

Advanced Econometrics

Jean-Paul Renne

2022-09-19

Contents

1 Before starting	5
2 Linear Regressions	7
2.1 Hypotheses	7
2.2 Least square estimation	10
2.3 Common pitfalls in linear regressions	25
2.4 Instrumental Variables	27
2.5 General Regression Model (GRM) and robust covariance matrices	33
2.6 Shrinkage methods	45
3 Panel regressions	51
3.1 Specification and notations	51
3.2 Estimation of Fixed-Effects Models	54
3.3 Estimation of random effects models	58
3.4 Dynamic Panel Regressions	63
3.5 Introduction to program evaluation	68
4 Estimation Methods	75
4.1 Generalized Method of Moments (GMM)	75
4.2 Maximum Likelihood Estimation	81
4.3 Bayesian approach	93

5 Microeconometrics	103
5.1 Binary-choice models	103
5.2 Multiple Choice Models	117
5.3 Tobit models	131
5.4 Sample Selection Models	139
5.5 Models of Count Data	146
6 Time Series	155
6.1 Introduction to time series	155
6.2 Univariate processes	161
6.3 Multivariate models	191
6.4 Forecasting	239
7 Appendix	247
7.1 Principal component analysis (PCA)	247
7.2 Linear algebra: definitions and results	249
7.3 Statistical analysis: definitions and results	252
7.4 Some properties of Gaussian variables	257
7.5 Proofs	259
7.6 Additional codes	267
7.7 Statistical Tables	269

Chapter 1

Before starting

This course covers various econometric topics, including linear regression models, discrete-choice models, and time series analysis. It provides examples or simulations based on R codes.

The R codes use various packages that can be obtained from CRAN. Several pieces of code also involve procedures and data from my `AEC` package. The latter is available on GitHub. To install it, one need to employ the `devtools` library:

```
library(devtools)
install_github("jrenne/AEC")
library(AEC)
```

Useful (R) links:

- Download R:
 - R software: <https://cran.r-project.org> (the basic R software)
 - RStudio: <https://www.rstudio.com> (a convenient R editor)
- Tutorials:
 - Rstudio: <https://dss.princeton.edu/training/RStudio101.pdf> (by Oscar Torres-Reyna)
 - R: https://cran.r-project.org/doc/contrib/Paradis-rdebut_en.pdf (by Emmanuel Paradis)
 - My own tutorial: https://jrenne.shinyapps.io/Rtuto_publiShiny/

Chapter 2

Linear Regressions

Definition 2.1. A linear regression is a model defined through:

$$y_i = \beta' \mathbf{x}_i + \varepsilon_i, \quad (2.1)$$

where $\mathbf{x}_i = [x_{i,1}, \dots, x_{i,K}]'$ is a vector of dimension $K \times 1$.

For entity i , the $x_{i,k}$'s, for $k \in \{1, \dots, K\}$, are explanatory variables, regressors, or covariates. The variable of interest, y_i , is often called dependent variable, or regressand. The last term of the specification, namely ε_i , is called error, or disturbance.

The researcher is usually interested in the components of vector β , denoted by β_k , $k \in \{1, \dots, K\}$. She usually aims at estimating these coefficients based on observations of $\{y_i, \mathbf{x}_i\}$, $i \in \{1, \dots, n\}$, which constitutes a **sample** (of size n).

To have an intercept in the specification (2.1), one has to set $x_{i,1} = 1$ for all i ; β_1 then corresponds to the intercept.

2.1 Hypotheses

In the following, we introduce different assumptions regarding the covariates and/or the errors. The properties of the estimators used by the researcher depend on which of these assumptions are satisfied.

Hypothesis 2.1 (Full rank). There is no exact linear relationship among the independent variables (the $x_{i,k}$'s, for a given $i \in \{1, \dots, n\}$).

Intuitively, when Hypothesis 2.1 is satisfied, then the estimation of the model parameters is unfeasible since, for any value of β , some changes in the explanatory variables will be exactly compensated by other changes in another set of explanatory variables, preventing the identification of these effects.

Let us denote by \mathbf{X} the matrix containing all explanatory variables, of dimension $n \times K$. (That is, row i of \mathbf{X} is \mathbf{x}'_i .) The following hypothesis concerns the relationship between the errors (gathered in ε , a n -dimensional vector) and the explanatory variables \mathbf{X} :

Hypothesis 2.2 (Conditional mean-zero assumption).

$$\mathbb{E}(\varepsilon|\mathbf{X}) = 0. \quad (2.2)$$

The following proposition states some implications of Hypothesis 2.2:

Proposition 2.1. *Under Hypothesis 2.2:*

- i. $\mathbb{E}(\varepsilon_i) = 0$;
- ii. the x_{ij} 's and the ε_i 's are uncorrelated, i.e. $\forall i, j \quad \text{Corr}(x_{ij}, \varepsilon_i) = 0$.

Proof. Let us prove (i) and (ii):

- i. By the law of iterated expectations:

$$\mathbb{E}(\varepsilon) = \mathbb{E}(\mathbb{E}(\varepsilon|\mathbf{X})) = \mathbb{E}(0) = 0.$$

$$\text{ii. } \mathbb{E}(x_{ij}\varepsilon_i) = \mathbb{E}(\mathbb{E}(x_{ij}\varepsilon_i|\mathbf{X})) = \mathbb{E}(x_{ij} \underbrace{\mathbb{E}(\varepsilon_i|\mathbf{X})}_{=0}) = 0. \square$$

□

Let us now present two hypotheses (?? and ??) concerning the stochastic properties of the errors ε_i :

Hypothesis 2.3 (Homoskedasticity).

$$\forall i, \quad \text{Var}(\varepsilon_i|\mathbf{X}) = \sigma^2.$$

The following lines of code generate a figure comparing two situations: Panel (a) of Figure 2.1 corresponds to a situation of homoskedasticity, and Panel (b) corresponds to a situation of heteroskedasticity. Let us be more specific. In the two plots, we have $X_i \sim \mathcal{N}(0, 1)$ and $\varepsilon_i^* \sim \mathcal{N}(0, 1)$. In Panel (a) (homoskedasticity): $Y_i = 2 + 2X_i + \varepsilon_i^*$. In Panel (b) (heteroskedasticity): $Y_i = 2 + 2X_i + (2\mathbb{1}_{\{X_i < 0\}} + 0.2\mathbb{1}_{\{X_i \geq 0\}})\varepsilon_i^*$.

Figure 2.2 shows a situation of heteroskedasticity, based on data taken from the Swiss Household Panel. The sample is restricted to persons younger than 35 year in 2019, and that have completed at least 19 years of study. The figure shows that the dispersion of yearly income increases with age.




Figure 2.1: Homoskedasticity vs heteroskedasticity.

```

library(AEC)
table(shp$edyear19)

##
##      8     9    10    12    13    14    16    19    21
##    70   325   350  1985   454   117   990  1263   168

shp_higherEd <- subset(shp, (edyear19>18)&age19<35)
plot(i19wyg/1000~age19, data=shp_higherEd, pch=19, las=1, xlab="Age", ylab="Yearly work income")
abline(lm(i19wyg/1000~age19, data=shp_higherEd), col="red", lwd=2)

```

The next assumption concerns the correlation of the errors across entities.

Hypothesis 2.4 (Uncorrelated errors).

$$\forall i \neq j, \quad \text{Cov}(\varepsilon_i, \varepsilon_j | \mathbf{X}) = 0.$$

We will often work with covariance matrices. Proposition 2.2 give the specific form of the conditional covariance of the errors when both Hypotheses 2.3 and 2.4 are satisfied.

Proposition 2.2. *If Hypotheses 2.3 and 2.4 hold, then:*

$$\text{Var}(\varepsilon | \mathbf{X}) = \sigma^2 \text{Id},$$

where Id is the $n \times n$ identity matrix.




Figure 2.2: Income versus age. Data are from the Swiss Household Panel. The sample is restricted to persons that have completed at least 19 years of study. The figure shows that the dispersion of yearly income increases with age.

We will sometimes assume that errors are Gaussian — or normal. We will then work under Hypothesis 2.5:

Hypothesis 2.5 (Normal distribution).

$$\forall i, \quad \varepsilon_i \sim \mathcal{N}(0, \sigma^2).$$

2.2 Least square estimation

2.2.1 Derivation of the OLS formula

In this section, we will present and study the properties of the very popular estimation approach called **Ordinary Least Squares (OLS)**. As suggested by its name, the OLS estimator of β is defined as the vector \mathbf{b} that minimizes the sum of squared residuals. (The **residuals** are the estimates of the **errors** ε_i .)

For a given vector of coefficients $\mathbf{b} = [b_1, \dots, b_K]'$, the sum of squared residuals is:

$$f(\mathbf{b}) = \sum_{i=1}^n \left(y_i - \sum_{j=1}^K x_{i,j} b_j \right)^2 = \sum_{i=1}^n (y_i - \mathbf{x}_i' \mathbf{b})^2.$$

Minimizing this sum amounts to minimizing:

$$f(\mathbf{b}) = (\mathbf{y} - \mathbf{X}\mathbf{b})'(\mathbf{y} - \mathbf{X}\mathbf{b}).$$

Since:

$$\frac{\partial f}{\partial \mathbf{b}}(\mathbf{b}) = -2\mathbf{X}'\mathbf{y} + 2\mathbf{X}'\mathbf{X}\mathbf{b},$$

it comes that a necessary first-order condition (FOC) is:

$$\mathbf{X}'\mathbf{X}\mathbf{b} = \mathbf{X}'\mathbf{y}. \quad (2.3)$$

Under Assumption 2.1, $\mathbf{X}'\mathbf{X}$ is invertible. Hence:

$$\boxed{\mathbf{b} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}.}$$

Vector \mathbf{b} minimizes the sum of squared residuals. (f is a non-negative quadratic function, it therefore admits a minimum.)

We have:

$$\mathbf{y} = \underbrace{\mathbf{X}\mathbf{b}}_{\text{fitted values } (\mathbf{y})} + \underbrace{\mathbf{e}}_{\text{residuals}}$$

The estimated residuals are:

$$\mathbf{e} = \mathbf{y} - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} = \mathbf{M}\mathbf{y} \quad (2.4)$$

where $\mathbf{M} := \mathbf{I} - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ is called the **residual maker matrix**.

Moreover, the fitted values $\hat{\mathbf{y}}$ are given by:

$$\hat{\mathbf{y}} = \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} = \mathbf{P}\mathbf{y}, \quad (2.5)$$

where $\mathbf{P} = \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ is a **projection matrix**.

These matrices \mathbf{M} and \mathbf{P} are such that:

- $\mathbf{M}\mathbf{X} = \mathbf{0}$: if one regresses one of the explanatory variables on \mathbf{X} , the residuals are null.
- $\mathbf{M}\mathbf{y} = \mathbf{M}\varepsilon$ (because $\mathbf{y} = \mathbf{X}\beta + \varepsilon$ and $\mathbf{M}\mathbf{X} = \mathbf{0}$).

Here are some additional properties of \mathbf{M} and \mathbf{P} :

- \mathbf{M} is symmetric ($\mathbf{M} = \mathbf{M}'$) and **idempotent** ($\mathbf{M} = \mathbf{M}^2 = \mathbf{M}^k$ for $k > 0$).
- \mathbf{P} is symmetric and idempotent.
- $\mathbf{P}\mathbf{X} = \mathbf{X}$.
- $\mathbf{P}\mathbf{M} = \mathbf{M}\mathbf{P} = \mathbf{0}$.
- $\mathbf{y} = \mathbf{Py} + \mathbf{My}$ (decomposition of \mathbf{y} into two orthogonal parts).

It is easily checked that $\mathbf{X}'\mathbf{e} = 0$. Each column of \mathbf{X} is therefore orthogonal to \mathbf{e} . In particular, if an intercept is included in the regression ($x_{i,1} \equiv 1$, i.e., the first column of \mathbf{X} is filled with ones), the average of the residuals is null.

Example 2.1 (Bivariate case). Consider a bivariate situation, where we regress y_i on a constant and an explanatory variable w_i . We have $K = 2$, and \mathbf{X} is a $n \times 2$ matrix whose i^{th} row is $[x_{i,1}, x_{i,2}]$, with $x_{i,1} = 1$ (to account for the intercept) and with $w_i = x_{i,2}$ (say).

We have:

$$\begin{aligned}\mathbf{X}'\mathbf{X} &= \begin{bmatrix} n & \sum_i w_i \\ \sum_i w_i & \sum_i w_i^2 \end{bmatrix}, \\ (\mathbf{X}'\mathbf{X})^{-1} &= \frac{1}{n \sum_i w_i^2 - (\sum_i w_i)^2} \begin{bmatrix} \sum_i w_i^2 & -\sum_i w_i \\ -\sum_i w_i & n \end{bmatrix}, \\ (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} &= \frac{1}{n \sum_i w_i^2 - (\sum_i w_i)^2} \begin{bmatrix} \sum_i w_i^2 \sum_i y_i - \sum_i w_i \sum_i w_i y_i \\ -\sum_i w_i \sum_i y_i + n \sum_i w_i y_i \end{bmatrix} \\ &= \frac{1}{\frac{1}{n} \sum_i (w_i - \bar{w})^2} \begin{bmatrix} \frac{\bar{y}}{n} \sum_i w_i^2 - \frac{\bar{w}}{n} \sum_i w_i y_i \\ \frac{1}{n} \sum_i (w_i - \bar{w})(y_i - \bar{y}) \end{bmatrix}.\end{aligned}$$

It can be seen that the second element of $\mathbf{b} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ is:

$$b_2 = \frac{\overline{\text{Cov}(W, Y)}}{\overline{\text{Var}(W)}},$$

where $\overline{\text{Cov}(W, Y)}$ and $\overline{\text{Var}(W)}$ are sample estimates.

Since there is a constant in the regression, we have $b_1 = \bar{y} - b_2 \bar{w}$.

2.2.2 Properties of the OLS estimate (small sample)

Proposition 2.3 states first properties of the OLS estimator:

Proposition 2.3 (Properties of the OLS estimator). *We have:*

- i. Under Assumptions 2.1 and 2.2, the OLS estimator is linear and unbiased.
- ii. Under Hypotheses 2.1 to 2.4, the conditional covariance matrix of \mathbf{b} is:
 $\text{Var}(\mathbf{b}|\mathbf{X}) = \sigma^2(\mathbf{X}'\mathbf{X})^{-1}$.

Proof. Under Hypothesis 2.1, $\mathbf{X}'\mathbf{X}$ can be inverted. We have:

$$\mathbf{b} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y} = \beta + (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\epsilon.$$

- i. Let us consider the expectation of the last term, i.e. $\mathbb{E}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\epsilon)$. Using the law of iterated expectations, we obtain:

$$\mathbb{E}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}') = \mathbb{E}(\mathbb{E}[(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'|\mathbf{X}]) = \mathbb{E}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}[\epsilon|\mathbf{X}]).$$

By Hypothesis 2.2, we have $\mathbb{E}[\epsilon|\mathbf{X}] = 0$. Hence $\mathbb{E}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}') = 0$ and result (i) follows.

- ii. $\text{Var}(\mathbf{b}|\mathbf{X}) = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbb{E}(\varepsilon\varepsilon'|\mathbf{X})\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}$. By Prop. 2.2, if 2.3 and 2.4 hold, then we have $\mathbb{E}(\varepsilon\varepsilon'|\mathbf{X}) = \text{Var}(\varepsilon|\mathbf{X}) = \sigma^2 I_d$.

□

Together, Hypotheses 2.1 to 2.4 form the Gauss-Markov set of assumptions. Under these assumptions, the OLS estimator feature the lowest possible variance:

Theorem 2.1 (Gauss-Markov Theorem). *Under Assumptions 2.1 to 2.4, for any vector w , the minimum-variance linear unbiased estimator of $w'\beta$ is $w'\mathbf{b}$, where \mathbf{b} is the least squares estimator. (BLUE: Best Linear Unbiased Estimator.)*

Proof. Consider $\mathbf{b}^* = C\mathbf{y}$, another linear unbiased estimator of β . Since it is unbiased, we must have $\mathbb{E}(C\mathbf{y}|\mathbf{X}) = \mathbb{E}(C\mathbf{X}\beta + C\varepsilon|\mathbf{X}) = \beta$. We have $\mathbb{E}(C\varepsilon|\mathbf{X}) = C\mathbb{E}(\varepsilon|\mathbf{X}) = 0$ (by 2.2).

Therefore \mathbf{b}^* is unbiased if $\mathbb{E}(C\mathbf{X})\beta = \beta$. This has to be the case for any β , which implies that we must have $C\mathbf{X} = \mathbf{I}$.

Let us compute $\text{Var}(\mathbf{b}^*|\mathbf{X})$. For this, we introduce $D = C - (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$, which is such that $D\mathbf{y} = \mathbf{b}^* - \mathbf{b}$. The fact that $C\mathbf{X} = \mathbf{I}$ implies that $D\mathbf{X} = \mathbf{0}$.

We have $\text{Var}(\mathbf{b}^*|\mathbf{X}) = \text{Var}(C\mathbf{y}|\mathbf{X}) = \text{Var}(C\varepsilon|\mathbf{X}) = \sigma^2 CC'$ (by Assumptions 2.3 and 2.4, see Prop. 2.2). Using $C = D + (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ and exploiting the fact that $D\mathbf{X} = \mathbf{0}$ leads to:

$$\text{Var}(\mathbf{b}^*|\mathbf{X}) = \sigma^2 [(D + (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}')(D + (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}')'] = \text{Var}(\mathbf{b}|\mathbf{X}) + \sigma^2 \mathbf{D}\mathbf{D}'.$$

Therefore, we have $\text{Var}(w'\mathbf{b}^*|\mathbf{X}) = w'\text{Var}(\mathbf{b}|\mathbf{X})w + \sigma^2 w'\mathbf{D}\mathbf{D}'w \geq w'\text{Var}(\mathbf{b}|\mathbf{X})w = \text{Var}(w'\mathbf{b}|\mathbf{X})$. □

The Frisch-Waugh theorem (Theorem 2.2) reveals the relationship between the OLS estimator and the notion of partial correlation coefficient. Consider the linear least square regression of \mathbf{y} on \mathbf{X} . We introduce the notations:

- $\mathbf{b}^{\mathbf{y}/\mathbf{X}}$: OLS estimates of β ,
- $\mathbf{M}^{\mathbf{X}}$: residual-maker matrix of any regression on \mathbf{X} ,
- $\mathbf{P}^{\mathbf{X}}$: projection matrix of any regression on \mathbf{X} .

Let us split the set of explanatory variables into two: $\mathbf{X} = [\mathbf{X}_1, \mathbf{X}_2]$. With obvious notations: $\mathbf{b}^{\mathbf{y}/\mathbf{X}} = [\mathbf{b}'_1, \mathbf{b}'_2]'$.

Theorem 2.2 (Frisch-Waugh Theorem). *We have:*

$$\mathbf{b}_2 = \mathbf{b}^{\mathbf{M}^{\mathbf{X}_1}\mathbf{y}/\mathbf{M}^{\mathbf{X}_1}\mathbf{X}_2}.$$

Proof. The minimization of the least squares leads to (these are first-order conditions, see Eq. (2.3)):

$$\begin{bmatrix} \mathbf{X}'_1 \mathbf{X}_1 & \mathbf{X}'_1 \mathbf{X}_2 \\ \mathbf{X}'_2 \mathbf{X}_1 & \mathbf{X}'_2 \mathbf{X}_2 \end{bmatrix} \begin{bmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{X}'_1 \mathbf{y} \\ \mathbf{X}'_2 \mathbf{y} \end{bmatrix}.$$

Use the first-row block of equations to solve for \mathbf{b}_1 first; it comes as a function of \mathbf{b}_2 . Then use the second set of equations to solve for \mathbf{b}_2 , which leads to:

$$\mathbf{b}_2 = [\mathbf{X}'_2 \mathbf{X}_2 - \mathbf{X}'_2 \mathbf{X}_1 (\mathbf{X}'_1 \mathbf{X}_1) \mathbf{X}'_1 \mathbf{X}_2]^{-1} \mathbf{X}'_2 (Id - \mathbf{X}_1 (\mathbf{X}'_1 \mathbf{X}_1) \mathbf{X}'_1) \mathbf{y} = [\mathbf{X}'_2 \mathbf{M}^{\mathbf{X}_1} \mathbf{X}_2]^{-1} \mathbf{X}'_2 \mathbf{M}^{\mathbf{X}_1} \mathbf{y}.$$

Using the fact that $\mathbf{M}^{\mathbf{X}_1}$ is idempotent and symmetric leads to the result. \square

This suggests a second way of estimating \mathbf{b}_2 :

1. Regress Y on X_1 , regress X_2 on X_1 .
2. Regress the residuals associated with the former regression on the ones associated with the latter regressions.

This is illustrated by the following code, where we run different regressions involving the number of Google searches for “parapluie” (*umbrella* in French). In the broad specification, we regress it on French precipitations and month dummies. Next, we deseasonalize both the dependent variable and the precipitations by regressing them on the month dummies. As stated by Theorem 2.2, regressing deseasonalized Google searches on deseasonalized precipitations give the same coefficient as in the baseline regression.

```
library(AEC)
dummies <- as.matrix(parapluie[, 4:14])
eq_all <- lm(parapluie ~ precip + dummies, data=parapluie)
deseas_parapluie <- lm(parapluie ~ dummies, data=parapluie)$residuals
deseas_precip <- lm(precip ~ dummies, data=parapluie)$residuals
eq_frac <- lm(deseas_parapluie ~ deseas_precip)
rbind(eq_all$coefficients,
      c(eq_frac$coefficients, rep(NaN, 11)))

##           (Intercept)      precip   dummies1   dummies2   dummies3   dummies4   dummies5
## [1,] -2.953448e+00  0.1300055 -8.599068 -13.32904 -7.982958 -3.392353 -3.703816
## [2,] -1.038345e-15  0.1300055       NaN       NaN       NaN       NaN       NaN
##           dummies6   dummies7   dummies8   dummies9   dummies10  dummies11
## [1,] -3.360641 -7.315881 -7.717277 -4.6492 -5.109199  1.98077
## [2,]       NaN       NaN       NaN       NaN       NaN       NaN
```

When b_2 is scalar (and then \mathbf{X}_2 is of dimension $n \times 1$), Theorem 2.2 gives the expression of the **partial regression coefficient** b_2 :

$$b_2 = \frac{\mathbf{X}'_2 M^{\mathbf{X}_1} \mathbf{y}}{\mathbf{X}'_2 M^{\mathbf{X}_1} \mathbf{X}_2}.$$

2.2.3 Goodness of fit

Define the total variation in y as the sum of squared deviations:

$$TSS = \sum_{i=1}^n (y_i - \bar{y})^2.$$

We have:

$$\mathbf{y} = \mathbf{X}\mathbf{b} + \mathbf{e} = \hat{\mathbf{y}} + \mathbf{e}$$

In the following, we assume that the regression includes a constant (i.e. for all i , $x_{i,1} = 1$). Denote by \mathbf{M}^0 the matrix that transforms observations into deviations from sample means. Using that $\mathbf{M}^0\mathbf{e} = \mathbf{e}$ and that $\mathbf{X}'\mathbf{e} = 0$, we have:

$$\begin{aligned} \underbrace{\mathbf{y}'\mathbf{M}^0\mathbf{y}}_{\text{Total sum of sq.}} &= (\mathbf{X}\mathbf{b} + \mathbf{e})'\mathbf{M}^0(\mathbf{X}\mathbf{b} + \mathbf{e}) \\ &= \underbrace{\mathbf{b}'\mathbf{X}'\mathbf{M}^0\mathbf{X}\mathbf{b}}_{\text{"Explained" sum of sq.}} + \underbrace{\mathbf{e}'\mathbf{e}}_{\text{Sum of sq. residuals}} \\ TSS &= Expl.SS + SSR. \end{aligned}$$

We can now define the coefficient of determination:

$$\text{Coefficient of determination} = \frac{Expl.SS}{TSS} = 1 - \frac{SSR}{TSS} = 1 - \frac{\mathbf{e}'\mathbf{e}}{\mathbf{y}'\mathbf{M}^0\mathbf{y}}. \quad (2.6)$$

It can be shown (Greene, 2003, Section 3.5) that:

$$\text{Coefficient of determination} = \frac{[\sum_{i=1}^n (y_i - \bar{y})(\hat{y}_i - \bar{y})]^2}{\sum_{i=1}^n (y_i - \bar{y})^2 \sum_{i=1}^n (\hat{y}_i - \bar{y})^2}.$$

That is, the R^2 is the sample squared correlation between y and the (regression-implied) y 's predictions.

The higher the R^2 , the higher the goodness of fit of a model. One however has to be cautious with R^2 . Indeed, it is easy to increase it: it suffices to add explanatory variables. As stated by Proposition 2.5, adding an explanatory variable (even if it does not truly relate to the dependent variable) results in an increase in the R^2 . In the limit, taking any set of n non-linearly-dependent explanatory variables (i.e., variables satisfying Hypothesis 2.1) results in a R^2 equal to one.

Proposition 2.4 (Change in SSR when a variable is added). *We have:*

$$\mathbf{u}'\mathbf{u} = \mathbf{e}'\mathbf{e} - c^2(\mathbf{z}^*\mathbf{z}^*) \quad (\leq \mathbf{e}'\mathbf{e}) \quad (2.7)$$

where (i) \mathbf{u} and \mathbf{e} are the residuals in the regressions of \mathbf{y} on $[\mathbf{X}, \mathbf{z}]$ and of \mathbf{y} on \mathbf{X} , respectively, (ii) c is the regression coefficient on \mathbf{z} in the former regression and where \mathbf{z}^* are the residuals in the regression of \mathbf{z} on \mathbf{X} .

Proof. The OLS estimates $[\mathbf{d}', \mathbf{c}']'$ in the regression of \mathbf{y} on $[\mathbf{X}, \mathbf{z}]$ satisfies (first-order cond., Eq. (2.3))

$$\begin{bmatrix} \mathbf{X}'\mathbf{X} & \mathbf{X}'\mathbf{z} \\ \mathbf{z}'\mathbf{X} & \mathbf{z}'\mathbf{z} \end{bmatrix} \begin{bmatrix} \mathbf{d} \\ \mathbf{c} \end{bmatrix} = \begin{bmatrix} \mathbf{X}'\mathbf{y} \\ \mathbf{z}'\mathbf{y} \end{bmatrix}.$$

Hence, in particular $\mathbf{d} = \mathbf{b} - (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{z}\mathbf{c}$, where \mathbf{b} is the OLS of \mathbf{y} on \mathbf{X} . Substituting in $\mathbf{u} = \mathbf{y} - \mathbf{X}\mathbf{d} - \mathbf{z}\mathbf{c}$, we get $\mathbf{u} = \mathbf{e} - \mathbf{z}^*c$. We therefore have:

$$\mathbf{u}'\mathbf{u} = (\mathbf{e} - \mathbf{z}^*c)(\mathbf{e} - \mathbf{z}^*c) = \mathbf{e}'\mathbf{e} + c^2(\mathbf{z}^{*'}\mathbf{z}^*) - 2c\mathbf{z}^{*'}\mathbf{e}. \quad (2.8)$$

Now $\mathbf{z}^{*'}\mathbf{e} = \mathbf{z}^{*'}(\mathbf{y} - \mathbf{X}\mathbf{b}) = \mathbf{z}^{*'}\mathbf{y}$ because \mathbf{z}^* are the residuals in an OLS regression on \mathbf{X} . Since $c = (\mathbf{z}^{*'}\mathbf{z}^*)^{-1}\mathbf{z}^{*'}\mathbf{y}^*$ (by an application of Theorem 2.2), we have $(\mathbf{z}^{*'}\mathbf{z}^*)c = \mathbf{z}^{*'}\mathbf{y}^*$ and, therefore, $\mathbf{z}^{*'}\mathbf{e} = (\mathbf{z}^{*'}\mathbf{z}^*)c$. Inserting this in Eq. (2.8) leads to the results. \square

Proposition 2.5 (Change in the coefficient of determination when a variable is added). *Denoting by R_W^2 the coefficient of determination in the regression of \mathbf{y} on some variable \mathbf{W} , we have:*

$$R_{\mathbf{X}, \mathbf{z}}^2 = R_{\mathbf{X}}^2 + (1 - R_{\mathbf{X}}^2)(r_{yz}^{\mathbf{X}})^2,$$

where $r_{yz}^{\mathbf{X}}$ is the coefficient of partial correlation (see Definition 7.5).

Proof. Let's use the same notations as in Prop. @ref{prp:chgeR2}. Theorem 2.2 implies that $c = (\mathbf{z}^{*'}\mathbf{z}^*)^{-1}\mathbf{z}^{*'}\mathbf{y}^*$. Using this in Eq. (2.7) gives $\mathbf{u}'\mathbf{u} = \mathbf{e}'\mathbf{e} - (\mathbf{z}^{*'}\mathbf{y}^*)^2/(\mathbf{z}^{*'}\mathbf{z}^*)$. Using the definition of the partial correlation (Eq. (7.2)), we get $\mathbf{u}'\mathbf{u} = \mathbf{e}'\mathbf{e}(1 - (r_{yz}^{\mathbf{X}})^2)$. The results is obtained by dividing both sides of the previous equation by $\mathbf{y}'\mathbf{M}_0\mathbf{y}$. \square

Figure 2.3, below, illustrates the fact that one can obtain an R^2 of one by regressing a sample of length n on any set of n linearly-independent variables.

```

n <- 30; Y <- rnorm(n); X <- matrix(rnorm(n^2), n, n)
all_R2 <- NULL; all_adjR2 <- NULL
for(j in 0:(n-1)){
  if(j==0){eq <- lm(Y~1)
  }else{eq <- lm(Y~X[,1:j])}
  all_R2 <- c(all_R2, summary(eq)$r.squared)
  all_adjR2 <- c(all_adjR2, summary(eq)$adj.r.squared)
}
plot(all_R2, pch=19, ylim=c(min(all_adjR2, na.rm = TRUE), 1), xlab="number of regressors", ylab="Adjusted R-squared")
points(all_adjR2, pch=3); abline(h=0, col="light grey", lwd=2)
legend("topleft", c("R2", "Adjusted R2"),
      lty=NaN, col=c("black"), pch=c(19, 3), lwd=2)

```




Figure 2.3: This figure illustrates the monotonous increase in the R^2 as a function of the number of explanatory variables. In the true model, there is no explanatory variables, i.e., $y_i = \varepsilon_i$. We then take (independent) regressors and regress y on the latter, progressively increasing the set of regressors.

In order to address the risk of adding irrelevant explanatory variables, measures of **adjusted** R^2 have been proposed. Compared to the standard R^2 , these measures add penalties that depend on the number of covariates employed in the regression. A common adjusted R^2 measure, denoted by \bar{R}^2 , is the following:

$$\bar{R}^2 = 1 - \frac{\mathbf{e}'\mathbf{e}/(n-K)}{\mathbf{y}'\mathbf{M}^0\mathbf{y}/(n-1)} = 1 - \frac{n-1}{n-K}(1-R^2).$$

2.2.4 Inference and confidence intervals (in small sample)

Under the normality assumption (Assumption 2.5), we know the distribution of \mathbf{b} (conditional on \mathbf{X}). Indeed, $(\mathbf{b}|\mathbf{X}) \equiv (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ is multivariate Gaussian:

$$\mathbf{b}|\mathbf{X} \sim \mathcal{N}(\beta, \sigma^2(\mathbf{X}'\mathbf{X})^{-1}). \quad (2.9)$$

Eq. (2.9) can be used to conduct inference and tests. However, in practice, we do not know σ^2 (which is a population parameter). The following proposition gives an unbiased estimate of σ^2 .

Proposition 2.6. *Under 2.1 to 2.4, an unbiased estimate of σ^2 is given by:*

$$s^2 = \frac{\mathbf{e}'\mathbf{e}}{n-K}. \quad (2.10)$$

(It is sometimes denoted by σ_{OLS}^2 .)

Proof. $\mathbb{E}(\mathbf{e}'\mathbf{e}|\mathbf{X}) = \mathbb{E}(\varepsilon'\mathbf{M}\varepsilon|\mathbf{X}) = \mathbb{E}(\text{Tr}(\varepsilon'\mathbf{M}\varepsilon)|\mathbf{X}) = \text{Tr}(\mathbf{M}\mathbb{E}(\varepsilon\varepsilon'|\mathbf{X})) = \sigma^2\text{Tr}(\mathbf{M})$. (Note that we have $\mathbb{E}(\varepsilon\varepsilon'|\mathbf{X}) = \sigma^2\text{Id}$ by Assumptions 2.3 and 2.4, see Prop. 2.2.) Finally:

$$\begin{aligned}\text{Tr}(\mathbf{M}) &= n - \text{Tr}(\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}') \\ &= n - \text{Tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}) = n - \text{Tr}(Id_{K \times K}),\end{aligned}$$

which leads to the result. \square

Two results will prove important to produce inference:

- i. We know the distribution of s^2 (Prop. 2.7).
- ii. s^2 and \mathbf{b} are independent random variables (Prop. 2.8).

Proposition 2.7. *Under 2.1 to 2.5, we have: $\frac{s^2}{\sigma^2}|\mathbf{X} \sim \chi^2(n-K)/(n-K)$.*

Proof. We have $\mathbf{e}'\mathbf{e} = \varepsilon'\mathbf{M}\varepsilon$. \mathbf{M} is an idempotent symmetric matrix. Therefore it can be decomposed as PDP' where D is a diagonal matrix and P is an orthogonal matrix. As a result $\mathbf{e}'\mathbf{e} = (P'\varepsilon)'D(P'\varepsilon)$, i.e. $\mathbf{e}'\mathbf{e}$ is a weighted sum of independent squared Gaussian variables (the entries of $P'\varepsilon$ are independent because they are Gaussian —under 2.5— and uncorrelated). The variance of each of these i.i.d. Gaussian variable is σ^2 . Because \mathbf{M} is an idempotent symmetric matrix, its eigenvalues are either 0 or 1, and its rank equals its trace (see Propositions 7.3 and 7.4). Further, its trace is equal to $n - K$ (see proof of Eq. (2.10)). Therefore D has $n - K$ entries equal to 1 and K equal to 0. Hence, $\mathbf{e}'\mathbf{e} = (P'\varepsilon)'D(P'\varepsilon)$ is a sum of $n - K$ squared independent Gaussian variables of variance σ^2 . Therefore $\frac{\mathbf{e}'\mathbf{e}}{\sigma^2} = (n - K)\frac{s^2}{\sigma^2}$ is a sum of $n - K$ squared i.i.d. standard normal variables. \square

Proposition 2.8. *Under Hypotheses 2.1 to 2.5, \mathbf{b} and s^2 are independent.*

Proof. We have $\mathbf{b} = \beta + [\mathbf{X}'\mathbf{X}]^{-1}\mathbf{X}\varepsilon$ and $s^2 = \varepsilon'\mathbf{M}\varepsilon/(n - K)$. Hence \mathbf{b} is an affine combination of ε and s^2 is a quadratic combination of the same Gaussian shocks. One can write s^2 as $s^2 = (\mathbf{M}\varepsilon)'(\mathbf{M}\varepsilon)/(n - K)$ and \mathbf{b} as $\beta + \mathbf{T}\varepsilon$. Since $\mathbf{T}\mathbf{M} = 0$, $\mathbf{T}\varepsilon$ and $\mathbf{M}\varepsilon$ are independent (because two uncorrelated Gaussian variables are independent), therefore \mathbf{b} and s^2 , which are functions of two sets of independent variables, are independent. \square

Consistently with Eq. (2.9), under Hypotheses 2.1 to 2.5, the k^{th} entry of \mathbf{b} satisfies:

$$b_k|\mathbf{X} \sim \mathcal{N}(\beta_k, \sigma^2 v_k),$$

where v_k is the k^{th} component of the diagonal of $(\mathbf{X}'\mathbf{X})^{-1}$.

Moreover, we have (Prop. 2.7):

$$\frac{(n-K)s^2}{\sigma^2}|\mathbf{X} \sim \chi^2(n-K).$$

As a result (using Propositions 2.7 and 2.8), we have:

$$t_k = \frac{\frac{b_k - \beta_k}{\sqrt{\sigma^2 v_k}}}{\sqrt{\frac{(n-K)s^2}{\sigma^2(n-K)}}} = \frac{b_k - \beta_k}{\sqrt{s^2 v_k}} \sim t(n-K), \quad (2.11)$$

where $t(n-K)$ denotes a t distribution with $n-K$ degrees of freedom.¹

Note that $s^2 v_k$ is not exactly the conditional variance of b_k : The variance of b_k conditional on \mathbf{X} is $\sigma^2 v_k$. However $s^2 v_k$ is an unbiased estimate of $\sigma^2 v_k$ (by Prop. 2.6).

The previous result (Eq. (2.11)) can be extended to any linear combinations of elements of \mathbf{b} (Eq. (2.11) is for its k^{th} component only).

Let us consider $\alpha' \mathbf{b}$, the OLS estimate of $\alpha' \beta$. From Eq. (2.9), we have:

$$\alpha' \mathbf{b} |\mathbf{X} \sim \mathcal{N}(\alpha' \beta, \sigma^2 \alpha' (\mathbf{X}' \mathbf{X})^{-1} \alpha).$$

Therefore:

$$\frac{\alpha' \mathbf{b} - \alpha' \beta}{\sqrt{\sigma^2 \alpha' (\mathbf{X}' \mathbf{X})^{-1} \alpha}} |\mathbf{X} \sim \mathcal{N}(0, 1).$$

Using the same approach as the one used to derive Eq. (2.11), one can show that Props. 2.7 and 2.8 also imply that:

$$\frac{\alpha' \mathbf{b} - \alpha' \beta}{\sqrt{s^2 \alpha' (\mathbf{X}' \mathbf{X})^{-1} \alpha}} \sim t(n-K). \quad (2.12)$$

What precedes is widely exploited for statistical inference in the context of linear regressions. Indeed, Eq. (2.11) gives a sense of the distances between b_k and β_k that can be deemed as “likely”. For instance, it implies that, if $\sqrt{v_k s^2}$ is equal to 1 (say), then the probability to obtain b_k smaller than $\beta_k - 4.587 \times \sqrt{v_k s^2}$ or larger than $\beta_k + 4.587 \times \sqrt{v_k s^2}$ is equal to 0.1% when $n-K=10$.

That means for instance that, under the assumption that $\beta_k = 0$, it would be extremely unlikely to have obtained $b_k / \sqrt{v_k s^2}$ smaller than 4.587 or larger than 4.587. More generally, this shows that the **t-statistic**, i.e., the ratio $b_k / \sqrt{v_k s^2}$, is the test statistic associated with the null hypothesis:

$$H_0 : \beta_k = 0.$$

¹We have $\frac{b_k - \beta_k}{\sqrt{\sigma^2 v_k}} |\mathbf{X} \sim \mathcal{N}(0, 1)$ and $\frac{(n-K)s^2}{\sigma^2} |\mathbf{X} \sim \chi^2(n-K)$. These two distributions do not depend on \mathbf{X} \Rightarrow the *marginal distribution* of t_k is also t .




Figure 2.4: The higher the degree of freedom, the closer the distribution of $t(\nu)$ gets to the normal distribution. (Convergence in distribution.)

Under the null hypothesis, the test statistic follows a Student-t distribution with $n - K$ degrees of freedom. The **t-statistic** is therefore of particular importance, and, as a result, it is routinely reported in regression outputs. This is illustrated below, in a regression that aims at determining covariates of households' income. This example makes use of data from the Swiss Household Panel (SHP). `edyear19` is the number of years of education and `age19` is the age of the respondent, as of 2019.

```
library(AEC)
library(sandwich)
shp$income <- shp$i19ptotn/1000
shp$female <- 1*(shp$sex19==2)
eq <- lm(income ~ edyear19 + age19 + I(age19^2) + female, data=shp)
```

The last two columns give the **t-statistic** and p-values associated with the t-test, whose **critical region** for the test of size α is:

$$\left[-\infty, -\Phi_{t(n-K)}^{-1} \left(1 - \frac{\alpha}{2} \right) \right] \cup \left[\Phi_{t(n-K)}^{-1} \left(1 - \frac{\alpha}{2} \right), +\infty \right].$$

We recall that the **p-value** is defined as the probability that $|Z| > |t|$, where t is the (computed) t statistics and where $Z \sim t(n - K)$. That is, the p-value is given by $2(1 - \Phi_{t(n-K)}(|t_k|))$. See this webpage for details regarding the link between critical regions, p-value, and test outcomes.

Now, suppose we want to compute a (symmetrical) **confidence interval** $[I_{d,1-\alpha}, I_{u,1-\alpha}]$ that is such that $\mathbb{P}(\beta_k \in [I_{d,1-\alpha}, I_{u,1-\alpha}]) = 1 - \alpha$. In particular, we want to have: $\mathbb{P}(\beta_k < I_{d,1-\alpha}) = \frac{\alpha}{2}$.

For this purpose, we make use of Eq. (2.11), i.e., $t_k = \frac{b_k - \beta_k}{\sqrt{s^2 v_k}} \sim t(n - K)$. We have:

$$\begin{aligned} \mathbb{P}(\beta_k < I_{d,1-\alpha}) = \frac{\alpha}{2} &\Leftrightarrow \\ \mathbb{P}\left(\frac{b_k - \beta_k}{\sqrt{s^2 v_k}} > \frac{b_k - I_{d,1-\alpha}}{\sqrt{s^2 v_k}}\right) = \frac{\alpha}{2} &\Leftrightarrow \mathbb{P}\left(t_k > \frac{b_k - I_{d,1-\alpha}}{\sqrt{s^2 v_k}}\right) = \frac{\alpha}{2} \Leftrightarrow \\ 1 - \mathbb{P}\left(t_k \leq \frac{b_k - I_{d,1-\alpha}}{\sqrt{s^2 v_k}}\right) = \frac{\alpha}{2} &\Leftrightarrow \frac{b_k - I_{d,1-\alpha}}{\sqrt{s^2 v_k}} = \Phi_{t(n-K)}^{-1} \left(1 - \frac{\alpha}{2} \right), \end{aligned}$$

where $\Phi_{t(n-K)}(\alpha)$ is the c.d.f. of the $t(n - K)$ distribution (Table 7.2).

Doing the same for $I_{u,1-\alpha}$, we obtain:

$$\begin{aligned} [I_{d,1-\alpha}, I_{u,1-\alpha}] &= \\ \left[b_k - \Phi_{t(n-K)}^{-1} \left(1 - \frac{\alpha}{2} \right) \sqrt{s^2 v_k}, b_k + \Phi_{t(n-K)}^{-1} \left(1 - \frac{\alpha}{2} \right) \sqrt{s^2 v_k} \right]. \end{aligned}$$

Using the results of the previous regression, we compute lower and upper bounds of 95% confidence intervals for the estimated parameters as follows:

```

n <- length(eq$residuals); K <- length(eq$coefficients)
lower.b <- eq$coefficients - pt(.025,df=n-K)*sqrt(diag(vcov(eq)))
upper.b <- eq$coefficients + pt(.025,df=n-K)*sqrt(diag(vcov(eq)))
cbind(lower.b,upper.b)

##           lower.b      upper.b
## (Intercept) -74.8848532 -69.06276141
## edyear19     4.7334839   4.95504836
## age19        3.1272532   3.34998983
## I(age19^2)   -0.0300164  -0.02788323
## female       -32.5523381 -31.06546311

```

2.2.5 Testing a set of linear restrictions

We sometimes want to test if a set of restrictions is *jointly* consistent with the data at hand. Let us formalize such a set of (J) linear restrictions:

$$\begin{aligned} r_{1,1}\beta_1 + \cdots + r_{1,K}\beta_K &= q_1 \\ &\vdots \\ r_{J,1}\beta_1 + \cdots + r_{J,K}\beta_K &= q_J. \end{aligned} \tag{2.13}$$

In matrix form, we get:

$$\mathbf{R}\beta = \mathbf{q}. \tag{2.14}$$

Define the **Discrepancy vector** $\mathbf{m} = \mathbf{R}\mathbf{b} - \mathbf{q}$. Under the null hypothesis:

$$\begin{aligned} \mathbb{E}(\mathbf{m}|\mathbf{X}) &= \mathbf{R}\beta - \mathbf{q} = 0 \quad \text{and} \\ \mathbb{V}ar(\mathbf{m}|\mathbf{X}) &= \mathbf{R}\mathbb{V}ar(\mathbf{b}|\mathbf{X})\mathbf{R}'. \end{aligned}$$

With these notations, the assumption to test is:

$$H_0 : \mathbf{R}\beta - \mathbf{q} = 0 \text{ against } H_1 : \mathbf{R}\beta - \mathbf{q} \neq 0. \tag{2.15}$$

Under Hypotheses 2.1 to 2.4, we have $\mathbb{V}ar(\mathbf{m}|\mathbf{X}) = \sigma^2 \mathbf{R}(\mathbf{X}'\mathbf{X})^{-1} \mathbf{R}'$ (see Prop. 2.3). If we add the normality assumption (Hypothesis 2.5), we have:

$$W = \mathbf{m}'\mathbb{V}ar(\mathbf{m}|\mathbf{X})^{-1}\mathbf{m} \sim \chi^2(J). \tag{2.16}$$

If σ^2 was known, we could then conduct a **Wald test**. But this is not the case in practice and we cannot compute W . We can, however, approximate it by replacing σ^2 by s^2 . The distribution of this new statistic is not $\chi^2(J)$ any more; it is an **F distribution** (whose quantiles are shown in Table 7.4), and the test is called **F test**.

Proposition 2.9. *Under Hypotheses 2.1 to 2.5 and if Eq. (2.15) holds, we have:*

$$F = \frac{W}{J} \frac{\sigma^2}{s^2} = \frac{\mathbf{m}'(\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}')^{-1}\mathbf{m}}{s^2 J} \sim \mathcal{F}(J, n - K), \quad (2.17)$$

where \mathcal{F} is the distribution of the F-statistic.

Proof. According to Eq. (2.16), $W/J \sim \chi^2(J)/J$. Moreover, the denominator (s^2/σ^2) is $\sim \chi^2(n-K)$. Therefore, F is the ratio of a r.v. distributed as $\chi^2(J)/J$ and another distributed as $\chi^2(n-K)/(n-K)$. It remains to verify that these r.v. are independent.

Under H_0 , we have $\mathbf{m} = \mathbf{R}(\mathbf{b} - \boldsymbol{\beta}) = \mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\boldsymbol{\varepsilon}$. Therefore $\mathbf{m}'(\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}')^{-1}\mathbf{m}$ is of the form $\boldsymbol{\varepsilon}'\mathbf{T}\boldsymbol{\varepsilon}$ with $\mathbf{T} = \mathbf{D}'\mathbf{C}\mathbf{D}$ where $\mathbf{D} = \mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ and $\mathbf{C} = (\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}')^{-1}$. Under Hypotheses 2.1 to 2.4, the covariance between $\mathbf{T}\boldsymbol{\varepsilon}$ and $\mathbf{M}\boldsymbol{\varepsilon}$ is $\sigma^2\mathbf{T}\mathbf{M} = \mathbf{0}$. Therefore, under 2.5, these variables are Gaussian variables with 0 covariance. Hence they are independent. \square

For large $n - K$, the $\mathcal{F}_{J,n-K}$ distribution converges to $\mathcal{F}_{J,\infty} = \chi^2(J)/J$.

The following proposition proposes another computation of the F-statistic, based on the R^2 of the restricted and unrestricted linear models.

Proposition 2.10. *The F-statistic defined by Eq. (2.17) is also equal to:*

$$F = \frac{(R^2 - R_*^2)/J}{(1 - R^2)/(n - K)} = \frac{(SSR_{restr} - SSR_{unrestr})/J}{SSR_{unrestr}/(n - K)}, \quad (2.18)$$

where R_*^2 is the coef. of determination (Eq. (2.6)) of the “restricted regression” (SSR: sum of squared residuals.)

Proof. Let's denote by $\mathbf{e}_* = \mathbf{y} - \mathbf{X}\mathbf{b}_*$ the vector of residuals associated to the *restricted regression* (i.e. $\mathbf{R}\mathbf{b}_* = \mathbf{q}$). We have $\mathbf{e}_* = \mathbf{e} - \mathbf{X}(\mathbf{b}_* - \mathbf{b})$. Using $\mathbf{e}'\mathbf{X} = 0$, we get $\mathbf{e}'_*\mathbf{e}_* = \mathbf{e}'\mathbf{e} + (\mathbf{b}_* - \mathbf{b})'\mathbf{X}'\mathbf{X}(\mathbf{b}_* - \mathbf{b}) \geq \mathbf{e}'\mathbf{e}$.

By Proposition 7.5 (in Appendix ??), we have: $\mathbf{b}_* - \mathbf{b} = -(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}'\{\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}'\}^{-1}(\mathbf{R}\mathbf{b} - \mathbf{q})$. Therefore:

$$\mathbf{e}'_*\mathbf{e}_* - \mathbf{e}'\mathbf{e} = (\mathbf{R}\mathbf{b} - \mathbf{q})'[\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}']^{-1}(\mathbf{R}\mathbf{b} - \mathbf{q}).$$

This implies that the F statistic defined in Prop. 2.9 is also equal to:

$$\frac{(\mathbf{e}'_*\mathbf{e}_* - \mathbf{e}'\mathbf{e})/J}{\mathbf{e}'\mathbf{e}/(n - K)},$$

which leads to the result. \square

The null hypothesis H_0 (Eq. (2.15)) of the F-test is rejected if F –defined by Eq. (2.17) or (2.18)– is higher than $\mathcal{F}_{1-\alpha}(J, n - K)$. (Hence, this test is a one-sided test.)

2.2.6 Large Sample Properties

Even if we relax the normality assumption (Hypothesis 2.5), we can approximate the finite-sample behavior of the estimators by using *large-sample* or *asymptotic properties*.

To begin with, we proceed under Hypothesis 2.1 to 2.4. (We will see, later on, how to deal with –partial– relaxations of Hypothesis 2.3 and 2.4.)

Under regularity assumptions, and under Hypotheses 2.1 to 2.4, even if the residuals are not normally-distributed, the least square estimators can be *asymptotically normal* and inference can be performed as in small samples when Hypotheses 2.1 to 2.5 hold. This derives from Prop. 2.11 (below). The F-test (Prop. 2.10) and the t-test (Eq. (2.11)) can then be performed.

Proposition 2.11. *Under Assumptions 2.1 to 2.4, and assuming further that:*

$$Q = \text{plim}_{n \rightarrow \infty} \frac{\mathbf{X}'\mathbf{X}}{n}, \quad (2.19)$$

and that the $(\mathbf{x}_i, \varepsilon_i)$'s are independent (across entities i), we have:

$$\sqrt{n}(\mathbf{b} - \beta) \xrightarrow{d} \mathcal{N}(0, \sigma^2 Q^{-1}). \quad (2.20)$$

Proof. Since $\mathbf{b} = \beta + \left(\frac{\mathbf{X}'\mathbf{X}}{n}\right)^{-1} \left(\frac{\mathbf{X}'\varepsilon}{n}\right)$, we have: $\sqrt{n}(\mathbf{b} - \beta) = \left(\frac{\mathbf{X}'\mathbf{X}}{n}\right)^{-1} \left(\frac{1}{\sqrt{n}}\right) \mathbf{X}'\varepsilon$. Since $f : A \rightarrow A^{-1}$ is a continuous function (for $A \neq \mathbf{0}$), $\text{plim}_{n \rightarrow \infty} \left(\frac{\mathbf{X}'\mathbf{X}}{n}\right)^{-1} = \mathbf{Q}^{-1}$ (see Prop. 7.12). Let us denote by V_i the vector $\mathbf{x}_i\varepsilon_i$. Because the $(\mathbf{x}_i, \varepsilon_i)$'s are independent, the V_i 's are independent as well. Their covariance matrix is $\sigma^2 \mathbb{E}(\mathbf{x}_i \mathbf{x}_i') = \sigma^2 Q$. Applying the multivariate central limit theorem on vectors V_i gives $\sqrt{n} \left(\frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \varepsilon_i\right) = \left(\frac{1}{\sqrt{n}}\right) \mathbf{X}'\varepsilon \xrightarrow{d} \mathcal{N}(0, \sigma^2 Q)$. An application of Slutsky's theorem (Prop. 7.12) then leads to the results. \square

In practice, σ^2 is estimated with $\frac{\mathbf{e}'\mathbf{e}}{n-K}$ (Eq. (2.10)) and \mathbf{Q}^{-1} with $\left(\frac{\mathbf{X}'\mathbf{X}}{n}\right)^{-1}$. That is, the covariance matrix of the estimator is approximated by:

$$\widehat{\text{Var}}(\mathbf{b}) = s^2 (\mathbf{X}'\mathbf{X})^{-1}. \quad (2.21)$$

Eqs. (2.19) and (2.20) respectively correspond to convergences in probability and in distribution (see Definitions 7.14 and 7.17, respectively).

2.3 Common pitfalls in linear regressions

2.3.1 Multicollinearity

Consider the model: $y_i = \beta_1 x_{i,1} + \beta_2 x_{i,2} + \varepsilon_i$, where all variables are zero-mean and $\text{Var}(\varepsilon_i) = \sigma^2$. We have

$$\mathbf{X}'\mathbf{X} = \begin{bmatrix} \sum_i x_{i,1}^2 & \sum_i x_{i,1}x_{i,2} \\ \sum_i x_{i,1}x_{i,2} & \sum_i x_{i,2}^2 \end{bmatrix},$$

therefore:

$$(\mathbf{X}'\mathbf{X})^{-1} = \frac{1}{\sum_i x_{i,1}^2 \sum_i x_{i,2}^2 - (\sum_i x_{i,1}x_{i,2})^2} \begin{bmatrix} \sum_i x_{i,2}^2 & -\sum_i x_{i,1}x_{i,2} \\ -\sum_i x_{i,1}x_{i,2} & \sum_i x_{i,1}^2 \end{bmatrix}.$$

The inverse of the upper-left parameter of $(\mathbf{X}'\mathbf{X})^{-1}$ is:

$$\sum_i x_{i,1}^2 - \frac{(\sum_i x_{i,1}x_{i,2})^2}{\sum_i x_{i,2}^2} = \sum_i x_{i,1}^2 (1 - \text{correl}_{1,2}^2), \quad (2.22)$$

where $\text{correl}_{1,2}$ is the sample correlation between \mathbf{x}_1 and \mathbf{x}_2 .

Hence, the closer to one $\text{correl}_{1,2}$, the higher the variance of b_1 (recall that the variance of b_1 is the upper-left component of $\sigma^2(\mathbf{X}'\mathbf{X})^{-1}$).

2.3.2 Omitted variables

Consider the following model (the “True model”):

$$\mathbf{y} = \underbrace{\mathbf{X}_1}_{n \times K_1} \underbrace{\beta_1}_{K_1 \times 1} + \underbrace{\mathbf{X}_2}_{n \times K_2} \underbrace{\beta_2}_{K_2 \times 1} + \varepsilon$$

If one computes \mathbf{b}_1 by regressing \mathbf{y} on \mathbf{X}_1 only, one gets:

$$\mathbf{b}_1 = (\mathbf{X}'_1 \mathbf{X}_1)^{-1} \mathbf{X}'_1 \mathbf{y} = \beta_1 + (\mathbf{X}'_1 \mathbf{X}_1)^{-1} \mathbf{X}'_1 \mathbf{X}_2 \beta_2 + (\mathbf{X}'_1 \mathbf{X}_1)^{-1} \mathbf{X}'_1 \varepsilon.$$

This results in the omitted-variable formula:

$$\boxed{\mathbb{E}(\mathbf{b}_1 | \mathbf{X}) = \beta_1 + \underbrace{(\mathbf{X}'_1 \mathbf{X}_1)^{-1} (\mathbf{X}'_1 \mathbf{X}_2)}_{K_1 \times K_2} \beta_2}$$

(each column of $(\mathbf{X}'_1 \mathbf{X}_1)^{-1} (\mathbf{X}'_1 \mathbf{X}_2)$ are the OLS regressors obtained when regressing the columns of \mathbf{X}_2 on \mathbf{X}_1). Unless the variables included in \mathbf{X}_1 are orthogonal to those in \mathbf{X}_2 , we obtain a bias.

Example 2.2. Let us use the California Test Score dataset (in the package AER). Assume we want to measure the effect of the students-to-teacher ratio (`str`) on student test scores (`testscr`). The following regressions show that the effect is lower when controls are added.

```
library(AER); data("CASchools")
CASchools$str <- CASchools$students/CASchools$teachers
CASchools$testscr <- .5 * (CASchools$math + CASchools$read)
eq1 <- lm(testscr~str,data=CASchools)
eq2 <- lm(testscr~str+lunch,data=CASchools)
eq3 <- lm(testscr~str+lunch+english,data=CASchools)
stargazer::stargazer(eq1,eq2,eq3,type="text",no.space = TRUE)

##
## -----
##                               Dependent variable:
##                               -----
##                               testscr
##                               (1)          (2)          (3)
## -----
## str           -2.280***      -1.117***     -0.998***
##                 (0.480)        (0.240)        (0.239)
## lunch          -0.600***     -0.547***     -0.547***
##                 (0.017)        (0.022)        (0.022)
## english         -0.122**     -0.122**      (0.032)
##                 (0.032)
## Constant       698.933***    702.911***    700.150...
##                 (9.467)        (4.700)        (4.686)
## -----
## Observations      420          420          420
## R2              0.051         0.767         0.775
## Adjusted R2      0.049         0.766         0.773
## Residual Std. Error 18.581 (df = 418)    9.222 (df = 417)    9.080 (df ...
## F Statistic      22.575*** (df = 1; 418) 685.756*** (df = 2; 417) 476.306*** (df ...
## -----
## Note:           *p<0.1; **p<0.05;
```

2.3.3 Irrelevant variable

Consider the *True model*:

$$\mathbf{y} = \mathbf{X}_1\beta_1 + \varepsilon,$$

while the *Estimated model* is:

$$\mathbf{y} = \mathbf{X}_1\beta_1 + \mathbf{X}_2\beta_2 + \varepsilon$$

The estimates are unbiased. However, adding irrelevant explanatory variables increases the variance of the estimate of β_1 (compared to the case where one uses the correct explanatory variables). This is the case unless the correlation between \mathbf{X}_1 and \mathbf{X}_2 is null, see Eq. (2.22).

In other words, the estimator is *inefficient*, i.e., there exists an alternative consistent estimator whose variance is lower. The inefficiency problem can have serious consequences when testing hypotheses of type $H_0 : \beta_1 = 0$ due to the loss of power, so we might infer that they are no relevant variables when they truly are (Type-II error; False Negative).

2.4 Instrumental Variables

Nice of interpretation of tests

The conditional mean zero assumption (Hypothesis 2.2), according to which $\mathbb{E}(\varepsilon|\mathbf{X}) = 0$ —which implies in particular that \mathbf{x}_i and ε_i are uncorrelated— is sometimes not consistent with the considered economic framework. When it is the case, the parameters of interest may still be estimated consistently by resorting to instrumental variable techniques.

Consider the following model:

$$y_i = \mathbf{x}_i' \beta + \varepsilon_i, \quad \text{where } \mathbb{E}(\varepsilon_i) = 0 \text{ and } \mathbf{x}_i \perp \!\!\! \perp \varepsilon_i. \quad (2.23)$$

Let us illustrate how this situation may result in biased OLS estimate. Consider for instance the situation where:

$$\mathbb{E}(\varepsilon_i) = 0 \quad \text{and} \quad \mathbb{E}(\varepsilon_i \mathbf{x}_i) = \gamma, \quad (2.24)$$

in which case we have $\mathbf{x}_i \perp \!\!\! \perp \varepsilon_i$ (consistently with Eq. (2.23)).

By the law of large numbers, $\text{plim}_{n \rightarrow \infty} \mathbf{X}' \varepsilon / n = \gamma$. If $\mathbf{Q}_{xx} := \text{plim} \mathbf{X}' \mathbf{X} / n$, the OLS estimator is not consistent because

$$\mathbf{b} = \beta + (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \varepsilon \xrightarrow{P} \beta + \mathbf{Q}_{xx}^{-1} \gamma \neq \beta.$$

Let us now introduce the notion of instruments.

Definition 2.2 (Instrumental variables). The L -dimensional random variable \mathbf{z}_i is a valid set of instruments if:

- a. \mathbf{z}_i is correlated to \mathbf{x}_i ;
- b. we have $\mathbb{E}(\varepsilon|\mathbf{Z}) = 0$ and
- c. the orthogonal projections of the \mathbf{x}_i 's on the \mathbf{z}_i 's are not multicollinear.

If \mathbf{z}_i is a valid set of instruments, we have:

$$\text{plim} \left(\frac{\mathbf{Z}'\mathbf{y}}{n} \right) = \text{plim} \left(\frac{\mathbf{Z}'(\mathbf{X}\beta + \varepsilon)}{n} \right) = \text{plim} \left(\frac{\mathbf{Z}'\mathbf{X}}{n} \right) \beta$$

Indeed, by the law of large numbers, $\frac{\mathbf{Z}'\varepsilon}{n} \xrightarrow{P} \mathbb{E}(\mathbf{z}_i\varepsilon_i) = 0$.

If $L = K$, the matrix $\frac{\mathbf{Z}'\mathbf{X}}{n}$ is of dimension $K \times K$ and we have:

$$\left[\text{plim} \left(\frac{\mathbf{Z}'\mathbf{X}}{n} \right) \right]^{-1} \text{plim} \left(\frac{\mathbf{Z}'\mathbf{y}}{n} \right) = \beta.$$

By continuity of the inverse funct.: $\left[\text{plim} \left(\frac{\mathbf{Z}'\mathbf{X}}{n} \right) \right]^{-1} = \text{plim} \left(\left(\frac{\mathbf{Z}'\mathbf{X}}{n} \right)^{-1} \right)$. The Slutsky Theorem (Prop. 7.12) further implies that:

$$\text{plim} \left(\frac{\mathbf{Z}'\mathbf{X}}{n} \right)^{-1} \text{plim} \left(\frac{\mathbf{Z}'\mathbf{y}}{n} \right) = \text{plim} \left(\left(\frac{\mathbf{Z}'\mathbf{X}}{n} \right)^{-1} \frac{\mathbf{Z}'\mathbf{y}}{n} \right).$$

Hence \mathbf{b}_{iv} is consistent if it is defined by:

$$\boxed{\mathbf{b}_{iv} = (\mathbf{Z}'\mathbf{X})^{-1}\mathbf{Z}'\mathbf{y}.}$$

Proposition 2.12 (Asymptotic distribution of the IV estimator). *If \mathbf{z}_i is a L -dimensional random variable that constitutes a valid set of instruments (see Def. 2.2) and if $L = K$, then the asymptotic distribution of \mathbf{b}_{iv} is:*

$$\mathbf{b}_{iv} \xrightarrow{d} \mathcal{N} \left(\beta, \frac{\sigma^2}{n} [Q_{xz} Q_{zz}^{-1} Q_{zx}]^{-1} \right)$$

where $\text{plim } \mathbf{Z}'\mathbf{Z}/n =: \mathbf{Q}_{zz}$, $\text{plim } \mathbf{Z}'\mathbf{X}/n =: \mathbf{Q}_{zx}$, $\text{plim } \mathbf{X}'\mathbf{Z}/n =: \mathbf{Q}_{xz}$.

Proof. The proof is very similar to that of Prop. 2.11, the starting point being that $\mathbf{b}_{iv} = \beta + (\mathbf{Z}'\mathbf{X})^{-1}\mathbf{Z}'\varepsilon$. \square

When $L = K$, we have:

$$[Q_{xz} Q_{zz}^{-1} Q_{zx}]^{-1} = Q_{zx}^{-1} Q_{zz} Q_{xz}^{-1}$$

In practice, to estimate $\text{Var}(\mathbf{b}_{iv}) = \frac{\sigma^2}{n} Q_{zx}^{-1} Q_{zz} Q_{xz}^{-1}$, we replace σ^2 by:

$$s_{iv}^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{x}'_i \mathbf{b}_{iv})^2.$$

What about when $L > K$? In this case, we proceed as follows:

1. Regress \mathbf{X} on the space spanned by \mathbf{Z} and

2. Regress \mathbf{y} on the fitted values $\hat{\mathbf{X}} := \mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{X}$.

The results in:

$$\mathbf{b}_{iv} = [\mathbf{X}'\mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{X}]^{-1}\mathbf{X}'\mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{Y}. \quad (2.25)$$

In this case, Prop. 2.12 still holds, with \mathbf{b}_{iv} given by Eq. (2.25).

\mathbf{b}_{iv} is also the result of the regression of \mathbf{y} on \mathbf{X}^* , where the columns of \mathbf{X}^* are the (orthogonal) projections of those of \mathbf{X} on \mathbf{Z} , i.e. $\mathbf{X}^* = \mathbf{P}^Z\mathbf{X}$ (using the notations introduced in Eq. (2.5)). Hence the other names of this estimator: **Two-Stage Least Squares (TSLS)**.

If the instruments do not properly satisfy Condition (a) in Def. 2.2 (i.e. if \mathbf{x}_i and \mathbf{z}_i are only loosely related), the instruments are said to be **weak** (see, e.g., Stock and Yogo (2005), available here or Andrews et al. (2019)). A simple standard way to test for weak instruments consist in looking at the F-statistic associated with the first stage of the estimation. The easier it is to reject the null hypothesis (large test statistic), the less weak the instruments.

The Durbin-Wu-Hausman test (Durbin (1954), Wu (1973), Hausman (1978)) can be used to test if IV necessary. IV techniques are required if $\text{plim}_{n \rightarrow \infty} \mathbf{X}'\varepsilon/n \neq 0$. Hausman (1978) proposes a test of the efficiency of estimators. Under the null hypothesis two estimators, \mathbf{b}_0 and \mathbf{b}_1 , are consistent but \mathbf{b}_0 is (asymptotically) efficient relative to \mathbf{b}_1 . Under the alternative hypothesis, \mathbf{b}_1 (IV in the present case) remains consistent but not \mathbf{b}_0 (OLS in the present case). That is, when we reject the null hypothesis, it means that the OLS estimator is not consistent, potentially due to endogeneity issue.

The test statistic is:

$$H = (\mathbf{b}_1 - \mathbf{b}_0)'MPI(\mathbb{V}ar(\mathbf{b}_1) - \mathbb{V}ar(\mathbf{b}_0))(\mathbf{b}_1 - \mathbf{b}_0),$$

where MPI is the Moore-Penrose pseudo-inverse. Under the null hypothesis, $H \sim \chi^2(q)$, where q is the rank of $\mathbb{V}ar(\mathbf{b}_1) - \mathbb{V}ar(\mathbf{b}_0)$.

Example 2.3 (Estimation of price elasticity). See e.g. WHO and estimation of tobacco price elasticity of demand.

We want to estimate what is the effect on demand of an *exogenous increase* in prices of cigarettes (say).

The model is:

$$\begin{aligned} \log(\text{demand}) &= \alpha_0 + \alpha_1 \underbrace{\times p_t}_{\text{log(price)}} + \alpha_2 \underbrace{\times w_t}_{\text{income}} + \varepsilon_t^d \\ \log(\text{supply}) &= \gamma_0 + \gamma_1 \times p_t + \gamma_2 \underbrace{\times \mathbf{y}_t}_{\text{cost factors}} + \varepsilon_t^s, \end{aligned}$$

where $\mathbf{y}_t, w_t, \varepsilon_t^s \sim \mathcal{N}(0, \sigma_s^2)$ and $\varepsilon_t^d \sim \mathcal{N}(0, \sigma_d^2)$ are independent.

Equilibrium: $q_t^d = q_t^s$. This implies that prices are **endogenous**:

$$p_t = \frac{\alpha_0 + \alpha_2 w_t + \varepsilon_t^d - \gamma_0 - \gamma_2 \mathbf{y}_t - \varepsilon_t^s}{\gamma_1 - \alpha_1}.$$

In particular we have $\mathbb{E}(p_t \varepsilon_t^d) = \frac{\sigma_d^2}{\gamma_1 - \alpha_1} \neq 0 \Rightarrow$ Regressing by OLS q_t^d on p_t gives biased estimates (see Eq. (2.24)).




Figure 2.5: This figure illustrates the situation prevailing when estimating a price-elasticity (and the price is endogenous).

Let us use IV regressions to estimate the price elasticity of cigarette demand. For that purpose, we use the **CigarettesSW** dataset of package **AER** (these data are used by Stock and Watson (2003)). This panel dataset documents cigarette consumption for the 48 continental US States from 1985–1995. The instrument is the real tax on cigarettes arising from the state’s general sales tax. The rationale is that larger general sales tax drives cigarette prices up, but the general tax is not determined by other forces affecting ε_t^d .

```
data("CigarettesSW", package = "AER")
CigarettesSW$rprice <- with(CigarettesSW, price/cpi)
CigarettesSW$rincome <- with(CigarettesSW, income/population/cpi)
CigarettesSW$tdiff <- with(CigarettesSW, (taxs - tax)/cpi)
```

```

## model
eq.IV1 <- ivreg(log(packs) ~ log(rprice) + log(rincome) | log(rincome) + tdiff + I(tax/cpi),
                   data = CigarettesSW, subset = year == "1995")
eq.IV2 <- ivreg(log(packs) ~ log(rprice) | tdiff,
                   data = CigarettesSW, subset = year == "1995")
eq.no.IV <- lm(log(packs) ~ log(rprice) + log(rincome),
                   data = CigarettesSW, subset = year == "1995")
stargazer::stargazer(eq.no.IV, eq.IV1, eq.IV2, type="text", no.space = TRUE)

##
## =====
##                               Dependent variable:
##                               -----
##                               log(packs)
##                               OLS          instrumental
##                               variable
##                               (1)          (2)          (3)
## -----
## log(rprice)           -1.407***      -1.277***      -1.084***  

##                         (0.251)        (0.263)        (0.317)  

## log(rincome)          0.344          0.280  

##                         (0.235)        (0.239)  

## Constant              10.342***     9.895***     9.720***  

##                         (1.023)        (1.059)        (1.514)
## -----
## Observations          48            48            48
## R2                    0.433          0.429          0.401
## Adjusted R2           0.408          0.404          0.388
## Residual Std. Error   0.187 (df = 45)    0.188 (df = 45)  0.190 (df = 46)
## F Statistic           17.165*** (df = 2; 45)
## =====
## Note: *p<0.1; **p<0.05; ***p<0.01

summary(eq.IV1, diagnostics = TRUE)$diagnostics

##             df1 df2 statistic      p-value
## Weak instruments 2  44  244.7337536 1.444054e-24
## Wu-Hausman       1  44   3.0678163 8.682505e-02
## Sargan           1  NA   0.3326221 5.641191e-01

```

Example 2.4 (Education and wage). In this example, we make use of another dataset proposed by Stock and Watson (2003), namely the `CollegeDistance` dataset.² the objective is to estimate the effect of education on wages. Education

²Cross-section data from the High School and Beyond survey conducted by the Department of Education in the 80s. The survey includes students from approximately 1,100 high schools.

choice is suspected to be an endogenous variable, which calls for an IV strategy. The instrumental variable is the distance to college.

```

library(sem)
data("CollegeDistance", package = "AER")
eq.1st.stage <- lm(education ~ urban + gender + ethnicity + unemp + distance,
                    data = CollegeDistance)
CollegeDistance$ed.pred<- predict(eq.1st.stage)
eq.2nd.stage <- lm(wage ~ urban + gender + ethnicity + unemp + ed.pred ,
                     data = CollegeDistance)
eqOLS <- lm(wage ~ urban + gender + ethnicity + unemp + education,
              data=CollegeDistance)
eqTSLS <- ivreg(wage ~ urban + gender + ethnicity + unemp + education |
                  urban + gender + ethnicity + unemp + distance,
                  data=CollegeDistance)
stargazer::stargazer(eq.1st.stage, eq.2nd.stage, eqTSLS, eqOLS, type="text", no.space = TRUE)

## 
## =====
##                               Dependent variable:
##                               -----
##                               education          wage
##                               OLS      OLS   instrumental    OLS
##                               variable
##                               (1)      (2)      (3)      (4)
## -----
## urbanyes                   -0.092    0.046    0.046    0.070
##                               (0.065)  (0.045)  (0.060)  (0.045)
## genderfemale                -0.025   -0.071*   -0.071   -0.085**
##                               (0.052)  (0.037)  (0.050)  (0.037)
## ethnicityafam               -0.524*** -0.227*** -0.227** -0.556***
##                               (0.072)  (0.073)  (0.099)  (0.052)
## ethnicityhispanic            -0.275*** -0.351*** -0.351*** -0.544***
##                               (0.068)  (0.057)  (0.077)  (0.049)
## unemp                       0.010    0.139*** 0.139*** 0.133***
##                               (0.010)  (0.007)  (0.009)  (0.007)
## distance                     -0.087*** 
##                               (0.012)
## ed.pred                      0.647*** 
##                               (0.101)
## education                     0.647*** 0.005
##                               (0.136) (0.010)
## Constant                      14.061*** -0.359   -0.359   8.641***
##                               (0.083) (1.412) (1.908) (0.157)
## -----

```

```

## Observations           4,739      4,739      4,739      4,739
## R2                   0.023      0.117     -0.612      0.110
## Adjusted R2          0.022      0.116     -0.614      0.109
## Residual Std. Error (df = 4732) 1.770      1.263      1.706      1.268
## F Statistic (df = 6; 4732)    18.552*** 104.971*** 97.274*** 
## =====
## Note:                  *p<0.1; **p<0.05; ***p<0.01

```

2.5 General Regression Model (GRM) and robust covariance matrices

The statistical inference presented above relies on strong assumptions regarding the stochastic properties of the errors. Namely, they are assumed to be mutually uncorrelated (Hypothesis @ref(hyp:noncorrel_resid)) and homoskedastic (Hypothesis 2.3).

The objective of this section is to present approaches aimed at adjusting the estimate of the covariance matrix of the OLS estimator ($(\mathbf{X}'\mathbf{X})^{-1}s^2$, see Eq. (2.21)), when the previous hypotheses do not hold.

2.5.1 Presentation of the General Regression Model (GRM)

It will prove useful to introduce the following notation:

$$\text{Var}(\varepsilon|\mathbf{X}) = \mathbb{E}(\varepsilon\varepsilon'|\mathbf{X}) = \Sigma. \quad (2.26)$$

Note that Eq. ((2.26)) is more general than Hypothesis 2.3 and @ref(hyp:noncorrel_resid) because the diagonal entries of Σ may be different (as opposed to under Hypothesis 2.3), and the non-diagonal entries of Σ can be non-null (as opposed to under Hypothesis 2.4).

Definition 2.3 (General Regression Model (GRM)). Hypothesis 2.1 and 2.2, together with Eq. (2.26), form the General Regression Model (GRM) framework.

Naturally, a regression model where Hypotheses 2.1 to 2.4 hold is a specific case of the GRM framework.

The GRM context notably encompasses situations of heteroskedasticity and autocorrelation:

- Heteroskedasticity:

$$\Sigma = \begin{bmatrix} \sigma_1^2 & 0 & \dots & 0 \\ 0 & \sigma_2^2 & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \dots & 0 & \sigma_n^2 \end{bmatrix}. \quad (2.27)$$

- Autocorrelation:

$$\Sigma = \sigma^2 \begin{bmatrix} 1 & \rho_{2,1} & \dots & \rho_{n,1} \\ \rho_{2,1} & 1 & & \vdots \\ \vdots & & \ddots & \rho_{n,n-1} \\ \rho_{n,1} & \rho_{n,2} & \dots & 1 \end{bmatrix}. \quad (2.28)$$

Example 2.5 (Auto-regressive processes). Autocorrelation is, in particular, a recurrent problem when time-series data are used (see Section 6).

In a time-series context, subscript i refers to a date. Assume for instance that:

$$y_i = \mathbf{x}'_i \beta + \varepsilon_i \quad (2.29)$$

with

$$\varepsilon_i = \rho \varepsilon_{i-1} + v_i, \quad v_i \sim \mathcal{N}(0, \sigma_v^2). \quad (2.30)$$

In this case, we are in the GRM context, with:

$$\Sigma = \frac{\sigma_v^2}{1 - \rho^2} \begin{bmatrix} 1 & \rho & \dots & \rho^{n-1} \\ \rho & 1 & & \vdots \\ \vdots & & \ddots & \rho \\ \rho^{n-1} & \rho^{n-2} & \dots & 1 \end{bmatrix}. \quad (2.31)$$

In some cases —in particular when one assumes a parametric formulation for Σ — one can determine a better (more accurate) estimator than the OLS one. This approach is called Generalized Least Squares (GLS), which we present below.

2.5.2 Generalized Least Squares

Assume Σ is known (“feasible GLS”). Because Σ is symmetric positive, it admits a spectral decomposition of the form $\Sigma = \mathbf{C}\Lambda\mathbf{C}'$, where \mathbf{C} is an orthogonal matrix (i.e. $\mathbf{C}\mathbf{C}' = Id$) and Λ is a diagonal matrix (the diagonal entries are the eigenvalues of Σ).

We have $\Sigma = (\mathbf{P}\mathbf{P}')^{-1}$ with $\mathbf{P} = \mathbf{C}\Lambda^{-1/2}$. Consider the transformed model:

$$\mathbf{P}'\mathbf{y} = \mathbf{P}'\mathbf{X}\beta + \mathbf{P}'\varepsilon \quad \text{or} \quad \mathbf{y}^* = \mathbf{X}^*\beta + \varepsilon^*.$$

The variance of ε^* is \mathbf{I} . In the transformed model, OLS is BLUE (Gauss-Markow Theorem 2.1).

The **Generalized least squares** estimator of β is:

$$\mathbf{b}_{GLS} = (\mathbf{X}'\Sigma^{-1}\mathbf{X})^{-1}\mathbf{X}'\Sigma^{-1}\mathbf{y}. \quad (2.32)$$

We have:

$$\text{Var}(\mathbf{b}_{GLS}|\mathbf{X}) = (\mathbf{X}'\Sigma^{-1}\mathbf{X})^{-1}.$$

However, in general, Σ is unknown. The GLS estimator is then said to be *infeasible*. Some structure is required. Assume Σ admits a parametric form $\Sigma(\theta)$. The estimation becomes *feasible* (FGLS) if one replaces $\Sigma(\theta)$ by $\Sigma(\hat{\theta})$, where $\hat{\theta}$ is a consistent estimator of θ . In that case, the FGLS is asymptotically efficient (see Example ??).

When Σ has no obvious structure: the OLS (or IV) is the only estimator available. Under regularity assumptions, it remains unbiased, consistent, and asymptotically normally distributed, but not efficient. Standard inference procedures are no longer appropriate.

Example 2.6 (GLS in the auto-correlation case). Consider the case presented in Example 2.5. Because the OLS estimate \mathbf{b} of β is consistent, the estimates e_i of the ε_i 's also are. Consistent estimators of ρ and σ_v are then obtained by regressing the e_i 's on the e_{i-1} s. Using these estimates in Eq. (2.31) provides a consistent estimate of Σ . Applying these steps recursively gives an efficient estimator of β (Cochrane and Orcutt (1949)).

2.5.3 Asymptotic properties of the OLS estimator in the GRM framework

Since $\mathbf{b} = \beta + (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\varepsilon$ and $\text{Var}(\varepsilon|\mathbf{X}) = \Sigma$, we have:

$$\text{Var}(\mathbf{b}|\mathbf{X}) = \frac{1}{n} \left(\frac{1}{n} \mathbf{X}'\mathbf{X} \right)^{-1} \left(\frac{1}{n} \mathbf{X}'\Sigma\mathbf{X} \right) \left(\frac{1}{n} \mathbf{X}'\mathbf{X} \right)^{-1}. \quad (2.33)$$

The conditional covariance matrix of the OLS estimator is therefore not $\sigma^2(\mathbf{X}'\mathbf{X})^{-1}$ any longer. Therefore, using $s^2(\mathbf{X}'\mathbf{X})^{-1}$ for inference may be misleading. Below, we will see appropriate how to construct appropriate estimates of the covariance matrix of \mathbf{b} . Before, that, let us prove that the OLS estimator remains consistent in the GRM framework.

Proposition 2.13 (Consistency of the OLS estimator in the GRM framework). *If $\text{plim}(\mathbf{X}'\mathbf{X}/n)$ and $\text{plim}(\mathbf{X}'\Sigma\mathbf{X}/n)$ are finite positive definite matrices, then $\text{plim}(\mathbf{b}) = \beta$.*

Proof. We have $\text{Var}(\mathbf{b}) = \mathbb{E}[\text{Var}(\mathbf{b}|\mathbf{X})] + \text{Var}[\mathbb{E}(\mathbf{b}|\mathbf{X})]$. Since $\mathbb{E}(\mathbf{b}|\mathbf{X}) = \beta$, $\text{Var}[\mathbb{E}(\mathbf{b}|\mathbf{X})] = 0$. Eq. (2.33) implies that $\text{Var}(\mathbf{b}|\mathbf{X}) \rightarrow 0$. Hence \mathbf{b} converges in mean square, and therefore in probability (see Prop. ??). \square

Prop. 2.14 gives the asymptotic distribution of the OLS estimator in the GRM framework.

Proposition 2.14 (Asymptotic distribution of the OLS estimator in the GRM framework). *If $Q_{xx} = \text{plim}(\mathbf{X}'\mathbf{X}/n)$ and $Q_{x\Sigma x} = \text{plim}(\mathbf{X}'\Sigma\mathbf{X}/n)$ are finite positive definite matrices, then:*

$$\sqrt{n}(\mathbf{b} - \beta) \xrightarrow{d} \mathcal{N}(0, Q_{xx}^{-1}Q_{x\Sigma x}Q_{xx}^{-1}).$$

The IV estimator also features a normal asymptotic distribution:

Proposition 2.15 (Asymptotic distribution of the IV estimator in the GRM framework). *If regressors and IV variables are “well-behaved”, then:*

$$\mathbf{b}_{iv} \xrightarrow{a} \mathcal{N}(\beta, \mathbf{V}_{iv}),$$

where

$$\mathbf{V}_{iv} = \frac{1}{n}(\mathbf{Q}^*) \text{plim} \left(\frac{1}{n} \mathbf{Z}' \Sigma \mathbf{Z} \right) (\mathbf{Q}^*)',$$

with

$$\mathbf{Q}^* = [\mathbf{Q}_{xz} \mathbf{Q}_{zz}^{-1} \mathbf{Q}_{zx}]^{-1} \mathbf{Q}_{xz} \mathbf{Q}_{zz}^{-1}.$$

For practical purposes, one needs to have estimates of Σ in Props. 2.14 or 2.15. The complication comes from the fact that Σ is of dimension $n \times n$, and its estimation —based on a sample of length n — is therefore infeasible in the general case. Notwithstanding, looking at Eq. (2.33), it appears that one can focus on the estimation of $Q_{x\Sigma x} = \text{plim}(\mathbf{X}'\Sigma\mathbf{X}/n)$ (or $\text{plim}(\frac{1}{n}\mathbf{Z}'\Sigma\mathbf{Z})$ in the IV case). This matrix being of dimension $K \times K$, its estimation is easier.

We have:

$$\frac{1}{n} \mathbf{X}' \Sigma \mathbf{X} = \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n \sigma_{i,j} \mathbf{x}_i \mathbf{x}_j' \quad (2.34)$$

The so-called **robust covariance matrices** are estimates of the previous matrix. Their computation is based on the fact that if \mathbf{b} is consistent, then the e_i 's are consistent (pointwise) estimators of the ε_i 's.

In the following sections (2.5.4 and 2.5.5), we present such robust covariance matrices in two broad cases.

2.5.4 HAC-robust covariance matrices

When only heteroskedasticity prevails, i.e., when matrix Σ is as in Eq. (2.27), then one can use the formula proposed by White (1980) to estimate $\frac{1}{n}\mathbf{X}'\Sigma\mathbf{X}$ (see Example 2.7). When the residuals feature both heteroskedasticity and auto-correlation, then one can use the Newey and West (1987) approach (see Example 2.8).

Example 2.7 (Heteroskedasticity). This is the case of Eq. (2.27). In this case, we have $\sigma_{i,j} = 0$ for $i \neq j$. Hence, in this case, we then need to estimate $\frac{1}{n} \sum_{i=1}^n \sigma_i^2 \mathbf{x}_i \mathbf{x}'_i$. White (1980) has shown that, under general conditions:

$$\text{plim} \left(\frac{1}{n} \sum_{i=1}^n \sigma_i^2 \mathbf{x}_i \mathbf{x}'_i \right) = \text{plim} \left(\frac{1}{n} \sum_{i=1}^n e_i^2 \mathbf{x}_i \mathbf{x}'_i \right). \quad (2.35)$$

The estimator of $\frac{1}{n} \mathbf{X}' \Sigma \mathbf{X}$ therefore is:

$$M_{HC0} = \frac{1}{n} \mathbf{X}' \begin{bmatrix} e_1^2 & 0 & \dots & 0 \\ 0 & e_2^2 & & \\ \vdots & & \ddots & 0 \\ 0 & \dots & 0 & e_n^2 \end{bmatrix} \mathbf{X}. \quad (2.36)$$

where the e_i are the OLS residuals of the regression. The previous estimator is often called **HC0**. The **HC1** estimator, due to MacKinnon and White (1985), is obtained by applying an adjustment factor $n/(n-K)$ for the number of degrees of freedom (as in Prp. 2.6). That is:

$$M_{HC1} = \frac{n}{n-K} M_{HC0}. \quad (2.37)$$

We can illustrate the influence of heteroskedasticity using simulations. Consider the following model:

$$y_i = x_i + \varepsilon_i, \quad \varepsilon_i \sim \mathcal{N}(0, x_i^2),$$

where the x_i 's are i.i.d. $t(6)$.

Here is a simulated sample ($n = 200$) of this model:

```
n <- 200
x <- rt(n, df=6)
y <- x + x*rnorm(n)
par(plt=c(.1,.95,.1,.95))
plot(x,y,pch=19)
```

We simulate 1000 samples of the same model with $n = 200$. For each sample, we compute the OLS estimate of $\beta (=1)$. For each of the 1000 OLS estimations, we employ (a) the standard OLS variance formula ($s^2(\mathbf{X}'\mathbf{X})^{-1}$) and (b) the White formula to estimate the variance of b . For each formula, we compute the average of the 1000 resulting standard deviations and compare these with the standard deviation of the 1000 OLS estimate of β .

```
n <- 200 # sample size
N <- 1000 # number of simulated samples
XX <- matrix(rt(n*N, df=6), n, N)
```




Figure 2.6: Situation of heteroskedasticity. The model is $y_i = x_i + \varepsilon_i$, $\varepsilon_i \sim \mathcal{N}(0, x_i^2)$, where the x_i 's are i.i.d. $t(6)$.

```

YY <- matrix(XX + XX*rnorm(n), n, N)
all_b      <- NULL; all_V_OLS    <- NULL; all_V_White <- NULL
for(j in 1:N){
  Y <- matrix(YY[,j], ncol=1)
  X <- matrix(XX[,j], ncol=1)
  b <- solve(t(X) %*% X) %*% t(X) %*% Y
  e <- Y - X %*% b
  S <- 1/n * t(X) %*% diag(c(e^2)) %*% X
  V_OLS   <- solve(t(X) %*% X) * var(e)
  V_White <- 1/n * (solve(1/n*t(X) %*% X)) %*% S %*% (solve(1/n*t(X) %*% X))
  all_b    <- c(all_b, b)
  all_V_OLS <- c(all_V_OLS, V_OLS)
  all_V_White <- c(all_V_White, V_White)
}
c(sd(all_b), mean(sqrt(all_V_OLS)), mean(sqrt(all_V_White)))
## [1] 0.13711930 0.06659173 0.13676988

```

The results show that the White formula yields, on average, an estimated standard deviation that is much closer to the “true” value than the standard OLS formula. The latter underestimate the standard deviation of b .

In the following example, we regress GDP growth rates from the Jordà et al. (2017) database on a systemic financial crisis dummy. We compute the HC0-

and HC1-based standard deviations of the parameter estimate, and compare it to the one based on the standard OLS formula.

```
library(lmtest)
library(sandwich)
nT <- dim(JST)[1]
JST$growth <- NaN
JST$growth[2:nT] <- log(JST$rgdpbarro[2:nT]/JST$rgdpbarro[1:(nT-1)])
JST.red <- subset(JST, year>1950)
JST.red$iso <- as.factor(JST.red$iso)
JST.red$year <- as.factor(JST.red$year)
eq <- lm(growth~crisisJST+iso, data=JST.red)
rbind(coefest(eq)[2,],
      coefest(eq, vcov = vcovHC(eq, type = "HC0"))[2,],
      coefest(eq, vcov = vcovHC(eq, type = "HC1"))[2,])

##           Estimate Std. Error   t value   Pr(>|t|) 
## [1,] -0.02481424 0.005490411 -4.519560 6.789284e-06
## [2,] -0.02481424 0.005605452 -4.426804 1.040602e-05
## [3,] -0.02481424 0.005648268 -4.393247 1.212105e-05
```

Example 2.8 (Heteroskedasticity and Autocorrelation (HAC)). Newey and West (1987) have proposed a formula to address both heteroskedasticity and auto-correlation of the residuals (Eqs. (2.27) and (2.28)). They show that, if the correlation between terms i and j gets sufficiently small when $|i-j|$ increases:

$$\begin{aligned} \text{plim} \left(\frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n \sigma_{i,j} \mathbf{x}_i \mathbf{x}'_j \right) &\approx \\ \text{plim} \left(\frac{1}{n} \sum_{t=1}^n e_t^2 \mathbf{x}_t \mathbf{x}'_t + \frac{1}{n} \sum_{\ell=1}^L \sum_{t=\ell+1}^n w_\ell e_t e_{t-\ell} (\mathbf{x}_t \mathbf{x}'_{t-\ell} + \mathbf{x}_{t-\ell} \mathbf{x}'_t) \right) \end{aligned} \quad (2.38)$$

where $w_\ell = 1 - \ell/(L+1)$ (with L large).

Let us illustrate the influence of autocorrelation using simulations. We consider the following model:

$$y_i = x_i + \varepsilon_i, \quad \varepsilon_i \sim \mathcal{N}(0, x_i^2), \quad (2.39)$$

where the x_i 's and the ε_i 's are such that:

$$x_i = 0.8x_{i-1} + u_i \quad \text{and} \quad \varepsilon_i = 0.8\varepsilon_{i-1} + v_i, \quad (2.40)$$

where the u_i 's and the v_i 's are i.i.d. $\mathcal{N}(0, 1)$.

We simulate 500 samples of the same model with $n = 200$. For each sample, we compute the OLS estimate of β (=1). For each of the 1000 OLS estimations, we employ (a) the standard OLS variance formula ($s^2(\mathbf{X}'\mathbf{X})^{-1}$), (b) the White

formula, and (c) the Newey-West formula to estimate the variance of b . For each formula, using we compute the average of the 500 resulting standard deviations and compare these with the standard deviation of the 500 OLS estimate of β .

```

library(AEC)
n <- 100 # sample length
nb.sim <- 500 # number of simulated samples
all.b <- NULL; all.OLS.stdv.b <- NULL; all.Whi.stdv.b <- NULL; all.NW.stdv.b <- NULL
for(i in 1:nb.sim){
  eps <- rnorm(n); x <- rnorm(n)
  for(i in 2:n){
    eps[i] <- eps[i] + .8*eps[i-1]
    x[i] <- x[i] + .8*x[i-1]
  }
  y <- x + eps
  eq <- lm(y~x)
  all.b <- c(all.b,eq$coefficients[2])
  all.OLS.stdv.b <- c(all.OLS.stdv.b,summary(eq)$coefficients[2,2])
  X <- cbind(rep(1,n),x)
  XX <- t(X) %*% X; XX_1 <- solve(XX)
  E2 <- diag(eq$residuals^2)
  White.V <- XX_1 %*% (t(X) %*% E2 %*% X) %*% XX_1
  all.Whi.stdv.b <- c(all.Whi.stdv.b,sqrt(White.V[2,2]))
  # HAC:
  U <- X * cbind(eq$residuals,eq$residuals)
  XSigmaX <- NW.LongRunVariance(U,5)
  NW.V <- 1/n * (n*XX_1) %*% XSigmaX %*% (n*XX_1)
  all.NW.stdv.b <- c(all.NW.stdv.b,sqrt(NW.V[2,2]))
}
cbind(sd(all.b),mean(all.OLS.stdv.b),mean(all.Whi.stdv.b),mean(all.NW.stdv.b))

##          [,1]      [,2]      [,3]      [,4]
## [1,] 0.22045 0.1014711 0.09696198 0.1490803

```

The results show that the Newey-West formula yields, on average, an estimated standard deviation that is closer to the “true” value than the standard OLS formula. The latter underestimate the standard deviation of b .

What precedes suggest that, when the residuals feature autocorrelation, it is important to use appropriately adjusted covariance matrices to make statistical inference. How to detect autocorrelation in residuals? A popular test has been proposed by Durbin and Watson (1950) and Durbin and Watson (1951). The

Durbin-Watson test statistic is:

$$DW = \frac{\sum_{i=2}^n (e_i - e_{i-1})^2}{\sum_{i=1}^n e_i^2} = 2(1 - r) - \frac{e_1^2 + e_n^2}{\sum_{i=1}^n e_i^2},$$

$\xrightarrow{p} 0$

where r is the slope in the regression of the e_i 's on the e_{i-1} 's, i.e.:


$$r = \frac{\sum_{i=2}^n e_i e_{i-1}}{\sum_{i=1}^{n-1} e_i^2}.$$

(r is a consistent estimator of $\text{Cor}(\varepsilon_i, \varepsilon_{i-1})$, i.e. ρ in Eq. (2.30).)

The one-sided test for $H_0: \rho = 0$ against $H_1: \rho > 0$ is carried out by comparing DW to values $d_L(T, K)$ and $d_U(T, K)$:

$$\begin{cases} \text{If } DW < d_L, & \text{the null hypothesis is rejected;} \\ \text{if } DW > d_U, & \text{the hypothesis is not rejected;} \\ \text{If } d_L \leq DW \leq d_U, & \text{no conclusion is drawn.} \end{cases}$$

```
library(car)
data <- subset(JST, iso=="USA"); T <- dim(data)[1]
data$infl <- c(NaN, 100*log(data$cpi[2:T]/data$cpi[1:(T-1)]))
data$infl[(data$infl < -5) | (data$infl > 10)] <- NaN
par(mfrow=c(1,2))
plot(data$year, data$stir, ylim=c(-10, 20), type="l", lwd=2, xlab="", ylab="", main="Nominal rate and infl")
lines(data$year, data$infl, col="red", lwd=2)
eq <- lm(stir~infl, data=data)
plot(eq$residuals, type="l", col="blue", main="Residuals", xlab="", ylab="")
```



```
durbinWatsonTest(eq)
```

```
##   lag Autocorrelation D-W Statistic p-value
##     1          0.7321902    0.4984178      0
## Alternative hypothesis: rho != 0
```

2.5.5 Cluster-robust covariance matrices

The present section, based on MacKinnon et al. (2022), explains how to compute estimates of $Q_{x\Sigma_x} = \text{plim}(\mathbf{X}'\Sigma\mathbf{X}/n)$ (see Prop. 2.14) when the dataset can be decomposed into clusters. These clusters are such that residuals may be correlated within a given cluster, but not across different clusters. A cluster may, e.g., gather entities from a given geographical area, from the same industry, or same age cohort. Another useful reference is Cameron and Miller (2014).

The OLS estimator satisfies:

$$\mathbf{b} = \beta + (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\varepsilon. \quad (2.41)$$

Consider a set $\{n_1, n_2, \dots, n_G\}$ s.t. $n = \sum_g n_g$, on which is based the following decomposition of \mathbf{X} :

$$\mathbf{X} = \begin{bmatrix} \mathbf{X}_1 \\ \mathbf{X}_2 \\ \vdots \\ \mathbf{X}_G \end{bmatrix}.$$

With these notations, Eq. (2.41) rewrites:

$$\mathbf{b} - \beta = \left(\sum_{g=1}^G \mathbf{X}'_g \mathbf{X}_g \right)^{-1} \mathbf{X}' \sum_{g=1}^G \mathbf{s}_g, \quad (2.42)$$

where

$$\mathbf{s}_g = \mathbf{X}'_g \varepsilon_g \quad (2.43)$$

denotes the score vector (of dimension $K \times 1$) associated with the g^{th} cluster.

If the model is correctly specified then $\mathbb{E}(\mathbf{s}_g) = 0$ for all clusters g . Note that Eq. (2.42) is valid for any partition of $\{1, \dots, n\}$. Dividing the sample into clusters becomes meaningful if we further assume that the following hypothesis holds:

Hypothesis 2.6 (Clusters). We have:

$$(i) \mathbb{E}(\mathbf{s}_g \mathbf{s}'_g) = \Sigma_g, \quad (ii) \mathbb{E}(\mathbf{s}_g \mathbf{s}'_q) = 0, \quad g \neq q,$$

where s_g is defined in Eq. (2.43).

The real assumption here is (ii); the first one simply gives a notation for the covariance matrix of the score associated with the g^{th} cluster. Remark that these covariance matrices can differ across clusters. That is, **cluster-based inference is robust against both heteroskedasticity and intra-cluster dependence without imposing any restrictions on the (unknown) form of either of them**.

Naturally, matrix Σ_g depends on the covariance structure of the ε 's. In particular, if $\Omega_g = \mathbb{E}(\varepsilon_g \varepsilon'_g | \mathbf{X}_g)$, then we have $\Sigma_g = \mathbb{E}(\mathbf{X}'_g \Omega_g \mathbf{X}_g)$.

Under Hypothesis 2.6, it comes that the conditional covariance matrix of \mathbf{b} is:

$$(\mathbf{X}' \mathbf{X})^{-1} \left(\sum_{g=1}^G \Sigma_g \right) (\mathbf{X}' \mathbf{X})^{-1} \quad (2.44)$$

Let us denote by $\varepsilon_{g,i}$ the error associated with the i^{th} component of vector ε_g . Consider the special case where $\mathbb{E}(\varepsilon_{g,i} \varepsilon_{g,j} | \mathbf{X}_g) = \sigma^2 \mathbb{I}_{\{i=j\}}$, then Eq. (2.44) gives the standard expression $\sigma^2 (\mathbf{X}' \mathbf{X})^{-1}$ (see Eq. (2.9)).

If we have $\mathbb{E}(\varepsilon_{gi} \varepsilon_{gj} | \mathbf{X}_g) = \sigma_{gi}^2 \mathbb{I}_{\{i=j\}}$, then we fall in the case addressed by the White formula (see Example (??)), i.e.:

$$(\mathbf{X}' \mathbf{X})^{-1} \left(\mathbf{X}' \begin{bmatrix} \sigma_1^2 & 0 & \dots & 0 \\ 0 & \sigma_2^2 & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \dots & 0 & \sigma_n^2 \end{bmatrix} \mathbf{X} \right) (\mathbf{X}' \mathbf{X})^{-1}.$$

As in White (1980), the natural way to approach the conditional covariance given in Eq. (2.44) consists in replacing the Σ_g matrices by their sample equivalent, i.e. $\widehat{\Sigma}_g = \mathbf{X}'_g \mathbf{e}_g \mathbf{e}'_g \mathbf{X}_g$. Adding corrections for the degrees of freedom, this leads to the following estimate of the covariance matrix of \mathbf{b} :

$$\frac{G(n-1)}{(G-1)(n-K)} (\mathbf{X}'\mathbf{X})^{-1} \left(\sum_{g=1}^G \widehat{\Sigma}_g \right) (\mathbf{X}'\mathbf{X})^{-1}. \quad (2.45)$$

The previous estimate is CRCV1 in MacKinnon et al. (2022). Note that we indeed find the White-MacKinnon estimator (Eq. (2.37)) when $G = n$.

Remark that, if there was only one cluster ($G = 1$), and neglecting the degree-of-freedom correction, we would have:

$$(\mathbf{X}'\mathbf{X})^{-1} (\mathbf{X}'\mathbf{e}\mathbf{e}'\mathbf{X}) (\mathbf{X}'\mathbf{X})^{-1} = 0$$

because $\mathbf{X}'\mathbf{e} = 0$. Hence, having large clusters does not necessarily increase variance.

Often, when working with panel data (see Chapter 3), we want to cluster in different dimensions. A typical case is when the data are indexed by both individuals (i) and time (t). In that case, we may indeed suspect that: (a) the residuals are correlated across clusters of dates (e.g., with monthly data, a cluster may be one year) and (b) the residuals are correlated across clusters of individuals (e.g., with data at the county level a cluster may be a state). In this case, one can employ **two-way clustering**.

Formally, consider two distinct partitions of the data: one through index g , with $g \in \{1, \dots, G\}$, and the other through index h , with $h \in \{1, \dots, H\}$. Accordingly, we denote by $\mathbf{X}_{g,h}$ the submatrix of \mathbf{X} that contains the explanatory variables corresponding to clusters g and h (e.g., the firms of a given country g at a given date h). We also denote by $\mathbf{X}_{g,\bullet}$ (respectively $\mathbf{X}_{\bullet,h}$) the submatrix of \mathbf{X} containing all explanatory variables pertaining to cluster g , for all possible values of h (resp. to cluster h , for all possible values of g).

We will make the following assumption:

Hypothesis 2.7 (Two-way clusters). We have:

$$\begin{aligned} \mathbb{E}(\mathbf{s}_{g,\bullet} \mathbf{s}'_{g,\bullet}) &= \Sigma_g, & \mathbb{E}(\mathbf{s}_{\bullet,h} \mathbf{s}'_{\bullet,h}) &= \Sigma_h^*, & \mathbb{E}(\mathbf{s}_{g,h} \mathbf{s}'_{g,h}) &= \Sigma_{g,h}, \\ \mathbb{E}(\mathbf{s}_{g,h} \mathbf{s}'_{q,k}) &= 0 \text{ if } g \neq q \text{ and } h \neq k. \end{aligned}$$

Proposition 2.16 (Covariance of scores in the two-way-cluster setup). *Under this assumption, the matrix of covariance of the scores is given by:*

$$\Sigma = \sum_{g=1}^G \Sigma_g + \sum_{h=1}^H \Sigma_h^* - \sum_{g=1}^G \sum_{h=1}^H \Sigma_{g,h}.$$

(The last term on the right-hand side must be subtracted in order to avoid double counting.)

Proof. We have:

$$\begin{aligned}\Sigma &= \sum_{g=1}^G \sum_{q=1}^G \sum_{h=1}^H \sum_{k=1}^H \mathbf{s}_{g,h} \mathbf{s}'_{q,k} \\ &= \sum_{g=1}^G \underbrace{\left(\sum_{h=1}^H \sum_{k=1}^H \mathbf{s}_{g,h} \mathbf{s}'_{g,k} \right)}_{=\Sigma_g} + \sum_{h=1}^H \underbrace{\left(\sum_{g=1}^G \sum_{q=1}^G \mathbf{s}_{g,h} \mathbf{s}'_{q,h} \right)}_{=\Sigma_h^*} - \sum_{g=1}^G \sum_{h=1}^H \mathbf{s}_{g,h} \mathbf{s}'_{g,h},\end{aligned}$$

which gives the result. \square

The asymptotic theory can be based on two different approaches: (i) large number of clusters (common case), and (ii) fixed number of clusters but large number of observations in each cluster (see Subsections 4.1 and 4.2 in MacKinnon et al. (2022)). The more variable the N_g 's, the less reliable asymptotic inference based on Eq. (2.45), especially when a very few clusters are unusually large, or when the distribution of the data is heavy-tailed. These issues are somehow mitigated when the clusters have an approximate factor structure.

In practice, Σ is estimated by:

$$\widehat{\Sigma} = \sum_{g=1}^G \widehat{\mathbf{s}}_{g,\bullet} \widehat{\mathbf{s}}'_{g,\bullet} + \sum_{h=1}^H \widehat{\mathbf{s}}_{\bullet,h} \widehat{\mathbf{s}}'_{\bullet,h} - \sum_{g=1}^G \sum_{h=1}^H \widehat{\mathbf{s}}_{g,h} \widehat{\mathbf{s}}'_{g,h},$$

and we use:

$$\widehat{\text{Var}}(\mathbf{b}) = (\mathbf{X}' \mathbf{X})^{-1} \widehat{\Sigma} (\mathbf{X}' \mathbf{X})^{-1}.$$

As an alternative to the asymptotic approximation to the distribution of a statistic of interest, one can resort to bootstrap approximation (see Section 5 of MacKinnon et al. (2022)). In R, the package **fwildclusterboot** allows to implement such approaches.³

Let us come back to the analysis of the effect of systemic financial crises on GDP growth. Clustering the data at the country level and, further, at both the country and time levels gives the following:

```
eq <- lm(growth~crisisJST+iso, data=JST.red)
rbind(coeftest(eq)[2,],
      coeftest(eq, vcov = vcovHC(eq, type = "HC1"))[2,],
      coeftest(eq, vcov = vcovCL(eq, cluster = JST.red[, c("iso")]))[2,],
      coeftest(eq, vcov = vcovCL(eq, cluster = JST.red[, c("iso", "year")]))[2,])

##           Estimate Std. Error   t value   Pr(>|t|) 
## [1,] -0.02481424 0.005490411 -4.519560 6.789284e-06
## [2,] -0.02481424 0.005648268 -4.393247 1.212105e-05
## [3,] -0.02481424 0.005847708 -4.243413 2.365508e-05
## [4,] -0.02481424 0.006546931 -3.790209 1.577572e-04
```

³See, e.g., this tutorial by Alexander Fischer).

2.6 Shrinkage methods

Choosing the appropriate explanatory variables is often complicated, especially in the presence of many potentially relevant covariates. Keeping a large number of covariates results in large standard deviations for the estimated parameters (see Section 2.3.3). In order to address this issue, shrinkage methods have been designed. The objective of these methods is to help to select a limited number of variables (by shrinking the regression coefficients of the less useful variables towards zero). The two best-known shrinkage techniques are **ridge regression** and the **lasso** approach.⁴

In both cases (ridge and lasso), the OLS minimization problem (see Section 2.2), that is

$$\mathbf{b} = \underset{\beta}{\operatorname{argmin}} \sum_{i=1}^n (y_i - \mathbf{x}'_i \beta) \quad (2.46)$$

is replaced by the following:

$$\mathbf{b}_\lambda = \underset{\beta}{\operatorname{argmin}} \sum_{i=1}^n (y_i - \mathbf{x}'_i \beta) + \lambda f(\beta), \quad (2.47)$$

where $\lambda f(\beta)$ is a penalty term that positively depends on the “size” of the components of β . This term is called the *shrinkage penalty* term.

Specifically, assuming that the vector \mathbf{x}_i of potential covariates is of dimension $K \times 1$, we have:

$$\begin{aligned} f(\beta) &= \sum_{j=1}^K \beta_j^2 \quad \text{in the ridge case } (\ell_2 \text{ norm}), \\ f(\beta) &= \sum_{j=1}^K |\beta_j| \quad \text{in the lasso case } (\ell_1 \text{ norm}). \end{aligned}$$

In most cases, we do not want to involve the intercept in the set of parameters to shrink, and the preceding equations are respectively replaced with:

$$\begin{aligned} f(\beta) &= \sum_{j=2}^K \beta_j^2 \quad (\text{ridge}), \\ f(\beta) &= \sum_{j=2}^K |\beta_j| \quad (\text{lasso}). \end{aligned}$$

The nature of the penalty (based on the ℓ_1 or ℓ_2 norms) implies a different behaviour of the parameter estimates when λ –the *tuning parameter*– grows.

⁴See Tibshirani (2011) for a review of the lasso approach. See also Section 6.2 of James et al. (2013).

In the ridge regression, coefficient estimates go to zero (shrinkage); in the lasso case, some coefficients reach zero when λ reach some values. In other words, while ridge regression achieve shrinkage, lasso regressions achieve shrinkage *and* variable selection.

Parameter λ has to be determined separately from the minimization problem of Eq. (2.47). One can combine standard criteria (e.g., BIC or Akaike) with lasso regressions to help determine λ .

In R, one can use the `glmnet` package to run ridge and lasso regressions. In the following example, we employ this package to model the interest rates proposed to debtors. The data come from the Lending Club.

To begin with, let us define the variables we want to consider:

```
library(AEC)
library(glmnet)
credit$owner <- 1*(credit$home_ownership=="OWN")
credit$renter <- 1*(credit$home_ownership=="MORTGAGE")
credit$verification_status <- 1*(credit$verification_status=="Not Verified")
credit$emp_length_10 <- 1*(credit$emp_length_10)
credit$log_annual_inc <- log(credit$annual_inc)
credit$log_funded_amnt <- log(credit$funded_amnt)
credit$annual_inc2 <- (credit$annual_inc)^2
credit$funded_amnt2 <- (credit$funded_amnt)^2
x <- subset(credit,select = c(delinq_2yrs, annual_inc, annual_inc2, log_annual_inc, dti, installm
```

Let us standardize the data:


```
y <- credit$int_rate/sd(credit$int_rate,na.rm = TRUE)
stdv.x <- apply(x,2,function(a){sd(a,na.rm = TRUE)})
x <- x/t(matrix(stdv.x,dim(x)[2],dim(x)[1]))
```

Next, we define the set of λ we will use, and run the ridge and lasso regressions:

```
grid.lambda <- seq(0,.2,by=.005)
result.ridge <- glmnet(x, y, alpha = 0, lambda = grid.lambda)
result.lasso <- glmnet(x, y, alpha = 1, lambda = grid.lambda)
```

The following figure shows how estimated parameters depend on λ :

```
variab <- 6
plot(result.ridge$lambda,coef(result.ridge)[variab,],type="l",
      ylim=c(min(coef(result.ridge)[variab,]),max(coef(result.ridge)[variab,],coef(result.lasso)[variab,])),
      xlab=expression(lambda),ylab="Estimated parameter")
lines(result.lasso$lambda,coef(result.lasso)[variab,],col="red")
```



Let us take two values of λ and see the associated estimated parameters in the context of lasso regressions:

```
i <- 20; j <- 40
cbind(result.lasso$lambda[i],result.lasso$lambda[j])

##      [,1]  [,2]
## [1,] 0.105 0.005

cbind(coef(result.lasso)[,i],coef(result.lasso)[,j])

##                               [,1]          [,2]
## (Intercept)            3.24385583  6.4716266208
## delinq_2yrs           0.06308870  0.0689352682
## annual_inc            0.00000000  0.0045956535
## annual_inc2           0.00000000  0.0000000000
## log_annual_inc        0.00000000 -0.0361238200
## dti                   0.00000000  0.0224224582
## installment          0.14767959  8.2287289816
## verification_status   0.00000000 -0.0009750047
## funded_amnt           0.00000000 -7.3091694210
## funded_amnt2          0.00000000 -0.4711846250
## log_funded_amnt       0.00000000 -0.2460932367
## pub_rec                0.03390816  0.0599725219
```

```

## emp_length_10      0.00000000 -0.0192494122
## owner              0.00000000 -0.0244459908
## renter             -0.03882640 -0.0624308746
## pub_rec_bankruptcies 0.00000000  0.000000000000
## revol_util         0.00000000  0.000000000000
## revol_bal           0.00000000  0.0024022685

# Compute values of y predicted by the model, for all lambdas:
pred1 <- predict(result.lasso,as.matrix(x))
# Compute values of y predicted by the model, for a specific value:
pred2 <- predict(result.lasso,as.matrix(x),s=0.085)


```

The `glmnet` package (see Hastie et al. (2021)) also offers tools to implement cross-validation:

```

# cross validation:
cvglmnet <- cv.glmnet(as.matrix(x),y)
plot(cvglmnet)

```



```

cvglmnet$lambda.min # value of lambda.min, that is the value of lambda that gives minimum mean cr
## [1] 0.004091039

```

```

cvglmnet$lambda.1se # largest value of lambda that is such that the sample cost is within one standard error of the minimum

## [1] 0.00448991

coef(cvglmnet, s = "lambda.min") # associated parameters

## 18 x 1 sparse Matrix of class "dgCMatrix"
##                                     s1
## (Intercept)      6.583140504
## delinq_2yrs     0.066258597
## annual_inc      0.006054923
## annual_inc2     .
## log_annual_inc -0.037951509
## dti              0.020652724
## installment      8.518960621
## verification_status -0.002894435
## funded_amnt     -7.560852965
## funded_amnt2    -0.504192252
## log_funded_amnt -0.253756726
## pub_rec          0.058043332
## emp_length_10   -0.018946660
## owner            -0.024841917
## renter           -0.060398310
## pub_rec_bankruptcies .
## revol_util       .
## revol_bal        0.003093752

predict(cvglmnet, newx = as.matrix(x)[1:5,], s = "lambda.min") # predicted values of y

##      lambda.min
## 21529 3.807030
## 21547 3.416528
## 21579 4.026621
## 21583 3.255096
## 21608 3.340412

```

Chapter 3

Panel regressions

3.1 Specification and notations

A standard panel situation is as follows: the sample covers a lot of “entities”, indexed by $i \in \{1, \dots, n\}$, with n large, and, for each entity, we observe different variables over a small number of periods ($t \in \{1, \dots, T\}$). This is a *longitudinal dataset*.

The linear panel regression model is:

$$y_{i,t} = \underbrace{\mathbf{x}'_{i,t}}_{K \times 1} \underbrace{\beta}_{\text{Individual effects}} + \underbrace{\mathbf{z}'_i \alpha}_{\text{Individual effects}} + \varepsilon_{i,t}. \quad (3.1)$$

Our objective is to estimate the previous equation.

Figure 3.1 illustrates a panel-data situation. The model is $y_i = \alpha_i + \beta x_{i,t} + \varepsilon_{i,t}$, $t \in \{1, 2\}$. On Panel (b), blue dots are for $t = 1$, red dots are for $t = 2$. The lines relate the dots associated with the same entity i . What is remarkable in the simulated model is that, while the unconditional correlation between y and x is negative, the conditional correlation (conditional on α_i) is positive. Indeed, the sign of this conditional correlation is the sign of β , which is positive in the simulated example ($\beta = 5$). In other words, if one did not know the panel nature of the data, that would be tempting to say that $\beta < 0$, but this is not the case, due to **fixed effects** (the α_i 's) that are negatively correlated to the x_i 's.

```
T <- 2; n <- 12 # 2 periods and 12 entities
alpha <- 5*rnorm(n) # draw fixed effects
x.1 <- rnorm(n) - .5*alpha # note: x_i's correlate to alpha_i's
x.2 <- rnorm(n) - .5*alpha
beta <- 5; sigma <- .3
```

```

y.1 <- alpha + x.1 + sigma*rnorm(n);y.2 <- alpha + x.2 + sigma*rnorm(n)
x <- c(x.1,x.2) # pooled x
y <- c(y.1,y.2) # pooled y
par(mfrow=c(1,2))
plot(x,y,col="black",pch=19,xlab="x",ylab="y",main="(a)")
plot(x,y,col="black",pch=19,xlab="x",ylab="y",main="(b)")
points(x.1,y.1,col="blue",pch=19);points(x.2,y.2,col="red",pch=19)
for(i in 1:n){lines(c(x.1[i],x.2[i]),c(y.1[i],y.2[i]))}

```




Figure 3.1: The data are the same for both panels. On Panel (b), blue dots are for $t = 1$, red dots are for $t = 2$. The lines relate the dots associated to the same entity i .

Figure 3.2 presents the same type of plot based on the Cigarette Consumption Panel dataset (`CigarettesSW` dataset, used in Stock and Watson (2003)). This dataset documents the average consumption of cigarettes in 48 continental US states for two dates (1985 and 1995).

Notations:

$$\mathbf{y}_i = \begin{bmatrix} y_{i,1} \\ \vdots \\ y_{i,T} \end{bmatrix}_{T \times 1}, \quad \boldsymbol{\varepsilon}_i = \underbrace{\begin{bmatrix} \varepsilon_{i,1} \\ \vdots \\ \varepsilon_{i,T} \end{bmatrix}}_{T \times 1}, \quad \mathbf{x}_i = \underbrace{\begin{bmatrix} \mathbf{x}'_{i,1} \\ \vdots \\ \mathbf{x}'_{i,T} \end{bmatrix}}_{T \times K}, \quad \mathbf{X} = \underbrace{\begin{bmatrix} \mathbf{x}_1 \\ \vdots \\ \mathbf{x}_n \end{bmatrix}}_{(nT) \times K}.$$

$$\tilde{\mathbf{y}}_i = \begin{bmatrix} y_{i,1} - \bar{y}_i \\ \vdots \\ y_{i,T} - \bar{y}_i \end{bmatrix}, \quad \tilde{\boldsymbol{\varepsilon}}_i = \begin{bmatrix} \varepsilon_{i,1} - \bar{\varepsilon}_i \\ \vdots \\ \varepsilon_{i,T} - \bar{\varepsilon}_i \end{bmatrix},$$




Figure 3.2: Cigarette consumption versus real price in the CigarettesSW panel dataset.

$$\tilde{\mathbf{x}}_i = \begin{bmatrix} \mathbf{x}'_{i,1} - \bar{\mathbf{x}}'_i \\ \vdots \\ \mathbf{x}'_{i,T} - \bar{\mathbf{x}}'_i \end{bmatrix}, \quad \tilde{\mathbf{X}} = \begin{bmatrix} \tilde{\mathbf{x}}_1 \\ \vdots \\ \tilde{\mathbf{x}}_n \end{bmatrix}, \quad \tilde{\mathbf{Y}} = \begin{bmatrix} \tilde{y}_1 \\ \vdots \\ \tilde{y}_n \end{bmatrix},$$

where

$$\bar{y}_i = \frac{1}{T} \sum_{t=1}^T y_{i,t}, \quad \bar{\varepsilon}_i = \frac{1}{T} \sum_{t=1}^T \varepsilon_{i,t} \quad \text{and} \quad \bar{\mathbf{x}}_i = \frac{1}{T} \sum_{t=1}^T \mathbf{x}_{i,t}.$$

3.1.1 Three standard cases

- **Pooled regression:** $\mathbf{z}_i \equiv 1$. This case amounts to the case studied in Chapter 2.
- **Fixed Effects** (Section 3.2): \mathbf{z}_i is unobserved, but correlates with $\mathbf{x}_i \Rightarrow \mathbf{b}$ is biased and inconsistent in the OLS regression of \mathbf{y} on \mathbf{X} (omitted variable, see Section 2.3.2).
- **Random Effects** (Section 3.3): \mathbf{z}_i is unobserved, but uncorrelated with \mathbf{x}_i . The model writes:

$$y_{i,t} = \mathbf{x}'_{i,t}\beta + \alpha + \underbrace{u_i + \varepsilon_{i,t}}_{\text{compound disturbance}},$$

where $\alpha = \mathbb{E}(\mathbf{z}'_i\alpha)$ and $u_i = \mathbf{z}'_i\alpha - \mathbb{E}(\mathbf{z}'_i\alpha) \perp \mathbf{x}_i$. In that case, the OLS is consistent, but not efficient.

3.2 Estimation of Fixed-Effects Models

Hypothesis 3.1 (Fixed-effect model). We assume that:

- i. There is no perfect multicollinearity among the regressors.
- ii. $\mathbb{E}(\varepsilon_{i,t} | \mathbf{X}) = 0$, for all i, t .
- iii. We have:

$$\mathbb{E}(\varepsilon_{i,t} \varepsilon_{j,s} | \mathbf{X}) = \begin{cases} \sigma^2 & \text{if } i = j \text{ and } s = t, \\ 0 & \text{otherwise.} \end{cases}$$

These assumptions are analogous to those introduced in the standard linear regression:

$$(i) \leftrightarrow 2.1, (ii) \leftrightarrow 2.2, (iii) \leftrightarrow 2.3 + 2.4.$$

In matrix form, for a given i , the model writes:

$$\mathbf{y}_i = \mathbf{X}_i \beta + \mathbf{1} \alpha_i + \varepsilon_i,$$

where $\mathbf{1}$ is a T -dimensional vector of ones.

This is the **Least Square Dummy Variable (LSDV)** model:

$$\boxed{\mathbf{y} = [\mathbf{X} \quad \mathbf{D}] \begin{bmatrix} \beta \\ \alpha \end{bmatrix} + \varepsilon}, \quad (3.2)$$

with:

$$\mathbf{D} = \underbrace{\begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & & & \\ 0 & 0 & \dots & 1 \end{bmatrix}}_{(nT \times n)}.$$

The linear regression (Eq. (3.2)) –with the dummy variables– satisfies the Gauss-Markov conditions (Theorem 2.1). Hence, in this context, the OLS estimator is the **best linear unbiased estimator**.

Denoting by \mathbf{Z} the matrix $[\mathbf{X} \quad \mathbf{D}]$, and by \mathbf{b} and \mathbf{a} the respective OLS estimates of β and of α , we have:

$$\boxed{\begin{bmatrix} \mathbf{b} \\ \mathbf{a} \end{bmatrix} = [\mathbf{Z}' \mathbf{Z}]^{-1} \mathbf{Z}' \mathbf{y}.} \quad (3.3)$$

The asymptotical distribution of $[\mathbf{b}', \mathbf{a}']'$ derives from the standard OLS context: Prop. 2.11 can be used after having replaced \mathbf{X} by $\mathbf{Z} = [\mathbf{X} \quad \mathbf{D}]$.

We have:

$$\begin{bmatrix} \mathbf{b} \\ \mathbf{a} \end{bmatrix} \xrightarrow{d} \mathcal{N} \left(\begin{bmatrix} \beta \\ \alpha \end{bmatrix}, \sigma^2 \frac{Q^{-1}}{nT} \right) \quad (3.4)$$

where

$$Q = \text{plim}_{nT \rightarrow \infty} \frac{1}{nT} \mathbf{Z}' \mathbf{Z}.$$

In practice, an estimator of the covariance matrix of $[\mathbf{b}', \mathbf{a}']'$ is:

$$s^2 (\mathbf{Z}' \mathbf{Z})^{-1} \quad \text{with} \quad s^2 = \frac{\mathbf{e}' \mathbf{e}}{nT - K - n},$$

where \mathbf{e} is the $(nT) \times 1$ vector of OLS residuals.

There is an alternative way of expressing the LSDV estimators.

It is based on matrix $\mathbf{M}_D = \mathbf{I} - \mathbf{D}(\mathbf{D}'\mathbf{D})^{-1}\mathbf{D}'$, which acts as an operator that removes entity-specific means, i.e.:

$$\tilde{\mathbf{Y}} = \mathbf{M}_D \mathbf{Y}, \quad \tilde{\mathbf{X}} = \mathbf{M}_D \mathbf{X} \quad \text{and} \quad \tilde{\varepsilon} = \mathbf{M}_D \varepsilon.$$

With these notations, using Theorem 2.2, we get:

$$\mathbf{b} = [\mathbf{X}' \mathbf{M}_D \mathbf{X}]^{-1} \mathbf{X}' \mathbf{M}_D \mathbf{y}. \quad (3.5)$$

This amounts to regressing the $\tilde{y}_{i,t}$'s ($= y_{i,t} - \bar{y}_i$) on the $\tilde{\mathbf{x}}_{i,t}$'s ($= \mathbf{x}_{i,t} - \bar{\mathbf{x}}_i$).

The estimate of α is given by:

$$\mathbf{a} = (\mathbf{D}'\mathbf{D})^{-1} \mathbf{D}' (\mathbf{y} - \mathbf{X}\mathbf{b}), \quad (3.6)$$

which is obtained by developing the second row of

$$\begin{bmatrix} \mathbf{X}' \mathbf{X} & \mathbf{X}' \mathbf{D} \\ \mathbf{D}' \mathbf{X} & \mathbf{D}' \mathbf{D} \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ \mathbf{a} \end{bmatrix} = \begin{bmatrix} \mathbf{X}' \mathbf{Y} \\ \mathbf{D}' \mathbf{Y} \end{bmatrix},$$

which are the first-order conditions resulting from the least squares problem (see Eq. (2.3)).

One can use different types of fixed effects in the same regression. Typically, one can have time and entity fixed effects. In that case, the model writes:

$$y_{i,t} = \mathbf{x}'_i \beta + \alpha_i + \gamma_t + \varepsilon_{i,t}.$$

The LSDV approach (Eq. (3.2)) can still be resorted to. It suffices to extend the \mathbf{Z} matrix with additional columns (*time dummies*):

$$\mathbf{y} = [\mathbf{X} \quad \mathbf{D} \quad \mathbf{C}] \begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix} + \varepsilon, \quad (3.7)$$

with:

$$\mathbf{C} = \begin{bmatrix} \delta_1 & \delta_2 & \dots & \delta_{T-1} \\ \vdots & \vdots & & \vdots \\ \delta_1 & \delta_2 & \dots & \delta_{T-1} \end{bmatrix},$$

where the T -dimensional vector δ_t (the *time dummy*) is

$$[0, \dots, 0, \underset{\text{t^{th} entry}}{1}, 0, \dots, 0]'$$

Using state and year fixed effects in the `CigarettesSW` panel dataset yields the following results:

```
CigarettesSW$rincome <- with(CigarettesSW, income/population/cpi)
eq.pooled <- lm(log(packs)~log(rprice)+log(rincome), data=CigarettesSW)
eq.LSDV <- lm(log(packs)~log(rprice)+log(rincome)+state,
                 data=CigarettesSW)
CigarettesSW$year <- as.factor(CigarettesSW$year)
eq.LSDV2 <- lm(log(packs)~log(rprice)+log(rincome)+state+year,
                  data=CigarettesSW)
stargazer::stargazer(eq.pooled, eq.LSDV, eq.LSDV2, type="text", no.space = TRUE,
                      omit=c("state", "year"),
                      add.lines=list(c('State FE', 'No', 'Yes', 'Yes'),
                                    c('Year FE', 'No', 'No', 'Yes')),
                      omit.stat=c("f", "ser"))

##
## =====
##             Dependent variable:
##             -----
##                   log(packs)
##                   (1)      (2)      (3)
## ----- 
## log(rprice) -1.334*** -1.210*** -1.056*** 
##             (0.135)   (0.114)   (0.149)
## log(rincome)  0.318**   0.121     0.497 
##             (0.136)   (0.190)   (0.304)
## Constant     10.067*** 9.954*** 8.360*** 
##             (0.516)   (0.264)   (1.049)
## ----- 
## State FE      No       Yes      Yes
## Year FE       No       No       Yes
## Observations   96      96      96
## R2            0.552    0.966    0.967
## Adjusted R2   0.542    0.929    0.931
## =====
## Note:          *p<0.1; **p<0.05; ***p<0.01
```

Example 3.1 (Housing prices and interest rates). In this example, we want to estimate the effect of short and long-term interest rate on housing prices. The data come from the Jordà et al. (2017) dataset see this website.

```

library(AEC)
JST <- subset(JST, year>1950); N <- dim(JST)[1]
JST$hpreal <- JST$hpnom/JST$cpi # real house price index
JST$chge_ctry <- c(NaN, JST$iso[2:N] != JST$iso[1:(N-1)])# will be use to put NA's when computing ch
JST$dhpreal <- 100*log(JST$hpreal/c(NaN, JST$hpreal[1:(N-1)]))
JST$dhpreal[JST$chge_ctry==1] <- NaN
JST$dhpreal[abs(JST$dhpreal)>30] <- NaN # remove extreme price change
JST$YEAR <- as.factor(JST$year) # to have time fixed effects
eq1 <- lm(dhpreal ~ stir + ltrate + iso + YEAR, data=JST)
eq2 <- lm(dhpreal ~ I(ltrate-stir) + iso + YEAR, data=JST)
library(sandwich); library(lmtest)
vcov_cluster1 <- vcovHC(eq1, cluster = JST[, c("iso", "YEAR")])
vcov_cluster2 <- vcovHC(eq2, cluster = JST[, c("iso", "YEAR")])
robust_se_FE1 <- sqrt(diag(vcov_cluster1))
robust_se_FE2 <- sqrt(diag(vcov_cluster2))
stargazer::stargazer(eq1, eq1, eq2, type = "text",
                      column.labels = c("no Cluster", "Clusters", "no Cluster", "Clusters"),
                      omit = c("iso", "YEAR"), no.space = TRUE,
                      omit.labels = c("Country FE", "Year FE"), keep.stat = "n",
                      se = list(NULL, robust_se_FE1, NULL, robust_se_FE2))

## 
## =====
##          Dependent variable:
## -----
##                  dhpreal
##          no Cluster Clusters no Cluster Clusters
##                  (1)      (2)      (3)      (4)
## -----
## stir           0.532***  0.532***  

##                 (0.134)   (0.170)  

## ltrate        -0.384*** -0.384**  

##                 (0.145)   (0.182)  

## I(ltrate - stir)           -0.475*** -0.475***  

##                               (0.125)   (0.159)  

## Constant       -1.662    -1.662    -0.996    -0.996  

##                 (2.141)   (2.843)   (2.073)   (2.780)
## -----
## Country FE     Yes      Yes      Yes      Yes  

## Year FE       Yes      Yes      Yes      Yes
## -----

```

```

## Observations      1,141      1,141      1,141      1,141
## =====
## Note:           *p<0.1; **p<0.05; ***p<0.01

```

3.3 Estimation of random effects models

Here, the individual effects are assumed to be not correlated to other variables (the \mathbf{x}_i 's).

Random-effect models write:

$$y_{i,t} = \mathbf{x}'_{it}\beta + (\alpha + \underbrace{u_i}_{\text{Random heterogeneity}}) + \varepsilon_{i,t}$$

with

$$\begin{aligned} \mathbb{E}(\varepsilon_{i,t}|\mathbf{X}) &= \mathbb{E}(u_i|\mathbf{X}) = 0, \\ \mathbb{E}(\varepsilon_{i,t}\varepsilon_{j,s}|\mathbf{X}) &= \begin{cases} \sigma_\varepsilon^2 & \text{if } i = j \text{ and } s = t, \\ 0 & \text{otherwise.} \end{cases} \\ \mathbb{E}(u_i u_j|\mathbf{X}) &= \begin{cases} \sigma_u^2 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases} \\ \mathbb{E}(\varepsilon_{i,t} u_j|\mathbf{X}) &= 0 \quad \text{for all } i, j \text{ and } t. \end{aligned}$$

Let us introduce the following notations: $\eta_{i,t} = u_i + \varepsilon_{i,t}$ and $\eta_i = [\eta_{i,1}, \eta_{i,2}, \dots, \eta_{i,T}]'$.

We have $\mathbb{E}(\eta_i|\mathbf{X}) = \mathbf{0}$ and $\text{Var}(\eta_i|\mathbf{X}) = \Gamma$ where

$$\Gamma = \begin{bmatrix} \sigma_\varepsilon^2 + \sigma_u^2 & \sigma_u^2 & \sigma_u^2 & \dots & \sigma_u^2 \\ \sigma_u^2 & \sigma_\varepsilon^2 + \sigma_u^2 & \sigma_u^2 & \dots & \sigma_u^2 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \sigma_u^2 & \sigma_u^2 & \sigma_u^2 & \dots & \sigma_\varepsilon^2 + \sigma_u^2 \end{bmatrix} = \sigma_\varepsilon^2 \mathbf{I} + \sigma_u^2 \mathbf{1}\mathbf{1}'.$$

Denoting by Σ the covariance matrix of $\eta = [\eta'_1, \dots, \eta'_n]'$, we have:

$$\Sigma = \mathbf{I} \otimes \Gamma.$$

If we knew Σ , we would apply (feasible) GLS (Eq. (2.32), in Section 2.5.2):

$$\beta = (\mathbf{X}' \Sigma^{-1} \mathbf{X})^{-1} \mathbf{X}' \Sigma^{-1} \mathbf{y}.$$

(As explained in Section 2.5.2, this amounts to regressing $\Sigma^{-1/2}' \mathbf{y}$ on $\Sigma^{-1/2}' \mathbf{X}$.)

It can be checked that $\Sigma^{-1/2} = \mathbf{I} \otimes (\Gamma^{-1/2})$ where

$$\Gamma^{-1/2} = \frac{1}{\sigma_\varepsilon} \left(\mathbf{I} - \frac{\theta}{T} \mathbf{1}\mathbf{1}' \right),$$

with

$$\theta = 1 - \frac{\sigma_\varepsilon}{\sqrt{\sigma_\varepsilon^2 + T\sigma_u^2}}.$$

Hence, if we knew Σ , we would transform the data as follows:

$$\Gamma^{-1/2} \mathbf{y}_i = \frac{1}{\sigma_\varepsilon} \begin{bmatrix} y_{i,1} - \theta \bar{y}_i \\ y_{i,2} - \theta \bar{y}_i \\ \vdots \\ y_{i,T} - \theta \bar{y}_i \end{bmatrix}.$$

What about when Σ is unknown? One can take deviations from group means to remove heterogeneity:

$$y_{i,t} - \bar{y}_i = [\mathbf{x}_{i,t} - \bar{\mathbf{x}}_i]' \beta + (\varepsilon_{i,t} - \bar{\varepsilon}_i).$$

The previous equation can be consistently estimated by OLS. (Although the residuals are correlated across t 's for the observations pertaining to a given entity, the OLS remain consistent; see Prop. @ref{prp:XXX}).

We have $\mathbb{E} \left[\sum_{i=1}^T (\varepsilon_{i,t} - \bar{\varepsilon}_i)^2 \right] = (T-1)\sigma_\varepsilon^2$.

The $\varepsilon_{i,t}$'s are not observed but \mathbf{b} is a consistent estimator of β . Using an adjustment for the degrees of freedom, we have:

$$\hat{\sigma}_e^2 = \frac{1}{nT - n - K} \sum_{i=1}^n \sum_{t=1}^T (e_{i,t} - \bar{e}_i)^2.$$

What about σ_u^2 ? We can exploit the fact that OLS are consistent in the pooled regression:

$$\text{plim } s_{\text{pooled}}^2 = \text{plim } \frac{\mathbf{e}' \mathbf{e}}{nT - K - 1} = \sigma_u^2 + \sigma_\varepsilon^2,$$

and therefore use $s_{\text{pooled}}^2 - \hat{\sigma}_e^2$ as an approximation to σ_u^2 .

Let us come back to Example 3.1. In the following, we use the random effect specification; and compare the results with those from the pooled regression and the fixed-effect model. For that, we use the function `plm` of the package of the same name. Note that `eq.FE` is similar to `eq1` in Example 3.1.

```
library(plm)
eq.RE <- plm(dhpreal ~ stir + ltrate, data=JST, index=c("iso", "YEAR"), model="random",
               effect="twoways")
eq.FE <- plm(dhpreal ~ stir + ltrate, data=JST, index=c("iso", "YEAR"), model="within",
               effect="twoways")
eq0 <- plm(dhpreal ~ stir + ltrate, data=JST, index=c("iso", "YEAR"), model="pooling")
stargazer::stargazer(eq0, eq.RE, eq.FE, type = "text", no.space = TRUE,
                      column.labels = c("Pooled", "Random Effect", "Fixed Effects"),
                      add.lines=list(c('State FE', 'No', 'Yes', 'Yes'),
                                    c('Year FE', 'No', 'Yes', 'Yes')), omit.stat=c("f", "ser"))
```

```

## 
## =====
##          Dependent variable:
## -----
##                  dhpreal
##          Pooled   Random Effect Fixed Effects
##          (1)      (2)      (3)
## -----
## stir       0.485***    0.456***    0.532***  

##           (0.114)     (0.019)     (0.134)  

## ltrate     -0.690***   -0.541***   -0.384***  

##           (0.127)     (0.020)     (0.145)  

## Constant   4.103***    3.341***  

##           (0.421)     (0.096)
## -----
## State FE    No        Yes        Yes  

## Year FE    No        Yes        Yes  

## Observations 1,141    1,141    1,141  

## R2          0.027    0.024    0.015  

## Adjusted R2 0.025    0.022   -0.067
## =====
## Note: *p<0.1; **p<0.05; ***p<0.01

```

One can run an Hausman (1978) test in order to check whether or not the fixed-effect model is needed. Indeed, if this is not the case (i.e., if the covariates are not correlated to the disturbances), then it is preferable to use the random-effect estimation as the latter is more efficient.

```
phtest(eq.FE, eq.RE)
```

```

## 
## Hausman Test
## 
## data: dhpreal ~ stir + ltrate
## chisq = 3.8386, df = 2, p-value = 0.1467
## alternative hypothesis: one model is inconsistent

```

The p-value being high, we do not reject the null hypothesis according to which the covariates and the errors are uncorrelated. We should therefore prefer the random-effect model.

Example 3.2 (Spatial data). This example makes use of Airbnb prices (Zürich, 22 June 2017), collected from Tom Slee's website. The covariates are the number of bedrooms and the number of people that can be accommodated. We consider the use of district fixed effects. Figure 3.3 shows the price to explain. (The size of the circles is proportional to prices.)




Figure 3.3: Airbnb prices for the Zurich area, 22 June 2017. The size of the circles is proportional to prices.

Let us regress prices on the covariates as well as on district dummies:

```

eq_noFE <- lm(price~bedrooms+accommodates,data=airbnb)
eq_FE    <- lm(price~bedrooms+accommodates+neighborhood,data=airbnb)
# Adjust standard errors:
cov_FE      <- vcovHC(eq_FE, cluster = airbnb[, c("neighborhood")])
robust_se_FE <- sqrt(diag(cov_FE))
cov_noFE    <- vcovHC(eq_noFE, cluster = airbnb[, c("neighborhood")])
robust_se_noFE <- sqrt(diag(cov_noFE))
stargazer::stargazer(eq_FE, eq_noFE, eq_FE, type = "text",
                      column.labels = c("FE (no HAC)", "No FE (no HAC)",
                      "FE (with HAC)", "No FE (with HAC)"),
                      omit = c("neighborhood"), no.space = TRUE,
                      omit.labels = c("District FE"), keep.stat = "n",
                      se = list(NULL, NULL, robust_se_FE, robust_se_noFE))

##
## =====
##                               Dependent variable:
## -----
##                               price
## FE (no HAC) No FE (no HAC) FE (with HAC) No FE (with HAC)

```

```

##              (1)          (2)          (3)          (4)
## -----
## bedrooms    7.229***   5.629**   7.229***   5.629***
##             (2.135)    (2.194)    (2.052)    (2.073)
## accommodates 16.426*** 17.449*** 16.426*** 17.449*** 
##             (1.284)    (1.323)    (1.431)    (1.428)
## Constant     95.118***  68.417***  95.118***  68.417*** 
##             (5.323)    (3.223)    (5.664)    (3.527)
## -----
## District FE Yes        No        Yes        No
## -----
## Observations 1,321      1,321      1,321      1,321
## =====
## Note: *p<0.1; **p<0.05; ***p<0.01

```

Figure 3.4 compares the residuals with and without fixed effects. The sizes of the circles are proportional to the absolute values of the residuals, the color indicates the sign (blue for negative).




Figure 3.4: Regression residuals. The sizes of the circles are proportional to the absolute values of the residuals, the color indicates the sign (blue for negative).

With fixed effects, the colors are better balanced within each district.

3.4 Dynamic Panel Regressions

In what precedes, it has been assumed that there is no correlation between the observations indexed by (i, t) and those indexed by (j, s) as long as $j \neq i$ or $t \neq s$. If one suspects that the errors $\varepsilon_{i,t}$ are correlated (across entities i for a given date t , or across dates for a given entity, or both), then one should employ a robust covariance matrix (see Section ??).

In several cases, auto-correlation in the variable of interest may stem from an auto-regressive specification. That is, Eq. (3.1) is then replaced by:

$$y_{i,t} = \rho y_{i,t-1} + \mathbf{x}'_{i,t} \begin{matrix} \beta \\ \alpha_i \\ K \times 1 \end{matrix} + \varepsilon_{i,t}. \quad (3.8)$$

Individual effects

In that case, even if the explanatory variables $\mathbf{x}_{i,t}$ are uncorrelated to the errors $\varepsilon_{i,t}$, we have that the additional *explanatory variable* $y_{i,t-1}$ correlates to the errors $\varepsilon_{i,t-1}, \varepsilon_{i,t-2}, \dots, \varepsilon_{i,1}$. As a result, the LSDV estimate of the model parameters $\{\rho, \beta\}$ may be biased, even if n is large. To see this, notice that the LSDV regression amounts to regressing $\tilde{\mathbf{y}}$ on $\tilde{\mathbf{X}}$ (see Eq. (3.5)), where the elements of $\tilde{\mathbf{X}}$ are the explanatory variables to which we subtract their within-sample means. In particular, we have:

$$\tilde{y}_{i,t-1} = y_{i,t-1} - \frac{1}{T} \sum_{s=1}^T y_{i,s-1},$$

which correlates to the corresponding error, that is:

$$\tilde{\varepsilon}_{i,t} = \varepsilon_{i,t} - \frac{1}{T} \sum_{s=1}^T \varepsilon_{i,s}.$$

The previous equation shows that the *within-group* estimator (LSDV) introduces all realizations of the $\varepsilon_{i,t}$ errors into the transformed error term ($\tilde{\varepsilon}_{i,t}$). As a result, in large n , fixed T panels, it is consistent only if all the right-hand-side variables of the regression are strictly exogenous (i.e., do not correlate to past, present, and future errors $\varepsilon_{i,t}$).¹ This is not the case when there are lags of $y_{i,t}$ on the right-hand side of the regression formula.

The following simulation illustrate this bias. The x -coordinates of the dots are the fixed effects α_i 's, and the y -coordinates are their LSDV estimates. The blue line is the 45-degree line.

In the previous example, the estimate of ρ (whose true value is 0.9) is 0.666.

To address this, one can resort to instrumental-variable regressions. Anderson and Hsiao (1982) have, in particular, proposed a first-differenced Two Stage

¹The bias may vanish for large T 's, though.




Figure 3.5: illustration of the bias pertaining to the LSDV estimation approach in the presence of auto-correlation of the depend variable.

Least Squares (2SLS) estimator. This estimation is based on the following transformation of the model:

$$\Delta y_{i,t} = \rho \Delta y_{i,t-1} + (\Delta \mathbf{x}_{i,t})' \beta + \Delta \varepsilon_{i,t}. \quad (3.9)$$

The OLS estimates of the parameters are biased because ε_{i,t_1} —which is part of the error $\Delta \varepsilon_{i,t}$ — is correlated to $y_{i,t-1}$ —which is part of the “explanatory variable”, namely $\Delta y_{i,t-1}$. But consistent estimates can be obtained using 2SLS with instrumental variables that are both correlated with $\Delta y_{i,t}$ and orthogonal to $\Delta \varepsilon_{i,t}$. One can for instance use $\{y_{i,t-2}, \mathbf{x}_{i,t-2}\}$ as instruments. Note that this approach can be implemented only if there are more than 3 time observations per entity i .

If the explanatory variables $\mathbf{x}_{i,t}$ are assumed to be predetermined (i.e., do not contemporaneous correlate with the errors $\varepsilon_{i,t}$), then $\mathbf{x}_{i,t-1}$ can be added to the instruments associated with $\Delta y_{i,t}$. Further, if these variables (the $\mathbf{x}_{i,t}$'s) are exogenous (i.e., do not contemporaneous correlate with any of the errors $\varepsilon_{i,s}$, $\forall s$), then $\mathbf{x}_{i,t}$ also constitute a valid instrument.

Using the previous (simulated) example, this approach consists in the following steps:

```

Dy <- c(all_y[3:T,]) - c(all_y[2:(T-1),])
Dy_1 <- c(all_y[2:(T-1),]) - c(all_y[1:(T-2),])
y_2 <- c(all_y[1:(T-2),])
Z <- matrix(y_2, ncol=1)
Pz <- Z %*% solve(t(Z) %*% Z) %*% t(Z)
Dy_hat <- Pz %*% Dy; Dy_1hat <- Pz %*% Dy_1
rho_TSLS <- solve(t(Dy_1hat) %*% Dy_1hat) %*% t(Dy_1hat) %*% Dy_hat

```

While the OLS estimate of ρ (whose true value is 0.9) was 0.666, we obtain here $\text{rho_TSLS} = 1.617$.

Let us come back to the general case (with covariates $\mathbf{x}_{i,k}$'s). For $t = 3$, $y_{i,1}$ (and $\mathbf{x}_{i,1}$) is the only possible instrument. However, for $t = 4$, one could use $y_{i,2}$ and $y_{i,1}$ (as well as $\mathbf{x}_{i,2}$ and $\mathbf{x}_{i,1}$). More generally, defining matrix Z_i as follows:

$$Z_i = \begin{bmatrix} \mathbf{z}'_{i,1} & 0 & \dots & & & & \\ 0 & \mathbf{z}'_{i,1} & \mathbf{z}'_{i,2} & 0 & \dots & & \\ 0 & 0 & 0 & \mathbf{z}'_{i,1} & \mathbf{z}'_{i,2} & \mathbf{z}'_{i,3} & 0 & \dots \\ \vdots & & & & & & & \\ 0 & \dots & & & & & 0 & \mathbf{z}'_{i,1} & \dots & \mathbf{z}'_{i,T-2} \end{bmatrix},$$

where $\mathbf{z}_{i,t} = [y_{i,t}, \mathbf{x}'_{i,t}]'$, we have the moment conditions:²

$$\mathbb{E}(Z'_i \Delta \mathbf{v}_i) = 0,$$

with $\Delta \mathbf{v}_i = [\Delta v_{i,3}, \dots, \Delta v_{i,T}]'$.

These restrictions are used in the GMM approach employed by Arellano and Bond (1991). Specifically, a GMM estimator of the model parameters is given by:

$$\operatorname{argmin} \left(\frac{1}{n} \sum_{i=1}^n Z'_i \Delta \mathbf{v}_i \right)' W_n \left(\frac{1}{n} \sum_{i=1}^n Z'_i \Delta \mathbf{v}_i \right),$$

using the weighting matrix

$$W_n = \left(\frac{1}{n} \sum_{i=1}^n Z'_i \widehat{\Delta \mathbf{v}_i} \widehat{\Delta \mathbf{v}_i}' Z_i \right)^{-1},$$

where the $\widehat{\Delta \mathbf{v}_i}$'s are consistent estimates of the $\Delta \mathbf{v}_i$'s that result from a preliminary estimation. In this sense, this estimator is a two-step GMM one.

If the disturbances are homoskedastic, then it can be shown that an asymptotically equivalent (efficient) GMM estimator can be obtained by using:

$$W_{1,n} = \left(\frac{1}{n} Z'_i H Z_i \right)^{-1},$$

²If $\mathbf{x}_{i,t}$ is predetermined (exogenous), we can use $\mathbf{z}_{i,t} = [y_{i,t}, \mathbf{x}_{i,t+1}, \mathbf{x}'_{i,t}]'$ (respectively $\mathbf{z}_{i,t} = [y_{i,t}, \mathbf{x}_{i,t+2}, \mathbf{x}_{i,t+1}, \mathbf{x}'_{i,t}]'$).

where H is is $(T - 2) \times (T - 2)$ matrix of the form:

$$H = \begin{bmatrix} 2 & -1 & 0 & \dots & 0 \\ -1 & 2 & -1 & & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & & -1 & 2 & -1 \\ 0 & \dots & 0 & -1 & 2 \end{bmatrix}.$$

It is straightforward to extend these GMM methods to cases where there is more than one lag of the dependent variable on the right-hand side of the equation or in cases where disturbances feature limited moving-average serial correlation.

The `pdynmc` package allows to run these GMM approaches (see Fritsch et al. (2019)). The following lines of code allow to replicate the results of Arellano and Bond (1991):

```
library(pdynmc)
data(EmplUK, package = "plm")
dat <- EmplUK
dat[,c(4:7)] <- log(dat[,c(4:7)])
m1 <- pdynmc(dat = dat, # name of the dataset
               varname.i = "firm", # name of the cross-section identifier
               varname.t = "year", # name of the time-series identifiers
               use.mc.diff = TRUE, # use moment conditions from equations in differences
               use.mc.lev = FALSE, # use moment conditions from equations in levels? (i.e., use.mc.nonlin = FALSE)
               use.mc.nonlin = FALSE, # use nonlinear (quadratic) moment conditions?
               include.y = TRUE, # instruments should be derived from the lags of the dependent variable
               varname.y = "emp", # name of the dependent variable in the dataset
               lagTerms.y = 2, # number of lags of the dependent variable
               fur.con = TRUE, # further control variables (covariates) are included?
               fur.con.diff = TRUE, # include further control variables in equations from differences
               fur.con.lev = FALSE, # include further control variables in equations from levels
               varname.reg.fur = c("wage", "capital", "output"), # covariate(s) -in the regressions-
               lagTerms.reg.fur = c(1,2,2), # number of lags of the further controls
               include.dum = TRUE, # A logical variable indicating whether dummy variables are included
               dum.diff = TRUE, # A logical variable indicating whether dummy variables are included in differences
               dum.lev = FALSE, # A logical variable indicating whether dummy variables are included in levels
               varname.dum = "year",
               w.mat = "iid.err", # One of the character strings c('"iid.err"', '"identif"')
               std.err = "corrected",
               estimation = "onestep", # One of the character strings c('"onestep"', '"two-step"')
               opt.meth = "none" # numerical optimization procedure. When no nonlinear method is specified, this is used
)
summary(m1)

##
```

```

## Dynamic linear panel estimation (onestep)
## Estimation steps: 1
##
## Coefficients:
##              Estimate Std.Err.rob z-value.rob Pr(>|z.rob|)
## L1.emp      0.686226  0.144594    4.746   < 2e-16 ***
## L2.emp     -0.085358  0.056016   -1.524    0.12751
## L0.wage    -0.607821  0.178205   -3.411   0.00065 ***
## L1.wage     0.392623  0.167993    2.337   0.01944 *
## L0.capital  0.356846  0.059020    6.046   < 2e-16 ***
## L1.capital -0.058001  0.073180   -0.793    0.42778
## L2.capital -0.019948  0.032713   -0.610    0.54186
## L0.output   0.608506  0.172531    3.527   0.00042 ***
## L1.output   -0.711164  0.231716   -3.069   0.00215 **
## L2.output   0.105798  0.141202    0.749    0.45386
## 1979        0.009554  0.010290    0.929    0.35289
## 1980        0.022015  0.017710    1.243    0.21387
## 1981       -0.011775  0.029508   -0.399    0.68989
## 1982       -0.027059  0.029275   -0.924    0.35549
## 1983       -0.021321  0.030460   -0.700    0.48393
## 1976       -0.007703  0.031411   -0.245    0.80646
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## 41 total instruments are employed to estimate 16 parameters
## 27 linear (DIF)
## 8 further controls (DIF)
## 6 time dummies (DIF)
##
## J-Test (overid restrictions): 70.82 with 25 DF, pvalue: <0.001
## F-Statistic (slope coeff): 528.06 with 10 DF, pvalue: <0.001
## F-Statistic (time dummies): 14.98 with 6 DF, pvalue: 0.0204

```

We generate novel results (`m2`) by replacing “`onestep`” with “`twostep`” (in the `estimation` field). The resulting estimated coefficients are:

```

##      L1.emp      L2.emp      L0.wage      L1.wage      L0.capital  L1.capital
##  0.62870890 -0.06518800 -0.52575951  0.31128961  0.27836190  0.01409950
##      L2.capital  L0.output  L1.output      L2.output          1979          1980
## -0.04024847  0.59192286 -0.56598515  0.10054264  0.01121551  0.02306871
##          1981          1982          1983          1976
## -0.02135806 -0.03111604 -0.01799335 -0.02336762

```

Arellano and Bond (1991) have proposed a specification test. If the model is correctly specified, then the errors of Eq. (3.9) –that is the first-difference

equation– should feature non-zero first-order auto-correlation, but zero higher-order autocorrelation.

Function `mtest.fct` of package `pdynmc` implements this test. Here is its result in the present case:

```
mtest.fct(m1, order=2)

##
##  Arellano and Bond (1991) serial correlation test of degree 2
##
## data: 1step GMM Estimation
## normal = -0.16523, p-value = 0.8688
## alternative hypothesis: serial correlation of order 2 in the error terms
```

One can also implement the Hansen J-test of the over-identifying restrictions (see Section 4.1.3):

```
jtest.fct(m1)

##
##  J-Test of Hansen
##
## data: 1step GMM Estimation
## chisq = 70.82, df = 25, p-value = 2.905e-06
## alternative hypothesis: overidentifying restrictions invalid

jtest.fct(m2)

##
##  J-Test of Hansen
##
## data: 2step GMM Estimation
## chisq = 31.381, df = 25, p-value = 0.1767
## alternative hypothesis: overidentifying restrictions invalid
```

3.5 Introduction to program evaluation

This section briefly introduces the econometrics of program evaluation. A comprehensive review is proposed by Abadie and Cattaneo (2018). A seminal book on the subject is that of Angrist and Pischke (2008).

Program evaluation refer to the analysis of the causal effects of some “treatments” in a broad sense. These treatment can, e.g., correspond to the implementation (or announcement) of policy measures.

3.5.1 Presentation of the problem

To begin with, let us consider a single entity, so let us drop the entity index (i). Let us denote by Y the outcome variable (for the variable of interest), by W is a binary variable indicating whether the considered entity has received treatment ($W = 1$) or not ($W = 0$), and by X a vector of covariates, assumed to be predetermined relative to the treatment. That is, W and X could be correlated, but the values of X have been determined before that of W (in such a way that the realization of W does not affect X). Typically, X contains characteristics of the considered entity.

We are interested in the following:

$$Y_1 - Y_0,$$

where Y_1 correspond to the outcome obtained under treatment, while Y_0 is the outcome obtained without it. That is, we have:

$$Y = (1 - W)Y_0 + WY_1.$$

The problem is that observing (Y, W, X) is not sufficient to observe $Y_1 - Y_0$, that is the treatment effect. Additional assumptions are needed to estimate, in particular, the average treatment effect:

$$ATE = \mathbb{E}(Y_1 - Y_0).$$

Importantly, ATE is different from the following quantity:

$$\alpha = \underbrace{\mathbb{E}(Y|W=1)}_{=\mathbb{E}(Y_1|W=1)} - \underbrace{\mathbb{E}(Y|W=0)}_{=\mathbb{E}(Y_0|W=0)},$$

that is easier to estimate. Indeed, we won't have a lot of direct information regarding $\mathbb{E}(Y_0|W=1)$ and $\mathbb{E}(Y_1|W=0)$. However, these two conditional expectations are part of ATE . Indeed, $ATE = \mathbb{E}(Y_1) - \mathbb{E}(Y_0)$, and:

$$\begin{aligned} \mathbb{E}(Y_1) &= \mathbb{E}(Y_1|W=0)\mathbb{P}(W=0) + \mathbb{E}(Y_1|W=1)\mathbb{P}(W=1) \\ \mathbb{E}(Y_0) &= \mathbb{E}(Y_0|W=0)\mathbb{P}(W=0) + \mathbb{E}(Y_0|W=1)\mathbb{P}(W=1). \end{aligned}$$

3.5.2 Randomized controlled trials (RCTs)

In the context of Randomized controlled trials (RCTs), entities are randomly assigned to receive the treatment. As a result, we have $\mathbb{E}(Y_1) = \mathbb{E}(Y_1|W=0) = \mathbb{E}(Y_1|W=1)$ and $\mathbb{E}(Y_0) = \mathbb{E}(Y_0|W=0) = \mathbb{E}(Y_0|W=1)$. Hence, in this context, $ATE = \alpha$.

Therefore, in this context, estimating $\mathbb{E}(Y_1 - Y_0)$ amounts to computing the difference between two sample means, namely (a) the sample mean of the subset

of Y_i 's corresponding to the entities for which $W_i = 1$, and (b) the one for which $W_i = 0$.

More accurate estimates can be obtained through regressions. Assume that the model reads:

$$Y_i = W_i\beta_1 + Z'_i\beta_z + \varepsilon_i,$$

where $\mathbb{E}(\varepsilon_i|Z_i) = 0$ (and W_i is independent from Z_i and ε_i). In this case, we obtain a consistent estimate of β_0 by regressing \mathbf{y} on $\mathbf{X} = [\mathbf{w}, \mathbf{Z}]$.

3.5.3 Difference-in-Difference (DiD) approach

The DiD approach is a popular methodology implemented in cases where W cannot be considered as an independent variable. It exploits two dimensions: entities (i), and time (t). To simplify the exposition, we consider only two periods here ($t = 0$ and $t = 1$).

Consider the following model:

$$Y_{i,t} = W_{i,t}\beta_1 + \mu_i + \delta_t + \varepsilon_{i,t} \quad (3.10)$$

The parameter of interest is β_1 , which is the treatment effect. Usually, for all entities i , we have $W_{i,t=0} = 0$. But only some of them are treated on date 1, i.e., $W_{i,1} = 0 \in \{0, 1\}$.

The disturbance $\varepsilon_{i,t}$ affects the outcome, but it does not relate the selection for treatment; therefore, $\mathbb{E}(\varepsilon_{i,t}|W_{i,t}) = 0$. By contrast, we do not exclude some correlation between $W_{i,t}$ (for $t = 1$) and μ_i ; hence, μ_i may constitute a **confounder**.

We have:


$$\begin{aligned} \mathbb{E}(Y_{i,1}|W_{i,1} = 1) &= \beta_1 + \mathbb{E}(\mu_i|W_{i,1} = 1) + \mathbb{E}(\delta_1|W_{i,1} = 1) + \mathbb{E}(\varepsilon_{i,1}) \\ \mathbb{E}(Y_{i,0}|W_{i,1} = 1) &= \mathbb{E}(\mu_i|W_{i,1} = 1) + \mathbb{E}(\delta_0|W_{i,1} = 1) + \mathbb{E}(\varepsilon_{i,0}) \\ \mathbb{E}(Y_{i,1}|W_{i,1} = 0) &= \mathbb{E}(\mu_i|W_{i,1} = 0) + \mathbb{E}(\delta_1|W_{i,1} = 0) + \mathbb{E}(\varepsilon_{i,1}) \\ \mathbb{E}(Y_{i,0}|W_{i,1} = 0) &= \mathbb{E}(\mu_i|W_{i,1} = 0) + \mathbb{E}(\delta_0|W_{i,1} = 0) + \mathbb{E}(\varepsilon_{i,0}). \end{aligned}$$

and, therefore, it can be checked that:

$$\beta_1 = \mathbb{E}(\Delta Y_{i,1}|W_{i,1} = 1) - \mathbb{E}(\Delta Y_{i,1}|W_{i,1} = 0),$$

where $\Delta Y_{i,1} = Y_{i,1} - Y_{i,0}$. Therefore, in this context, the treatment effect appears to be a difference (of two conditional expectations) of difference (of the outcome variable, through time).

This is illustrated by Figure ??, which represents the generic DiD framework.

**Figure 5**

Identification in a difference-in-differences model. The dashed line represents the outcome that the treated units would have experienced in the absence of the treatment.

Figure 3.6: Source: Abadie et al., (1998).

In practice, implementing this approach consists in running a linear regression of the type of Eq. (3.10). These regressions also usually involve controls on top of the fixed effects μ_i . As illustrated in the next subsection, the parameter of interest (β_1) is often associated with an interaction term.

3.5.4 Application of the DiD approach

This example is based on the data used in Meyer et al. (1995). This dataset is part of the `wooldridge` package. This paper examines the effect of workers' compensation for injury on time out of work. It exploits a **natural experiment** approach of comparing individuals injured before and after increases in the maximum weekly benefit amount. Specifically, in 1980, the cap on weekly earnings covered by worker's compensation was increased. Let us check whether this new policy was followed by an increase in the amount of time workers spent unemployed (for example, higher compensation may reduce workers' incentives to avoid injury).

As shown in Figure ??, the measure has only affected high-earning workers. The idea exploited by Meyer et al. (1995) was to compare the increase in time out of work before-after 1980 for higher-earnings workers on the one hand (entities who received the treatment) and low-earnings workers on the other hand (control group).

The next lines of codes replicate some of their results. The dependent variable is the logarithm of the duration of benefits. For more information use `?injury`, after having loaded the `wooldridge` library.




FIGURE 1. TEMPORARY TOTAL BENEFIT SCHEDULE
BEFORE AND AFTER AN INCREASE IN
THE MAXIMUM WEEKLY BENEFIT

Figure 3.7: Source: Meyer et al., (1995).

In the table of results below, the parameter of interest is the one associated with the interaction term `afchng:highearn`. Columns 2 and 3 correspond to the first two column of Table 6 in Meyer et al. (1995).

```
library(wooldridge)
data(injury)
injury <- subset(injury,ky==1)
injury$indust <- as.factor(injury$indust)
injury$injtype <- as.factor(injury$injtype)
#names(injury)
eq1 <- lm(log(durat) ~ afchng + highearn + afchng*highearn,data=injury)
eq2 <- lm(log(durat) ~ afchng + highearn + afchng*highearn + lprewage*highearn + male
eq3 <- lm(log(durat) ~ afchng + highearn + afchng*highearn + lprewage*highearn + male
stargazer::stargazer(eq1,eq2,eq3,type="text",
                      order = c(1,2,18,3:17,19,20))

##
## =====
##                                     Dependent variable:
##                                     -----
##                                     log(durat)
##                                     (1)          (2)
## -----
## ## afchng           0.008          -0.004
## ##                         (0.045)        (0.038)
## ##
```

```

##          0.256***      -0.595      -1.522
##          (0.047)      (0.930)      (1.099)
##
## afchnge:highearn      0.191***      0.162***      0.215*** 
##          (0.069)      (0.059)      (0.069)
##
## lprewage                  0.207**      0.258** 
##          (0.088)      (0.104)
##
## male                     -0.070*      -0.072
##          (0.039)      (0.046)
##
## married                  0.055       0.051
##          (0.035)      (0.041)
##
## lage                     0.244***      0.252*** 
##          (0.044)      (0.052)
##
## ltotmed                  0.361***      0.249*** 
##          (0.011)      (0.059)
##
## hosp                     0.252***      0.173*** 
##          (0.044)      (0.042)
##
## indust2                  0.197***      0.249*** 
##          (0.050)      (0.059)
##
## indust3                  0.153***      0.173*** 
##          (0.035)      (0.042)
##
## injtype2                 0.790***      0.780*** 
##          (0.132)      (0.156)
##
## injtype3                 0.564***      0.348*** 
##          (0.078)      (0.092)
##
## injtype4                 0.575***      0.634*** 
##          (0.085)      (0.101)
##
## injtype5                 0.617***      0.501*** 
##          (0.078)      (0.092)
##
## injtype6                 0.616***      0.395*** 
##          (0.079)      (0.093)
##

```

```

## injtype7          0.958***   0.781*** (0.175) (0.175)
## 
## 
## injtype8          0.432***   0.511*** (0.109) (0.109)
## 
## 
## highearn:lprewage 0.065     0.065 (0.158) (0.158)
## 
## 
## Constant          1.126***  -3.536*** (-1.126) (-3.536)
## 
## 
## -----
## Observations      5,626       5,347      5,347
## R2                0.021       0.319      0.319
## Adjusted R2        0.020       0.316      0.316
## Residual Std. Error 1.269 (df = 5622) 1.054 (df = 5327) 1.246 (df = 5327)
## F Statistic        39.540*** (df = 3; 5622) 131.260*** (df = 19; 5327) 16.089*** (df = 19; 5327)
## 
## =====
## Note: *p<0.1; **p<0.05; ***p<0.01

```

Chapter 4

Estimation Methods

This chapter presents three approaches to estimate parametric models: the General Method of Moments (GMM), the Maximum Likelihood approach (ML), and the Bayesian approach. The general context is the following: You observe a sample $\mathbf{y} = \{y_1, \dots, y_n\}$; you assume that these data have been generated by a model parameterized by $\theta \in \mathbb{R}^K$, and you want to estimate this vector θ_0 .

4.1 Generalized Method of Moments (GMM)

4.1.1 Definition of the GMM estimator

We denote by y_i a $p \times 1$ vector of variables; by θ an $K \times 1$ vector of parameters, and by $h(y_i; \theta)$ a continuous $r \times 1$ vector-valued function.

We denote by θ_0 the true value of θ and we assume that θ_0 satisfies:

$$\mathbb{E}[h(y_i; \theta_0)] = 0.$$

We denote by y_i the information contained in the current and past observations of y_i , that is: $\underline{y}_i = \{y_i, y_{i-1}, \dots, y_1\}$. We denote by $g(\underline{y}_n; \theta)$ the sample average of $h(y_i; \theta)$, i.e.:

$$g(\underline{y}_n; \theta) = \frac{1}{n} \sum_{i=1}^n h(y_i; \theta).$$

The intuition behind the GMM estimator is the following: choose θ so as to make the sample moment as close as possible to their population values, that is 0.

Definition 4.1. A GMM estimator of θ_0 is given by:

$$\hat{\theta}_n = \operatorname{argmin}_{\theta} g(\underline{y}_n; \theta)' W_n g(\underline{y}_n; \theta),$$

where W_n is a positive definite matrix (that may depend on \underline{y}_n).

In the specific case where $K = r$ (the dimension of θ is the same as that of $h(y_i; \theta)$ —or of $g(\underline{y}_n; \theta)$ — then $\hat{\theta}_n$ is such that:

$$g(\underline{y}_n; \hat{\theta}_n) = 0.$$

Under regularity and identification conditions, this estimator is consistent, that is $\hat{\theta}_n$ converges towards θ_0 in probability, which we denote by:

$$\hat{\theta}_n \xrightarrow{P} \theta_0, \quad (4.1)$$

i.e. $\forall \varepsilon > 0, \lim_{n \rightarrow \infty} \mathbb{P}(|\hat{\theta}_n - \theta_0| > \varepsilon) = 0$ (this is Definition 7.14).

Definition 4.1 involves a positive definite matrix W_n . While one can take any positive definite matrix to have consistency (Eq. (4.1)), it can be shown that the GMM estimator achieves the minimum asymptotic variance when W_n is the inverse of matrix S , the latter being defined by:

$$S = \text{Asy.Var}(\sqrt{n}g(\underline{y}_n; \hat{\theta}_n)).$$

In this case, W_n is said to be the **optimal weighting matrix**.

The intuition behind this result is the same that underlies Generalized Least Squares (see Section 2.5.2), that is: it is beneficial to use a criterion in which the weights are inversely proportional to the variances of the moments.

If $h(x_i; \theta_0)$ is not correlated to $h(x_j; \theta_0)$, for $i \neq j$, then we have:

$$S = \text{Var}(h(x_i; \theta_0)),$$

which can be approximated by

$$\hat{\Gamma}_{0,n} = \frac{1}{n} \sum_{i=1}^n h(x_i; \hat{\theta}_n) h(x_i; \hat{\theta}_n)'.$$

In a time series context, we often have correlation between x_i and x_{i+k} , especially for small k 's. In this case, and if the time series $\{y_i\}$ is covariance stationary (see Def. 6.4), then we have:

$$S := \sum_{\nu=-\infty}^{\infty} \Gamma_{\nu},$$

where $\Gamma_{\nu} := \mathbb{E}[h(x_i; \theta_0)h(x_{i-\nu}; \theta_0)']$. Matrix S is called the **long-run variance** of process $\{y_i\}$ (see Def. 6.9).

For $\nu \geq 0$, let us define $\hat{\Gamma}_{\nu,n}$ by:

$$\hat{\Gamma}_{\nu,n} = \frac{1}{n} \sum_{i=\nu+1}^n h(x_i; \hat{\theta}_n) h(x_{i-\nu}; \hat{\theta}_n)',$$

then S can be approximated by the Newey and West (1987) formula (similar to Eq. (6.6)):

$$\hat{\Gamma}_{0,n} + \sum_{\nu=1}^q \left[1 - \frac{\nu}{q+1} \right] (\hat{\Gamma}_{\nu,n} + \hat{\Gamma}'_{\nu,n}), \quad (4.2)$$

4.1.2 Asymptotic distribution of the GMM estimator

We have:

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} \mathcal{N}(0, V), \quad (4.3)$$

where $V = (DS^{-1}D')^{-1}$, with

$$D := \mathbb{E} \left(\frac{\partial h(x_i; \theta)}{\partial \theta'} \right) \Big|_{\theta=\theta_0}.$$

V can be approximated by

$$\hat{V}_n = (\hat{D}_n \hat{S}_n^{-1} \hat{D}'_n)^{-1}, \quad (4.4)$$

where \hat{S}_n is given by Eq. (4.2) and

$$\hat{D}'_n := \frac{\partial g(y_n; \theta)}{\partial \theta'} \Big|_{\theta=\hat{\theta}_n}.$$

In practice, the previous matrix is computed numerically.

4.1.3 Testing hypotheses in the GMM framework

A first important test is the one concerning the validity of the moment restrictions (Sargan-Hansen test; Sargan (1958) and Hansen (1982)). Assume that the number of restrictions imposed is larger than the number of parameters to estimate ($r > K$). In this case, the restrictions are said to be over-identifying.

Under correct specification, we asymptotically have:

$$\sqrt{n}g(\underline{y}_n; \theta_0) \sim \mathcal{N}(0, S).$$

As a result, it comes that:

$$J_n = (\sqrt{n}g(\underline{y}_n; \theta_0))' S^{-1} (\sqrt{n}g(\underline{y}_n; \theta_0)) \quad (4.5)$$

asymptotically follows a χ^2 distribution. The number of degrees of freedom is equal to $r - K$. (Note that, for $r = K$, we have, as expected, $J = 0$.) That is, asymptotically:

$$J_n \sim \chi^2(r - K).$$

The GMM framework also allows to easily test linear restrictions on the parameters. First, given Eq. (4.3), Wald tests (see Eq. (2.16) in Section 2.2.5) are readily available. Second, one can also resort to a test equivalent to the *likelihood ratio tests* (see Definition 4.8). More precisely, consider an unconstrained model and a constrained version of this model, the number of restrictions being equal to k . If the two models are estimated by considering the same moment constraints, and the same weighting matrix (using Eq. (4.4), based on the unrestricted model), then we have that:

$$n \left[(g(\underline{y}_n; \hat{\theta}_n^*) - g(\underline{y}_n; \hat{\theta}_n)) \right] \sim \chi^2(k),$$

where $\hat{\theta}_n^*$ is the constrained estimate of θ_0 .

4.1.4 Example: Estimation of the Stochastic Discount Factor (s.d.f.)

Under the no-arbitrage assumption, there exists a random variable $\mathcal{M}_{t,t+1}$ such that

$$\mathbb{E}_t(\mathcal{M}_{t,t+1} R_{t+1}) = 1$$

for any (gross) asset return R_t . In the following, R_t denotes a n_r -dimensional vector of gross returns.

We consider the following specification of the s.d.f.:

$$\mathcal{M}_{t,t+1} = 1 - \mathbf{b}'_M(F_{t+1} - \mathbb{E}_t(F_{t+1})), \quad (4.6)$$

where F_t is a vector of factors. Eq. (4.6) then reads:

$$\mathbb{E}_t([1 - \mathbf{b}'_M(F_{t+1} - \mathbb{E}_t(F_{t+1}))] R_{t+1}) = 1.$$

Assume that the date- t information set is $\mathcal{I}_t = \{\mathbf{z}_t, \mathcal{I}_{t-1}\}$, where \mathbf{z}_t is a vector of variables observed on date t . (We then have $\mathbb{E}_t(\bullet) \equiv \mathbb{E}(\bullet | \mathcal{I}_t)$.)

We can use \mathbf{z}_t as an instrument. Indeed, we have:

$$\begin{aligned} & \mathbb{E}(z_{i,t} [\mathbf{b}'_M \{F_{t+1} - \mathbb{E}_t(F_{t+1})\} R_{t+1} - R_{t+1} + 1]) \\ &= \mathbb{E}(\mathbb{E}_t \{z_{i,t} [\mathbf{b}'_M \{F_{t+1} - \mathbb{E}_t(F_{t+1})\} R_{t+1} - R_{t+1} + 1]\}) \\ &= \mathbb{E}(z_{i,t} \underbrace{\mathbb{E}_t \{\mathbf{b}'_M \{F_{t+1} - \mathbb{E}_t(F_{t+1})\} R_{t+1} - R_{t+1} + 1\}}_{=0}) = 0. \end{aligned} \quad (4.7)$$

We have then converted a conditional moment condition into an unconditional one (which we need to implement the theory above). However, at that stage, we cannot still not directly use the GMM formulas because of the conditional expectation $\mathbb{E}_t(F_{t+1})$ that appears in $\mathbb{E}(z_{i,t}[\mathbf{b}'_M\{F_{t+1} - \mathbb{E}_t(F_{t+1})\}R_{t+1} - R_{t+1} + 1]) = 0$.

To go further, let us assume that:

$$\mathbb{E}_t(F_{t+1}) = \mathbf{b}_F \mathbf{z}_t.$$

We can then easily estimate matrix \mathbf{b}_F (of dimension $n_F \times n_z$) by OLS. Note here that these OLS can be seen as a special GMM case. Indeed, as was done in Eq. (4.7), we can show that, for the j^{th} component of F_t , we have:

$$\mathbb{E}([F_{j,t+1} - \mathbf{b}_{F,j} \mathbf{z}_t] \mathbf{z}_t) = 0,$$

where $\mathbf{b}_{F,j}$ denotes the j^{th} row of \mathbf{b}_F . This yields the OLS formula.

At that stage, we count on the following moment restrictions to estimate \mathbf{b}_M :

$$\mathbb{E}(z_{i,t}[\mathbf{b}'_M\{F_{t+1} - \mathbf{b}_F \mathbf{z}_t\}R_{t+1} - R_{t+1} + 1]) = 0.$$

Specifically, the number of restrictions is $n_R \times n_z$. Let us implement this approach in the U.S. context, using data extracted from the FRED database. In factor F_t , we use the changes in the VIX and in the personal consumption expenditures. The returns (R_t) are based on the Wilshire 5000 Price Index (a stock price index) and on the ICE BofA BBB US Corporate Index Total Return Index (a bond return index).

```
library(fredr)
fredr_set_key("df65e14c054697a52b4511e77fcfa1f3")
start_date <- as.Date("1990-01-01"); end_date <- as.Date("2022-01-01")
f <- function(ticker){
  fredr(series_id = ticker,
        observation_start = start_date, observation_end = end_date,
        frequency = "m", aggregation_method = "avg")
}
vix <- f("VIXCLS") # VIX
pce <- f("PCE") # Personal consumption expenditures
sto <- f("WILL5000PRFC") # Wilshire 5000 Full Cap Price Index
bdr <- f("BAMLCCOA4BBTRIV") # ICE BofA BBB US Corporate Index Total Return Index
T <- dim(vix)[1]
dvix <- c(vix$value[3:T]/vix$value[2:(T-1)]) # change in VIX t+1
dpce <- c(pce$value[3:T]/pce$value[2:(T-1)]) # change in PCE t+1
dsto <- c(sto$value[3:T]/sto$value[2:(T-1)]) # return t+1
dbdr <- c(bdr$value[3:T]/bdr$value[2:(T-1)]) # return t+1
dvix_1 <- c(vix$value[2:(T-1)]/vix$value[1:(T-2)]) # change in VIX t
dpce_1 <- c(pce$value[2:(T-1)]/pce$value[1:(T-2)]) # change in PCE t
```

```

dsto_1 <- c(sto$value[2:(T-1)]/sto$value[1:(T-2)]) # return t
dbdr_1 <- c(bdr$value[2:(T-1)]/bdr$value[1:(T-2)]) # return t

```

Define the matrices containing the F_{t+1} , \mathbf{z}_t , and R_{t+1} vectors:

```

F_tp1 <- cbind(dvix,dpce)
Z      <- cbind(1,dvix_1,dpce_1,dsto_1,dbdr_1)
b_F    <- t(solve(t(Z) %*% Z) %*% t(Z) %*% F_tp1)
F_innov <- F_tp1 - Z %*% t(b_F)
R_tp1  <- cbind(dsto,dbdr)
n_F    <- dim(F_tp1)[2]; n_R <- dim(R_tp1)[2]; n_z <- dim(Z)[2]

```

Function `f_aux` compute the $h(x_t; \theta)$ and the $g(y_T; \theta)$; function `f2beMin` is the function to be minimized.

```

f_aux <- function(theta){
  b_M <- matrix(theta[1:n_F],ncol=1)
  R_aux <- matrix(F_innov %*% b_M,T-2,n_R) * R_tp1 - R_tp1 + 1
  H <- (R_aux %x% matrix(1,1,n_z)) * (matrix(1,1,n_R) %x% Z)
  g <- matrix(apply(H,2,mean),ncol=1)
  return(list(g=g,H=H))
}
f2beMin <- function(theta,W){# function to be minimized
  res <- f_aux(theta)
  return(t(res$g) %*% W %*% res$g)
}

```

Now, let's minimize this function. We run 5 iterations (where W is updated).

```

library(AEC)
theta <- c(rep(0,n_F)) # initial value
for(i in 1:10){# recursion on W
  res <- f_aux(theta)
  W <- solve(NW.LongRunVariance(res$H,q=6))
  res.optim <- optim(theta,f2beMin,W=W,
                      method="BFGS", # could be "Nelder-Mead"
                      control=list(trace=FALSE,maxit=200),hessian=TRUE)
  theta <- res.optim$par
}

```

Finally, let's compute the standard deviation of the parameter estimates, using Eq. (4.4):

```

eps <- .0001
g0 <- f_aux(theta)$g
D <- NULL
for(i in 1:length(theta)){
  theta.i <- theta
  theta.i[i] <- theta.i[i] + eps
  gi <- f_aux(theta.i)$g
  D <- cbind(D,(gi-g0)/eps)
}
V <- 1/T * solve(t(D) %*% W %*% D)
std.dev <- sqrt(diag(V));t.stud <- theta/std.dev
cbind(theta, std.dev, t.stud)

##           theta     std.dev      t.stud
## [1,] -0.7355469  0.4653449 -1.5806490
## [2,] -10.9859162 16.8406714 -0.6523443

```

The Hansen statistic can be used to test the model (see Eq. (4.5)). If the model is correct, we have:

$$T g(\underline{y}_T; \theta)' S^{-1} g(\underline{y}_T; \theta) \sim i.i.d. \chi^2(J - K),$$

where J is the number of moment constraints ($n_z \times n_r$ here) and K is the number of estimated parameters ($= n_F$ here).

```

g <- f_aux(theta)$g
Hansen_stat <- T * t(g) %*% W %*% g
pvalue <- pchisq(q = Hansen_stat, df = n_R*n_z - n_F)
pvalue

##           [,1]
## [1,] 0.8796173

```

4.2 Maximum Likelihood Estimation

4.2.1 Intuition

Intuitively, the Maximum Likelihood Estimation consists in looking for the value of θ that is such that the probability of having observed \mathbf{y} (the sample at hand) is the highest possible.

To set an example, assume that the time periods between the arrivals of two customers in a shop, denoted by y_i , are i.i.d. and follow an exponential distribution, i.e. $y_i \sim i.i.d. \mathcal{E}(\lambda)$. You have observed these arrivals for some time,

thereby constituting a sample $\mathbf{y} = \{y_1, \dots, y_n\}$. You want to estimate λ (i.e. in that case, the vector of parameters is simply $\theta = \lambda$).

The density of Y (one observation) is $f(y; \lambda) = \frac{1}{\lambda} \exp(-y/\lambda)$. Fig. 4.1 represents that density functions for different values of λ .

Your 200 observations are reported at the bottom of Fig. 4.1 (red bars). You build the histogram and report it on the same chart.




Figure 4.1: The red ticks, at the bottom, indicate observations (there are 200 of them). The histogram is based on these 200 observations

What is your estimate of λ ? Intuitively, one is led to take the λ for which the (theoretical) distribution is the closest to the histogram (that can be seen as an “empirical distribution”). This approach is consistent with the idea of picking the λ for which the probability of observing the values included in \mathbf{y} is the highest.

Let us be more formal. Assume that you have only four observations: $y_1 = 1.1$, $y_2 = 2.2$, $y_3 = 0.7$ and $y_4 = 5.0$. What was the probability of observing, for a small ε ,

- $1.1 - \varepsilon \leq Y_1 < 1.1 + \varepsilon$,
- $2.2 - \varepsilon \leq Y_2 < 2.2 + \varepsilon$,
- $0.7 - \varepsilon \leq Y_3 < 0.7 + \varepsilon$ and
- $5.0 - \varepsilon \leq Y_4 < 5.0 + \varepsilon$?

Because the y_i s are i.i.d., this probability is $\prod_{i=1}^4 (2\varepsilon f(y_i, \lambda))$. The next plot shows the probability (divided by $16\varepsilon^4$, which does not depend on λ) as a function of λ .




Figure 4.2: Proba. that $y_i - \varepsilon \leq Y_i < y_i + \varepsilon$, $i \in \{1, 2, 3, 4\}$. The vertical red line indicates the maximum of the function.

The value of λ that maximizes the probability is 2.26.

Let us come back to the example with 200 observations:

In that case, the value of λ that maximizes the probability is 3.42.

4.2.2 Notations

$f(y; \theta)$ denotes the probability density function (p.d.f.) of a random variable Y which depends on a set of parameters θ .

Density of n independent and identically distributed (i.i.d.) observations of Y :

$$f(y_1, \dots, y_n; \theta) = \prod_{i=1}^n f(y_i; \theta).$$

\mathbf{y} denotes the vector of observations; $\mathbf{y} = \{y_1, \dots, y_n\}$.

Definition 4.2 (Likelihood function). $\mathcal{L} : \theta \rightarrow \mathcal{L}(\theta; \mathbf{y}) = f(y_1, \dots, y_n; \theta)$ is the **likelihood function**.

We often work with $\log \mathcal{L}$, the **log-likelihood function**.

Example 4.1 (Gaussian distribution). If $y_i \sim \mathcal{N}(\mu, \sigma^2)$, then

$$\log \mathcal{L}(\theta; \mathbf{y}) = -\frac{1}{2} \sum_{i=1}^n \left(\log \sigma^2 + \log 2\pi + \frac{(y_i - \mu)^2}{\sigma^2} \right).$$




Figure 4.3: Log-likelihood function associated with the 200 i.i.d. observations. The vertical red line indicates the maximum of the function.

Definition 4.3 (Score). The score $S(y; \theta)$ is given by $\frac{\partial \log f(y; \theta)}{\partial \theta}$.

If $y_i \sim \mathcal{N}(\mu, \sigma^2)$ (Example 4.1), then

$$\frac{\partial \log f(y; \theta)}{\partial \theta} = \begin{bmatrix} \frac{\partial \log f(y; \theta)}{\partial \mu} \\ \frac{\partial \log f(y; \theta)}{\partial \sigma^2} \end{bmatrix} = \begin{bmatrix} \frac{y - \mu}{\sigma^2} \\ \frac{1}{2\sigma^2} \left(\frac{(y - \mu)^2}{\sigma^2} - 1 \right) \end{bmatrix}.$$

Proposition 4.1 (Score expectation). *The expectation of the score is zero.*

Proof. We have:

$$\begin{aligned} \mathbb{E} \left(\frac{\partial \log f(Y; \theta)}{\partial \theta} \right) &= \int \frac{\partial \log f(y; \theta)}{\partial \theta} f(y; \theta) dy \\ &= \int \frac{\partial f(y; \theta)/\partial \theta}{f(y; \theta)} f(y; \theta) dy = \frac{\partial}{\partial \theta} \int f(y; \theta) dy = \partial 1 / \partial \theta = 0, \end{aligned}$$

which gives the result. \square

Definition 4.4 (Fisher information matrix). The **information matrix** is (minus) the expectation of the second derivatives of the log-likelihood function:

$$\mathcal{I}_Y(\theta) = -\mathbb{E} \left(\frac{\partial^2 \log f(Y; \theta)}{\partial \theta \partial \theta'} \right).$$

Proposition 4.2. We have $\mathcal{I}_Y(\theta) = \mathbb{E} \left[\left(\frac{\partial \log f(Y; \theta)}{\partial \theta} \right) \left(\frac{\partial \log f(Y; \theta)}{\partial \theta} \right)' \right] = \text{Var}[S(Y; \theta)]$.

Proof. We have $\frac{\partial^2 \log f(Y; \theta)}{\partial \theta \partial \theta'} = \frac{\partial^2 f(Y; \theta)}{\partial \theta \partial \theta'} \frac{1}{f(Y; \theta)} - \frac{\partial \log f(Y; \theta)}{\partial \theta} \frac{\partial \log f(Y; \theta)}{\partial \theta'}$. The expectation of the first right-hand side term is $\partial^2 1 / (\partial \theta \partial \theta') = \mathbf{0}$, which gives the result. \square

Example 4.2. If $y_i \sim i.i.d. \mathcal{N}(\mu, \sigma^2)$, let $\theta = [\mu, \sigma^2]'$ then

$$\frac{\partial \log f(y; \theta)}{\partial \theta} = \left[\frac{y - \mu}{\sigma^2} \quad \frac{1}{2\sigma^2} \left(\frac{(y - \mu)^2}{\sigma^2} - 1 \right) \right]'$$

and

$$\mathcal{J}_Y(\theta) = \mathbb{E} \left(\frac{1}{\sigma^4} \begin{bmatrix} \sigma^2 & y - \mu \\ y - \mu & \frac{(y - \mu)^2}{\sigma^2} - \frac{1}{2} \end{bmatrix} \right) = \begin{bmatrix} 1/\sigma^2 & 0 \\ 0 & 1/(2\sigma^4) \end{bmatrix}.$$

Proposition 4.3 (Additive property of the Info. mat.). *The information matrix resulting from two independent experiments is the sum of the information matrices:*

$$\mathcal{J}_{X,Y}(\theta) = \mathcal{J}_X(\theta) + \mathcal{J}_Y(\theta).$$

Proof. Immediately obtained from the definition (see Def. 4.4). \square

Theorem 4.1 (Frechet-Darmois-Cramer-Rao bound). *Consider an unbiased estimator of θ denoted by $\hat{\theta}(Y)$.*

The variance of the random variable $\omega' \hat{\theta}$ (which is a linear combination of the components of $\hat{\theta}$) is larger than:

$$(\omega' \omega)^2 / (\omega' \mathcal{J}_Y(\theta) \omega).$$

Proof. The Cauchy-Schwarz inequality implies that $\sqrt{\text{Var}(\omega' \hat{\theta}(Y)) \text{Var}(\omega' S(Y; \theta))} \geq |\omega' \text{Cov}[\hat{\theta}(Y), S(Y; \theta)] \omega|$. Now, $\text{Cov}[\hat{\theta}(Y), S(Y; \theta)] = \int_y \hat{\theta}(y) \frac{\partial \log f(y; \theta)}{\partial \theta} f(y; \theta) dy = \frac{\partial}{\partial \theta} \int_y \hat{\theta}(y) f(y; \theta) dy = \mathbf{I}$ because $\hat{\theta}$ is unbiased.

Therefore $\text{Var}(\omega' \hat{\theta}(Y)) \geq \text{Var}(\omega' S(Y; \theta))^{-1} (\omega' \omega)^2$. Prop. 4.2 leads to the result. \square

Definition 4.5 (Identifiability). The vector of parameters θ is identifiable if, for any other vector θ^* :

$$\theta^* \neq \theta \Rightarrow \mathcal{L}(\theta^*; \mathbf{y}) \neq \mathcal{L}(\theta; \mathbf{y}).$$

Definition 4.6 (Maximum Likelihood Estimator (MLE)). The maximum likelihood estimator (MLE) is the vector θ that maximizes the likelihood function. Formally:

$$\theta_{MLE} = \arg \max_{\theta} \mathcal{L}(\theta; \mathbf{y}) = \arg \max_{\theta} \log \mathcal{L}(\theta; \mathbf{y}). \quad (4.8)$$

Definition 4.7 (Likelihood equation). Necessary condition for maximizing the likelihood function:

$$\frac{\partial \log \mathcal{L}(\theta; \mathbf{y})}{\partial \theta} = \mathbf{0}. \quad (4.9)$$

Hypothesis 4.1 (Regularity assumptions). We have:

- i. $\theta \in \Theta$ where Θ is compact.
- ii. θ_0 is identified.
- iii. The log-likelihood function is continuous in θ .
- iv. $\mathbb{E}_{\theta_0}(\log f(Y; \theta))$ exists.
- v. The log-likelihood function is such that $(1/n) \log \mathcal{L}(\theta; \mathbf{y})$ converges almost surely to $\mathbb{E}_{\theta_0}(\log f(Y; \theta))$, uniformly in $\theta \in \Theta$.
- vi. The log-likelihood function is twice continuously differentiable in an open neighborhood of θ_0 .
- vii. The matrix $\mathbf{I}(\theta_0) = -\mathbb{E}_0 \left(\frac{\partial^2 \log \mathcal{L}(\theta; \mathbf{y})}{\partial \theta \partial \theta'} \right)$ —the Fisher Information matrix— exists and is nonsingular.

Proposition 4.4 (Properties of MLE). *Under regularity conditions (Assumptions 4.1), the MLE is:*

- a. **Consistent:** $\text{plim } \theta_{MLE} = \theta_0$ (θ_0 is the true vector of parameters).
 - b. **Asymptotically normal:** $\theta_{MLE} \sim \mathcal{N}(\theta_0, \mathbf{I}(\theta_0)^{-1})$, where
- $$\mathbf{I}(\theta_0) = -\mathbb{E}_0 \left(\frac{\partial^2 \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta \partial \theta'} \right) = n \mathcal{J}_Y(\theta_0). \quad (\text{Fisher Info. matrix})$$
- c. **Asymptotically efficient:** θ_{MLE} is asymptotically efficient and achieves the Fr'echet-Darmois-Cram'er-Rao lower bound for consistent estimators.
 - d. **Invariant:** The MLE of $g(\theta_0)$ is $g(\theta_{MLE})$ if g is a continuous and continuously differentiable function.

Proof. See Appendix 7.5. □

Note that (b) also writes:

$$\sqrt{n}(\theta_{MLE} - \theta_0) \xrightarrow{d} \mathcal{N}(0, \mathcal{J}_Y(\theta_0)^{-1}). \quad (4.10)$$

The asymptotic covariance matrix of the MLE is:

$$[\mathbf{I}(\theta_0)]^{-1} = \left[-\mathbb{E}_0 \left(\frac{\partial^2 \log \mathcal{L}(\theta; \mathbf{y})}{\partial \theta \partial \theta'} \right) \right]^{-1}.$$

A direct (analytical) evaluation of this expectation is often out of reach.

It can however be estimated by, either:

$$\hat{\mathbf{I}}_1^{-1} = \left(-\frac{\partial^2 \log \mathcal{L}(\theta_{MLE}; \mathbf{y})}{\partial \theta \partial \theta'} \right)^{-1}, \quad (4.11)$$

$$\hat{\mathbf{I}}_2^{-1} = \left(\sum_{i=1}^n \frac{\partial \log \mathcal{L}(\theta_{MLE}; y_i)}{\partial \theta} \frac{\partial \log \mathcal{L}(\theta_{MLE}; y_i)}{\partial \theta'} \right)^{-1}. \quad (4.12)$$

Asymptotically, we have $(\hat{\mathbf{I}}_1^{-1})\hat{\mathbf{I}}_2 = Id$, that is, the two formulas provide the same result.

In case of (suspected) misspecification, one can use the so-called *sandwich estimator* of the covariance matrix.¹ This covariance matrix is given by:

$$\hat{\mathbf{I}}_3^{-1} = \hat{\mathbf{I}}_2^{-1} \hat{\mathbf{I}}_1 \hat{\mathbf{I}}_2^{-1}. \quad (4.13)$$

4.2.3 To sum up – MLE in practice

- A parametric model (depending on the vector of parameters θ whose “true” value is θ_0) is specified.
- i.i.d. sources of randomness are identified.
- The density associated to one observation y_i is computed analytically (as a function of θ): $f(y; \theta)$.
- The log-likelihood is $\log \mathcal{L}(\theta; \mathbf{y}) = \sum_i \log f(y_i; \theta)$.
- The MLE estimator results from the optimization problem (this is Eq. (4.8)):

$$\theta_{MLE} = \arg \max_{\theta} \log \mathcal{L}(\theta; \mathbf{y}). \quad (4.14)$$

- We have: $\theta_{MLE} \sim \mathcal{N}(\theta_0, \mathbf{I}(\theta_0)^{-1})$, where $\mathbf{I}(\theta_0)^{-1}$ is estimated by means of Eq. (4.11), Eq. (4.12), or Eq. (4.13). Most of the time, this computation is numerical.

4.2.4 Example: MLE estimation of a mixture of Gaussian distribution

Consider the returns of the SMI index. Let's assume that these returns are independently drawn from a mixture of Gaussian distributions. The p.d.f. $f(x; \theta)$, with $\theta = [\mu_1, \sigma_1, \mu_2, \sigma_2, p]'$, is given by:

$$p \frac{1}{\sqrt{2\pi\sigma_1^2}} \exp\left(-\frac{(x-\mu_1)^2}{2\sigma_1^2}\right) + (1-p) \frac{1}{\sqrt{2\pi\sigma_2^2}} \exp\left(-\frac{(x-\mu_2)^2}{2\sigma_2^2}\right).$$

¹See, e.g., Charles Geyer's lectures notes.

(See p.d.f. of mixtures of Gaussian dist..)

```
library(AEC)
T <- dim(smi)[1]
h <- 5 # holding period (one week)
smi$r <- c(rep(NaN,h),
           100*c(log(smi$Close[(1+h):T]/smi$Close[1:(T-h)])))
indic.dates <- seq(1,T,by=5) # weekly returns
smi <- smi[indic.dates,]
smi <- smi[complete.cases(smi),]
par(mfrow=c(1,1))
plot(smi$Date,smi$r,type="l",xlab="",ylab="in percent")
abline(h=0,col="blue")
abline(h=mean(smi$r,na.rm = TRUE)+2*sd(smi$r,na.rm = TRUE),lty=3,col="blue")
abline(h=mean(smi$r,na.rm = TRUE)-2*sd(smi$r,na.rm = TRUE),lty=3,col="blue")
```




Figure 4.4: Time series of SMI weekly returns (source: Yahoo Finance).

```
f <- function(theta,y){ # Likelihood function
  mu.1 <- theta[1]; mu.2 <- theta[2]
  sigma.1 <- theta[3]; sigma.2 <- theta[4]
  p <- exp(theta[5])/(1+exp(theta[5]))
  res <- p*1/sqrt(2*pi*sigma.1^2)*exp(-(y-mu.1)^2/(2*sigma.1^2)) +
    (1-p)*1/sqrt(2*pi*sigma.2^2)*exp(-(y-mu.2)^2/(2*sigma.2^2))
  return(res)
}
log.f <- function(theta,y){ #log-Likelihood function
  return(-sum(log(f(theta,y))))
}
```

```

}

res.optim <- optim(c(0,0,0.5,1.5,.5),
                    log.f,
                    y=smi$r,
                    method="BFGS", # could be "Nelder-Mead"
                    control=list(trace=FALSE,maxit=100),hessian=TRUE)
theta <- res.optim$par
theta

## [1] 0.3012379 -1.3167476 1.7715072 4.8197596 1.9454889

```

Now, let us compute estimates of the covariance matrix of the MLE:

```

# Hessian approach:
I.1 <- solve(res.optim$hessian)
# Outer-product of gradient approach:
log.f.0 <- log(f(theta,smi$r))
epsilon <- .00000001
d.log.f <- NULL
for(i in 1:length(theta)){
  theta.i <- theta
  theta.i[i] <- theta.i[i] + epsilon
  log.f.i <- log(f(theta.i,smi$r))
  d.log.f <- cbind(d.log.f,
                     (log.f.i - log.f.0)/epsilon)
}
I.2 <- solve(t(d.log.f) %*% d.log.f)
# Misspecification-robust approach (sandwich formula):
I.3 <- I.1 %*% solve(I.2) %*% I.1
cbind(diag(I.1),diag(I.2),diag(I.3))

##          [,1]      [,2]      [,3]
## [1,] 0.003683422 0.003199481 0.00586160
## [2,] 0.226892824 0.194283391 0.38653389
## [3,] 0.005764271 0.002769579 0.01712255
## [4,] 0.194081311 0.047466419 0.83130838
## [5,] 0.092114437 0.040366005 0.31347858

```

According to the first (respectively third) type of estimate for the covariance matrix, a 95% confidence interval for μ_1 is [0.182, 0.42] (resp. [0.151, 0.451]).

Note that we have not directly estimated parameter p but $\nu = \log(p/(1-p))$ (in such a way that $p = \exp(\nu)/(1+\exp(\nu))$). In order to get an estimate of the standard deviation of our esitmate of p , we can implement the **Delta method**.

This method is based on the fact that, for a function g that is continuous in the neighborhood of θ_0 and for large n , we have:

$$\text{Var}(g(\hat{\theta}_n)) \approx \frac{\partial g(\hat{\theta}_n)}{\partial \theta'} \text{Var}(\hat{\theta}_n) \frac{\partial g(\hat{\theta}_n)'}{\partial \theta}. \quad (4.15)$$

```

g <- function(theta){
  mu.1 <- theta[1]; mu.2 <- theta[2]
  sigma.1 <- theta[3]; sigma.2 <- theta[4]
  p <- exp(theta[5])/(1+exp(theta[5]))
  return(c(mu.1,mu.2,sigma.1,sigma.2,p))
}
# Computation of g's gradient around estimated theta:
eps <- .00001
g.theta <- g(theta)
g.gradient <- NULL
for(i in 1:5){
  theta.perturb <- theta
  theta.perturb[i] <- theta[i] + eps
  g.gradient <- cbind(g.gradient,(g(theta.perturb)-g.theta)/eps)
}
Var <- g.gradient %*% t(g.gradient)
stdv.g.theta <- sqrt(diag(Var))
stdv.theta <- sqrt(diag(I.3))
cbind(theta,stdv.theta,g.theta,stdv.g.theta)

##           theta stdv.theta   g.theta stdv.g.theta
## [1,]  0.3012379 0.07656108  0.3012379   0.07656108
## [2,] -1.3167476 0.62171850 -1.3167476   0.62171850
## [3,]  1.7715072 0.13085316  1.7715072   0.13085316
## [4,]  4.8197596 0.91176114  4.8197596   0.91176114
## [5,]  1.9454889 0.55989158  0.8749539   0.06125726

```

To finish with, let us draw the estimated parametric p.d.f. (the mixture of Gaussian distribution), and compare it to a non-parametric (kernel-based) estimate of this p.d.f. (using function `density`):

```

x <- seq(-5,5,by=.01)
par(plt=c(.1,.95,.1,.95))
plot(x,f(theta,x),type="l",lwd=2,xlab="returns, in percent",ylab="",
      ylim=c(0,1.4*max(f(theta,x))))
lines(density(smi$r),type="l",lwd=2,lty=3)
lines(x,dnorm(x,mean=mean(smi$r),sd = sd(smi$r)),col="red",lty=2,lwd=2)
rug(smi$r,col="blue")
legend("topleft",

```

```

c("Kernel estimate (non-parametric)", "Estimated mixture of Gaussian distr. (MLE, parametric",
  lty=c(3,1,2), # gives the legend appropriate symbols (lines)
  lwd=c(2), # line width
  col=c("black","black","red"), # gives the legend lines the correct color and width
  pt.bg=c(1),
  pt.cex = c(1),
  bg="white",
  seg.len = 4
)

```




Figure 4.5: Comparison of different estimates of the distribution of returns.

4.2.5 Test procedures

The objective is to test the following parameter restrictions:

$$H_0 : \underbrace{h(\theta)}_{r \times 1} = 0. \quad (4.16)$$

In the context of MLE, three tests are largely used:

- Likelihood Ratio (LR) test,
- Wald (W) test,
- Lagrange Multiplier (LM) test.

Here is the rationale behind these three tests:²

²An interesting graphical presentation of the tests is proposed in Buse (1982).

- LR: If $h(\theta) = 0$, then imposing this restriction during the estimation (restricted estimator) should not result in a large decrease in the likelihood function (w.r.t the unrestricted estimation).
- Wald: If $h(\theta) = 0$, then $h(\hat{\theta})$ should not be far from 0 (even if these restrictions are not imposed during the MLE).
- LM: If $h(\theta) = 0$, then the gradient of the likelihood function should be small when evaluated at the restricted estimator.

In terms of implementation, while the LR necessitates to estimate both restricted and unrestricted models, the Wald test requires the estimation of the unrestricted model only, and the LM tests requires the estimation of the restricted model only.

As shown below, the three test statistics associated with these three tests coincide asymptotically. (Therefore, they naturally have the same asymptotic distribution, that are χ^2).

Proposition 4.5 (Asymptotic distribution of the Wald statistic). *Under regularity conditions (Assumptions 4.1) and under $H_0 : h(\theta) = 0$, the Wald statistic, defined by:*

$$\xi^W = h(\hat{\theta})' \text{Var}[h(\hat{\theta})]^{-1} h(\hat{\theta}),$$

where

$$\text{Var}[h(\hat{\theta})] = \left(\frac{\partial h(\hat{\theta})}{\partial \theta'} \right) \text{Var}[\hat{\theta}] \left(\frac{\partial h(\hat{\theta})}{\partial \theta} \right)', \quad (4.17)$$

is asymptotically $\chi^2(r)$, where the number of degrees of freedom r corresponds to the dimension of $h(\theta)$. (Note that Eq. (4.17) is the same as the one used in the Delta method, see Eq. (4.15).)

The Wald test, defined by the critical region

$$\{\xi^W \geq \chi_{1-\alpha}^2(r)\},$$

where $\chi_{1-\alpha}^2(r)$ denotes the quantile of level $1 - \alpha$ of the $\chi^2(r)$ distribution, has asymptotic level α and is consistent.³

Proof. See Appendix 7.5. □

In practice, in Eq. (4.17), $\text{Var}[\hat{\theta}]$ is replaced by an estimate given, e.g., by Eq. (4.11), Eq. (4.12), or Eq. (4.13).

Proposition 4.6 (Asymptotic distribution of the LM test statistic). *Under regularity conditions (Assumptions 4.1) and under $H_0 : h(\theta) = 0$, the LM statistic*

$$\xi^{LM} = \left(\frac{\partial \log \mathcal{L}(\theta)}{\partial \theta'} \Big|_{\theta=\hat{\theta}^0} \right) [\mathbf{I}(\hat{\theta}^0)]^{-1} \left(\frac{\partial \log \mathcal{L}(\theta)}{\partial \theta} \Big|_{\theta=\hat{\theta}^0} \right), \quad (4.18)$$

³See Defs. 7.7 and 7.8 for definitions of the asymptotic levels and consistency of tests.

(where $\hat{\theta}^0$ is the restricted MLE estimator) is $\chi^2(r)$.

The test defined by the critical region:

$$\{\xi^{LM} \geq \chi^2_{1-\alpha}(r)\}$$

has asymptotic level α and is consistent (see Defs. 7.7 and 7.8).

This test is called Score or Lagrange Multiplier (LM) test.

Proof. See Appendix 7.5. □

Definition 4.8 (Likelihood Ratio test statistics). The likelihood ratio associated to a restriction of the form $H_0 : h(\theta) = 0$ is given by:

$$LR = \frac{\mathcal{L}_R(\theta; \mathbf{y})}{\mathcal{L}_U(\theta; \mathbf{y})} \quad (\in [0, 1]),$$

where \mathcal{L}_R (respectively \mathcal{L}_U) is the likelihood function that imposes (resp. that does not impose) the restriction. The likelihood ratio test statistic is given by $-2 \log(LR)$, that is:

$$\boxed{\xi^{LR} = 2(\log \mathcal{L}_U(\theta; \mathbf{y}) - \log \mathcal{L}_R(\theta; \mathbf{y})).}$$

Proposition 4.7 (Asymptotic equivalence of LR, LM, and Wald tests). Under the null hypothesis H_0 , we have, asymptotically:

$$\xi^{LM} = \xi^{LR} = \xi^W.$$

Proof. See Appendix 7.5. □

4.3 Bayesian approach

4.3.1 Introduction

An excellent introduction to Bayesian methods is proposed by Martin Haugh, 2017.

As suggested by the name of this approach, the starting point is the Bayes formula:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \& B)}{\mathbb{P}(B)},$$

where A and B are two “events”. For instance, A may be: parameter α (conceived as something stochastic) lies in interval $[a, b]$. Assume that you are interested in the probability of occurrence of A . Without any specific information (or “unconditionally”), this probability is $\mathbb{P}(A)$. Your evaluation of this probability

can only be better if you are provided with any additional form of information. Typically, if the event B tends to occur simultaneously with A , then knowledge of B can be useful. The Bayes formula says how this additional information (on B) can be used to “update” the probability of event A .

In our case, this intuition will work as follows: assume that you know the form of the data-generating process (DGP). That is, you know the structure of the model used to draw some stochastic data; you also know the type of distributions used to generate these data. However, you do not know the numerical values of all the parameters characterizing the DGP. Let us denote by θ the vector of unknown parameters. While these parameters are not known exactly, assume that we have –even without having observed any data– some **priors** on their distribution. Then, as was the case in the example above (with A and B), the observation of data generated by the model can only reduce the uncertainty associated with θ . Loosely speaking, combining the priors and the observations of data generated by the model should result in “thinner” distributions for the components of θ . The latter distributions are called the **posterior distributions**.⁴

Let us formalize this intuition. Define the prior by $f_\theta(\theta)$ and the model realizations (the “data”) by vector \mathbf{y} . The joint distribution of (\mathbf{y}, θ) is given by:

$$f_{Y,\theta}(\mathbf{y}, \theta) = f_{Y|\theta}(\mathbf{y}, \theta)f_\theta(\theta),$$

and, symmetrically, by

$$f_{Y,\theta}(\mathbf{y}, \theta) = f_{\theta|Y}(\theta, \mathbf{y})f_Y(\mathbf{y}),$$

where $f_{\theta|Y}(\cdot, \mathbf{y})$, the distribution of the parameters conditional on the observations, is the **posterior** distribution.

The last two equations imply that:

$$f_{\theta|Y}(\theta, \mathbf{y}) = \frac{f_{Y|\theta}(\mathbf{y}, \theta)f_\theta(\theta)}{f_Y(\mathbf{y})}. \quad (4.19)$$

Note that f_Y is the marginal (or unconditional) distribution of \mathbf{y} , that can be written:

$$f_Y(\mathbf{y}) = \int f_{Y|\theta}(\mathbf{y}, \theta)f_\theta(\theta)d\theta. \quad (4.20)$$

Eq. (4.19) is sometimes rewritten as follows:

$$f_{\theta|Y}(\theta, \mathbf{y}) \propto f_{\theta,Y}(\theta, \mathbf{y}) := f_{Y|\theta}(\mathbf{y}, \theta)f_\theta(\theta), \quad (4.21)$$

where \propto means, loosely speaking, “*proportional to*”. In rare instances, starting from given priors, one can analytically compute the posterior distribution

⁴The output of the Bayesian approach will be the (posterior) distribution of the vector of parameters (θ). When we speak about the *distributions of the components of θ* , we mean the marginal distributions of each component of the vector.

$f_\theta(\theta, \mathbf{y})$. However, in most cases, this is out of reach. One then has to resort to numerical approaches to compute the posterior distribution. Monte Carlo Markov Chains (MCMC) is one of them.

According to the Bernstein-von Mises Theorem, Bayesian and MLE estimators have the same large sample properties. (In particular, the Bayesian approach also achieves the FDCR bound, see Theorem 4.1.) The intuition behind this result is that the influence of the prior diminishes with increasing sample sizes.

4.3.2 Monte-Carlo Markov Chains

MCMC techniques aim at using simulations to approach a distribution whose distribution is difficult to obtain analytically. Indeed, in some circumstances, one can draw in a distribution even if we do not know its analytical expression.

Definition 4.9 (Markov Chain). The sequence $\{z_i\}$ is said to be a (first-order) Markovian process if it satisfies:

$$f(z_i|z_{i-1}, z_{i-2}, \dots) = f(z_i|z_{i-1}).$$

The Metropolis-Hastings (MH) algorithm is a specific MCMC approach that allows to generate samples of θ 's whose distribution approximately corresponds to the posterior distribution of Eq. (4.19).

The MH algorithm is a recursive algorithm. That is, one can draw the i^{th} value of θ , denoted by θ_i , if one has already drawn θ_{i-1} . Assume we have θ_{i-1} . We obtain a value for θ_i by implementing the following steps:

1. Draw $\tilde{\theta}_i$ from the conditional distribution $Q_{\tilde{\theta}|\theta}(\cdot, \theta_{i-1})$, called **proposal distribution**.
2. Draw u in a uniform distribution on $[0, 1]$.
3. Compute

$$\alpha(\tilde{\theta}_i, \theta_{i-1}) := \min \left(\frac{f_{\theta,Y}(\tilde{\theta}_i, \mathbf{y})}{f_{\theta,Y}(\theta_{i-1}, \mathbf{y})} \times \frac{Q_{\tilde{\theta}|\theta}(\theta_{i-1}, \tilde{\theta}_i)}{Q_{\tilde{\theta}|\theta}(\tilde{\theta}_i, \theta_{i-1})}, 1 \right), \quad (4.22)$$

where $f_{\theta,Y}$ is given in Eq. (4.21).

4. If $u < \alpha(\tilde{\theta}_i, \theta_{i-1})$, then take $\theta_i = \tilde{\theta}_i$, otherwise we leave θ_i equal to θ_{i-1} .

It can be shown that, the distribution of the draws converges to the posterior distribution. That is, after a sufficiently large number of iterations, the draws can be considered to be drawn from the posterior distribution.⁵

⁵The proof of this claim is based on the fact that, if θ_{i-1} is drawn from the posterior distribution, then it is also the case for θ_i .

To get some insights into the algorithm, consider the case of a **symmetric proposal distribution**, that is:

$$Q_{\tilde{\theta}|\theta}(\tilde{\theta}_i, \theta_{i-1}) = Q_{\tilde{\theta}|\theta}(\theta_{i-1}, \tilde{\theta}_i). \quad (4.23)$$

We then have:

$$\alpha(\tilde{\theta}, \theta_{i-1}) = \min \left(\frac{q(\tilde{\theta}, y)}{q(\theta_{i-1}, y)}, 1 \right). \quad (4.24)$$

Remember that, up to the marginal distribution of the data ($f_Y(\mathbf{y})$), $f_{\theta,Y}(\tilde{\theta}, \mathbf{y})$ is the probability of observing \mathbf{y} conditional on having a model parameterized by $\tilde{\theta}$. Then, under Eq. (4.24), it appears that if this probability is larger for $\tilde{\theta}$ than for θ_{i-1} (in which case $\tilde{\theta}$ seems “more consistent with the observations \mathbf{y} ” than θ_{i-1}), we accept θ_i . By contrast, if $f_{\theta,Y}(\tilde{\theta}, \mathbf{y}) < f_{\theta,Y}(\theta_{i-1}, \mathbf{y})$, then we do not necessarily accept the proposed value $\tilde{\theta}$, especially if $f_{\theta,Y}(\tilde{\theta}, \mathbf{y}) \ll f_{\theta,Y}(\theta_{i-1}, \mathbf{y})$ (in which case $\tilde{\theta}$ seems far less consistent with the observations \mathbf{y} than θ_{i-1} , and, accordingly, the acceptance probability, namely $\alpha(\tilde{\theta}, \theta_{i-1})$, is small).

The choice of the **proposal distribution** $Q_{\tilde{\theta}|\theta}$ is crucial to get a rapid convergence of the algorithm. Looking at Eq. (4.22), it is easily seen that the optimal choice would be $Q_{\tilde{\theta}|\theta}(\cdot, \theta_i) = f_{\theta|Y}(\cdot, \mathbf{y})$. In that case, we would have $\alpha(\tilde{\theta}_i, \theta_{i-1}) \equiv 1$ (see Eq. (4.22)). We would then accept all draws from the proposal distribution, as this distribution would directly be the posterior distribution. Of course, this situation is not realistic as the objective of the algorithm is precisely to approximate the posterior distribution.

A common choice for Q is a multivariate normal distribution. If θ is of dimension K , we can for instance use:

$$Q(\tilde{\theta}, \theta) = \frac{1}{(\sqrt{2\pi\sigma^2})^K} \exp \left(-\frac{1}{2} \sum_{j=1}^K \frac{(\tilde{\theta}_j - \theta_j)^2}{\sigma^2} \right),$$

which is an example of symmetric proposal distribution (see Eq. (4.23)). Equivalently, we then have:

$$\tilde{\theta} = \theta + \varepsilon,$$

where ε is a K -dimensional vector of independent zero-mean normal disturbances of variance σ^2 .⁶ One then has to determine an appropriate value for σ . If it is too low, then α will be close to 1 (as $\tilde{\theta}_i$ will be close to θ_{i-1}), and we will accept very often the proposed value ($\tilde{\theta}_i$). This seems to be a favourable situation. But it may not be. Indeed, it means that it will take a large number of iterations to explore the whole distribution of θ . What if σ is very large? In

⁶We could also have different variances for the different components of θ . However, this may lead to complicated settings. A useful practice consists in looking for model (re)parametrization-based, e.g., on exponential and/or logistic functions- that are such that the components of θ are all expected to be of the order of magnitude of the unity.

this case, it is likely that the proposed values ($\tilde{\theta}_i$) will often result in poor likelihoods; The probability of acceptance will then be low and the Markov chain may be blocked at its initial value. Therefore, intermediate values of σ^2 have to be determined. The acceptance rate (i.e., the average value of $\alpha(\tilde{\theta}, \theta_{i-1})$) can be used as a guide for that. Indeed, a literature explores the optimal values for such acceptance rate (in order to obtain the best possible fit of the posterior for a minimum number of algorithm iterations). In particular, following Roberts et al. (1997), people often target acceptance rate of the order of magnitude of 20%.

It is important to note that, to implement this approach, one only has to be able to compute the joint p.d.f. $q(\theta, \mathbf{y}) = f_{Y|\theta}(\mathbf{y}, \theta)f_\theta(\theta)$ (Eq. (4.21)). That is, as soon as one can evaluate the likelihood ($f_{Y|\theta}(\mathbf{y}, \theta)$) and the prior ($f_\theta(\theta)$), we can employ this methodology.


4.3.3 Example: AR(1) specification

In the following example, we employ MCMC in order to estimate the posterior distributions of the three parameters defining an AR(1) model (see Section 6.2.2). The specification is as follows:

$$y_t = \mu + \rho y_{t-1} + \sigma \varepsilon_t, \quad \varepsilon_t \sim i.i.d. \mathcal{N}(0, 1).$$

Hence, we have $\theta = [\mu, \rho, \sigma]$. Let us first simulate the process on T periods:

```
mu <- .6; rho <- .8; sigma <- 2.5 # true model specification
T <- 20 # number of observations
y0 <- mu/(1-rho)
Y <- NULL
for(t in 1:T){
  if(t==1){
    y <- y0
  }
  y <- mu + rho*y + sigma * rnorm(1)
  Y <- c(Y,y)
}
plot(Y,type="l",xlab="time t",ylab=expression(y[t]))
```



Next, let us write the likelihood function, i.e. $f_{Y|\theta}(\mathbf{y}, \theta)$. For ρ (which is expected to be between 0 and 1), we use a logistic transformation. For σ , that is expected to be positive, we use an exponential transformation.

```
likelihood <- function(param,Y){
  mu  <- param[1]
  rho <- exp(param[2])/(1+exp(param[2]))
  sigma <- exp(param[3])
  MU <- mu/(1-rho)
  SIGMA2 <- sigma^2/(1-rho^2)
  L <- 1/sqrt(2*pi*SIGMA2)*exp(-(Y[1]-MU)^2/(2*SIGMA2))
  Y1 <- Y[2:length(Y)]
  Y0 <- Y[1:(length(Y)-1)]
  aux <- 1/sqrt(2*pi*sigma^2)*exp(-((Y1-mu-rho*Y0)^2/(2*sigma^2)))
  L <- L * prod(aux)
  return(L)
}
```

Next define function `rQ` that draws from the (Gaussian) proposal distribution, as well as function `Q`, that computes $Q_{\tilde{\theta}|\theta}(\tilde{\theta}, \theta)$:

```
rQ <- function(x,a){
  n <- length(x)
  y <- x + a * rnorm(n)
  return(y)
```

```

}

Q <- function(y,x,a){
  q <- 1/sqrt(2*pi*a^2)*exp(-(y - x)^2/(2*a^2))
  return(prod(q))
}

```

We consider Gaussian priors:

```

prior <- function(param,means_prior,stdv_prior){
  f <- 1/sqrt(2*pi*stdv_prior^2)*exp(-(param - means_prior)^2/(2*stdv_prior^2))
  return(prod(f))
}

```

Function `p_tilde` corresponds to $f_{\theta,Y}$:

```

p_tilde <- function(param,Y,means_prior,stdv_prior){
  p <- likelihood(param,Y) * prior(param,means_prior,stdv_prior)
  return(p)
}

```

We can now define function α (Eq. (4.22)):

```

alpha <- function(y,x,means_prior,stdv_prior,a){
  aux <- p_tilde(y,Y,means_prior,stdv_prior)/
    p_tilde(x,Y,means_prior,stdv_prior) * Q(y,x,a)/Q(x,y,a)
  alpha_proba <- min(aux,1)
  return(alpha_proba)
}

```

Now, all is set for us to write the MCMC function:

```

MCMC <- function(Y,means_prior,stdv_prior,a,N){
  x <- means_prior
  all_theta <- NULL
  count_accept <- 0
  for(i in 1:N){
    y <- rQ(x,a)
    alph <- alpha(y,x,means_prior,stdv_prior,a)
    #print(alph)
    u <- runif(1)
    if(u < alph){
      count_accept <- count_accept + 1
      x <- y
    }
  }
}

```

```

    all_theta <- rbind(all_theta,x)
}
print(paste("Acceptance rate:",toString(round(count_accept/N,3))))
return(all_theta)
}

```

let us specify the Gaussian priors:

```

true_values <- c(mu,log(rho/(1-rho)),log(sigma))
means_prior <- c(1,0,0) # as if we did not know the true values
stdv_prior <- rep(2,3)
resultMCMC <- MCMC(Y,means_prior,stdv_prior,a=.45,N=20000)

## [1] "Acceptance rate: 0.301"

par(mfrow=c(2,3))
for(i in 1:length(means_prior)){
  m <- means_prior[i]
  s <- stdv_prior[i]
  x <- seq(m-3*s,m+3*s,length.out = 100)
  par(mfg=c(1,i))
  aux <- density(resultMCMC[,i])
  par(plt=c(.15,.95,.15,.85))
  plot(x,dnorm(x,m,s),type="l",xlab="",ylab="",main=paste("Parameter",i),
       ylim=c(0,max(aux$y)))
  lines(aux$x,aux$y,col="red",lwd=2)
  abline(v=true_values[i],lty=2,col="blue")
  par(mfg=c(2,i))
  plot(resultMCMC[,i],1:length(resultMCMC[,i]),xlim=c(min(x),max(x)),
       type="l",xlab="",ylab="")
}

```




Figure 4.6: The upper line of plot compares prior (black) and posterior (red) distributions. The vertical dashed blue lines indicate the true values of the parameters. The second row of plots show the sequence of θ_i 's generated by the MCMC algorithm. These sequences are the ones used to produce the posterior distributions (red lines) in the upper plots.

Chapter 5

Microeconomics

5.1 Binary-choice models

In many instances, the variables to be explained (y_i s) have only two possible values (0 and 1, say).

Assume we suspect some variable \mathbf{x}_i ($K \times 1$) to be able to account for the probability that $y_i = 1$.

The model reads:

$$y_i | \mathbf{X} \sim \mathcal{B}(g(\mathbf{x}_i; \theta)), \quad (5.1)$$

where $g(\mathbf{x}_i; \theta)$ is the parameter of the Bernoulli distribution. In other words, conditionally on \mathbf{X} :

$$y_i = \begin{cases} 1 & \text{with probability } g(\mathbf{x}_i; \theta) \\ 0 & \text{with probability } 1 - g(\mathbf{x}_i; \theta), \end{cases}$$

where θ is a vector of parameters to be estimated.

The objective is to estimate the vector of population parameters θ .

Binary-choice models can be used to account for...

- any binary decisions (e.g. in referendums, being owner or renter, living in the city or in the countryside, in/out of the labour force,...),
- contamination (disease or default),
- success/failure (exams).

A possibility is to run a linear regression (a situation called **Linear Probability Model, LPM**):

$$y_i = \theta' \mathbf{x}_i + \varepsilon_i.$$

Such a regression could be consistent with the *conditional-mean-zero assumption* (Hypothesis 2.2) and with the *assumption of non-correlated residuals* (Hypothesis 2.4), but more difficultly with the **homoskedasticity assumption** (Hypothesis 2.3). Moreover, the ε_i s cannot be Gaussian (because $y_i \in \{0, 1\}$). Therefore, using a linear regression to study the relationship between \mathbf{x}_i and y_i can be consistent but it is inefficient.




Figure 5.1: Fitting a binary variable with a linear model (Linear Probability Model, LPM). The model is $\mathbb{P}(y_i = 1|x_i) = \Phi(0.5 + 2x_i)$, where Φ is the c.d.f. of the normal distribution and where $x_i \sim i.i.d. \mathcal{N}(0, 1)$.

Table 5.1: This table provides examples of function g , s.t. $\mathbb{P}(y_i = 1|\mathbf{x}_i; \theta) = g(\theta' \mathbf{x}_i)$. The “linear” case is given for comparison, but note that it does not satisfy $g(\theta' \mathbf{x}_i)$ for any value of $\theta' \mathbf{x}_i$.

Model	Function g	Derivative
Probit	Φ	ϕ
Logit	$\frac{\exp(x)}{1 + \exp(x)}$	$\frac{\exp(x)}{(1 + \exp(x))^2}$
log-	$1 - \exp(-\exp(x))$	$\exp(-\exp(x)) \exp(x)$
log		
linear	x	1

Two prominent models to tackle this situation. In both models, we have:

$$\mathbb{P}(y_i = 1|\mathbf{x}_i; \theta) = g(\theta' \mathbf{x}_i).$$

In the **Probit model**, we have

$$g(z) = \Phi(z), \quad (5.2)$$

where Φ is the c.d.f. of the normal distribution.

And for the **logit model**:

$$g(z) = \frac{1}{1 + \exp(-z)}. \quad (5.3)$$

Figure 5.2 displays the functions g appearing in Table 5.1.




Figure 5.2: Probit, Logit, and Log-log functions.

5.1.1 Interpretation in terms of latent variable, and utility-based models

The probit model has an interpretation in terms of **latent variables**. In this model, we indeed have:

$$\mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta) = \Phi(\theta' \mathbf{x}_i) = \mathbb{P}(-\varepsilon_i < \theta' \mathbf{x}_i),$$

where $\varepsilon_i \sim \mathcal{N}(0, 1)$. That is:

$$\mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta) = \mathbb{P}(0 < y_i^*),$$

where $y_i^* = \theta' \mathbf{x}_i + \varepsilon_i$, with $\varepsilon_i \sim \mathcal{N}(0, 1)$. Variable y_i^* can be interpreted as a (latent) variable that determines y_i since $y_i = \mathbb{I}_{\{y_i^* > 0\}}$.




Figure 5.3: The model is $\mathbb{P}(y_i = 1|x_i) = \Phi(0.5 + 2x_i)$, where Φ is the c.d.f. of the normal distribution and where $x_i \sim i.i.d. \mathcal{N}(0, 1)$. Crosses give the model-implied probabilities of having $y_i = 1$.




Figure 5.4: Distribution of y_i^* conditional on x_i .

Random Utility Models (RUM) are based on such a view of probit models. Assume that agent (i) chooses $y_i = 1$ if the utility associated with this choice ($U_{i,1}$) is higher than the one associated with $y_i = 0$ ($U_{i,0}$). Assume further that the utility of agent i , if she chooses outcome j ($\in \{0, 1\}$), is given by

$$U_{i,j} = V_{i,j} + \varepsilon_{i,j},$$

where $V_{i,j}$ is the deterministic component of the utility associated with choice and $\varepsilon_{i,j}$ is a random (agent-specific) component.

Moreover, posit $V_{i,j} = \theta'_j \mathbf{x}_i$. We then have:

$$\begin{aligned} \mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta) &= \mathbb{P}(\theta'_1 \mathbf{x}_i + \varepsilon_{i,1} > \theta'_0 \mathbf{x}_i + \varepsilon_{i,0}) \\ &= F(\theta'_1 \mathbf{x}_i - \theta'_0 \mathbf{x}_i) = F([\theta_1 - \theta_0]' \mathbf{x}_i), \end{aligned} \quad (5.4)$$

where F is the c.d.f. of $\varepsilon_{i,0} - \varepsilon_{i,1}$.

Note that only the difference $\theta_1 - \theta_0$ is identifiable (as opposed to θ_1 AND θ_0). Indeed, replacing U with aU ($a > 0$) gives the same model \Leftrightarrow scaling issue, solved by fixing the variance of $\varepsilon_{i,0} - \varepsilon_{i,1}$.

Other types of structural models –based on the comparison of marginal costs and benefits– give rise to the existence of latent variable and to probit models. An example is Nakosteen and Zimmer (1980). The main ingredients of their approach are as follows:

- Wage that can be earned at the present location: $y_p^* = \theta'_p \mathbf{x}_p + \varepsilon_p$.
- Migration cost: $C^* = \theta'_c \mathbf{x}_c + \varepsilon_c$.
- Wage earned elsewhere: $y_m^* = \theta'_m \mathbf{x}_m + \varepsilon_m$.

In this context, agents decision to migrate if $y_m^* > y_p^* + C^*$, i.e. if

$$y^* = y_m^* - y_p^* - C^* = \theta' \mathbf{x} + \underbrace{\xi}_{=\varepsilon_m - \varepsilon_c - \varepsilon_p} > 0,$$

where \mathbf{x} is the union of the \mathbf{x}_i s, for $i \in \{p, m, c\}$.

5.1.2 Alternative-Varying Regressors

In some cases, the regressors may depend on the considered alternative (0 or 1). For instance:

- When modeling the decision to participate in the labour force (or not), the wage depends on the alternative. It cannot be included among the regressors given it is not observed if the considered agent has decided not to work.

- In the context of the choice of transportation mode, the *time cost* depends on the considered transportation mode.

In terms of utility, we then have:

$$V_{i,j} = \theta_j^{(u)'} \mathbf{u}_{i,j} + \theta_j^{(v)'} \mathbf{v}_i,$$

where the $\mathbf{u}_{i,j}$ s are regressors associated with agent i , but taking different values for the different choices ($j = 0$ or $j = 1$).

In that case, Eq. (5.4) becomes:

$$\mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta) = F\left(\theta_1^{(u)'} \mathbf{u}_{i,1} - \theta_0^{(u)'} \mathbf{u}_{i,0} + [\theta_1^{(v)} - \theta_0^{(v)}]'\mathbf{v}_i\right), \quad (5.5)$$

and, if $\theta_1^{(u)} = \theta_0^{(u)} = \theta^{(u)}$ –as is customary– we get:

$$\mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta) = F\left(\theta_1^{(u)'} (\mathbf{u}_{i,1} - \mathbf{u}_{i,0}) + [\theta_1^{(v)} - \theta_0^{(v)}]'\mathbf{v}_i\right). \quad (5.6)$$

The fishing-mode dataset used in Cameron and Trivedi (2005) (Chapter 14 and 15) contains alternative-specific variables. Specifically, for each individual, the price and catch rate depend on the fishing model. In the table reported below, lines `price` and `catch` correspond to the prices and catch rates associated with the chosen alternative.

```
library(mlogit)
data("Fishing", package="mlogit")
stargazer::stargazer(Fishing, type="text")

##
## =====
## Statistic      N    Mean    St. Dev.     Min     Max
## -----
## price.beach   1,182  103.422  103.641   1.290  843.186
## price.pier    1,182  103.422  103.641   1.290  843.186
## price.boat    1,182   55.257   62.713   2.290  666.110
## price.charter 1,182   84.379   63.545   27.290  691.110
## catch.beach   1,182    0.241    0.191   0.068   0.533
## catch.pier    1,182    0.162    0.160   0.001   0.452
## catch.boat    1,182    0.171    0.210   0.0002  0.737
## catch.charter 1,182    0.629    0.706   0.002   2.310
## income         1,182 4,099.337 2,461.964 416.667 12,500.000
## -----
```

5.1.3 Estimation

These models can be estimated by Maximum Likelihood approaches (see Section 4.2).

To simplify the exposition, we consider the \mathbf{x}_i to be deterministic. Also, we assume that the r.v. are independent across entities i . How to write the likelihood here? It can be seen that:

$$\begin{aligned} f(y_i|\mathbf{x}_i; \theta) &= y_i g(\theta' \mathbf{x}_i) + (1 - y_i)(1 - g(\theta' \mathbf{x}_i)) \\ &= g(\theta' \mathbf{x}_i)^{y_i} (1 - g(\theta' \mathbf{x}_i))^{1-y_i}. \end{aligned} \quad (5.7)$$

Therefore, if the observations (\mathbf{x}_i, y_i) are independent across entities i , then:

$$\log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n y_i \log[g(\theta' \mathbf{x}_i)] + (1 - y_i) \log[1 - g(\theta' \mathbf{x}_i)].$$

The likelihood equation reads (FOC of the optimization program, see Def. 4.7):

$$\frac{\partial \log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X})}{\partial \theta} = \mathbf{0},$$

that is:

$$\sum_{i=1}^n y_i \mathbf{x}_i \frac{g'(\theta' \mathbf{x}_i)}{g(\theta' \mathbf{x}_i)} - (1 - y_i) \mathbf{x}_i \frac{g'(\theta' \mathbf{x}_i)}{1 - g(\theta' \mathbf{x}_i)} = \mathbf{0}.$$

This is a nonlinear equation that generally has to be numerically solved. Under regularity conditions (Hypotheses 4.1), we have (Prop. 4.4):

$$\theta_{MLE} \sim \mathcal{N}(\theta_0, \mathbf{I}(\theta_0)^{-1}),$$

where

$$\mathbf{I}(\theta_0) = -\mathbb{E}_0 \left(\frac{\partial^2 \log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X})}{\partial \theta \partial \theta'} \right) = n \mathcal{J}_Y(\theta_0).$$

For finite samples, we can e.g. approximate $\mathbf{I}(\theta_0)^{-1}$ by (Eq. (4.11)):

$$\mathbf{I}(\theta_0)^{-1} \approx - \left(\frac{\partial^2 \log \mathcal{L}(\theta_{MLE}; \mathbf{y}, \mathbf{X})}{\partial \theta \partial \theta'} \right)^{-1}.$$

In the Probit case (see Table 5.1), it can be shown that we have:

$$\begin{aligned} \frac{\partial^2 \log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X})}{\partial \theta \partial \theta'} &= - \sum_{i=1}^n g'(\theta' \mathbf{x}_i) [\mathbf{x}_i \mathbf{x}_i'] \times \\ &\left[y_i \frac{g'(\theta' \mathbf{x}_i) + \theta' \mathbf{x}_i g(\theta' \mathbf{x}_i)}{g(\theta' \mathbf{x}_i)^2} + (1 - y_i) \frac{g'(\theta' \mathbf{x}_i) - \theta' \mathbf{x}_i (1 - g(\theta' \mathbf{x}_i))}{(1 - g(\theta' \mathbf{x}_i))^2} \right]. \end{aligned}$$

In the Logit case (see Table 5.1), it can be shown that we have:

$$\frac{\partial^2 \log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X})}{\partial \theta \partial \theta'} = - \sum_{i=1}^n g'(\theta' \mathbf{x}_i) \mathbf{x}_i \mathbf{x}'_i,$$

$$\text{where } g'(x) = \frac{\exp(-x)}{(1 + \exp(-x))^2}.$$

Since $g'(x) > 0$, it can be checked that $-\partial^2 \log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X})/\partial \theta \partial \theta'$ is positive definite.

5.1.4 Marginal effectss

How to measure marginal effects, i.e. the effect on the probability that $y_i = 1$ of a marginal increase of $x_{i,k}$? This object is given by:

$$\frac{\partial \mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta)}{\partial x_{i,k}} = \underbrace{g'(\theta' \mathbf{x}_i)}_{>0} \theta_k,$$

which is of the same sign as θ_k .

It can be estimated by $g'(\theta'_{MLE} \mathbf{x}_i) \theta_{MLE,k}$. It is important to see that the marginal effect depends on \mathbf{x}_i : increases by 1 unit of $x_{i,k}$ (entity i) and of $x_{j,k}$ (entity j) do not necessarily have the same effects on $\mathbb{P}(y_i = 1 | \mathbf{x}_i; \theta)$ and on $\mathbb{P}(y_j = 1 | \mathbf{x}_j; \theta)$, respectively.

To address this issue, one can compute some measures of “average” marginal effect. There are two main solutions. For each explanatory variable k :

- i. Denoting by $\hat{\mathbf{x}}$ the sample average of the \mathbf{x}_i s, compute $g'(\theta'_{MLE} \hat{\mathbf{x}}) \theta_{MLE,k}$.
- ii. Compute the average (across i) of $g'(\theta'_{MLE} \mathbf{x}_i) \theta_{MLE,k}$.

5.1.5 Goodness of fit

There is no obvious version of “ R^2 ” for binary-choice models. Existing measures are called **pseudo- R^2** measures.

Denoting by $\log \mathcal{L}_0(\mathbf{y})$ the (maximum) log-likelihood that would be obtained for a model containing only a constant term (i.e. with $\mathbf{x}_i = 1$ for all i), the **McFadden's pseudo- R^2** is given by:


$$R_{MF}^2 = 1 - \frac{\log \mathcal{L}(\theta; \mathbf{y})}{\log \mathcal{L}_0(\mathbf{y})}.$$

Intuitively, $R_{MF}^2 = 0$ if the explanatory variables do not allow to predict the decision (y).

5.1.6 Example: Credit data

This example makes use of the `credit` data of package `AEC`. The objective is to model the default probabilities of lenders.

```
library(AEC)
credit$Default <- 0
credit$Default[credit$loan_status == "Charged Off"] <- 1
credit$Default[credit$loan_status == "Does not meet the credit policy. Status:Charged Off"] <- 1
credit$amt2income <- credit$loan_amnt/credit$annual_inc
plot(as.factor(credit$Default)~log(credit$annual_inc),
     ylevels=2:1,ylab="Default status",xlab="log(annual income)")
```



We consider three specifications. The first one, with no explanatory variables, is trivial. It will just be used to compute the pseudo- R^2 .

```
eq0 <- glm(Default ~ 1,data=credit,family=binomial(link="probit"))
eq1 <- glm(Default ~ log(loan_amnt) + amt2income + delinq_2yrs + log(annual_inc)+ I(log(annual_inc)^2),
            data=credit,family=binomial(link="probit"))
eq2 <- glm(Default ~ grade + log(loan_amnt) + amt2income + delinq_2yrs + log(annual_inc)+ I(log(annual_inc)^2),
            data=credit,family=binomial(link="probit"))
logL0 <- logLik(eq0)
logL1 <- logLik(eq1)
logL2 <- logLik(eq2)
1 - logL1/logL0 # pseudo R2
```

```

## 'log Lik.' 0.01173993 (df=6)

1 - logL2/logL0 # pseudo R2

## 'log Lik.' 0.0558487 (df=12)

stargazer::stargazer(eq0,eq1,eq2,type="text")

## 
## =====
##                               Dependent variable:
##                               -----
##                               Default
## (1)          (2)          (3)
## -----
## gradeB           0.400***  

##                   (0.055)
## 
## gradeC           0.587***  

##                   (0.057)
## 
## gradeD           0.820***  

##                   (0.061)
## 
## gradeE           0.874***  

##                   (0.091)
## 
## gradeF           1.230***  

##                   (0.147)
## 
## gradeG           1.439***  

##                   (0.227)
## 
## log(loan_amnt)   -0.149** -0.194***  

##                   (0.060)   (0.061)
## 
## amt2income        1.266***  1.222***  

##                   (0.383)   (0.393)
## 
## delinq_2yrs       0.096***  0.009  

##                   (0.034)   (0.035)
## 
## log(annual_inc)   -1.444** -0.874  

##                   (0.569)   (0.586)

```

```

## 
## I(log(annual_inc)^2)          0.064**   0.038
##                               (0.025)   (0.026)
## 
## Constant        -1.231***  7.937***  4.749
##                   (0.017)   (3.060)   (3.154)
## 
## -----
## Observations      9,156     9,156     9,156
## Log Likelihood   -3,157.696 -3,120.625 -2,981.343
## Akaike Inf. Crit. 6,317.392  6,253.250  5,986.686
## =====
## Note:           *p<0.1; **p<0.05; ***p<0.01

```

Let us now compute marginal effects.

```
mean(dnorm(predict(eq2)), na.rm=TRUE)*eq2$coefficients
```

	(Intercept)	gradeB	gradeC
##	0.840731198	0.070747353	0.103944305
##	gradeD	gradeE	gradeF
##	0.145089219	0.154773742	0.217702041
##	gradeG	log(loan_amnt)	amt2income
##	0.254722161	-0.034289921	0.216251992
##	delinq_2yrs	log(annual_inc) I(log(annual_inc)^2)	
##	0.001574178	-0.154701321	0.006813694

There is, however, an issue for the `annual_inc` variable. Indeed, the previous computation does not realize that this variable appears twice among the explanatory variables. To address this, one can proceed as follows.

```

new_credit <- credit
new_credit$annual_inc <- 1.01 * new_credit$annual_inc # increase of income by 1%
bas_predict_eq2 <- predict(eq2, newdata = credit, type = "response")
# This is equivalent to pnorm(predict(eq2, newdata = credit))
new_predict_eq2 <- predict(eq2, newdata = new_credit, type = "response")
mean(new_predict_eq2 - bas_predict_eq2)

## [1] -6.562126e-05

```

This average effect is pretty low. To compare, let us compute the average effect associated with a unit increase in the number of delinquencies:

```

new_credit <- credit
new_credit$delinq_2yrs <- credit$delinq_2yrs + 1
new_predict_eq2 <- predict(eq2, newdata = new_credit, type = "response")
mean(new_predict_eq2 - bas_predict_eq2)

## [1] 0.001582332

```

We can employ a likelihood ratio test (see Def. 4.8) to see if the two variables associated with annual income are jointly statistically significant (in the context of eq1):

```

eq1restr <- glm(Default ~ log(loan_amnt) + amt2income + delinq_2yrs,
                  data=credit,family=binomial(link="probit"))
LRstat <- 2*(logL1 - logLik(eq1restr))
pvalue <- 1 - c(pchisq(LRstat,df=2))

```

The computation gives a p-value of 0.0436.

5.1.7 Replicating Table 14.2 of Cameron and Trivedi (2005)

The following lines of codes replicate Table 14.2 of Cameron and Trivedi (2005).

```

data.reduced <- subset(Fishing,mode %in% c("charter","pier"))
data.reduced$lnrelp <- log(data.reduced$price.charter/data.reduced$price.pier)
data.reduced$y <- 1*(data.reduced$mode=="charter")
# check first line of Table 14.1:
price.charter.y0 <- mean(data.reduced$pcharter[data.reduced$y==0])

## Warning in mean.default(data.reduced$pcharter[data.reduced$y == 0]): argument is
## not numeric or logical: returning NA

price.charter.y1 <- mean(data.reduced$pcharter[data.reduced$y==1])

## Warning in mean.default(data.reduced$pcharter[data.reduced$y == 1]): argument is
## not numeric or logical: returning NA

price.charter     <- mean(data.reduced$pcharter)

## Warning in mean.default(data.reduced$pcharter): argument is not numeric or
## logical: returning NA

```

```

# Run probit regression:
reg.probit <- glm(y ~ lnrelp,
                    data=data.reduced,
                    family=binomial(link="probit"))
# Run Logit regression:
reg.logit <- glm(y ~ lnrelp,
                    data=data.reduced,
                    family=binomial(link="logit"))
# Run OLS regression:
reg.OLS <- lm(y ~ lnrelp,
                    data=data.reduced)
# Replicates Table 14.2 of Cameron and Trivedi:
stargazer::stargazer(reg.logit, reg.probit, reg.OLS,
                      type="text")

## 
## =====
##                               Dependent variable:
##                               -----
##                               y
##          logistic    probit      OLS
##          (1)        (2)        (3)
## -----
## lnrelp           -1.823***   -1.056***     -0.243***
##                   (0.145)     (0.075)      (0.010)
## 
## Constant         2.053***    1.194***     0.784*** 
##                   (0.169)     (0.088)      (0.013)
## 
## -----
## Observations       630        630        630
## R2                  0.463      0.462
## Adjusted R2        0.462
## Log Likelihood    -206.827   -204.411
## Akaike Inf. Crit.  417.654   412.822
## Residual Std. Error                         0.330 (df = 628)
## F Statistic                                542.123*** (df = 1; 628)
## =====
## Note:                                         *p<0.1; **p<0.05; ***p<0.01

```

5.1.8 Predictions

How to define predicted outcomes? As is the case for y_i , predicted outcomes \hat{y}_i need to be valued in $\{0, 1\}$. A natural choice consists in considering that

$\hat{y}_i = 1$ if $\mathbb{P}(y_i = 1|\mathbf{x}_i; \theta) > 0.5$, i.e., in taking a cutoff of $c = 0.5$. However, we may have some models where all predicted probabilities are small, but some less than others. In this context, a model-implied probability of 10% (say) could characterize a “high-risk” entity. However, using a cutoff of 50% would not identify this level of riskiness.

The **receiver operating characteristics (ROC)** curve constitutes a more general approach. It works as follows:


For each potential cutoff $c \in [0, 1]$, compute (and plot):

- The fraction of $y = 1$ values correctly classified (*True Positive Rate*) against
- The fraction of $y = 0$ values incorrectly specified (*False Positive Rate*).

Such a curve mechanically starts at (0,0) [situation when $c = 1$] and terminates at (1,1) [situation when $c = 0$].

In the case of no predictive ability (worst situation), the ROC curve is a straight line between (0,0) and (1,1).

```
library(pROC)
predict_model <- predict.glm(reg.probit, type = "response")
roc(data.reduced$y, predict_model, percent=T,
    boot.n=1000, ci.alpha=0.9, stratified=T, plot=TRUE, grid=TRUE,
    show.thres=TRUE, legacy.axes = TRUE, reuse.auc = TRUE,
    print.auc = TRUE, print.thres.col = "blue", ci=TRUE,
    ci.type="bars", print.thres.cex = 0.7, col = 'red',
    main = paste("ROC curve using", "(N = ", nrow(data.reduced), ")") )
```



```
##  
## Call:  
## roc.default(response = data.reduced$y, predictor = predict_model, percent = T, ci = TRUE,  
##  
## Data: predict_model in 178 controls (data.reduced$y 0) < 452 cases (data.reduced$y 1).  
## Area under the curve: 91.69%  
## 95% CI: 89.5%-93.87% (DeLong)
```

5.2 Multiple Choice Models

We will now consider cases where the number of possible outcomes (or alternatives) is larger than two. Let us denote by J this number. We have $y_j \in \{1, \dots, J\}$. We also consider variables \mathbf{x}_i ($K \times 1$) that are suspected to account for the probabilities that $y_i = j$, $j \in \{1, \dots, J\}$.

We will assume that the y_i s are assumed to be independently distributed, with:

$$y_i = \begin{cases} 1 & \text{with probability } g_1(\mathbf{x}_i; \theta) \\ \vdots \\ J & \text{with probability } g_J(\mathbf{x}_i; \theta), \end{cases} \quad (5.8)$$

where θ is a vector of parameters to be estimated.

Of course, for all entities (i), we must have:

$$\sum_{j=1}^J g_j(\mathbf{x}_i; \theta) = 1.$$

Our objective is to estimate the vector of population parameters θ given functional forms for the g_j s.

Multiple-choice models can for instance be used to account for

- Opinions: strongly opposed / opposed / neutral / support (ranked choices),
- Occupational field: lawyer / farmer / engineer / doctor / ...,
- Alternative shopping areas,
- Transportation types.

In a few cases, the values associated with the choices will themselves be meaningful, for example, number of accidents per day: $y = 0, 1, 2, \dots$ (count data). In most cases, the values are meaningless.

5.2.1 Ordered case

Sometimes, there exist a natural order for the different alternatives. This is typically the case where respondents have to choose a level of agreement to a statement in five points: (1) Strongly disagree; (2) Disagree; (3) Neither agree nor disagree; (4) Agree; (5) Strongly agree. Another standard case is that of ratings (from A to F, say).

The ordered probit model consists in extending the binary case, considering the latent-variable view of the latter (see Section 5.1.1). Formally, the model is as follows:

$$\mathbb{P}(y_i = j | \mathbf{x}_i) = \mathbb{P}(\alpha_{j-1} < y_i^* < \alpha_j | \mathbf{x}_i), \quad (5.9)$$

where

$$y_i^* = \theta' \mathbf{x}_i + \varepsilon_i,$$

with $\varepsilon_i \sim i.i.d. \mathcal{N}(0, 1)$. The α_j , $j \in \{1, \dots, J - 1\}$, are important (new) parameters that have to be estimated, on top of θ . Naturally, we have $\alpha_1 < \alpha_2 < \dots < \alpha_{J-1}$. Moreover α_0 is $-\infty$ and α_J is $+\infty$, so that Eq. (5.9) is valid for $j \in \{1, \dots, J\}$.

We have:

$$\begin{aligned} g_j(\mathbf{x}_i; \theta, \alpha) = \mathbb{P}(y_i = j | \mathbf{x}_i) &= \mathbb{P}(\alpha_{j-1} < y_i^* < \alpha_j | \mathbf{x}_i) \\ &= \mathbb{P}(\alpha_{j-1} - \theta' \mathbf{x}_i < \varepsilon_i < \alpha_j - \theta' \mathbf{x}_i) \\ &= F(\alpha_j - \theta' \mathbf{x}_i) - F(\alpha_{j-1} - \theta' \mathbf{x}_i), \end{aligned}$$

where F is the c.d.f. of $\mathcal{N}(0, 1)$.

If one component of \mathbf{x}_i ($\forall i$) is 1 (intercept), then one of the α_j ($j \in \{1, \dots, J-1\}$) is not identified. (One can then arbitrarily set $\alpha_1 = 0$, which is done in the binary logit/probit cases.)

This model can be estimated by maximizing the likelihood function (see Section 4.2). This function is given by:

$$\log \mathcal{L}(\theta, \alpha; \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n \sum_{j=1}^J \mathbb{I}_{\{y_i=j\}} \log(g_j(\mathbf{x}_i; \theta, \alpha)). \quad (5.10)$$

Let us stress that we have two types of parameters to estimate: those included in vector θ , and the α_j 's, gathered in vector α .

Note that the estimated values of the θ_j 's are slightly more complicated to interpret (at least in term of sign) than in the binary case. Indeed, we have:

$$\mathbb{P}(y_i \leq j | \mathbf{x}_i) = F(\alpha_j - \theta' \mathbf{x}_i) \Rightarrow \frac{\partial \mathbb{P}(y_i \leq j | \mathbf{x}_i)}{\mathbf{x}_i} = -\underbrace{F'(\alpha_j - \theta' \mathbf{x}_i)}_{>0} \theta.$$

Hence the sign of θ_k indicates whether $\mathbb{P}(y_i \leq j | \mathbf{x}_i)$ increases or decreases w.r.t. $x_{i,k}$ (the k^{th} component of \mathbf{x}_i). By contrast:

$$\frac{\partial \mathbb{P}(y_i = j | \mathbf{x}_i)}{\mathbf{x}_i} = \underbrace{(-F'(\alpha_j + \theta' \mathbf{x}_i) + F'(\alpha_{j-1} + \theta' \mathbf{x}_i))}_{A} \theta.$$

Therefore the signs of the components of θ are not necessarily those of the marginal effects. (For the sign of A is a priori unknown.)

```
library(AEC)
library(MASS)
credit$emp_length_low5y <- credit$emp_length %in% c("< 1 year", "1 year", "2 years", "3 years", "4
credit$emp_length_high10y <- credit$emp_length=="10+ years"
credit$annual_inc <- credit$annual_inc/1000
credit$loan_amnt <- credit$loan_amnt/1000
credit$income2loan <- credit$annual_inc/credit$loan_amnt
training <- credit[1:20000,]
training <- subset(training, grade!=c("E", "F", "G"))
training <- droplevels(training)
training$grade.ordered <- factor(training$grade, ordered=TRUE,
                                    levels = c("D", "C", "B", "A"))
model1 <- polr(grade.ordered ~ log(loan_amnt) + log(income2loan) + delinq_2yrs,
                data=training, Hess=TRUE, method="probit")
model2 <- polr(grade.ordered ~ log(loan_amnt) + log(income2loan) + delinq_2yrs +
               emp_length_low5y + emp_length_high10y,
```

```

    data=training, Hess=TRUE, method="probit")
stargazer::stargazer(model1,model2,ord.intercepts = TRUE,type="text",
                      no.space = TRUE)

## 
## =====
##             Dependent variable:
## -----
##                  grade.ordered
##                  (1)          (2)
## -----
## # log(loan_amnt)      -0.014     -0.040*
## #                   (0.022)     (0.022)
## # log(income2loan)   0.115***   0.092***
## #                   (0.022)     (0.022)
## # delinq_2yrs        -0.399***  -0.404*** 
## #                   (0.025)     (0.025)
## # emp_length_low5y      -0.096*** 
## #                   (0.027)
## # emp_length_high10y    0.088**
## #                   (0.035)
## # D| C            -0.937***  -1.073*** 
## #                   (0.082)     (0.086)
## # C| B            -0.160**   -0.295*** 
## #                   (0.082)     (0.085)
## # B| A            0.696***   0.564*** 
## #                   (0.082)     (0.086)
## -----
## # Observations       8,695      8,695
## =====
## Note: *p<0.1; **p<0.05; ***p<0.01

pred.grade <- predict(model1,newdata = training) # predicted grade (the most likely ac-
pred.proba <- predict(model1,newdata = training, type="probs")

```

5.2.2 General multinomial model

Regressors may be **alternative-specific** or **alternative invariant** (see also Section 5.1.2).

Fishing data: prices and catch rates are alternative-specific, income is alternative-invariant.

Notations: as in Section 5.1.2, that is ...

- $\mathbf{u}_{i,j}$: vector of variables associated with agent i and alternative j (alternative-specific regressors).

Example: Travel time per type of transportation (transportation choice), wage per type of work, cost per type of car. * \mathbf{v}_i : vector of variables associated with agent i but alternative-invariant.

Example: age or gender of agent i ,

We may e.g. organize \mathbf{x}_i as follows:

$$\mathbf{x}_i = [\mathbf{u}'_{i,1}, \dots, \mathbf{u}'_{i,J}, \mathbf{v}'_i]' \quad (5.11)$$

General Multinomial Logit

The general formulation of a multinomial logit is as follows:

$$g_j(\mathbf{x}_i; \theta) = \frac{\exp(\theta'_j \mathbf{x}_i)}{\sum_{k=1}^J \exp(\theta'_k \mathbf{x}_i)}. \quad (5.12)$$

(natural extension of the binary logit model, see Table 5.1.)

Note: By construction, $g_j(\mathbf{x}_i; \theta) \in [0, 1]$ and $\sum_j g_j(\mathbf{x}_i; \theta) = 1$.

If \mathbf{x}_i organized as in Eq. ((5.11)), then, with obvious notations, θ_j is of the form:

$$\theta_j = [\theta_{1,j}^{(u)}, \dots, \theta_{J,j}^{(u)}, \theta_j^{(v)}]', \quad (5.13)$$

(and $\theta = [\theta'_1, \dots, \theta'_J]'$).

Classical specifications

Conditional logit (CL) with alternative-varying regressors:

$$\theta_j = [\mathbf{0}', \dots, \mathbf{0}', \underbrace{\beta'}_{j^{th} \text{ position}}, \mathbf{0}', \dots]', \quad (5.14)$$

i.e. we have $\beta = \theta_{1,1}^{(u)} = \dots = \theta_{J,J}^{(u)}$ and $\theta_{i,j}^{(u)} = 0$ for $i \neq j$.

Multinomial logit (MNL) with alternative-invariant regressors:

$$\theta_j = [\mathbf{0}', \dots, \mathbf{0}', \theta_j^{(v)}]', \quad (5.15)$$

(can embed intercepts.)

Mixed logit:

$$\theta_j = [\mathbf{0}', \dots, \mathbf{0}', \beta', \mathbf{0}', \dots, \mathbf{0}', \theta_j^{(v)}]'. \quad (5.16)$$

Remark: The labelling “CL” and “MNL” –used in the literature– are relatively *ad hoc* (see 15.4.1 in Cameron and Trivedi (2005)).

The following lines replicate Table 15.2 in Cameron and Trivedi (2005).

```

# Specify data organization:
library(mlogit)
library(stargazer)
data("Fishing", package="mlogit")
Fish <- mlogit.data(Fishing,
                     varying = c(2:9),
                     choice = "mode",
                     shape = "wide")
MNL1 <- mlogit(mode ~ price + catch, data = Fish)
MNL2 <- mlogit(mode ~ price + catch - 1, data = Fish)
MNL3 <- mlogit(mode ~ 0 | income, data = Fish)
MNL4 <- mlogit(mode ~ price + catch | income, data = Fish)
stargazer(MNL1,MNL2,MNL3,MNL4, type="text", no.space = TRUE)

## 
## =====
##                               Dependent variable:
## -----
##                                     mode
##             (1)          (2)          (3)          (4)
## 
## (Intercept):boat      0.871***      0.739***      0.527**  

##                      (0.114)        (0.197)        (0.223)  

## (Intercept):charter   1.499***      1.341***      1.694***  

##                      (0.133)        (0.195)        (0.224)  

## (Intercept):pier      0.307***      0.814***      0.778***  

##                      (0.115)        (0.229)        (0.220)  

## price                 -0.025***     -0.020***     -0.025***  

##                      (0.002)        (0.001)        (0.002)  

## catch                 0.377***      0.953***      0.358***  

##                      (0.110)        (0.089)        (0.110)  

## income:boat            0.0001**  

##                      (0.00004)        (0.0001)  

## income:charter        -0.00003  

##                      (0.00004)        (0.0001)  

## income:pier           -0.0001***  

##                      (0.0001)        (0.0001)  

## 
## Observations          1,182         1,182         1,182         1,182  

## R2                    0.178         0.014         0.189  

## Log Likelihood       -1,230.784    -1,311.980    -1,477.151    -1,215.138  

## LR Test               533.878*** (df = 5)      41.145*** (df = 6)  565.171*** (d  

## 
## Note: *p<0.1; **p<0.05; ***p<0.01

```

ML estimation

Consider the general model described in Eq. (5.8). It can be noted that:

$$f(y_i | \mathbf{x}_i; \theta) = \prod_{j=1}^J g_j(\mathbf{x}_i; \theta)^{\mathbb{I}_{\{y_i=j\}}},$$

which leads to

$$\log f(y_i | \mathbf{x}_i; \theta) = \sum_{j=1}^J \mathbb{I}_{\{y_i=j\}} \log(g_j(\mathbf{x}_i; \theta)).$$

The log-likelihood function is therefore given by:

$$\log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n \sum_{j=1}^J \mathbb{I}_{\{y_i=j\}} \log(g_j(\mathbf{x}_i; \theta)). \quad (5.17)$$

Numerical methods have to be employed in order to find the maximum-likelihood estimate of θ .

Marginal Effects

Let's use the notation $p_{i,j} \equiv \mathbb{P}(y_i = j | \mathbf{x}_i; \theta)$. According to the general logit formulation (Eq. (5.12)), we have:

$$\begin{aligned} \frac{\partial p_{i,j}}{\partial x_{i,s}} &= \frac{\theta_{j,s} \exp(\theta'_j \mathbf{x}_i) \sum_{k=1}^J \exp(\theta'_k \mathbf{x}_i)}{(\sum_{k=1}^J \exp(\theta'_k \mathbf{x}_i))^2} \\ &\quad - \frac{\exp(\theta'_j \mathbf{x}_i) \sum_{k=1}^J \theta_{k,s} \exp(\theta'_k \mathbf{x}_i)}{(\sum_{k=1}^J \exp(\theta'_k \mathbf{x}_i))^2} \\ &= \theta_{j,s} p_{i,j} - \sum_{k=1}^J \theta_{k,s} p_{i,j} p_{i,k} \\ &= p_{i,j} \times \left(\theta_{j,s} - \underbrace{\sum_{k=1}^J \theta_{k,s} p_{i,k}}_{=\bar{\theta}_s^{(i)}} \right), \end{aligned}$$

where $\bar{\theta}_s^{(i)}$ does not depend on j .

It then appears that the sign of the marginal effect is not necessarily that of $\theta_{j,s}$.

Utility models

More general models can be defined. Notably by resorting to **Additive Random Utility Models, ARUM** (see Section 5.1.1).

Let's drop the i subscript for simplicity. The utility derived from choosing j is given by $U_j = V_j + \varepsilon_j$.

We have (with obvious notations):

$$\begin{aligned}\mathbb{P}(y = j) &= \mathbb{P}(U_j > U_k, \forall k \neq j) \\ \mathbb{P}(y = j) &= \mathbb{P}(U_k - U_j < 0, \forall k \neq j) \\ \mathbb{P}(y = j) &= \mathbb{P}(\underbrace{\varepsilon_k - \varepsilon_j}_{=: \tilde{\varepsilon}_{k,j}} < \underbrace{V_j - V_k}_{=: -\tilde{V}_{k,j}}, \forall k \neq j).\end{aligned}$$

The last expression is an $(J-1)$ -variate integral. It has, in general, no analytical solution.

Utility Models and the Multinomial Logit

Definition 5.1 (Gumbel distribution). Gumbel distribution (\mathcal{W}), support: \mathbb{R} (see Theorem ??):

$$F(u) = \exp(-\exp(-u)), \quad f(u) = \exp(-u - \exp(u)).$$

If $X \sim \mathcal{W}$, then $\mathbb{E}(X) = 0.577$ (Euler constant)¹ and $\text{Var}(X) = \pi^2/6$.




Figure 5.5: C.d.f. of the Gumbel distribution ($F(x) = \exp(-\exp(-x))$).

Proposition 5.1 (Weibull). *In the context of the utility model described above, if $\varepsilon_j \sim$ i.i.d. \mathcal{W} , then*

$$\mathbb{P}(y = j) = \frac{\exp(V_j)}{\sum_{k=1}^J \exp(V_k)}.$$

¹The Euler constant γ satisfies $\gamma = \lim_{n \rightarrow \infty} (-\ln(n) + \sum_{k=1}^n \frac{1}{k})$.

Proof. We have:

$$\begin{aligned}\mathbb{P}(y = j) &= \mathbb{P}(\forall k \neq j, U_k < U_j) = \mathbb{P}(\forall k \neq j, \varepsilon_k < V_j - V_k + \varepsilon_j) \\ &= \int \prod_{k \neq j} F(V_j - V_k + \varepsilon) f(\varepsilon) d\varepsilon.\end{aligned}$$

After computation, it comes that

$$\prod_{k \neq j} F(V_j - V_k + \varepsilon) f(\varepsilon) = \exp[-\varepsilon - \exp(-\varepsilon + \lambda_j)],$$

where $\lambda_j = \log \left(1 + \frac{\sum_{k \neq j} \exp(V_k)}{\exp(V_j)} \right)$. We then have:

$$\begin{aligned}\mathbb{P}(y = j) &= \int \exp[-\varepsilon - \exp(-\varepsilon + \lambda_j)] d\varepsilon \\ &= \int \exp[-t - \lambda_j - \exp(-t)] d\varepsilon = \exp(-\lambda_j),\end{aligned}$$

which leads to the result. \square

Remarks on identification

We have:

$$\begin{aligned}\mathbb{P}(y = j) &= \frac{\exp(V_j)}{\sum_{k=1}^J \exp(V_k)} \\ &= \frac{\exp(V_j^*)}{1 + \sum_{k=2}^J \exp(V_k^*)},\end{aligned}$$

where $V_j^* = V_j - V_1$.

We can always assume that $V_1 = 0$.

Case where $V_{i,j} = \theta'_j \mathbf{x}_i = \beta' \mathbf{u}_{i,j} + \theta_j^{(v)'} \mathbf{v}_i$ (see Eqs., (??) and (5.16)): we can assume that

$$\begin{aligned}(A) \quad \mathbf{u}_{i,1} &= 0, \\ (B) \quad \theta_1^{(v)} &= 0.\end{aligned}$$

If (A) does not hold, we can replace $\mathbf{u}_{i,j}$ by $\mathbf{u}_{i,j} - \mathbf{u}_{i,1}$.

If $J = 2$ and $j \in \{0, 1\}$ (shift by one unit), we have $\mathbb{P}(y = 1 | \mathbf{x}) = \frac{\exp(\theta'_j \mathbf{x})}{1 + \exp(\theta'_j \mathbf{x})}$, this is the logit model (Table 5.1).

Limitations of logit models

In a Logit model:

$$\mathbb{P}(y = j | y \in \{k, j\}) = \frac{\exp(\theta'_j \mathbf{x})}{\exp(\theta'_j \mathbf{x}) + \exp(\theta'_k \mathbf{x})}. \quad (5.18)$$

This conditional probability does not depend on other alternatives (i.e. does not depend on θ_m , $m \neq j, k$). In particular, if $\mathbf{x} = [\mathbf{u}_1', \dots, \mathbf{u}_J', \mathbf{v}']'$, then changes in \mathbf{u}_m ($m \neq j, k$) have no impact on the object shown in Eq. (5.18).

That is, a Multinomial Logit can be seen as a series of pairwise comparisons that are unaffected by the characteristics of alternatives. Such a model is said to satisfy the **independence from irrelevant alternatives (IIA)** property.

IIA property: for any individual, the ratio of probabilities of choosing two alternatives is independent of the availability or attributes of any other alternatives.

In such a mode, the choice of commuting by car or (red) bus is independent of whether commuting using a blue bus is an option.

Red-Blue Buses

Assume one has a logit modelling the “car ($y = 1$) vs red bus ($y = 2$)” choice.

If a blue bus ($y = 3$) is exactly as a red bus, except for the color, then one would expect to have:

$$\mathbb{P}(y = 3|y \in \{2, 3\}) = 0.5,$$

i.e. $\theta_2 = \theta_3$.

Assume we had $V_1 = V_2$. We expect to have $V_2 = V_3$ (hence $p_2 = p_3$). The model will then imply $p_1 = p_2 = p_3 = 0.33$.

It would seem more reasonable to have $p_1 = p_2 + p_3 = 0.5$ and $p_2 = p_3 = 0.25$.

Sequential models

Naturally, other possible ad-hoc modelling strategy are possible, e.g. sequential modelling:

$$\begin{aligned}\mathbb{P}(y = 1|\mathbf{x}) &= F(\theta'_1 \mathbf{x}) \\ \mathbb{P}(y = 2|\mathbf{x}) &= (1 - F(\theta'_1 \mathbf{x})) \times F(\theta'_2 \mathbf{x}) \\ \mathbb{P}(y = 3|\mathbf{x}) &= (1 - F(\theta'_1 \mathbf{x})) \times (1 - F(\theta'_2 \mathbf{x})).\end{aligned}$$

Easily checked that $\sum_j \mathbb{P}(y = j|\mathbf{x}) = 1$. Example: car purchase; 1st choice = brand, 2nd choice = color.

Likelihood function:

$$\mathcal{L}(\theta_1, \theta_2; \mathbf{y}, \mathbf{X}) = \prod_{j \in \{1, 2, 3\}} \left(\prod_{i \text{ s.t. } y_i=j} \mathbb{P}(y_i = j|\mathbf{x}_i) \right).$$

The maximization of the log-likelihood can be performed sequentially (w.r.t. to θ_1 first, and next w.r.t. θ_2).

Weaknesses: results may depend on the chosen sequence + no clear mapping with utility framework.

Alternative: nested logit model (see below) where utilities of alternatives 2 and 3 are affected by correlated shocks.

5.2.3 Nested logits

Nested Logits are natural extensions of logit models when choices feature a nesting structure. The nested logit approach is based on the concept of sequential choices. That is, this approach is relevant when it makes sense to group some choices into the same *nest*, also called *limbs*. This framework is consistent with the idea according to which, for each agent, there exist unobserved nest-specific variables.

The setup is as follows: we consider J *limbs*. For each limb j , we have K_j *branches*. Let us denote by y_1 the limb choice (i.e., $y_1 \in \{1, \dots, J\}$) and by y_2 the branch choice (with $y_2 \in \{1, \dots, K_j\}$). The utility associated with the pair of choices (j, k) is given by

$$U_{j,k} = V_{j,k} + \varepsilon_{j,k}.$$

We have:

$$\mathbb{P}[(y_1, y_2) = (j, k) | \mathbf{x}] = \mathbb{P}(U_{j,k} > U_{l,m}, (l, m) \neq (j, k) | \mathbf{x}).$$

One usually make the following two assumptions:

- i. The deterministic part of the utility is given by:

$$V_{j,k} = \mathbf{u}'_j \alpha + \mathbf{v}'_{j,k} \beta_j,$$

where α is common to all nests and the β_j 's are nest-specific.

- ii. The disturbances ε follow the Generalized Extreme Value (GEV) distribution (see Def. 5.2).

Under (i) and (ii), we have:

$$\begin{aligned} \mathbb{P}[(y_1, y_2) = (j, k) | \mathbf{x}] &= \frac{\exp(\mathbf{u}'_j \alpha + \rho_j I_j)}{\sum_{m=1}^J \exp(\mathbf{u}'_m \alpha + \rho_m I_m)} \times \\ &\quad \underbrace{\frac{\exp(\mathbf{v}'_{j,k} \beta_j / \rho_j)}{\sum_{l=1}^{K_j} \exp(\mathbf{v}'_{j,l} \beta_j / \rho_j)}}_{=\mathbb{P}[y_2=k|y_1=j,\mathbf{x}]} \\ &= \mathbb{P}[y_1=j | \mathbf{x}] \mathbb{P}[y_2=k | y_1=j, \mathbf{x}] \end{aligned} \tag{5.19}$$

where I_j 's are **inclusive values** or **log sums**, given by:

$$I_j = \log \left(\sum_{l=1}^{K_j} \exp(\mathbf{v}'_{j,l} \beta_j / \rho_j) \right).$$

Definition 5.2 (Generalized Extreme Value (GEV) distribution). The vector of disturbances $\varepsilon = [\varepsilon_{1,1}, \dots, \varepsilon_{1,K_1}, \dots, \varepsilon_{J,1}, \dots, \varepsilon_{J,K_J}]'$ follows the Generalized Extreme Value (GEV) distribution if its c.d.f. is:

$$F(\varepsilon, \rho) = \exp(-G(e^{-\varepsilon_{1,1}}, \dots, e^{-\varepsilon_{J,K_J}}; \rho))$$

with

$$\begin{aligned} G(\mathbf{Y}; \rho) &\equiv G(Y_{1,1}, \dots, Y_{1,K_1}, \dots, Y_{J,1}, \dots, Y_{J,K_J}; \rho) \\ &= \sum_{j=1}^J \left(\sum_{k=1}^{K_j} Y_{jk}^{1/\rho_j} \right)^{\rho_j} \end{aligned}$$

Some remarks are in order:

- a. It can be shown that $\rho_j = \sqrt{1 - \text{Cor}(\varepsilon_{j,k}, \varepsilon_{j,l})}$, for $k \neq l$.
- b. $\rho_j = 1 \Rightarrow \varepsilon_{j,k}$ and $\varepsilon_{j,l}$ are uncorrelated (\Rightarrow Multinomial logit).
- c. When $J = 1$:

$$F([\varepsilon_1, \dots, \varepsilon_K]', \rho) = \exp \left(- \left(\sum_{k=1}^K \exp(-\varepsilon_k/\rho) \right)^\rho \right).$$

The following figure displays simulations of pairs $(\varepsilon_1, \varepsilon_2)$ drawn from GEV distributions for different values of ρ . The simulation approach is based on Bhat. The code used to produce this chart is in Appendix 7.6.1.




Figure 5.6: GEV simulations.

It can be shown that:

$$I_j = \mathbb{E}(\max_k(U_{j,k})) = \mathbb{E}(\max_k(V_{j,k} + \varepsilon_{j,k})),$$

The inclusive values can be seen as measures of the relative attractiveness of a nest.

This approach allows for some level of correlation across the $\varepsilon_{j,k}$ (for a given j). This can be interpreted as the existence of an (unobserved) *common error component* for the alternatives of a same nest. This component contributes to making the alternatives of a given nest more similar. In other words, this approach can accommodate a higher sensitivity (cross-elasticity) between the alternatives of a given nest.

Note that if the common component is reduced to zero (i.e. $\rho_i = 1$), the model boils down to the multinomial logit model with no covariance of error terms among the alternatives.

Contrary to the general multinomial model, nested logits can solve the Red-Blue problem described in Section 5.2.2. Assume you have estimated a model specifying $U_1 = V_1 + \varepsilon_1$ (car choice) and $U_2 = V_2 + \varepsilon_2$ (red bus choice). You can then assume that the blue-bus utility is of the form $U_3 = V_2 + \varepsilon_3$ where ε_3 is perfectly correlated to ε_2 . This is done by redefining the set of choices as follows:

$$\begin{aligned} j = 1 &\Leftrightarrow (j' = 1, k = 1) \\ j = 2 &\Leftrightarrow (j' = 2, k = 1) \\ j = 3 &\Leftrightarrow (j' = 2, k = 2), \end{aligned}$$

and by setting $\rho_2 \rightarrow 0$.

IIA holds within a nest, but not when considering alternatives in different nests. Indeed, using Eq.(5.19):

$$\frac{\mathbb{P}[y_1 = j, y_2 = k_A | \mathbf{x}]}{\mathbb{P}[y_1 = j, y_2 = k_B | \mathbf{x}]} = \frac{\exp(\mathbf{v}'_{j,k_A} \beta_j / \rho_j)}{\exp(\mathbf{v}'_{j,k_B} \beta_j / \rho_j)},$$

i.e. we have IIA in nest j .

By contrast:

$$\begin{aligned} \frac{\mathbb{P}[y_1 = j_A, y_2 = k_A | \mathbf{x}]}{\mathbb{P}[y_1 = j_B, y_2 = k_B | \mathbf{x}]} &= \frac{\exp(\mathbf{u}'_{j_A} \alpha + \rho_{j_A} I_{j_A}) \exp(\mathbf{v}'_{j_A,k_A} \beta_{j_A} / \rho_{j_A})}{\exp(\mathbf{u}'_{j_B} \alpha + \rho_{j_B} I_{j_B}) \exp(\mathbf{v}'_{j_B,k_B} \beta_{j_B} / \rho_{j_B})} \times \\ &\quad \frac{\sum_{l=1}^{K_{j_B}} \exp(\mathbf{v}'_{j_B,l} \beta_{j_B} / \rho_{j_B})}{\sum_{l=1}^{K_{j_A}} \exp(\mathbf{v}'_{j_A,l} \beta_{j_A} / \rho_{j_A})}, \end{aligned}$$

which depends on the expected utilities of all alternatives in nest j_A and j_B . So the IIA does not hold.

Travel model dataset used, e.g., by Hensher and Greene (2002) (see also Heiss (2002)).

```

library(mlogit)
library(stargazer)
data("TravelMode", package = "AER")
Prepared.TravelMode <- mlogit.data(TravelMode, chid.var = "individual",
                                    alt.var = "mode", choice = "choice",
                                    shape = "long")
# Fit a multinomial model:
hl <- mlogit(choice ~ wait + travel + vcost, Prepared.TravelMode,
              method = "bfgs", heterosc = TRUE, tol = 10)
## Fit a nested logit model:
TravelMode$avincome <- with(TravelMode, income * (mode == "air"))
TravelMode$time <- with(TravelMode, travel + wait)/60
TravelMode$timeair <- with(TravelMode, time * I(mode == "air"))
TravelMode$income <- with(TravelMode, income / 10)
# Hensher and Greene (2002), table 1 p.8-9 model 5
TravelMode$incomeother <- with(TravelMode, ifelse(mode %in% c('air', 'car'), income, 0))
nl1 <- mlogit(choice ~ gcost + wait + incomeother, TravelMode,
              shape='long', # Inidcates how the dataset is organized
              alt.var='mode', # This is the variable that defines the alternative choi
              nests=list(public=c('train', 'bus'),
                         car='car',
                         air='air'), # The different components of the list define the
              un.nest.el = TRUE
)
nl2 <- mlogit(choice ~ gcost + wait + time, TravelMode,
              shape='long', # Inidcates how the dataset is organized
              alt.var='mode', # This is the variable that defines the alternative choi
              nests=list(public=c('train', 'bus'),
                         car='car',
                         air='air'), # The different components of the list define the
              un.nest.el = TRUE
)
stargazer(nl1,nl2,type="text",no.space = TRUE)

##
## =====
##          Dependent variable:
##          -----
##          choice
##          (1)      (2)
## -----
## (Intercept):train    -0.211     -0.284
##                      (0.562)     (0.551)
## (Intercept):bus     -0.824     -0.712
##                      (0.708)     (0.690)

```

```

## (Intercept):car -5.237*** -3.845***  

##                   (0.785)   (0.844)  

## gcost          -0.013*** -0.004  

##                   (0.004)   (0.006)  

## wait           -0.088*** -0.089***  

##                   (0.011)   (0.011)  

## incomeother    0.430***  

##                   (0.113)  

## time            -0.202***  

##                   (0.060)  

## iv              0.835*** 0.877***  

##                   (0.192)   (0.198)  

## -----  

## Observations     210      210  

## R2              0.328      0.313  

## Log Likelihood  -190.779   -194.841  

## LR Test (df = 7) 185.959*** 177.836***  

## ======  

## Note:           *p<0.1; **p<0.05; ***p<0.01

```

5.3 Tobit models

In some situations, the dependent variable is incompletely observed, which may result in a non-representative sample. Typically, in some cases, observations of the dependent variable can have a lower and/or an upper limit, while the “true”, underlying, dependent variable has not. In this case, OLS regression may lead to inconsistent parameter estimates.

Tobit models have been designed to address some of these situations. This approach has been named after James Tobin, who developed this model in the late 50s (see Tobin (1956)).

Figure 5.7 illustrates the situation. The dots (white and black) represent the “true” observations. Now, assume that only the black are observed. If one uses these observations in an OLS regression to estimate the relationship between x and y , then one gets the red line. It is clear that the sensitivity of y to x is then underestimated. The blue line is the line one would obtain if white dots were also observed; the grey line represents the model used to generate the data ($y_i = x_i + \varepsilon_i$).

Assume that the (partially) observed dependent variable follows:

$$y^* = \beta' \mathbf{x} + \varepsilon,$$

with ε drawn from a distribution characterized by a p.d.f. denoted by f_γ^* and a c.d.f. denoted by F_γ^* ; these functions depend on a vector of parameters γ .




Figure 5.7: Bias in the case of sample selection. The grey line represents the population regression line. The model is $y_i = x_i + \varepsilon_i$, with $\varepsilon_{i,t} \sim \mathcal{N}(0, 1)$. The red line is the OLS regression line based on black dots only.

The observed dependent variable is:

$$\begin{aligned} \text{Censored case: } y &= \begin{cases} y^* & \text{if } y^* > L \\ L & \text{if } y^* \leq L, \end{cases} \\ \text{Truncated case: } y &= \begin{cases} y^* & \text{if } y^* > L \\ - & \text{if } y^* \leq L, \end{cases} \end{aligned}$$

where “-” stands for missing observations.

This formulation is easily extended to censoring from above ($L \rightarrow U$) or censoring from both below and above.

The model parameters: $\theta = [\beta', \gamma']'$.

Let us write the conditional p.d.f. of the observed variable:

$$\begin{aligned} \text{Censored case: } f(y|\mathbf{x}; \theta) &= \begin{cases} f_\gamma^*(y - \beta' \mathbf{x}) & \text{if } y > L \\ F_\gamma^*(L - \beta' \mathbf{x}) & \text{if } y = L, \end{cases} \\ \text{Truncated case: } f(y|\mathbf{x}; \theta) &= \frac{f_\gamma^*(y - \beta' \mathbf{x})}{1 - F_\gamma^*(L - \beta' \mathbf{x})} \quad \text{with } y > L. \end{aligned}$$

The (conditional) log-likelihood function is then given by:

$$\log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n \log f(y_i|\mathbf{x}_i; \theta).$$

In the censored case, we have:

$$\begin{aligned} \log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X}) &= \sum_{i=1}^n \left\{ \mathbb{I}_{\{y_i=L\}} \log [F_\gamma^*(L - \beta' \mathbf{x}_i)] + \right. \\ &\quad \left. \mathbb{I}_{\{y_i>0\}} \log [f_\gamma^*(y_i - \beta' \mathbf{x}_i)] \right\}. \end{aligned}$$

The **Tobit, or censored/truncated normal regression model** corresponds to the case described above, but with Gaussian errors ε . Specifically:

$$y^* = \beta' \mathbf{x} + \varepsilon,$$

with $\varepsilon \sim i.i.d. \mathcal{N}(0, \sigma^2)$ ($\Rightarrow \gamma = \sigma^2$).

Without loss of generality, we can assume that $L = 0$. (One can shift observed data if necessary.)

- The censored density (with $L = 0$) is given by:

$$f(y) = \left[\frac{1}{\sqrt{2\pi}\sigma^2} \exp\left(-\frac{1}{2\sigma^2}(y - \beta' \mathbf{x})^2\right) \right]^{\mathbb{I}_{\{y>0\}}} \left[1 - \Phi\left(\frac{\beta' \mathbf{x}}{\sigma}\right) \right]^{\mathbb{I}_{\{y=0\}}}.$$

- The truncated density (with $L = 0$) is given by:

$$f(y) = \frac{1}{\Phi(\beta' \mathbf{x})} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y - \beta' \mathbf{x})^2\right).$$

Results usually heavily rely on distributional assumptions (more than in uncensored/untruncated case). The framework is easy to extend to an heteroskedastic case, for instance by setting $\sigma_i^2 = \exp(\alpha' \mathbf{x}_i)$.




Figure 5.8: Censored dataset with heteroskedasticity. The model is $y_i = x_i + \varepsilon_i$, with $\varepsilon_{i,t} \sim \mathcal{N}(0, \sigma_i^2)$ where $\sigma_i = \exp(-1 + x_i)$.

Let us consider the conditional means of y in the general case, i.e., for any ε distribution. Assume \mathbf{x} is observed, such that expectations are conditional on \mathbf{x} .

- For data that are left-truncated at 0, we have:

$$\begin{aligned}\mathbb{E}(y) &= \mathbb{E}(y^* | y^* > 0) \\ &= \underbrace{\beta' \mathbf{x}}_{=\mathbb{E}(y^*)} + \underbrace{\mathbb{E}(\varepsilon | \varepsilon > -\beta' \mathbf{x})}_{> 0} > \mathbb{E}(y^*).\end{aligned}$$

- Consider data that are left-censored at 0. By Bayes, we have that

$$f_{y^*|y^*>0}(u) = \frac{f_{y^*}(u)}{\mathbb{P}(y^* > 0)} \mathbb{I}_{\{u>0\}}.$$

Therefore:

$$\begin{aligned} \mathbb{E}(y^*|y^* > 0) &= \frac{1}{\mathbb{P}(y^* > 0)} \int_{-\infty}^{\infty} u f_{y^*}(u) \mathbb{I}_{\{u>0\}} du \\ &= \frac{1}{\mathbb{P}(y^* > 0)} \mathbb{E}(y^* \underbrace{\mathbb{I}_{\{y^*>0\}}}_{=y}), \end{aligned}$$

and, further:

$$\begin{aligned} \mathbb{E}(y) &= \mathbb{P}(y^* > 0) \mathbb{E}(y^*|y^* > 0) \\ &> \mathbb{E}(y^*) = \mathbb{P}(y^* > 0) \mathbb{E}(y^*|y^* > 0) + \mathbb{P}(y^* < 0) \underbrace{\mathbb{E}(y^*|y^* < 0)}_{<0}. \end{aligned}$$

Now, let us come back to the Tobit (i.e., Gaussian case) case.

- For data that are left-truncated at 0:

$$\begin{aligned} \mathbb{E}(y) &= \beta' \mathbf{x} + \mathbb{E}(\varepsilon|\varepsilon > -\beta' \mathbf{x}) \\ &= \beta' \mathbf{x} + \sigma \underbrace{\frac{\phi(\beta' \mathbf{x}/\sigma)}{\Phi(\beta' \mathbf{x}/\sigma)}}_{=\lambda(\beta' \mathbf{x}/\sigma)} \\ &= \sigma \left(\frac{\beta' \mathbf{x}}{\sigma} + \lambda \left(\frac{\beta' \mathbf{x}}{\sigma} \right) \right). \end{aligned} \quad (5.20)$$

where the penultimate line is obtained by using Eq. (7.5).

- For data that are left-censored at 0:

$$\begin{aligned} \mathbb{E}(y) &= \mathbb{P}(y^* > 0) \mathbb{E}(y^*|y^* > 0) \\ &= \Phi \left(\frac{\beta' \mathbf{x}}{\sigma} \right) \sigma \left(\frac{\beta' \mathbf{x}}{\sigma} + \lambda \left(\frac{\beta' \mathbf{x}}{\sigma} \right) \right). \end{aligned}$$

Heckit regression

The previous formula can in particular be used in an alternative estimation approach, namely the Heckman two-step estimation. This approach is based on two steps:




Figure 5.9: Conditional means of y in Tobit models. The model is $y_i = x_i + \varepsilon_i$, with $\varepsilon_i \sim \mathcal{N}(0, 1)$.

- Using the complete sample, fit a Probit model of $\mathbb{I}_{\{y_i > 0\}}$ on \mathbf{x} . This provides a consistent estimate of $\frac{\beta}{\sigma}$, and therefore of $\lambda(\beta' \mathbf{x} / \sigma)$. (Indeed, if $z_i \equiv \mathbb{I}_{\{y_i > 0\}}$, then $\mathbb{P}(z_i = 1 | \mathbf{x}_i; \beta/\sigma) = \Phi(\beta' \mathbf{x}_i / \sigma)$.)
 - Using the truncated sample only: run an OLS regression of \mathbf{y} on $\{\mathbf{x}, \lambda(\beta' \mathbf{x} / \sigma)\}$. This provides a consistent estimate of (β, σ) .

The underlying specification is of the form:

Conditional mean + disturbance.

where “Conditional mean” comes from Eq. (5.20) and “disturbance” is an error of zero conditional mean.

This approach is also applied to the case of **sample selection models** (see Section, @ref(section:SSM)).

See 16.10.2 of Cameron and Trivedi (2005) for the derivation of asymptotic standard errors of β .

The following example are based on the dataset used in Mroz (1987) (this dataset is part of the `sampleSelection` package).

```

library(sampleSelection)
library(AER)
data("Mroz87")
Mroz87$lfp.yesno <- NaN
Mroz87$lfp.yesno[Mroz87$lfp==1] <- "yes"
Mroz87$lfp.yesno[Mroz87$lfp==0] <- "no"
Mroz87$lfp.yesno <- as.factor(Mroz87$lfp.yesno)
ols <- lm(wage ~ educ + exper + I(exper^2) + city, data=subset(Mroz87, lfp==1))
tobit <- tobit(wage ~ educ + exper + I(exper^2) + city,
                left = 0, right = Inf,
                data=Mroz87)
Heckit <- heckit(lfp ~ educ + exper + I(exper^2) + city, # selection equation
                  wage ~ educ + exper + I(exper^2) + city, # outcome equation
                  data=Mroz87)

stargazer(ols,Heckit,tobit,no.space = TRUE,type="text")

##  

## ======  

##                                     Dependent variable:  

## -----  

##                                     wage  

## OLS          Heckman        Tobit

```

		selection	
	(1)	(2)	(3)
<hr/>			
## educ	0.481*** (0.067)	0.759*** (0.270)	0.642*** (0.081)
## exper	0.032 (0.062)	0.430 (0.369)	0.461*** (0.068)
## I(exper2)	-0.0003 (0.002)	-0.008 (0.008)	-0.009*** (0.002)
## city	0.449 (0.318)	0.113 (0.522)	-0.087 (0.378)
## Constant	-2.561*** (0.929)	-12.251 (8.853)	-10.395*** (1.095)
<hr/>			
## Observations	428	753	753
## R2	0.125	0.128	
## Adjusted R2	0.117	0.117	
## Log Likelihood			-1,462.700
## rho		1.063	
## Inverse Mills Ratio		5.165 (4.594)	
## Residual Std. Error	3.111 (df = 423)		
## F Statistic	15.077*** (df = 4; 423)		
## Wald Test			153.892*** (df = 4)
<hr/>			
## Note:	*p<0.1; **p<0.05; ***p<0.01		

Figure 5.10 shows that, low wages, the OLS model tends to over-predict wages. The slope between observed and Tobit-predicted wages is closer to one (the adjustment line is closer to the 45-degree line.)

Two-part model

In the standard Tobit framework, the model determining censored –or truncated– data **censoring mechanism** is the same as the one determining non-censored –or observed– data **outcome mechanism**.

A **two-part model** adds flexibility by permitting the zeros and non-zeros to be generated by different densities.

Here, the second model characterizes the outcome *conditional on* the outcome being observed.

In a seminal paper, Duan et al. (1983) employ this methodology to account for individual annual hospital expenses. The two models are then as follows:

- 1st model: $\mathbb{P}(hosp = 1|\mathbf{x}) = \Phi(\mathbf{x}'_1 \boldsymbol{\beta}_1)$,
- 2nd model: $Expense = \exp(\mathbf{x}'_2 \boldsymbol{\beta}_2 + \eta)$, with $\eta \sim i.i.d. \mathcal{N}(0, \sigma^2_2)$.




Figure 5.10: Predicted versus observed wages.

We have:

$$\mathbb{E}(\text{Expense}|\mathbf{x}_1, \mathbf{x}_2) = \Phi(\mathbf{x}'_1 \beta_1) \exp\left(\mathbf{x}'_2 \beta_2 + \frac{\sigma^2_2}{2}\right).$$

In sample-selection models, studied in the next section, one specifies the joint distribution for the censoring and outcome mechanisms (while the two parts are independent here).

5.4 Sample Selection Models

The situation tackled by sample-selection models is the following. The dependent variable of interest, denoted by y_2 , depends on observed variables \mathbf{x}_2 . The fact that y_2 is observed depends on the value of a latent variable (y_1^*) that is correlated to observed variables \mathbf{x}_1 . The difference w.r.t. the two-part model sketched above is that, even conditionally on $(\mathbf{x}_1, \mathbf{x}_2)$, y_1^* and y_2 may be correlated.

As in the Tobit case, even in the simplest case of population conditional mean linear in regressors (i.e. $y_2 = \mathbf{x}'_2 \beta_2 + \varepsilon_2$), OLS regression leads to inconsistent parameter estimates because the sample is not representative of the population.

There are two latent variables: y_1^* and y_2^* . We observe y_1 and, if the considered entity ‘participates’, y_2 . More specifically:

$$\begin{aligned} y_1 &= \begin{cases} 1 & \text{if } y_1^* > 0 \\ 0 & \text{if } y_1^* \leq 0 \end{cases} \quad (\text{participation equation}) \\ y_2 &= \begin{cases} y_2^* & \text{if } y_1 = 1 \\ - & \text{if } y_1 = 0 \end{cases} \quad (\text{outcome equation}). \end{aligned}$$

Moreover:

$$\begin{aligned} y_1^* &= \mathbf{x}'_1 \beta_1 + \varepsilon_1 \\ y_2^* &= \mathbf{x}'_2 \beta_2 + \varepsilon_2, \end{aligned}$$

Note that the Tobit model (Section 5.3) is the special case where $y_1^* = y_2^*$.

Usually:

$$\begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \end{bmatrix} \sim \mathcal{N}\left(\mathbf{0}, \begin{bmatrix} 1 & \rho\sigma_2 \\ \rho\sigma_2 & \sigma_2^2 \end{bmatrix}\right).$$

Let us derive the likelihood associated with this model. We have:

$$f(\underbrace{0}_{=y_1}, \underbrace{-}_{=y_2} | \mathbf{x}; \theta) = \mathbb{P}(y_1^* \leq 0) = \Phi(-\mathbf{x}'_1 \beta_1) \quad (5.21)$$

$$\begin{aligned} f(1, y_2 | \mathbf{x}; \theta) &= f(y_2^* | \mathbf{x}; \theta) \mathbb{P}(y_1^* > 0 | y_2^*, \mathbf{x}; \theta) \\ &= \frac{1}{\sigma} \phi\left(\frac{y_2 - \mathbf{x}'_2 \beta_2}{\sigma}\right) \times \\ &\quad \mathbb{P}(y_1^* > 0 | y_2, \mathbf{x}; \theta). \end{aligned} \quad (5.22)$$

Let's compute $\mathbb{P}(y_1^* > 0 | y_2, \mathbf{x}; \theta)$. By Prop. ?? (in Appendix 7.4), applied to $(\varepsilon_1, \varepsilon_2)$, we have:

$$y_1^* | y_2 \sim \mathcal{N}\left(\mathbf{x}'_1 \beta_1 + \frac{\rho}{\sigma_2} (y_2 - \mathbf{x}'_2 \beta_2), 1 - \rho^2\right).$$

which leads to

$$\mathbb{P}(y_1^* > 0 | y_2, \mathbf{x}; \theta) = \Phi\left(\frac{\mathbf{x}'_1 \beta_1 + \frac{\rho}{\sigma_2} (y_2 - \mathbf{x}'_2 \beta_2)}{\sqrt{1 - \rho^2}}\right). \quad (5.23)$$

Figure 5.11 shows $\mathbb{P}(y_1^* > 0 | y_2, \mathbf{x}; \theta)$ for different values of y_2 and of ρ , in the case where $\beta_1 = \beta_2 = 0$.




Figure 5.11: Probability of observing y_2 depending on its value, for different values of conditional correlation between y_2 and y_1^* .

Using Eqs. (5.21), (5.22) and (5.23) gives the log-likelihood function:

$$\begin{aligned}\log \mathcal{L}(\theta; \mathbf{y}, \mathbf{X}) &= \sum_{i=1}^n (1 - y_{1,i}) \log \Phi(-\mathbf{x}'_{1,i} \beta_1) + \\ &\quad \sum_{i=1}^n y_{1,i} \log \left(\frac{1}{\sigma} \phi \left(\frac{y_{2,i} - \mathbf{x}'_{2,i} \beta_2}{\sigma} \right) \right) + \\ &\quad \sum_{i=1}^n y_{1,i} \log \left(\Phi \left(\frac{\mathbf{x}'_{1,i} \beta_1 + \frac{\rho}{\sigma_2} (y_{2,i} - \mathbf{x}'_{2,i} \beta_2)}{\sqrt{1-\rho^2}} \right) \right).\end{aligned}$$

Computation of conditional expectation:

$$\begin{aligned}\mathbb{E}(y_2^* | y_1 = 1, \mathbf{x}) &= \mathbb{E}(\mathbb{E}(y_2^* | y_1^*, \mathbf{x}) | y_1 = 1, \mathbf{x}) \\ &= \mathbb{E}(\mathbf{x}'_2 \beta_2 + \rho \sigma_2 (y_1^* - \mathbf{x}'_1 \beta_1) | y_1 = 1, \mathbf{x}) \\ &= \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \underbrace{\mathbb{E}(y_1^* - \mathbf{x}'_1 \beta_1 | \varepsilon_1 > -\mathbf{x}'_1 \beta_1, \mathbf{x})}_{=\varepsilon_1 \sim \mathcal{N}(0,1)} \\ &= \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \frac{\phi(-\mathbf{x}'_1 \beta_1)}{1 - \Phi(-\mathbf{x}'_1 \beta_1)} \\ &= \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \frac{\phi(\mathbf{x}'_1 \beta_1)}{\Phi(\mathbf{x}'_1 \beta_1)} \\ &= \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \lambda(\mathbf{x}'_1 \beta_1).\end{aligned}\tag{5.24}$$

Similarly:

$$\begin{aligned}\mathbb{E}(y_2^* | y_1 = 0, \mathbf{x}) &= \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \mathbb{E}(y_1^* - \mathbf{x}'_1 \beta_1 | \varepsilon_1 \leq -\mathbf{x}'_1 \beta_1, \mathbf{x}) \\ &= \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \frac{\phi(-\mathbf{x}'_1 \beta_1)}{1 - \Phi(-\mathbf{x}'_1 \beta_1)} \\ &= \mathbf{x}'_2 \beta_2 - \rho \sigma_2 \frac{\phi(-\mathbf{x}'_1 \beta_1)}{\Phi(-\mathbf{x}'_1 \beta_1)} = \mathbf{x}'_2 \beta_2 - \rho \sigma_2 \lambda(-\mathbf{x}'_1 \beta_1).\end{aligned}$$

Heckman procedure

In particular, Eq. ((5.24) shows that $\mathbb{E}(y_2^* | y_1 = 1, \mathbf{x}) \neq \mathbf{x}'_2 \beta_2$ when $\rho \neq 0 \Rightarrow$ biased OLS estimates based only on the sub-sample where $y_1 = 1$.

Heckman two-step procedure (or “Heckit’’): replace $\lambda(\mathbf{x}'_1 \beta_1)$ appearing in Eq. (5.24) by an estimate. More precisely:

1. Get an estimate $\widehat{\beta}_1$ of β_1 (probit regression of y_1 on \mathbf{x}_1).
2. Run the OLS regression (using only data associated with $y_1 = 1$):

$$y_2 = \mathbf{x}'_2 \beta_2 + \rho \sigma_2 \lambda(\mathbf{x}'_1 \widehat{\beta}_1) + \varepsilon_2,\tag{5.25}$$

considering $\lambda(\mathbf{x}'_1 \widehat{\beta}_1)$ as a regressor.

How to estimate σ_2^2 ? By Eq. (7.6), we have:

$$\mathbb{V}ar(y_2|y_1^* > 0, \mathbf{x}) = \mathbb{V}ar(\varepsilon_2|\varepsilon_1 > -\mathbf{x}'_1\beta_1, \mathbf{x}).$$

Using that ε_2 can be decomposed as $\rho\varepsilon_1 + \xi$, where $\xi \sim \mathcal{N}(0, \sigma_2^2(1 - \rho^2))$ is independent from ε_1 , we get:

$$\mathbb{V}ar(y_2|y_1^* > 0, \mathbf{x}) = \sigma_2^2(1 - \rho^2) + \rho^2\sigma_2^2\mathbb{V}ar(\varepsilon_1|\varepsilon_1 > -\mathbf{x}'_1\beta_1, \mathbf{x}).$$

Using Eq. (7.7), we get:

$$\mathbb{V}ar(\varepsilon_1|\varepsilon_1 > -\mathbf{x}'_1\beta_1, \mathbf{x}) = 1 - \mathbf{x}'_1\beta_1\lambda(\mathbf{x}'_1\beta_1) - \lambda(\mathbf{x}'_1\beta_1)^2.$$

$$\begin{aligned} & \mathbb{V}ar(y_2|y_1^* > 0, \mathbf{x}) \\ &= \sigma_2^2(1 - \rho^2) + \rho^2\sigma_2^2(1 - \mathbf{x}'_1\beta_1\lambda(\mathbf{x}'_1\beta_1) - \lambda(\mathbf{x}'_1\beta_1)^2) \\ &= \sigma_2^2 - \rho^2\sigma_2^2(\mathbf{x}'_1\beta_1\lambda(\mathbf{x}'_1\beta_1) + \lambda(\mathbf{x}'_1\beta_1)^2). \end{aligned}$$

Therefore:

$$\sigma_2^2 \approx \widehat{\mathbb{V}ar}(y_2|y_1^* > 0, \mathbf{x}) + \widehat{\rho\sigma_2}^2 \left(\mathbf{x}'_1\widehat{\beta}_1\lambda(\mathbf{x}'_1\widehat{\beta}_1) + \lambda(\mathbf{x}'_1\widehat{\beta}_1)^2 \right).$$

Heckman procedure is computationally simple. Although computational costs are no longer an issue, the two-step solution allows certain generalisations more easily than ML, and is more robust in certain circumstances.

The computation of parameter standard errors is complicated because of the two steps (see Cameron and Trivedi (2005)), Subsection 16.10.2). Bootstrap can be resorted to.

Let us use the Mroz (1987) dataset again, with the objective of explaining wage setting.

```
library(sampleSelection)
library(AER)
data("Mroz87")
Mroz87$lfp.yesno <- NaN
Mroz87$lfp.yesno[Mroz87$lfp==1] <- "yes"
Mroz87$lfp.yesno[Mroz87$lfp==0] <- "no"
Mroz87$lfp.yesno <- as.factor(Mroz87$lfp.yesno)

#Logit & Probit (selection equation)
logitW <- glm(lfp ~ age + I( age^2 ) + kids5 + huswage + educ,
               family = binomial(link = "logit"), data = Mroz87)
probitW <- glm(lfp ~ age + I( age^2 ) + kids5 + huswage + educ,
                family = binomial(link = "probit"), data = Mroz87)
# OLS for outcome:
```

```

ols1 <- lm(log(wage) ~ educ + exper + I(exper^2) + city, data=subset(Mroz87, lfp==1))
# Two-step Heckman estimation
heckvan <- heckit(lfp ~ age + I(age^2) + kids5 + huswage + educ, # selection equation
                  log(wage) ~ educ + exper + I(exper^2) + city, # outcome equation
                  data=Mroz87)
# Maximum likelihood estimation of selection model:
ml <- selection(lfp~age+I(age^2)+kids5+huswage+educ,
                  log(wage)~educ+exper+I(exper^2)+city, data = Mroz87)
# Print selection-equation estimates:
stargazer(logitW,probitW,heckvan,ml,type = "text",
           selection.equation = TRUE)

## 
## =====
##                               Dependent variable:
## -----
##                                     lfp
## logistic      probit      Heckman      selection
##                                     selection
##                                     (1)       (2)       (3)       (4)
## -----
## age            0.012     0.010     0.010     0.010
##             (0.114)   (0.069)   (0.069)   (0.069)
## 
## I(age2)        -0.001    -0.0005   -0.0005   -0.0005
##             (0.001)   (0.001)   (0.001)   (0.001)
## 
## kids5          -1.409***  -0.855***  -0.855***  -0.854*** 
##             (0.198)   (0.116)   (0.115)   (0.116)
## 
## huswage        -0.069***  -0.042***  -0.042***  -0.042*** 
##             (0.020)   (0.012)   (0.012)   (0.013)
## 
## educ           0.244***  0.148***  0.148***  0.148*** 
##             (0.040)   (0.024)   (0.023)   (0.024)
## 
## Constant       -0.938     -0.620     -0.620     -0.615
##             (2.508)   (1.506)   (1.516)   (1.518)
## 
## Observations    753       753       753       753
## R2              0.158
## Adjusted R2     0.148
## Log Likelihood -459.955  -459.901      -891.177
## Akaike Inf. Crit. 931.910  931.802

```


5.5 Models of Count Data

Count-data models aim at explaining dependent variables y_i that take integer values. Typically, one may want to account for the number of doctor visits, of customers, of hospital stays, of borrowers' defaults, of recreational trips, of accidents. Quite often, these data feature large proportion of zeros (see, e.g., Table 20.1 in Cameron and Trivedi (2005)), and/or are skewed to the right.

5.5.1 Poisson model

The most basic count-data model is the Poisson model. In this model, we have $y \sim \mathcal{P}(\mu)$, i.e.

$$\mathbb{P}(y = k) = \frac{\mu^k e^{-\mu}}{k!},$$

implying $\mathbb{E}(y) = \text{Var}(y) = \mu$.

the Poisson parameter, μ , is then assumed to depend on some observed variables, gathered in vector \mathbf{x}_i for entity i . To ensure that $\mu_i \geq 0$, it is common to take $\mu_i = \exp[\beta' \mathbf{x}_i]$, which gives: A standard assumption:

$$y_i \sim \mathcal{P}(\exp[\beta' \mathbf{x}_i]).$$

The Poisson regression is intrinsically heteroskedastic (since $\text{Var}(y_i) = \exp[\beta' \mathbf{x}_i]$).

Under the assumption of independence across entities, the log-likelihood is simple to derive:

$$\log \mathcal{L}(\beta; \mathbf{y}, \mathbf{X}) = \sum_{i=1}^n (y_i \beta' \mathbf{x}_i - \exp[\beta' \mathbf{x}_i] - \ln[y_i!]).$$

The first-order condition to get the MLE is:

$$\sum_{i=1}^n (y_i - \exp[\beta' \mathbf{x}_i]) \mathbf{x}_i = \underset{K \times 1}{\mathbf{0}}. \quad (5.26)$$

Eq. (5.26) is equivalent to what would define the **Pseudo Maximum-Likelihood** estimator of β in the (misspecified) model

$$y_i \sim i.i.d. \mathcal{N}(\exp[\beta' \mathbf{x}_i], \sigma^2).$$

That is, Eq. (5.26) also characterizes the (true) ML estimator of β in the previous model.

Since $\mathbb{E}(y_i | \mathbf{x}_i) = \exp(\beta' \mathbf{x}_i)$, we have

$$y_i = \exp(\beta' \mathbf{x}_i) + \varepsilon_i,$$

with $\mathbb{E}(\varepsilon_i | \mathbf{x}_i) = 0$. This notably implies that the (N)LS estimator of β is consistent.

How to interpret regression coefficients (the components of β)? We have:

$$\frac{\partial \mathbb{E}(y_i | \mathbf{x}_i)}{\partial x_{i,j}} = \beta_j \exp(\beta' \mathbf{x}_i),$$

which depends on the considered individual.

The average response is:

$$\hat{\beta}_j \frac{1}{n} \sum_{i=1}^n \exp(\beta' \mathbf{x}_i),$$

which is equal to $\hat{\beta}_j \bar{y}$ if the model includes a constant (e.g. if $x_{1,i} = 1$).

The limitation of the standard Poisson model is that the distribution of y_i conditional on \mathbf{x}_i depends on a single parameter (μ_i). Besides, there is often a tension between fitting the fraction of zeros, i.e. $\mathbb{P}(y_i = 0 | \mathbf{x}_i) = \exp[-\exp(\beta' \mathbf{x}_i)]$, and the distribution of $y_i | \mathbf{x}_i, y_i > 0$. The following models (negative binomial, or NB model, the Hurdle model, and the Zero-Inflated model) have been designed to address these points.

5.5.2 Negative binomial model

In the **negative binomial model**, we have:

$$y_i | \lambda_i \sim \mathcal{P}(\lambda_i),$$

but λ_i is now random. Specifically, it takes the form:

$$\lambda_i = \nu_i \times \exp(\beta' \mathbf{x}_i),$$

where $\nu_i \sim i.i.d. \Gamma(\underbrace{\delta}_{\text{shape}}, \underbrace{1/\delta}_{\text{scale}})$. That is, the p.d.f. of ν_i is:

$$g(\nu) = \frac{\nu^{\delta-1} e^{-\nu\delta} \delta^\delta}{\Gamma(\delta)},$$

where $\Gamma : z \mapsto \int_0^{+\infty} t^{z-1} e^{-t} dt$ (and $\Gamma(k+1) = k!$).

This notably implies that:

$$\mathbb{E}(\nu_i) = 1 \quad \text{and} \quad \text{Var}(\nu) = \frac{1}{\delta}.$$

Hence, the p.d.f. of y_i conditional on $\exp(\beta' \mathbf{x}_i) = \mu$ and δ is obtained as a mixture of densities:

$$\mathbb{P}(y_i = k | \exp(\beta' \mathbf{x}_i) = \mu; \delta) = \int_0^{\infty} \frac{e^{-\mu\nu} (\mu\nu)^k}{k!} \frac{\nu^{\delta-1} e^{-\nu\delta} \delta^\delta}{\Gamma(\delta)} d\nu.$$

It can be shown that:

$$\mathbb{E}(y|\mathbf{x}) = \mu \quad \text{and} \quad \text{Var}(y|\mathbf{x}) = \mu(1 + \alpha\mu),$$

where $\exp(\beta' \mathbf{x}_i) = \mu$ and $\alpha = 1/\delta$.

We have one additional degree of freedom w.r.t. the Poisson model (α).

Note that $\text{Var}(y|\mathbf{x}) > \mathbb{E}(y|\mathbf{x})$ (which is often called for by the data). Moreover, the conditional variance is quadratic in the mean:

$$\text{Var}(y|\mathbf{x}) = \mu + \alpha\mu^2.$$


the previous expression is the basis of the so-called **NB2** specification.

If δ is replaced by μ/γ , then we get the **NB1** model:

$$\text{Var}(y|\mathbf{x}) = \mu(1 + \gamma).$$

The following example compares different specifications, namely a linear regression model, a Poisson model, and a NB model, to account for the number of doctor visits. The dataset ('randdata') is the one used in Chapter 20 of Cameron and Trivedi (2005) (available on that page).

```
library(AEC)
#library(foreign)
library(COUNT)
library(pscl) # for predprob function and hurdle model
#randdata <- subset(randdata, year==1) # randdata included in AEC
plot(table(randdata$mdvis))
```



```

randdata$LC <- log(1 + randdata$coins)
model.OLS <- lm(mdvis ~ LC + idp + lpi + fmde + physlm + disea + hlthg + hlthf + hlthp - 1,
                  data=randdata)
model.poisson <- glm(mdvis ~ LC + idp + lpi + fmde + physlm + disea + hlthg + hlthf + hlthp - 1,
                      data=randdata,family = poisson)
model.neg.bin <- glm.nb(mdvis ~ LC + idp + lpi + fmde + physlm + disea + hlthg + hlthf + hlthp -
                        data=randdata)
model.neg.bin.with.intercept <- glm.nb(mdvis ~ LC + idp + lpi + fmde + physlm + disea + hlthg + hlthf + hlthp +
                                         data=randdata)
stargazer::stargazer(model.OLS,model.poisson,model.neg.bin, model.neg.bin.with.intercept,type="tex"
                      )

##
## =====
##                               Dependent variable:
##                               -----
##                               mdvis
##                               OLS          Poisson        negative
##                               (1)          (2)           (3)           binomial
##                               -----
## LC                   -0.155***      -0.051***      -0.057***      -0.058***  

## (0.020)                (0.003)       (0.006)       (0.006)  

## idp                  -0.546***      -0.183***      -0.212***      -0.268***  

## (0.075)                (0.011)       (0.023)       (0.023)

```

```

## lpi          0.230***      0.095***      0.088***      0
##                               (0.012)       (0.002)       (0.004)       (0)
## fmde        -0.073***     -0.029***     -0.030***     -0
##                               (0.012)       (0.002)       (0.004)       (0)
## physlm      0.945***      0.217***      0.229***      0
##                               (0.104)       (0.013)       (0.031)       (0)
## disea        0.177***      0.050***      0.062***      0
##                               (0.004)       (0.0005)     (0.001)       (0)
## hlthg        0.270***      0.126***      0.068***     -0
##                               (0.066)       (0.009)       (0.020)       (0)
## hlthf        0.455***      0.149***      0.084**      0
##                               (0.123)       (0.016)       (0.037)       (0)
## hlthp        1.537***      0.197***      0.185**      0
##                               (0.263)       (0.027)       (0.076)       (0)
## Constant
## 
## -----
## Observations           20,190          20,190          20,190          2
## R2                      0.322
## Adjusted R2            0.322
## Log Likelihood         -64,221.340    -43,745.860    -43
## theta                   0.732*** (0.010)  0.773
## Akaike Inf. Crit.      128,460.700   87,509.710   86
## Residual Std. Error     4.393 (df = 20181)
## F Statistic             1,066.623*** (df = 9; 20181)
## =====
## Note:                  *p<0.1; **p<0.05

```

Models' predictions can be obtained as follows:

```

predict_poisson.beta.x <- predict(model.poisson) # prediction of beta'x, equivalent to
predict_poisson <- predict(model.poisson,type="response") # prediction of the number of
predict_NB <- model.neg.bin$fitted.values

```

Let us now compute the model-implied probabilities, and let's compare them with the frequencies observed in the data.

```

prop.table.data <- prop.table(table(randdata$mdvis))
predprob.poisson <- predprob(model.poisson) # part of pscl package
predprob.nb      <- predprob(model.neg.bin)
print(rbind(
  prop.table.data[1:6],
  apply(predprob.poisson[,1:6],2,mean),
  apply(predprob.nb[,1:6],2,mean)
))

```

```

##          0         1         2         3         4         5
## [1,] 0.3124319 0.1890540 0.1385339 0.09331352 0.06661714 0.04794453
## [2,] 0.1220592 0.2230328 0.2271621 0.17173478 0.10888940 0.06244353
## [3,] 0.3486824 0.1884640 0.1219340 0.08385899 0.05968603 0.04350209

```

It appears that the NB model is better at capturing the relatively large number of zeros than the Poisson model. This will also be the case for the Hurdle and Zero-Inflation models:

5.5.3 Hurdle model

The main objective of this model, w.r.t. the Poisson model, is to generate more zeros in the data than predicted by the previous count models. The idea is to separate the modeling of the number of zeros and that of the number of non-zero counts. Specifically, the frequency of zeros is determined by f_1 , the (relative) frequencies of non-zero counts are determined by f_2 :

$$f(y) = \begin{cases} f_1(0) & \text{if } y = 0, \\ \frac{1-f_1(0)}{1-f_2(0)} f_2(y) & \text{if } y > 0. \end{cases}$$

Note that we are back to the standard Poisson model if $f_1 \equiv f_2$. This model is straightforwardly estimated by ML.

5.5.4 Zero-inflated model

The objective is the same as for the Hurdle model, the modeling is slightly different. It is based on a mixture of a binary process B (p.d.f. f_1) and a process Z (p.d.f. f_2). B and Z are independent. Formally:

$$y = BZ,$$

implying:

$$f(y) = \begin{cases} f_1(0) + (1-f_1(0))f_2(0) & \text{if } y = 0, \\ (1-f_1(0))f_2(y) & \text{if } y > 0. \end{cases}$$

Typically, f_1 corresponds to a logit model and f_2 is Poisson or negative binomial density. This model is easily estimated by ML techniques.

```

# Hurdle models
# See Subsection 2.2 of https://cran.r-project.org/web/packages/pscl/vignettes/countreg.pdf
model.hurdle <- hurdle(mdvis ~ LC + idp + lpi + fmde + physlm +
                      disea + hlthg + hlthf + hlthp, data=randdata,
                      dist = "poisson", zero.dist = "binomial", link = "logit")
model.zeroinfl <- zeroinfl(mdvis ~ LC + idp + lpi + fmde + physlm +

```

```

disea + hlthg + hlthf + hlthp, data=randdata,
dist = "poisson", link = "logit")
stargazer(model.hurdle,model.zeroinfl,zero.component=FALSE,no.space=TRUE,type="text")

## -----
## Dependent variable:
## -----
##          mdvis
##      hurdle   zero-inflated
##                  count data
##          (1)           (2)
## -----
## LC          -0.015***    -0.015***
##             (0.003)      (0.003)
## idp         -0.085***   -0.086*** 
##             (0.011)      (0.011)
## lpi          0.010***   0.010*** 
##             (0.002)      (0.002)
## fmde        -0.021***   -0.021*** 
##             (0.002)      (0.002)
## physlm      0.231***   0.231*** 
##             (0.012)      (0.012)
## disea        0.022***   0.022*** 
##             (0.001)      (0.001)
## hlthg        0.027***   0.026*** 
##             (0.010)      (0.010)
## hlthf        0.147***   0.146*** 
##             (0.016)      (0.016)
## hlthp        0.304***   0.303*** 
##             (0.026)      (0.026)
## Constant     1.133***   1.133*** 
##             (0.012)      (0.012)
## -----
## Observations   20,190    20,190
## Log Likelihood -54,772.100 -54,772.550
## -----
## Note:          *p<0.1; **p<0.05; ***p<0.01

stargazer(model.hurdle,model.zeroinfl,zero.component=TRUE,no.space=TRUE,type="text")

## -----
## Dependent variable:
## -----

```

```

## -----
##          mdvis
##      hurdle zero-inflated
##                  count data
##          (1)        (2)
## -----
## LC       -0.150***   0.154***  

##           (0.010)    (0.011)  

## idp     -0.631***   0.637***  

##           (0.038)    (0.040)  

## lpi      0.102***  -0.105***  

##           (0.007)    (0.007)  

## fmde    -0.062***   0.060***  

##           (0.006)    (0.006)  

## physlm   0.239***  -0.203***  

##           (0.056)    (0.058)  

## disea    0.062***  -0.059***  

##           (0.003)    (0.003)  

## hlthg    -0.142***   0.158***  

##           (0.034)    (0.036)  

## hlthf    -0.352***   0.396***  

##           (0.062)    (0.064)  

## hlthp    -0.181      0.233  

##           (0.149)    (0.151)  

## Constant 0.411***  -0.528***  

##           (0.044)    (0.047)
## -----
## Observations   20,190    20,190
## Log Likelihood -54,772.100 -54,772.550
## =====
## Note: *p<0.1; **p<0.05; ***p<0.01

```

Let us test the importance of LC in the model using a Wald test:

```

# Test whether LC is important in the model:  

model.hurdle.reduced <- update(model.hurdle,.~.-LC)  

lmtest::waldtest(model.hurdle, model.hurdle.reduced)

## Wald test
##  

## Model 1: mdvis ~ LC + idp + lpi + fmde + physlm + disea + hlthg + hlthf +  

##           hlthp  

## Model 2: mdvis ~ idp + lpi + fmde + physlm + disea + hlthg + hlthf + hlthp  

## Res.Df Df Chisq Pr(>Chisq)  

## 1  20170

```

```
## 2 20172 -2 247.64 < 2.2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Finally, we compare average model-implied probabilities with the frequencies observed in the data:

```
predprob.hurdle <- predprob(model.hurdle)
predprob.zeroinfl <- predprob(model.zeroinfl)
print(rbind(
  prop.table.data[1:6],
  apply(predprob.poisson[,1:6], 2, mean),
  apply(predprob.nb[,1:6], 2, mean),
  apply(predprob.hurdle[,1:6], 2, mean),
  apply(predprob.zeroinfl[,1:6], 2, mean)
))

##          0         1         2         3         4         5
## [1,] 0.3124319 0.18905399 0.1385339 0.09331352 0.06661714 0.04794453
## [2,] 0.1220592 0.22303277 0.2271621 0.17173478 0.10888940 0.06244353
## [3,] 0.3486824 0.18846395 0.1219340 0.08385899 0.05968603 0.04350209
## [4,] 0.3124319 0.06056959 0.1083120 0.13262624 0.12553899 0.09847017
## [5,] 0.3124684 0.06053026 0.1082799 0.13262562 0.12556531 0.09850218
```

Chapter 6

Time Series

6.1 Introduction to time series

A time series is an infinite sequence of random variables indexed by time: $\{y_t\}_{t=-\infty}^{+\infty} = \{\dots, y_{-2}, y_{-1}, y_0, y_1, \dots, y_t, \dots\}$, $y_i \in \mathbb{R}^k$. In practice, we only observe samples, typically: $\{y_1, \dots, y_T\}$.

Standard time series models are built using **shocks** that we will often denote by ε_t . Typically, $\mathbb{E}(\varepsilon_t) = 0$. In many models, the shocks are supposed to be i.i.d., but there exist other (less restrictive) notions of shocks. In particular, the definition of many processes is based on white noises:

Definition 6.1 (White noise). The process $\{\varepsilon_t\}_{t \in]-\infty, +\infty[}$ is a white noise if, for all t

- a. $\mathbb{E}(\varepsilon_t) = 0$,
- b. $\mathbb{E}(\varepsilon_t^2) = \sigma^2 < \infty$ and
- c. for all $s \neq t$, $\mathbb{E}(\varepsilon_t \varepsilon_s) = 0$.

Another type of shocks that are commonly used are Martingale Difference Sequences:

Definition 6.2 (Martingale Difference Sequence). The process $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ is a martingale difference sequence (MDS) if $\mathbb{E}(|\varepsilon_t|) < \infty$ and if, for all t ,

$$\underbrace{\mathbb{E}_{t-1}(\varepsilon_t)}_{\text{Expectation conditional on the past}} = 0.$$

By definition, if y_t is a martingale, then $y_t - y_{t-1}$ is a MDS.

Example 6.1 (ARCH process). The Autoregressive conditional heteroskedasticity (ARCH) process is an example of shock that satisfies the MDS definition but that is not i.i.d.:

$$\varepsilon_t = \sigma_t \times z_t,$$

where $z_t \sim i.i.d. \mathcal{N}(0, 1)$ and $\sigma_t^2 = w + \alpha \varepsilon_{t-1}^2$.

Example 6.2. A white noise process is not necessarily a MDS. This is for instance the following process:

$$\varepsilon_t = z_t + z_{t-1} z_{t-2},$$

where $z_t \sim i.i.d. \mathcal{N}(0, 1)$.

Let us now introduce the lag operator. The lag operator, denoted by L , is defined on the time series space and is defined by:

$$L : \{y_t\}_{t=-\infty}^{+\infty} \rightarrow \{w_t\}_{t=-\infty}^{+\infty} \quad \text{with} \quad w_t = y_{t-1}. \quad (6.1)$$

We have: $L^2 y_t = y_{t-2}$ and, more generally, $L^k y_t = y_{t-k}$.

Consider a time series y_t defined by $y_t = \mu + \phi y_{t-1} + \varepsilon_t$, where the ε_t 's are i.i.d. $\mathcal{N}(0, \sigma^2)$. Using the lag operator, the dynamics of y_t can be expressed as follows:

$$(1 - \phi L)y_t = \mu + \varepsilon_t.$$

It is easily checked that we have $L^2 y_t = y_{t-2}$ and, generally, $L^k y_t = y_{t-k}$.

If it exists, the **unconditional (or marginal) mean** of the random variable y_t is given by:

$$\mu_t := \mathbb{E}(y_t) = \int_{-\infty}^{\infty} y_t f_{Y_t}(y_t) dy_t,$$

where f_{Y_t} is the unconditional (or marginal) density of y_t . Similarly, if it exists, the **unconditional (or marginal) variance** of the random variable y_t is:

$$\text{Var}(y_t) = \int_{-\infty}^{\infty} (y_t - \mathbb{E}(y_t))^2 f_{Y_t}(y_t) dy_t.$$

Definition 6.3 (Autocovariance). The j^{th} autocovariance of y_t is given by:

$$\begin{aligned} \gamma_{j,t} &:= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} [y_t - \mathbb{E}(y_t)][y_{t-j} - \mathbb{E}(y_{t-j})] \times \\ &\quad f_{Y_t, Y_{t-1}, \dots, Y_{t-j}}(y_t, y_{t-1}, \dots, y_{t-j}) dy_t dy_{t-1} \dots dy_{t-j} \\ &= \mathbb{E}([y_t - \mathbb{E}(y_t)][y_{t-j} - \mathbb{E}(y_{t-j})]), \end{aligned}$$

where $f_{Y_t, Y_{t-1}, \dots, Y_{t-j}}(y_t, y_{t-1}, \dots, y_{t-j})$ is the joint distribution of $y_t, y_{t-1}, \dots, y_{t-j}$.

In particular, $\gamma_{0,t} = \text{Var}(y_t)$.

Definition 6.4 (Covariance stationarity). The process y_t is covariance stationary –or weakly stationary– if, for all t and j ,

$$\mathbb{E}(y_t) = \mu \quad \text{and} \quad \mathbb{E}\{(y_t - \mu)(y_{t-j} - \mu)\} = \gamma_j.$$

Figure 6.1 displays the simulation of a process that is not covariance stationary. This process follows $y_t = 0.1t + \varepsilon_t$, where $\varepsilon_t \sim i.i.d. \mathcal{N}(0, 1)$. Indeed, for such a process, we have: $\mathbb{E}(y_t) = 0.1t$, which depends on t .




Figure 6.1: Example of a process that is not covariance stationary ($y_t = 0.1t + \varepsilon_t$, where $\varepsilon_t \sim \mathcal{N}(0, 1)$).

Definition 6.5 (Strict stationarity). The process y_t is strictly stationary if, for all t and all sets of integers $J = \{j_1, \dots, j_n\}$, the distribution of $(y_t, y_{t+j_1}, \dots, y_{t+j_n})$ depends on J but not on t .

The following process is covariance stationary but not strictly stationary:

$$y_t = \mathbb{I}_{\{t < 1000\}} \varepsilon_{1,t} + \mathbb{I}_{\{t \geq 1000\}} \varepsilon_{2,t},$$

where $\varepsilon_{1,t} \sim \mathcal{N}(0, 1)$ and $\varepsilon_{2,t} \sim \sqrt{\frac{\nu-2}{\nu}} t(\nu)$ and $\nu = 4$.

Proposition 6.1. If y_t is covariance stationary, then $\gamma_j = \gamma_{-j}$.

Proof. Since y_t is covariance stationary, the covariance between y_t and y_{t-j} (i.e. γ_j) is the same as that between y_{t+j} and y_{t+j-j} (i.e. γ_{-j}). \square

Definition 6.6 (Auto-correlation). The j^{th} auto-correlation of a covariance-stationary process is:

$$\rho_j = \frac{\gamma_j}{\gamma_0}.$$




Figure 6.2: Example of a process that is covariance stationary but not strictly stationary. The red lines delineate the 99% confidence interval of the standard normal distribution (± 2.58).

Consider a long historical time series of the Swiss GDP growth, taken from the Jordà et al. (2017) dataset.¹

Definition 6.7 (Mean ergodicity). The covariance-stationary process y_t is ergodic for the mean if:

$$\text{plim}_{T \rightarrow +\infty} \frac{1}{T} \sum_{t=1}^T y_t = \mathbb{E}(y_t).$$

Definition 6.8 (Second-moment ergodicity). The covariance-stationary process y_t is ergodic for second moments if, for all j :

$$\text{plim}_{T \rightarrow +\infty} \frac{1}{T} \sum_{t=1}^T (y_t - \mu)(y_{t-j} - \mu) = \gamma_j.$$

It should be noted that ergodicity and stationarity are different properties. Typically if the process $\{x_t\}$ is such that, $\forall t$, $x_t \equiv y$, where $y \sim \mathcal{N}(0, 1)$ (say), then $\{x_t\}$ is stationary but not ergodic.

Theorem 6.1 (Central Limit Theorem for covariance-stationary processes). *If process y_t is covariance stationary and if the series of autocovariances is*

¹Version 6 of the dataset, available on this website.




Figure 6.3: Annual growth rate of Swiss GDP, based on the Jordà-Schularick-Taylor Macrohistory Database.




Figure 6.4: For order j , the slope of the blue line is, approximately, $\hat{\gamma}_j / \widehat{\text{Var}}(y_t)$, where hats indicate sample moments.

absolutely summable ($\sum_{j=-\infty}^{+\infty} |\gamma_j| < \infty$), then:

$$\bar{y}_T \xrightarrow{m.s.} \mu = \mathbb{E}(y_t) \quad (6.2)$$

$$\lim_{T \rightarrow +\infty} T \mathbb{E}[(\bar{y}_T - \mu)^2] = \sum_{j=-\infty}^{+\infty} \gamma_j \quad (6.3)$$

$$\sqrt{T}(\bar{y}_T - \mu) \xrightarrow{d} \mathcal{N}\left(0, \sum_{j=-\infty}^{+\infty} \gamma_j\right). \quad (6.4)$$

[Mean square (m.s.) and distribution (d.) convergences: see Definitions 7.17 and 7.15.]

Proof. By Proposition 7.8, Eq. (6.3) implies Eq. (6.2). For Eq. (6.3), see Appendix 7.5. For Eq. (6.4), see Anderson (1971), p. 429. \square

Definition 6.9 (Long-run variance). Under the assumptions of Theorem 6.1, the limit appearing in Eq. (6.3) exists and is called **long-run variance**. It is denoted by S , i.e.:

$$S = \sum_{j=-\infty}^{+\infty} \gamma_j = \lim_{T \rightarrow +\infty} T \mathbb{E}[(\bar{y}_T - \mu)^2].$$

If y_t is ergodic for second moments (see Def. 6.8), a natural estimator of S is:

$$\hat{\gamma}_0 + 2 \sum_{\nu=1}^q \hat{\gamma}_{\nu}, \quad (6.5)$$

where $\hat{\gamma}_{\nu} = \frac{1}{T} \sum_{t=1}^T (y_t - \bar{y})(y_{t-\nu} - \bar{y})$.

However, for small samples, Eq. (6.5) does not necessarily result in a positive definite matrix. Newey and West (1987) have proposed an estimator that does not have this defect. Their estimator is given by:

$$S^{NW} = \hat{\gamma}_0 + 2 \sum_{\nu=1}^q \left(1 - \frac{\nu}{q+1}\right) \hat{\gamma}_{\nu}. \quad (6.6)$$

Loosely speaking, Theorem 6.1 says that, for a given sample size, the higher the “persistency” of a process, the lower the accuracy of the sample mean as an estimate of the population mean. To illustrate, consider three processes that feature the same marginal variance (equal to one, say), but different autocorrelations: 0%, 70%, and 99.9%. Figure 6.5 displays simulated paths of such three processes. It indeed appears that, the larger the autocorrelation of the process, the further the sample mean (dashed red line) from the population mean (red solid line).

Identical simulations can be performed using this ShinyApp (use panel “AR(1)”).




Figure 6.5: The three samples have been simulated using the following data generating process: $x_t = \mu + \rho(x_{t-1} - \mu) + \sqrt{1 - \rho^2}\varepsilon_t$, where $\varepsilon_t \sim \mathcal{N}(0, 1)$. Case A: $\rho = 0$; Case B: $\rho = 0.7$; Case C: $\rho = 0.999$. In the three cases, $\mathbb{E}(x_t) = \mu = 2$ and $\text{Var}(x_t) = 1$.

6.2 Univariate processes

6.2.1 Moving Average (MA) processes

Definition 6.10. Consider a white noise process $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ (Def. 6.1). Then y_t is a first-order moving average process if, for all t :

$$y_t = \mu + \varepsilon_t + \theta\varepsilon_{t-1}.$$

If $\mathbb{E}(\varepsilon_t^2) = \sigma^2$, it is easily obtained that the unconditional mean and variances of y_t are:

$$\mathbb{E}(y_t) = \mu, \quad \text{Var}(y_t) = (1 + \theta^2)\sigma^2.$$

The first auto-covariance is:

$$\gamma_1 = \mathbb{E}\{(y_t - \mu)(y_{t-1} - \mu)\} = \theta\sigma^2.$$

Higher-order auto-covariances are zero ($\gamma_j = 0$ for $j > 1$). Therefore: An MA(1) process is covariance-stationary (Def. 6.4).

For a MA(1) process, the autocorrelation of order j (see Def. 6.6) is given by:

$$\rho_j = \begin{cases} 1 & \text{if } j = 0, \\ \theta/(1 + \theta^2) & \text{if } j = 1 \\ 0 & \text{if } j > 1. \end{cases}$$

Notice that process y_t defined through:

$$y_t = \mu + \varepsilon_t + \theta \varepsilon_{t-1},$$

where $\text{Var}(\varepsilon_t) = \sigma^2$, has the same mean and autocovariances as

$$y_t = \mu + \varepsilon_t^* + \frac{1}{\theta} \varepsilon_{t-1}^*,$$

where $\text{Var}(\varepsilon_t^*) = \theta^2 \sigma^2$.

Definition 6.11 (MA(q) process). A q^{th} order Moving Average process is defined through:

$$y_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}.$$

where $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ is a white noise process (Def. 6.1).

Proposition 6.2 (Covariance-stationarity of an MA(q) process). *Finite-order Moving Average processes are covariance-stationary.*

Moreover, the autocovariances of an MA(q) process (as defined in Def. 6.11) are given by:

$$\gamma_j = \begin{cases} \sigma^2(\theta_0 \theta_0 + \theta_1 \theta_1 + \cdots + \theta_q \theta_{q-j}) & \text{for } j \in \{0, \dots, q\} \\ 0 & \text{for } j > q, \end{cases} \quad (6.7)$$

where we use the notation $\theta_0 = 1$, and $\text{Var}(\varepsilon_t) = \sigma^2$.

Proof. The unconditional expectation of y_t does not depend on time, since $\mathbb{E}(y_t) = \mu$. Let's turn to autocovariances. We can extend the series of the θ_j 's by setting $\theta_j = 0$ for $j > q$. We then have:

$$\begin{aligned} \mathbb{E}((y_t - \mu)(y_{t-j} - \mu)) &= \mathbb{E}[(\theta_0 \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_j \varepsilon_{t-j} + \theta_{j+1} \varepsilon_{t-j-1} + \cdots) \times \\ &\quad (\theta_0 \varepsilon_{t-j} + \theta_1 \varepsilon_{t-j-1} + \cdots)]. \end{aligned}$$

Then use the fact that $\mathbb{E}(\varepsilon_t \varepsilon_s) = 0$ if $t \neq s$ (because $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ is a white noise process). \square

Figure 6.6 displays simulated paths of two MA processes (an MA(1) and an MA(4)). Such simulations can be produced by using panel “ARMA(p,q)” of this web interface.

```
library(AEC)
T <- 100; nb.sim <- 1
y.0 <- c(0)
c <- 1; phi <- c(0); sigma <- 1
theta <- c(1,1) # MA(1) specification
y.sim <- sim.arma(c,phi,theta,sigma,T,y.0,nb.sim)
```

```

par(mfrow=c(1,2))
par(plt=c(.2,.9,.2,.85))
plot(y.sim[,1],xlab="",ylab="",type="l",lwd=2,
      main=expression(paste(theta[0],"=1, ",theta[1],"=1",sep="")))
abline(h=c)
theta <- c(1,1,1,1,1) # MA(4) specification
y.sim <- sim.arma(c,phi,theta,sigma,T,y.0,nb.sim)
plot(y.sim[,1],xlab="",ylab="",type="l",lwd=2,
      main=expression(paste(theta[0],"=...=",theta[4],"=1",sep="")))
abline(h=c)

```



 $\theta_0=1, \theta_1=1$  $\theta_0=\dots=\theta_4=1$ 

Figure 6.6: Simulation of MA processes.

What if the order q of an MA(q) process gets infinite? The notion of **infinite-order Moving Average process** exists and is important in time series analysis. The (infinite) sequence of θ_j has to satisfy some conditions for such a process to be well-defined (see Theorem 6.2 below). These conditions relate to the “summability” of $\{\theta_i\}_{i \in \mathbb{N}}$ (see Definition 6.12).

Definition 6.12 (Absolute and square summability). The sequence $\{\theta_i\}_{i \in \mathbb{N}}$ is absolutely summable if $\sum_{i=0}^{\infty} |\theta_i| < +\infty$, and it is square summable if $\sum_{i=0}^{\infty} \theta_i^2 < +\infty$.

According to Theorem ??, absolute summability implies square summability.

Theorem 6.2 (Existence condition for an infinite MA process). *If $\{\theta_i\}_{i \in \mathbb{N}}$ is square summable (see Def. 6.12) and if $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ is a white noise process (see Def. 6.1), then*

$$\mu + \sum_{i=0}^{+\infty} \theta_i \varepsilon_{t-i}$$

defines a well-behaved [covariance-stationary] process, called infinite-order MA process ($MA(\infty)$).

Proof. See Appendix 3.A in Hamilton."Well behaved" means that $\sum_{i=0}^T \theta_{t-i} \varepsilon_{t-i}$ converges in mean square (Def. 7.15) to some random variable Z_t . The proof makes use of the fact that:

$$\mathbb{E} \left[\left(\sum_{i=N}^M \theta_i \varepsilon_{t-i} \right)^2 \right] = \sum_{i=N}^M |\theta_i|^2 \sigma^2,$$

and that, when $\{\theta_i\}$ is square summable, $\forall \eta > 0$, $\exists N$ s.t. the right-hand-side term in the last equation is lower than η for all $M \geq N$ (static Cauchy criterion, Theorem 7.2). This implies that $\sum_{i=0}^T \theta_i \varepsilon_{t-i}$ converges in mean square (stochastic Cauchy criterion, see Theorem 7.3). \square

Proposition 6.3 (First two moments of an infinite MA process). *If $\{\theta_i\}_{i \in \mathbb{N}}$ is absolutely summable, i.e. if $\sum_{i=0}^{\infty} |\theta_i| < +\infty$, then*

i. $y_t = \mu + \sum_{i=0}^{+\infty} \theta_i \varepsilon_{t-i}$ exists (Theorem ??) and is such that:

$$\begin{aligned} \mathbb{E}(y_t) &= \mu \\ \gamma_0 &= \mathbb{E}([y_t - \mu]^2) = \sigma^2(\theta_0^2 + \theta_1^2 + \dots) \\ \gamma_j &= \mathbb{E}([y_t - \mu][y_{t-j} - \mu]) = \sigma^2(\theta_0 \theta_j + \theta_1 \theta_{j+1} + \dots). \end{aligned}$$

ii. Process y_t has absolutely summable auto-covariances, which implies that the results of Theorem 6.1 (Central Limit) apply.

Proof. The absolute summability of $\{\theta_i\}$ and the fact that $\mathbb{E}(\varepsilon^2) < \infty$ imply that the order of integration and summation is interchangeable (see Hamilton, 1994, Footnote p. 52), which proves (i). For (ii), see end of Appendix 3.A in Hamilton (1994). \square

6.2.2 Auto-Regressive (AR) processes

Definition 6.13 (First-order AR process (AR(1))). Consider a white noise process $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ (see Def. 6.1). Process y_t is an AR(1) process if it is defined by the following difference equation:

$$y_t = c + \phi y_{t-1} + \varepsilon_t.$$

If $|\phi| \geq 1$, y_t is not stationary. Indeed, we have:

$$y_{t+k} = c + \varepsilon_{t+k} + \phi(c + \varepsilon_{t+k-1}) + \phi^2(c + \varepsilon_{t+k-2}) + \dots + \phi^{k-1}(c + \varepsilon_{t+1}) + \phi^k y_t.$$

Therefore, the conditional variance

$$\mathbb{V}ar_t(y_{t+k}) = \sigma^2(1 + \phi^2 + \phi^4 + \cdots + \phi^{2(k-1)})$$

does not converge for large k 's. This implies that $\mathbb{V}ar(y_t)$ does not exist.

By contrast, if $|\phi| < 1$, one can see that:

$$y_t = c + \varepsilon_t + \phi(c + \varepsilon_{t-1}) + \phi^2(c + \varepsilon_{t-2}) + \cdots + \phi^k(c + \varepsilon_{t-k}) + \dots$$

Hence, if $|\phi| < 1$, the unconditional mean and variance of y_t are:

$$\mathbb{E}(y_t) = \frac{c}{1 - \phi} =: \mu \quad \text{and} \quad \mathbb{V}ar(y_t) = \frac{\sigma^2}{1 - \phi^2}.$$

Let us compute the j^{th} autocovariance of the AR(1) process:

$$\begin{aligned} \mathbb{E}([y_t - \mu][y_{t-j} - \mu]) &= \mathbb{E}([\varepsilon_t + \phi\varepsilon_{t-1} + \phi^2\varepsilon_{t-2} + \cdots + \phi^j\varepsilon_{t-j} + \phi^{j+1}\varepsilon_{t-j-1} \dots] \times \\ &\quad [\varepsilon_{t-j} + \phi\varepsilon_{t-j-1} + \phi^2\varepsilon_{t-j-2} + \cdots + \phi^k\varepsilon_{t-j-k} + \dots]) \\ &= \mathbb{E}(\phi^j\varepsilon_{t-j}^2 + \phi^{j+2}\varepsilon_{t-j-1}^2 + \phi^{j+4}\varepsilon_{t-j-2}^2 + \dots) \\ &= \frac{\phi^j\sigma^2}{1 - \phi^2}. \end{aligned}$$

Therefore $\rho_j = \phi^j$.

By what precedes, we have:

Proposition 6.4 (Covariance-stationarity of an AR(1) process). *The AR(1) process, as defined in Def. 6.13, is covariance-stationary iff $|\phi| < 1$.*

Definition 6.14 (AR(p) process). Consider a white noise process $\{\varepsilon_t\}_{t=-\infty}^{+\infty}$ (see Def. 6.1). Process y_t is a p^{th} -order autoregressive process (AR(p)) if its dynamics is defined by the following difference equation (with $\phi_p \neq 0$):

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + \varepsilon_t. \quad (6.8)$$

As we will see, the covariance-stationarity of process y_t hinges on matrix F defined as:

$$F = \begin{bmatrix} \phi_1 & \phi_2 & \cdots & \phi_p \\ 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{bmatrix}. \quad (6.9)$$

Note that this matrix F is such that if y_t follows Eq. (6.8), then process \mathbf{y}_t follows:

$$\mathbf{y}_t = F\mathbf{y}_{t-1} + \xi_t$$

with

$$\xi_t = \begin{bmatrix} \varepsilon_t \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad \mathbf{y}_t = \begin{bmatrix} y_t \\ y_{t-1} \\ \vdots \\ y_{t-p+1} \end{bmatrix}.$$

Proposition 6.5 (The eigenvalues of matrix F). *The eigenvalues of F (defined by Eq. (6.9)) are the solutions of:*

$$\lambda^p - \phi_1 \lambda^{p-1} - \cdots - \phi_{p-1} \lambda - \phi_p = 0. \quad (6.10)$$

Proposition 6.6 (Covariance-stationarity of an AR(p) process). *These four statements are equivalent:*

- i. Process $\{y_t\}$, defined in Def. 6.14, is covariance-stationary.
- ii. The eigenvalues of F (as defined Eq. (6.9)) lie within the unit circle.
- iii. The roots of Eq. (6.11) (below) lie strictly outside the unit circle.

$$1 - \phi_1 z - \cdots - \phi_{p-1} z^{p-1} - \phi_p z^p = 0. \quad (6.11)$$

- iv. The roots of Eq. (6.12) (below) lie strictly inside the unit circle.

$$\lambda^p - \phi_1 \lambda^{p-1} - \cdots - \phi_{p-1} \lambda - \phi_p = 0. \quad (6.12)$$

Proof. We consider the case where the eigenvalues of F are distinct; Jordan decomposition can be used in the general case. When the eigenvalues of F are distinct, F admits the following spectral decomposition: $F = PDP^{-1}$, where D is diagonal. Using the notations introduced in Eq. (6.9), we have:

$$\mathbf{y}_t = \mathbf{c} + F\mathbf{y}_{t-1} + \xi_t.$$

Let's introduce $\mathbf{d} = P^{-1}\mathbf{c}$, $\mathbf{z}_t = P^{-1}\mathbf{y}_t$ and $\eta_t = P^{-1}\xi_t$. We have:

$$\mathbf{z}_t = \mathbf{d} + D\mathbf{z}_{t-1} + \eta_t.$$

Because D is diagonal, the different component of \mathbf{z}_t , denoted by $z_{i,t}$, follow AR(1) processes. The (scalar) autoregressive parameters of these AR(1) processes are the diagonal entries of D – which also are the eigenvalues of F – that we denote by λ_i .

Process y_t is covariance-stationary iff \mathbf{y}_t also is covariance-stationary, which is the case iff all $z_{i,t}$, $i \in [1, p]$, are covariance-stationary. By Prop. 6.4, process $z_{i,t}$ is covariance-stationary iff $|\lambda_i| < 1$. This proves that (i) is equivalent to (ii). Prop. 6.5 further proves that (ii) is equivalent to (iv). Finally, it is easily seen that (iii) is equivalent to (iv) (as long as $\phi_p \neq 0$). \square

Using the lag operator (see Eq (6.1)), if y_t is a covariance-stationary AR(p) process (Def. 6.14), we can write:

$$y_t = \mu + \psi(L)\varepsilon_t,$$

where

$$\psi(L) = (1 - \phi_1 L - \cdots - \phi_p L^p)^{-1} \quad \text{and} \quad \mu = \mathbb{E}(y_t) = \frac{c}{1 - \phi_1 - \cdots - \phi_p}. \quad (6.13)$$

In the following lines of codes, we compute the eigenvalues of the F matrices associated with the following processes (where ε_t is a white noise):

$$\begin{aligned} x_t &= 0.9y_{t-1} - 0.2x_{t-2} + \varepsilon_t \\ y_t &= 1.1y_{t-1} - 0.3y_{t-2} + \varepsilon_t \\ w_t &= 1.4y_{t-1} - 0.7w_{t-2} + \varepsilon_t \\ z_t &= 0.9y_{t-1} + 0.2z_{t-2} + \varepsilon_t \end{aligned}$$

```
F <- matrix(c(.9, 1, -.2, 0), 2, 2)
lambda_x <- eigen(F)$values
F[1,] <- c(1.1, -.3)
lambda_y <- eigen(F)$values
F[1,] <- c(1.4, -.7)
lambda_w <- eigen(F)$values
F[1,] <- c(.9, .2)
lambda_z <- eigen(F)$values
rbind(lambda_x, lambda_y, lambda_w, lambda_z)

## [,1] [,2]
## lambda_x 0.500000+0.0000000i 0.4000000+0.0000000i
## lambda_y 0.600000+0.0000000i 0.5000000+0.0000000i
## lambda_w 0.700000+0.4582576i 0.7000000-0.4582576i
## lambda_z 1.084429+0.0000000i -0.1844289+0.0000000i
```

The absolute values of the eigenvalues associated with process w_t are equal to 0.837. Therefore, according to Proposition 6.6, processes x_t , y_t , and w_t are covariance-stationary, but not z_t (because the absolute value of one of the eigenvalues of the F matrix associated with this process is larger than 1).

The computation of the autocovariances of y_t is based on the so-called **Yule-Walker equations** (Eq. (6.14)). Let's rewrite Eq. (6.8):

$$(y_t - \mu) = \phi_1(y_{t-1} - \mu) + \phi_2(y_{t-2} - \mu) + \cdots + \phi_p(y_{t-p} - \mu) + \varepsilon_t.$$

Multiplying both sides by $y_{t-j} - \mu$ and taking expectations leads to the (Yule-Walker) equations:

$$\gamma_j = \begin{cases} \phi_1\gamma_{j-1} + \phi_2\gamma_{j-2} + \cdots + \phi_p\gamma_{j-p} & \text{if } j > 0 \\ \phi_1\gamma_1 + \phi_2\gamma_2 + \cdots + \phi_p\gamma_p + \sigma^2 & \text{for } j = 0. \end{cases} \quad (6.14)$$

Using $\gamma_j = \gamma_{-j}$ (Prop. 6.1), one can express $(\gamma_0, \gamma_1, \dots, \gamma_p)$ as functions of $(\sigma^2, \phi_1, \dots, \phi_p)$.

6.2.3 AR-MA processes

Definition 6.15 (ARMA(p,q) process). $\{y_t\}$ is an ARMA(p,q) process if its dynamics is described by the following equation:

$$y_t = c + \underbrace{\phi_1 y_{t-1} + \cdots + \phi_p y_{t-p}}_{\text{AR part}} + \underbrace{\varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}}_{\text{MA part}}, \quad (6.15)$$

where $\{\varepsilon_t\}_{t \in [-\infty, +\infty[}$ is a white noise process (see Def. 6.1).

Proposition 6.7 (Stationarity of an ARMA(p,q) process). *The ARMA(p,q) process defined in 6.15 is covariance stationary iff the roots of*

$$1 - \phi_1 z - \cdots - \phi_p z^p = 0$$

lie strictly outside the unit circle or, equivalently, iff those of

$$\lambda^p - \phi_1 \lambda^{p-1} - \cdots - \phi_p = 0$$

lie strictly within the unit circle.

Proof. The proof of Prop. 6.6 can be adapted to the present case. □

We can write:

$$(1 - \phi_1 L - \cdots - \phi_p L^p) y_t = c + (1 + \theta_1 L + \cdots + \theta_q L^q) \varepsilon_t.$$

If the roots of $1 - \phi_1 z - \cdots - \phi_p z^p = 0$ lie outside the unit circle, we have:

$$y_t = \mu + \psi(L) \varepsilon_t, \quad (6.16)$$

where

$$\psi(L) = \frac{1 + \theta_1 L + \cdots + \theta_q L^q}{1 - \phi_1 L - \cdots - \phi_p L^p} \quad \text{and} \quad \mu = \frac{c}{1 - \phi_1 - \cdots - \phi_p}.$$

Eq. (6.16) is the **Wold representation** of this ARMA process (see Theorem 6.3 below).

The stationarity of the process depends only on the AR specification (or on the eigenvalues of matrix F , exactly as in Prop. 6.6). If the process is stationary, the weights in $\psi(L)$ decay at a geometric rate.

6.2.4 PACF approach to identify AR/MA processes

We have seen that the k^{th} -order auto-correlation of a MA(q) process is null if $k > q$. This is exploited, in practice, to determine the order of a MA process. Moreover, since this is not the case for an AR process, this can be used to distinguish an AR from an MA process.

There exists an equivalent approach to determine whether a process can be modeled as an AR process; it is based on partial auto-correlations (see Def. 6.17).

Definition 6.16 (Partial correlation). The partial correlation between X and Y , given Z , is the correlation between:

- a. the residuals of the linear regression of X on Z and
- b. the residuals of the linear regression of Y on Z .

Definition 6.17 (Partial auto-correlation). In a time series context, the partial auto-correlation ($\phi_{h,h}$) of process $\{y_t\}$ is defined as the partial correlation of y_{t+h} and y_t given $y_{t+h-1}, \dots, y_{t+1}$.

If $h > p$, the regression of y_{t+h} on $y_{t+h-1}, \dots, y_{t+1}$ is:

$$y_{t+h} = c + \phi_1 y_{t+h-1} + \dots + \phi_p y_{t+h-p} + \varepsilon_{t+h}.$$

The residuals of the latter regressions (ε_{t+h}) are uncorrelated to y_t . Then the partial autocorrelation is zero for $h > p$.

Besides, it can be shown that $\phi_{p,p} = \phi_p$. Hence $\phi_{p,p} = \phi_p$ but $\phi_{h,h} = 0$ for $h > p$. This can be used to determine the order of a AR process. By contrast (importantly) if y_t follows an MA(q) process, then $\phi_{k,k}$ asymptotically approaches zero instead of cutting off abruptly.

As illustrated below, functions `acf` and `pacf` can be conveniently used to employ the (P)ACF approach. (Note also the use of function `sim.arma` to simulate ARMA processes.)

```
library(AEC)
par(mfrow=c(3,2))
par(plt=c(.2,.9,.2,.95))
theta <- c(1,2,1);phi=0
y.sim <- sim.arma(c=0,phi,theta,sigma=1,T=1000,y.0=0,nb.sim=1)
par(mfg=c(1,1));plot(y.sim,type="l",lwd=2)
par(mfg=c(2,1));acf(y.sim)
par(mfg=c(3,1));pacf(y.sim)
theta <- c(1);phi=0.9
y.sim <- sim.arma(c=0,phi,theta,sigma=1,T=1000,y.0=0,nb.sim=1)
par(mfg=c(1,2));plot(y.sim,type="l",lwd=2)
```

```
par(mfg=c(2,2));acf(y.sim)
par(mfg=c(3,2));pacf(y.sim)
```




Figure 6.7: ACF/PACF analysis of two processes (MA process on the left, AR on the right).

6.2.5 Wold decomposition

The Wold decomposition is an important result in time series analysis:

Theorem 6.3 (Wold decomposition). *Any covariance-stationary process admits the following representation:*

$$y_t = \mu + \sum_0^{+\infty} \theta_i \varepsilon_{t-i} + \kappa_t,$$

where

- $\theta_0 = 1$, $\sum_{i=0}^{\infty} \theta_i^2 < +\infty$ (square summability, see Def. 6.12).
- $\{\varepsilon_t\}$ is a white noise (see Def. 6.1); ε_t is the error made when forecasting y_t based on a linear combination of lagged y_t 's ($\varepsilon_t = y_t - \hat{E}[y_t|y_{t-1}, y_{t-2}, \dots]$).
- For any $j \geq 1$, κ_t is not correlated with ε_{t-j} ; but κ_t can be perfectly forecasted based on a linear combination of lagged y_t 's (i.e. $\kappa_t = \hat{E}[\kappa_t|y_{t-1}, y_{t-2}, \dots]$). κ_t is called the **deterministic component** of y_t .

Proof. See Anderson (1971). Partial proof in L. Christiano. \square

For an ARMA process, the Wold representation is given by Eq. (6.16). As detailed in Prop. 6.8, it can be computed by recursively replacing the lagged y_t 's in Eq. (6.15). In this case, the deterministic component (κ) is null.

6.2.6 Impulse Response Functions (IRFs) in ARMA models

Consider the ARMA(p,q) process defined in Def. 6.15, whose associated sequence of white noise is $\{\varepsilon_t\}$. Let us construct a novel (counterfactual) sequence of shocks $\{\tilde{\varepsilon}_t^{(s)}\}$:

$$\tilde{\varepsilon}_t^{(s)} = \begin{cases} \varepsilon_t & \text{if } t \neq s, \\ \varepsilon_t + \delta & \text{if } t = s. \end{cases}$$

We denote by $\{\tilde{y}_t^{(s)}\}$ the process following Eq. (6.15) where $\{\varepsilon_t\}$ is replaced by $\{\tilde{\varepsilon}_t^{(s)}\}$. The time series $\{\tilde{y}_t^{(s)}\}$ is the (counterfactual) series $\{y_t\}$ that would have prevailed if ε_t had been shifted by δ on date s (only change).

What is the relationship between $\{y_t\}$ and $\{\tilde{y}_t^{(s)}\}$? By definition, we have:

$$\boxed{\tilde{y}_t^{(s)} = y_t + \frac{\partial y_t}{\partial \varepsilon_s} \delta.}$$

The object $\frac{\partial y_t}{\partial \varepsilon_s}$ is called the **dynamic multiplier** of y_t w.r.t. ε_t (of order $t-s$).

Let us consider the **Wold decomposition** of $\{y_t\}$ (see Theorem 6.3 and Eq. (6.16) for the application of this theorem to ARMA processes):

$$y_t = \mu + \sum_{i=0}^{+\infty} \psi_i \varepsilon_{t-i}.$$

For $t < s$, we have $y_t = \tilde{y}_t^{(s)}$ (because $\tilde{\varepsilon}_{t-i} = \varepsilon_{t-i}$ for all $i \geq 0$ if $t < s$).

For $t \geq s$:

$$\tilde{y}_t^{(s)} = \mu + \left(\sum_{i=0}^{t-s-1} \psi_i \varepsilon_{t-i} \right) + \psi_{t-s} (\varepsilon_s + \delta) + \left(\sum_{i=t-s+1}^{+\infty} \psi_i \varepsilon_{t-i} \right) = y_t + \frac{\partial y_t}{\partial \varepsilon_s} \delta.$$

Therefore, for $t \geq s$, we have:

$$\boxed{\frac{\partial y_t}{\partial \varepsilon_s} = \psi_{t-s}.}$$

That is, $\{y_t\}$'s dynamic multiplier of order k is the same object as the k^{th} loading ψ_k in the Wold decomposition of $\{y_t\}$. The sequence $\left\{ \frac{\partial y_{t+h}}{\partial \varepsilon_t} \right\}_{h \geq 0} \equiv \{\psi_h\}_{h \geq 0}$ defines the **impulse response function (IRF)** of y_t to the shock ε_t .

For ARMA processes, the computation of the IRFs is easy:

Proposition 6.8 (IRF of an ARMA(p,q) process). *The coefficients ψ_h , that define the IRF of process y_t to ε_t can be computed recursively as follows:*

1. Set $\psi_{-1} = \dots = \psi_{-p} = 0$.
2. For $h \geq 0$, (recursively) apply Eq. (6.15):

$$\psi_h = \phi_1 \psi_{h-1} + \dots + \phi_p \psi_{h-p} + \theta_h,$$

where $\theta_h = 0$ for $h > q$.

Proof. This is obtained by applying the operator $\frac{\partial}{\partial \varepsilon_t}$ on both sides of Eq. (6.15):

$$y_{t+h} = c + \phi_1 y_{t+h-1} + \dots + \phi_p y_{t+h-p} + \varepsilon_{t+h} + \theta_1 \varepsilon_{t+h-1} + \dots + \theta_q \varepsilon_{t+h-q}.$$

□

Note that Proposition 6.8 constitutes a simple way to compute the MA(∞) representation (or Wold representation) of an ARMA process.

One can use function `sim.arma`, with the argument `make.IRF = 1` to compute ARMA's IRFs:

```
T <- 21 # number of periods for IRF
theta <- c(1,1,1);phi <- c(0);c <- 0
y.sim <- sim.arma(c,phi,theta,sigma=1,T,y.0=rep(0,length(phi)),nb.sim=1,make.IRF = 1)
par(mfrow=c(1,3));par(plt=c(.25,.95,.2,.85))
plot(0:(T-1),y.sim[,1],type="l",lwd=2,
      main="(a) Process 1",xlab="Time after shock on epsilon",
      ylab="Dynamic multiplier (shock on epsilon at t=0)",col="red")
abline(h=0)
theta <- c(1,.5);phi <- c(0.6)
y.sim <- sim.arma(c,phi,theta,sigma=1,T,y.0=rep(0,length(phi)),nb.sim=1,make.IRF = 1)
plot(0:(T-1),y.sim[,1],type="l",lwd=2,
      main="(b) Process 2",xlab="Time after shock on epsilon",ylab="",col="red")
theta <- c(1,1,1);phi <- c(0,0,.5,.4)
y.sim <- sim.arma(c,phi,theta,sigma=1,T,y.0=rep(0,length(phi)),nb.sim=1,make.IRF = 1)
plot(0:(T-1),y.sim[,1],type="l",lwd=2,
      main="(c) Process 3",xlab="Time after shock on epsilon",ylab="",col="red")
```




Figure 6.8: IRFs of three processes. Process 1 (MA(2)): $y_t = \varepsilon_t + \varepsilon_{t-1} + \varepsilon_{t-2}$. Process 2 (ARMA(1,2)): $y_t = 0.6y_{t-1} + \varepsilon_t + 0.5\varepsilon_{t-1}$. Process 3 (ARMA(4,2)): $y_t = 0.5y_{t-3} + 0.4y_{t-4} + \varepsilon_t + \varepsilon_{t-1} + \varepsilon_{t-2}$.

Consider the annual Swiss GDP growth from the JST macro-history database. Let us first determine relevant orders for AR and MA processes using the (P)ACF approach.

```
library(AEC)
data <- subset(JST,iso=="CHE")
par(plt=c(.1,.95,.1,.95))
T <- dim(data)[1]
growth <- log(data$gdp[2:T]/data$gdp[1:(T-1)])
par(mfrow=c(3,1))
par(plt=c(.1,.95,.15,.95))
plot(data$year[2:T],growth,type="l",xlab="",ylab="",lwd=2)
abline(h=0,ity=2)
acf(growth)
pacf(growth)
```

The two bottom plots of Figure 6.9 suggest that either an MA(2) or an AR(1) could be used to model the GDP growth rate series. Figure 6.10 shows the IRFs based on these two respective specifications.

```
# Fit an AR process:
res <- arima(growth,order=c(1,0,0))
```




Figure 6.9: (P)ACF analysis of Swiss GDP growth.

```

phi <- res$coef[1]
T <- 11
y.sim <- sim.arma(c=0,phi,theta=1,sigma=1,T,y.0=rep(0,length(phi)),nb.sim=1,make.IRF = 1)
plot(0:(T-1),y.sim[,1],type="l",lwd=3,
     xlab="Time after shock on epsilon",
     ylab="Dynamic multiplier (shock on epsilon at t=0)",col="red")
# Fit a MA process:
res <- arima(growth,order=c(0,0,2))
phi <- 0
theta <- c(1,res$coef[1:2])
y.sim <- sim.arma(c=0,phi,theta,sigma=1,T,y.0=rep(0,length(phi)),nb.sim=1,make.IRF = 1)
lines(0:(T-1),y.sim[,1],lwd=3,col="red",lty=2)
abline(h=0)

```

The same kind of algorithm can be used to compute the impact of an increase in an exogenous variable x_t within an ARMAX(p,q,r) model (see next section).

6.2.7 ARMA processes with exogenous variables (ARMA-X)

ARMA processes do not allow to investigate the influence of an exogenous variable (say x_t) on the variable of interest (say y_t). When x_t and y_t have reciprocal influences, the Vector Autoregressive (VAR) model may be used (this tools will




Figure 6.10: Dynamic response of Swiss annual growth to a shock on the innovation ε_t at date $t = 0$. The solid line corresponds to an AR(1) specification; the dashed line corresponds to a MA(2) specification.

be studied later, in Section 6.3). However, when one suspects that x_t has an “exogenous” influence on y_t , then a simple extension of the ARMA processes may be considered. Loosely speaking, x_t has an “exogenous” influence on y_t if y_t does not affect x_t . This extension is called ARMAX(p,q,r).

To begin with, let us formalize this notion of exogeneity. Consider a white noise sequence $\{\varepsilon_t\}$ (Def. 6.1).

Definition 6.18 (Exogeneity). We say that x_t is (strictly) exogenous to $\{\varepsilon_t\}$ if

$$\mathbb{E}(\varepsilon_t \mid \underbrace{\dots, x_{t+1}, \dots}_{\text{future}}, \underbrace{x_t, x_{t-1}, \dots}_{\text{present and past}}) = 0.$$

For strictly exogenous processes $\{x_t\}$, past, present and future values of the exogenous variable do not allow to predict any of the ε_t 's.

In the following, we assume that $\{x_t\}$ is a covariance stationary process.

Definition 6.19 (ARMAX(p,q,r) model). The process $\{y_t\}$ is an ARMAX(p,q,r) if it follows a difference equation:

$$y_t = \underbrace{c + \phi_1 y_{t-1} + \dots + \phi_p y_{t-p}}_{\text{AR}(p) \text{ part}} + \underbrace{\beta_0 x_t + \dots + \beta_r x_{t-r}}_{\text{X}(r) \text{ part}} + \underbrace{\varepsilon_t + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q}}_{\text{MA}(q) \text{ part}}. \quad (6.17)$$

where $\{\varepsilon_t\}$ is an i.i.d. white noise sequence and $\{x_t\}$ is exogenous to y_t .

What is the effect of a one-unit increase in x_t on y_t ? To address this question, this notion of “effect” has to be formalized. Let us introduce two related sequences of values for $\{x\}$. Denote the first by $\{a\}^t$ and the second by $\{\tilde{a}\}^t$. Further, we posit $a_s = \tilde{a}_s$ for all $s \neq t$, and $\tilde{a}_t = a_t + 1$.

With these notations, we define $\frac{\partial y_{t+h}}{\partial x_t}$ as follows:

$$\frac{\partial y_{t+h}}{\partial x_t} := \mathbb{E}(y_{t+h} | \{x\} = \{\tilde{a}^t\}) - \mathbb{E}(y_{t+h} | \{x\} = \{a\}^t). \quad (6.18)$$

Under the exogeneity assumption, it is easily seen that

$$\frac{\partial y_t}{\partial x_t} = \beta_0.$$

Now, since

$$\begin{aligned} y_{t+1} &= c + \phi_1 y_t + \cdots + \phi_p y_{t+1-p} + \beta_0 x_{t+1} + \cdots + \beta_r x_{t+1-r} + \\ &\quad \varepsilon_{t+1} + \theta_1 \varepsilon_t + \cdots + \theta_q \varepsilon_{t+1-q}, \end{aligned}$$

and using the exogeneity assumption, we obtain:

$$\frac{\partial y_{t+1}}{\partial x_t} := \phi_1 \frac{\partial y_t}{\partial x_t} + \beta_1 = \phi_1 \beta_0 + \beta_1.$$

This can be applied recursively to give $\frac{\partial y_{t+h}}{\partial x_t}$ for any $h \geq 0$:

Proposition 6.9 (Dynamic multipliers in ARMAX models). *One can recursively compute the dynamic multipliers $\frac{\partial y_{t+h}}{\partial x_t}$ as follows:*

- i. *Initialization:* $\frac{\partial y_{t+h}}{\partial x_t} = 0$ for $h < 0$.
- ii. *For $h \geq 0$ and assuming that the first $h-1$ multipliers have been computed, we have:*

$$\frac{\partial y_{t+h}}{\partial x_t} = \phi_1 \frac{\partial y_{t+h-1}}{\partial x_t} + \cdots + \phi_p \frac{\partial y_{t+h-p}}{\partial x_t} + \beta_h, \quad (6.19)$$

where we use the notation $\beta_h = 0$ if $h > r$.


Remark that the resulting dynamic multipliers are the same as those obtained for an ARMA(p,r) model where the θ_i 's are replaced with β_i 's (see Proposition 6.8 in Section 6.2.7).

It has to be stressed that the definition of the dynamic multipliers (Eq. (6.19)) does not reflect a potential persistency of the shock occurring on date t in process

$\{x\}$ itself. Going in this direction would necessitate to model the dynamics of x_t (and, therefore, to go in the VAR direction , see Section 6.3).

The next example is based on data used in Stock and Watson (2003) (Chapter 16). The objective is to study the influence of the number of freezing days on the price of orange juice.

```
library(AEC)
library(AER)
data("FrozenJuice")
FJ <- as.data.frame(FrozenJuice)
date <- time(FrozenJuice)
price <- FJ$price/FJ$ppi
T <- length(price)
k <- 1
dprice <- 100*log(price[(k+1):T]/price[1:(T-k)])
fdd <- FJ$fdd[(k+1):T]
par(mfrow=c(3,1))
par(plt=c(.1,.95,.15,.75))
plot(date,price,type="l",xlab="",ylab="",main="(a) Price of orange Juice")
plot(date,c(NaN,dprice),type="l",xlab="",ylab="",main="(b) Monthly pct Change (y)")
plot(date,FJ$fdd,type="l",xlab="",ylab="",main="(c) Number of freezing days (x)")
```



```

nb.lags <- 12
FDD <- FJ$fdd[(nb.lags+1):T]
names.FDD <- NULL
for(i in 1:nb.lags){
  FDD <- cbind(FDD,FJ$fdd[(nb.lags+1-i):(T-i)])
  names.FDD <- c(names.FDD,paste(" Lag ",toString(i),sep=""))
}
colnames(FDD) <- c(" Lag 0",names.FDD)
dprice <- dprice[(length(dprice)-dim(FDD)[1]+1):length(dprice)]
eq <- lm(dprice~FDD)
# Compute the Newey-West std errors:
var.cov.mat <- NeweyWest(eq,lag = 7, prewhite = FALSE)
robust_se <- sqrt(diag(var.cov.mat))
# Stargazer output (with and without Robust SE)
stargazer::stargazer(eq, eq, type = "text",
                      column.labels = c("(no HAC)", "(HAC)"),keep.stat = "n",
                      se = list(NULL,robust_se),no.space = TRUE)

##
## =====
##             Dependent variable:
##             -----
##                   dprice
##             (no HAC)      (HAC)
##             (1)          (2)
## -----
## FDD Lag 0    0.496***   0.496***
##             (0.058)     (0.139)
## FDD Lag 1    0.150***   0.150*
##             (0.058)     (0.087)
## FDD Lag 2    0.046      0.046
##             (0.057)     (0.056)
## FDD Lag 3    0.062      0.062
##             (0.057)     (0.046)
## FDD Lag 4    0.024      0.024
##             (0.057)     (0.030)
## FDD Lag 5    0.036      0.036
##             (0.057)     (0.030)
## FDD Lag 6    0.037      0.037
##             (0.057)     (0.046)
## FDD Lag 7    0.019      0.019
##             (0.057)     (0.015)
## FDD Lag 8    -0.038     -0.038
##             (0.057)     (0.034)
## FDD Lag 9    -0.006     -0.006

```

```

##                      (0.057)      (0.050)
## FDD Lag 10        -0.112*     -0.112
##                      (0.057)      (0.069)
## FDD Lag 11        -0.063     -0.063
##                      (0.058)      (0.052)
## FDD Lag 12        -0.140**    -0.140*
##                      (0.058)      (0.078)
## Constant          -0.426*     -0.426*
##                      (0.238)      (0.243)
## -----
## Observations       600        600
## =====
## Note:           *p<0.1; **p<0.05; ***p<0.01

```

Let us now use function `estim.armax`, from package AECto estimate an ARMA-X(2,0,1) model:

```

nb.lags <- 1
FDD <- FJ$fd[(nb.lags+1):T]
names.FDD <- NULL
for(i in 1:nb.lags){
  FDD <- cbind(FDD,FJ$fd[(nb.lags+1-i):(T-i)])
  names.FDD <- c(names.FDD,paste(" Lag ",toString(i),sep=""))
}
colnames(FDD) <- c(" Lag 0",names.FDD)
dprice <- 100*log(price[(k+1):T]/price[1:(T-k)])
dprice <- dprice[(length(dprice)-dim(FDD)[1]+1):length(dprice)]
res.armax <- estim.armax(Y = dprice,p=3,q=0,X=FDD)

## [1] =====
## [1] " ESTIMATING"
## [1] =====
## [1] " END OF ESTIMATION"
## [1] =====
## [1] ""
## [1] " RESULTS:"
## [1] -----
##             THETA      st.dev   t.ratio
## c          -0.46556249 0.19554352 -2.380864
## phi      t-1  0.09788977 0.04025907  2.431496
## phi      t-2  0.05049849 0.03827488  1.319364
## phi      t-3  0.07155170 0.03764750  1.900570
## sigma     4.64917949 0.13300769 34.954215
## beta     t-0  0.47015552 0.05665344  8.298800
## beta     t-1  0.10015862 0.05972526  1.676989
## [1] =====

```

Figure 6.11 shows the IRF associated with each of the two models.

```
nb.periods <- 20
IRF1 <- sim.arma(c=0,phi=c(0),theta=eq$coefficients[2:13],sigma=1,
                   T=nb.periods,y.0=c(0),nb.sim=1,make.IRF=1)
IRF2 <- sim.arma(c=0,phi=res.armax$phi,theta=res.armax$beta,sigma=1,
                   T=nb.periods,y.0=rep(0,length(res.armax$phi)),nb.sim=1,make.IRF=1)
plot(IRF1,type="l",lwd=2,col="red",xlab="months after shock",ylab="Chge in price (percent")
lines(IRF2,lwd=2,col="red",lty=2)
abline(h=0,col="grey")
```




Figure 6.11: Response of changes in orange juice price (in percent) to the number of freezing days. The solid (respectively dashed) line corresponds to the ARMAX(0,0,12) (resp. ARMAX(3,0,1)) model. The first model is estimated by OLS, the second by MLE.

In the following example, we make use of monetary shocks identified through high-frequency data (see Gertler and Karadi (2015)). This dataset comes from Valerie Ramey's website (see Ramey (2016)).

```
library(AEC)
T <- dim(Ramey)[1]
# Construct growth series:
Ramey$growth <- Ramey$LIP - c(rep(NaN,12),Ramey$LIP[1:(length(Ramey$LIP)-12)])
# Prepare matrix of exogenous variables:
vec.lags <- c(9,12,18)
Matrix.of.Exog <- NULL
shocks <- Ramey$ED2_TC
```

```

for(i in 1:length(vec.lags)){
  Matrix.of.Exog <- cbind(Matrix.of.Exog,
                           c(rep(NaN,vec.lags[i]),shocks[1:(T-vec.lags[i])]))
}
# Look for dates where data are available:
indic.good.dates <- complete.cases(Matrix.of.Exog)

```




Figure 6.12: The blue line corresponds to monetary-policy shocks identified by means of the Gertler and Karadi (2015)'s approach (high-frequency change in Euro-dollar futures). The black solid line is the year-on-year growth rate of industrial production.

```

# Estimate ARMAX:
p <- 1; q <- 0
x <- estim.armax(Ramey$growth[indic.good.dates],p,q,
                   X=Matrix.of.Exog[indic.good.dates,])

## [1] "===="
## [1] "  ESTIMATING"
## [1] "===="
## [1] "  END OF ESTIMATION"
## [1] "===="
## [1] ""
## [1] "  RESULTS:"
## [1] "  -----"
## [1] "          THETA      st.dev    t.ratio"
## [1] "c      -0.0001716198 0.0005845907 -0.2935726"

```

```

## phi   t-1  0.9825608412 0.0120458531 81.5683897
## sigma        0.0087948724 0.0003211748 27.3834438
## beta  t-0 -0.0193570616 0.0087331529 -2.2165032
## beta  t-1 -0.0225707935 0.0086750938 -2.6017925
## beta  t-2 -0.0070131593 0.0086387440 -0.8118263
## [1] "====="

# Compute IRF:
irf <- sim.arma(0,x$phi,x$beta,x$sigma,T=60,y.0=rep(0,length(x$phi)),nb.sim=1,make.IRF=
                 X=NaN,beta=NaN)

```

Figure 6.13 displays the resulting IRF, with a 95% confidence band. The code used to produce the confidence bands (i.e., to compute the standard deviation of the dynamic multipliers for the different horizons) is based on the Delta method (see Eq. (4.15)). The codes are available in Appendix 7.6.2.




Figure 6.13: Response of industrial-production growth to monetary-policy shocks. Dashed lines correspond to the ± 2 -standard-deviation bands.

6.2.8 Maximum Likelihood Estimation of ARMA processes

Consider the general case (of any time series); assume we observe a sample $\mathbf{y} = [y_1, \dots, y_T]'$. In order to implement ML techniques (see Section 4.2), we need to evaluate the joint p.d.f. (or “likelihood”) of \mathbf{y} , i.e., $\mathcal{L}(\theta; \mathbf{y})$, where θ is a vector of parameters that characterizes the dynamics of y_t . The Maximum

Likelihood (ML) estimate of θ is then given by:

$$\boxed{\theta_{MLE} = \arg \max_{\theta} \mathcal{L}(\theta; \mathbf{y}) = \arg \max_{\theta} \log \mathcal{L}(\theta; \mathbf{y}).}$$

In the time series context, if process y_t is Markovian, then there exists a useful way to rewrite the likelihood $\mathcal{L}(\theta; \mathbf{y})$. Let us first recall the definition of a Markovian process (see also Def. 4.9):

Definition 6.20 (Markovian process). Process y_t is Markovian of order one if $f_{Y_t|Y_{t-1}, Y_{t-2}, \dots} = f_{Y_t|Y_{t-1}}$. More generally, it is Markovian of order k if $f_{Y_t|Y_{t-1}, Y_{t-2}, \dots} = f_{Y_t|Y_{t-1}, \dots, Y_{t-k}}$.

Now, remember Bayes' formula:

$$\mathbb{P}(X_2 = x, X_1 = y) = \mathbb{P}(X_2 = x|X_1 = y)\mathbb{P}(X_1 = y).$$

Using it leads to the following decomposition of our likelihood function:

$$\begin{aligned} f_{Y_T, \dots, Y_1}(y_T, \dots, y_1; \theta) &= f_{Y_T|Y_{T-1}, \dots, Y_1}(y_T, \dots, y_1; \theta) \times \\ &\quad f_{Y_{T-1}, \dots, Y_1}(y_{T-1}, \dots, y_1; \theta). \end{aligned}$$

Using the previous expression recursively, one obtains:

$$f_{Y_T, \dots, Y_1}(y_T, \dots, y_1; \theta) = f_{Y_1}(y_1; \theta) \prod_{t=2}^T f_{Y_t|Y_{t-1}, \dots, Y_1}(y_t, \dots, y_1; \theta). \quad (6.20)$$

Let us start with the Gaussian AR(1) process (which is Markovian of order one):

$$y_t = c + \phi_1 y_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim i.i.d. \mathcal{N}(0, \sigma^2).$$

For $t > 1$:

$$f_{Y_t|Y_{t-1}, \dots, Y_1}(y_t, \dots, y_1; \theta) = f_{Y_t|Y_{t-1}}(y_t, y_{t-1}; \theta)$$

and

$$f_{Y_t|Y_{t-1}}(y_t, y_{t-1}; \theta) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y_t - c - \phi_1 y_{t-1})^2}{2\sigma^2}\right).$$

These expressions can be plugged into Eq. (6.20). But what about $f_{Y_1}(y_1; \theta)$? There exist two possibilities:

1. **Case 1:** We use the marginal distribution: $y_1 \sim \mathcal{N}\left(\frac{c}{1-\phi_1}, \frac{\sigma^2}{1-\phi_1^2}\right)$.
2. **Case 2:** y_1 is considered to be deterministic. In a way, that means that the first observation is “sacrificed”.

In the Gaussian AR(1) process, we have:

1. **Case 1:** The (exact) log-likelihood is:

$$\begin{aligned}\log \mathcal{L}(\theta; \mathbf{y}) &= -\frac{T}{2} \log(2\pi) - T \log(\sigma) + \frac{1}{2} \log(1 - \phi_1^2) \\ &\quad - \frac{(y_1 - c/(1 - \phi_1))^2}{2\sigma^2/(1 - \phi_1^2)}\end{aligned}\tag{6.21}$$

$$-\sum_{t=2}^T \left[\frac{(y_t - c - \phi_1 y_{t-1})^2}{2\sigma^2} \right].\tag{6.22}$$

The Maximum Likelihood Estimator of $\theta = [c, \phi_1, \sigma^2]$ is obtained by numerical optimization.

2. **Case 2:** The (conditional) log-likelihood is:

$$\begin{aligned}\log \mathcal{L}^*(\theta; \mathbf{y}) &= -\frac{T-1}{2} \log(2\pi) - (T-1) \log(\sigma) \\ &\quad - \sum_{t=2}^T \left[\frac{(y_t - c - \phi_1 y_{t-1})^2}{2\sigma^2} \right].\end{aligned}\tag{6.23}$$

Exact MLE and conditional MLE have the same asymptotic (i.e. large-sample) distribution. Indeed, when the process is stationary, $f_{Y_1}(y_1; \theta)$ makes a relatively negligible contribution to $\log \mathcal{L}(\theta; \mathbf{y})$.

The conditional MLE has a substantial advantage: in the Gaussian case, the conditional MLE is simply obtained by OLS. Indeed, let us introduce the notations:

$$Y = \begin{bmatrix} y_2 \\ \vdots \\ y_T \end{bmatrix} \quad \text{and} \quad X = \begin{bmatrix} 1 & y_1 \\ \vdots & \vdots \\ 1 & y_{T-1} \end{bmatrix}.$$

Eq. (6.23) then rewrites:

$$\begin{aligned}\log \mathcal{L}^*(\theta; \mathbf{y}) &= -\frac{T-1}{2} \log(2\pi) - (T-1) \log(\sigma) \\ &\quad - \frac{1}{2\sigma^2} (Y - X[c, \phi_1]')' (Y - X[c, \phi_1]'),\end{aligned}\tag{6.24}$$

which is maximised for:

$$[\hat{c}, \hat{\phi}_1]' = (X'X)^{-1}X'Y\tag{6.25}$$

$$\begin{aligned}\hat{\sigma}^2 &= \frac{1}{T-1} \sum_{t=2}^T (y_t - \hat{c} - \hat{\phi}_1 y_{t-1})^2 \\ &= \frac{1}{T-1} Y' (I - X(X'X)^{-1}X') Y.\end{aligned}\tag{6.26}$$

Let us turn to the case of an AR(p) process. We use:

$$\begin{aligned}\log \mathcal{L}(\theta; \mathbf{y}) &= \log f_{Y_p, \dots, Y_1}(y_p, \dots, y_1; \theta) + \\ &\quad \underbrace{\sum_{t=p+1}^T \log f_{Y_t|Y_{t-1}, \dots, Y_{t-p}}(y_t, \dots, y_{t-p}; \theta)}_{\log \mathcal{L}^*(\theta; \mathbf{y})}.\end{aligned}$$

where $f_{Y_p, \dots, Y_1}(y_p, \dots, y_1; \theta)$ is the marginal distribution of $\mathbf{y}_{1:p} := [y_p, \dots, y_1]'$. The marginal distribution of $\mathbf{y}_{1:p}$ is Gaussian; it is therefore fully characterised by its mean and covariance matrix:

$$\begin{aligned}\mathbb{E}(\mathbf{y}_{1:p}) &= \frac{c}{1 - \phi_1 - \dots - \phi_p} \mathbf{1}_{p \times 1} \\ \mathbb{V}ar(\mathbf{y}_{1:p}) &= \begin{bmatrix} \gamma_0 & \gamma_1 & \dots & \gamma_{p-1} \\ \gamma_1 & \gamma_0 & \dots & \gamma_{p-2} \\ \vdots & & \ddots & \vdots \\ \gamma_{p-1} & \gamma_{p-2} & \dots & \gamma_0 \end{bmatrix},\end{aligned}$$

where the γ_i 's are computed using the Yule-Walker equations (Eq. (6.14)). Note that they depend, in a non-linear way, on the model parameters. Hence, the maximization of the exact log-likelihood necessitates numerical optimization procedures. By contrast, the maximization of the conditional log-likelihood $\log \mathcal{L}^*(\theta; \mathbf{y})$ only requires OLS, using Eqs. (6.25) and (6.26) with:

$$Y = \begin{bmatrix} y_{p+1} \\ \vdots \\ y_T \end{bmatrix} \quad \text{and} \quad X = \begin{bmatrix} 1 & y_p & \dots & y_1 \\ \vdots & \vdots & & \vdots \\ 1 & y_{T-1} & \dots & y_{T-p} \end{bmatrix}.$$

Again, for stationary processes, conditional and exact MLE have the same asymptotic (large-sample) distribution.

Consider the regression (where y_t follows an AR(p) process):

$$y_t = \beta' \mathbf{x}_t + \varepsilon_t, \tag{6.27}$$

with $\mathbf{x}_t = [1, y_{t-1}, \dots, y_{t-p}]'$ and $\beta = [c, \phi_1, \dots, \phi_p]'$.

The fact that \mathbf{x}_t correlates to the ε_s 's for $s < t$ implies that the OLS estimator \mathbf{b} of β is biased in small sample. Indeed, we have:

$$\mathbf{b} = \beta + (X'X)^{-1}X'\varepsilon, \tag{6.28}$$

and because of the specific form of X , we have non-zero correlation between \mathbf{x}_t and ε_s for $s < t$, therefore $\mathbb{E}[(X'X)^{-1}X'\varepsilon] \neq 0$. However, asymptotically, the previous expectation goes to zero, and we have:

Proposition 6.10 (Large-sample properties of the OLS estimator of AR(p) models). Assume $\{y_t\}$ follows the AR(p) process:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} + \varepsilon_t$$

where $\{\varepsilon_t\}$ is an i.i.d. white noise process. If \mathbf{b} is the OLS estimator of β (Eq. (6.27)), we have:

$$\sqrt{T}(\mathbf{b} - \beta) = \underbrace{\left[\frac{1}{T} \sum_{t=p}^T \mathbf{x}_t \mathbf{x}'_t \right]^{-1}}_{\xrightarrow{p} \mathbf{Q}^{-1}} \underbrace{\sqrt{T} \left[\frac{1}{T} \sum_{t=1}^T \mathbf{x}_t \varepsilon_t \right]}_{\xrightarrow{d} \mathcal{N}(0, \sigma^2 \mathbf{Q})},$$

where $\mathbf{Q} = \text{plim } \frac{1}{T} \sum_{t=p}^T \mathbf{x}_t \mathbf{x}'_t = \text{plim } \frac{1}{T} \sum_{t=1}^T \mathbf{x}_t \mathbf{x}'_t$ is given by:

$$\mathbf{Q} = \begin{bmatrix} 1 & \mu & \mu & \dots & \mu \\ \mu & \gamma_0 + \mu^2 & \gamma_1 + \mu^2 & \dots & \gamma_{p-1} + \mu^2 \\ \mu & \gamma_1 + \mu^2 & \gamma_0 + \mu^2 & \dots & \gamma_{p-2} + \mu^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \mu & \gamma_{p-1} + \mu^2 & \gamma_{p-2} + \mu^2 & \dots & \gamma_0 + \mu^2 \end{bmatrix}. \quad (6.29)$$

Proof. Rearranging Eq. (6.46), we have:

$$\sqrt{T}(\mathbf{b} - \beta) = (X' X / T)^{-1} \sqrt{T}(X' \varepsilon / T).$$

Let us consider the autocovariances of $\mathbf{v}_t = \mathbf{x}_t \varepsilon_t$, denoted by γ_j^v . Using the fact that \mathbf{x}_t is a linear combination of past ε_t 's and that ε_t is a white noise, we get that $\mathbb{E}(\varepsilon_t \mathbf{x}_t) = 0$. Therefore

$$\gamma_j^v = \mathbb{E}(\varepsilon_t \varepsilon_{t-j} \mathbf{x}_t \mathbf{x}'_{t-j}).$$

If $j > 0$, we have $\mathbb{E}(\varepsilon_t \varepsilon_{t-j} \mathbf{x}_t \mathbf{x}'_{t-j}) = \mathbb{E}(\mathbb{E}[\varepsilon_t \varepsilon_{t-j} \mathbf{x}_t \mathbf{x}'_{t-j} | \varepsilon_{t-j}, \mathbf{x}_t, \mathbf{x}_{t-j}]) = \mathbb{E}(\varepsilon_{t-j} \mathbf{x}_t \mathbf{x}'_{t-j} \mathbb{E}[\varepsilon_t | \varepsilon_{t-j}, \mathbf{x}_t, \mathbf{x}_{t-j}]) = 0$. Note that, for $j > 0$, we have $\mathbb{E}[\varepsilon_t | \varepsilon_{t-j}, \mathbf{x}_t, \mathbf{x}_{t-j}] = 0$ because $\{\varepsilon_t\}$ is an i.i.d. white noise sequence. If $j = 0$, we have:

$$\gamma_0^v = \mathbb{E}(\varepsilon_t^2 \mathbf{x}_t \mathbf{x}'_t) = \mathbb{E}(\varepsilon_t^2) \mathbb{E}(\mathbf{x}_t \mathbf{x}'_t) = \sigma^2 \mathbf{Q}.$$

The convergence in distribution of $\sqrt{T}(X' \varepsilon / T) = \sqrt{T} \frac{1}{T} \sum_{t=1}^T v_t$ results from Theorem 6.1 (applied on $\mathbf{v}_t = \mathbf{x}_t \varepsilon_t$), using the γ_j^v computed above. \square

These two cases (exact or conditional log-likelihoods) can be implemented when asking R to fit an AR process by means of function `arima`. Let us for instance use the output gap of the `US3var` dataset (US quarterly data, from 1959:2 to 2015:1, used in Gouriéroux et al. (2017)).

```

library(AEC)
y <- US3var$y.gdp.gap
ar3.Case1 <- arima(y,order = c(3,0,0),method="ML")
ar3.Case2 <- arima(y,order = c(3,0,0),method="CSS")
rbind(ar3.Case1$coef,ar3.Case2$coef)

##           ar1           ar2           ar3   intercept
## [1,] 1.191267 -0.08934705 -0.1781163 -0.9226007
## [2,] 1.192003 -0.08811150 -0.1787662 -1.0341696

```

The two sets of estimated coefficients appear to be very close to each other.

Let us now turn to Moving-Average processes. Start with the MA(1):

$$y_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1}, \quad \varepsilon_t \sim i.i.d.\mathcal{N}(0, \sigma^2).$$

The ε_t 's are easily computed recursively, starting with $\varepsilon_t = y_t - \mu - \theta_1 \varepsilon_{t-1}$. We obtain:

$$\varepsilon_t = y_t - \theta_1 y_{t-1} + \theta_1^2 y_{t-2}^2 + \cdots + (-1)^{t-1} \theta_1^{t-1} y_1 + (-1)^t \theta_1^t \varepsilon_0.$$

Assume that one wants to recover the sequence of $\{\varepsilon_t\}$'s based on observed values of y_t (from date 1 to date t). One can use the previous expression, but what value should be used for ε_0 ? If one does not use the true value of ε_0 but 0 (say), one does not obtain ε_t , but only an estimate of it ($\hat{\varepsilon}_t$, say), with:

$$\hat{\varepsilon}_t = \varepsilon_t - (-1)^t \theta_1^t \varepsilon_0.$$

Clearly, if $|\theta_1| < 1$, then the error becomes small for large t . Formally, when $|\theta_1| < 1$, we have:

$$\hat{\varepsilon}_t \xrightarrow{P} \varepsilon_t.$$

Hence, when $|\theta_1| < 1$, a consistent estimate of the conditional log-likelihood is given by:

$$\log \hat{\mathcal{L}}^*(\theta; \mathbf{y}) = -\frac{T}{2} \log(2\pi) - \frac{T}{2} \log(\sigma^2) - \sum_{t=1}^T \frac{\hat{\varepsilon}_t^2}{2\sigma^2}. \quad (6.30)$$

Loosely speaking, if $|\theta_1| < 1$ and if T is sufficiently large:

approximate conditional MLE \approx exact MLE.

Note that $\hat{\mathcal{L}}^*(\theta; \mathbf{y})$ is a complicated nonlinear function of μ and θ . Its maximization therefore has to be based on numerical optimization procedures.

Let us not consider the case of a Gaussian MA(q) process:

$$y_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}, \quad \varepsilon_t \sim i.i.d.\mathcal{N}(0, \sigma^2). \quad (6.31)$$

Let us assume that this process is an **invertible MA process**. That is, assume that the roots of:

$$\lambda^q + \theta_1\lambda^{q-1} + \cdots + \theta_{q-1}\lambda + \theta_q = 0 \quad (6.32)$$

lie strictly inside of the unit circle. In this case, the polynomial form $\Theta(L) = 1 + \theta_1L + \cdots + \theta_qL^q$ is *invertible* and Eq. (6.31) writes:

$$\varepsilon_t = \Theta(L)^{-1}(y_t - \mu),$$

which implies that, if we knew all past values of y_t , we would also know ε_t . In this case, we can consistently estimate the ε_t 's by recursively computing the $\hat{\varepsilon}_t$'s as follows (for $t > 0$):

$$\hat{\varepsilon}_t = y_t - \mu - \theta_1\hat{\varepsilon}_{t-1} - \cdots - \theta_q\hat{\varepsilon}_{t-q},$$

with

$$\hat{\varepsilon}_0 = \cdots = \hat{\varepsilon}_{-q+1} = 0. \quad (6.33)$$

In this context, a consistent estimate of the conditional log-likelihood is still given by Eq. (6.30).

Note that we could determine the exact likelihood of an MA process. Indeed, vector $Y = [y_1, \dots, y_T]'$ is a Gaussian-distributed vector of mean $\mu = [\mu, \dots, \mu]'$ and of variance:

$$\Omega = \begin{bmatrix} \gamma_0 & \gamma_1 & \cdots & \gamma_q & \color{red}{0} & \cdots & \color{red}{0} \\ \gamma_1 & \gamma_0 & \gamma_1 & & \ddots & \ddots & \vdots \\ \vdots & \gamma_1 & \ddots & \ddots & & \ddots & \color{red}{0} \\ \gamma_q & & \ddots & & & & \gamma_q \\ \color{red}{0} & & & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & & \gamma_1 & \gamma_0 & \gamma_1 \\ \color{red}{0} & \cdots & \color{red}{0} & \gamma_q & \cdots & \gamma_1 & \gamma_0 \end{bmatrix},$$

where the γ_j 's are given by Eq. (6.7). However, for large samples, the computation of the likelihood can be numerically demanding. Hence, the p.d.f. of Y is given by (see Prop. 7.18):

$$(2\pi)^{-T/2}|\Omega|^{-1/2} \exp\left(-\frac{1}{2}(Y - \mu)' \Omega^{-1} (Y - \mu)\right).$$

Finally, let us consider the MLE of an ARMA(p, q) processes:

$$y_t = c + \phi_1y_{t-1} + \cdots + \phi_py_{t-p} + \varepsilon_t + \theta_1\varepsilon_{t-1} + \cdots + \theta_q\varepsilon_{t-q}, \quad \varepsilon_t \sim i.i.d. \mathcal{N}(0, \sigma^2).$$

If the MA part of this process is invertible, the log-likelihood function can be consistently approximated by its conditional counterpart (of the form of Eq. (6.30)), using consistent estimates $\hat{\varepsilon}_t$ of the ε_t . The $\hat{\varepsilon}_t$'s are computed recursively as:

$$\hat{\varepsilon}_t = y_t - c - \phi_1y_{t-1} - \cdots - \phi_py_{t-p} - \theta_1\hat{\varepsilon}_{t-1} - \cdots - \theta_q\hat{\varepsilon}_{t-q}, \quad (6.34)$$

given some initial conditions, for instance:

- a. $\hat{\varepsilon}_0 = \dots = \hat{\varepsilon}_{-q+1} = 0$ and $y_0 = \dots = y_{-p+1} = \mathbb{E}(y_i) = \mu$. (Recursions in Eq. (6.34) then start for $t = 1$.)
- b. $\hat{\varepsilon}_p = \dots = \hat{\varepsilon}_{p-q+1} = 0$ and actual values of the y_i 's for $i \in [1, p]$. In that case, the first p observations of y_t will not be used. Recursions in Eq. (6.34) then start for $t = p + 1$.

6.2.9 Specification choice

The previous section explains how to fit a given ARMA specification. But how to choose an appropriate specification? A possibility is to employ the (P)ACF approach (see Figure 6.7). However, the previous approach leads to either an AR or a MA process (but not an ARMA process). If one wants to consider various ARMA(p, q) specifications, for $p \in \{1, \dots, P\}$ and $q \in \{1, \dots, Q\}$, say, then one can resort to **information criteria**.

In general, when choosing a specification, one faces the following dilemma:

- a. Too rich a specification may lead to “overfitting”/misspecification, implying additional estimation errors (in out-of-sample forecasts).
- b. Too simple a specification may lead to potential omission of valuable information (e.g., contained in older lags).

The lag selection approach based on the so-called **information criteria** consists in maximizing the fit of the data, but adding a penalty for the “richness” of the model. More precisely, using this approach amounts to minimizing a loss function that (a) negatively depends on the fitting errors and (b) positively depends on the number of parameters in the model.

Definition 6.21 (Information Criteria). The Akaike (AIC), Hannan-Quinn (HQ) and Schwarz information (BIC) criteria are of the form

$$c^{(i)}(k) = \underbrace{\frac{-2 \log \mathcal{L}(\hat{\theta}_T(k); \mathbf{y})}{T}}_{\text{decreases w.r.t. } k} + \underbrace{\frac{k \phi^{(i)}(T)}{T}}_{\text{increases w.r.t. } k}$$

with $(i) \in \{AIC, HQ, BIC\}$ and where $\hat{\theta}_T(k)$ denotes the ML estimate of $\theta_0(k)$, which is a vector of parameters of length k .

Criterion (i)		$\phi^{(i)}(T)$
Akaike	AIC	2
Hannan-Quinn	HQ	$2 \log(\log(T))$
Schwarz	BIC	$\log(T)$

The lag suggested by criterion (i) is then given by:

$$\hat{k}^{(i)} = \underset{k}{\operatorname{argmin}} \quad c^{(i)}(k).$$

In the case of an ARMA(p,q) process, $k = 2 + p + q$

Proposition 6.11 (Consistency of the criteria-based lag selection). *The lag selection procedure is consistent (see Def. 7.8) if*

$$\lim_{T \rightarrow \infty} \phi(T) = \infty \quad \text{and} \quad \lim_{T \rightarrow \infty} \phi(T)/T = 0.$$

This is notably the case of the HQ and the BIC criteria.

Proof. The true number of lags is denoted by k_0 . We will show that $\lim_{T \rightarrow \infty} \mathbb{P}(\hat{k}_T \neq k_0) = 0$.

- Case $k < k_0$: The model with k parameter is misspecified, therefore:

$$\text{plim}_{T \rightarrow \infty} \log \mathcal{L}(\hat{\theta}_T(k); \mathbf{y})/T < \text{plim}_{T \rightarrow \infty} \log \mathcal{L}(\hat{\theta}_T(k_0); \mathbf{y})/T.$$

Hence, if $\lim_{T \rightarrow \infty} \phi(T)/T = 0$, we have: $\lim_{T \rightarrow \infty} \mathbb{P}(c(k_0) \geq c(k)) \rightarrow 0$ and

$$\lim_{T \rightarrow \infty} \mathbb{P}(\hat{k} < k_0) \leq \lim_{T \rightarrow \infty} \mathbb{P}\{c(k_0) \geq c(k) \text{ for some } k < k_0\} = 0.$$

- Case $k > k_0$: under the null hypothesis, the likelihood ratio (LR) test statistic (see Def. 4.8) satisfies:

$$2 \left(\log \mathcal{L}(\hat{\theta}_T(k); \mathbf{y}) - \log \mathcal{L}(\hat{\theta}_T(k_0); \mathbf{y}) \right) \sim \chi^2(k - k_0).$$

If $\lim_{T \rightarrow \infty} \phi(T) = \infty$, we have: $\text{plim}_{T \rightarrow \infty} -2 \left(\log \mathcal{L}(\hat{\theta}_T(k); \mathbf{y}) - \log \mathcal{L}(\hat{\theta}_T(k_0); \mathbf{y}) \right) / \phi(T) = 0$. Hence $\text{plim}_{T \rightarrow \infty} T[c(k_0) - c(k)]/\phi(T) \leq -1$ and $\lim_{T \rightarrow \infty} \mathbb{P}(c(k_0) \geq c(k)) \rightarrow 0$, which implies, in the same spirit as before, that $\lim_{T \rightarrow \infty} \mathbb{P}(\hat{k} > k_0) = 0$.

Therefore, $\lim_{T \rightarrow \infty} \mathbb{P}(\hat{k} = k_0) = 1$. □

Example 6.3 (Linear regression). Consider a linear regression with normal disturbances:

$$y_t = \mathbf{x}'_t \beta + \varepsilon_t, \quad \varepsilon_t \sim i.i.d. \mathcal{N}(0, \sigma^2).$$

The associated log-likelihood is of the form of Eq. (6.30). In that case, we have:

$$\begin{aligned} c^{(i)}(k) &= \frac{-2 \log \mathcal{L}(\hat{\theta}_T(k); \mathbf{y})}{T} + \frac{k \phi^{(i)}(T)}{T} \\ &\approx \log(2\pi) + \log(\widehat{\sigma^2}) + \frac{1}{T} \sum_{t=1}^T \frac{\varepsilon_t^2}{\widehat{\sigma^2}} + \frac{k \phi^{(i)}(T)}{T}. \end{aligned}$$

For a large T , for all consistent estimation scheme, we have:

$$\widehat{\sigma^2} \approx \frac{1}{T} \sum_{t=1}^T \varepsilon_t^2 = SSR/T.$$

$$\text{Hence } \hat{k}^{(i)} \approx \underset{k}{\operatorname{argmin}} \log(SSR/T) + \frac{k \phi^{(i)}(T)}{T}.$$

Example 6.4 (Swiss GDP growth). Consider a long historical time series of the Swiss GDP growth (see Figure 6.3), taken from the Jordà et al. (2017) dataset. Let us look for the best ARMA specification using the AIC criteria:

```
library(AEC)
data <- subset(JST,iso=="CHE")
T <- dim(data)[1]
y <- c(NaN,log(data$gdp[2:T]/data$gdp[1:(T-1)]))
# Use AIC criteria to look for appropriate specif:
max.p <- 3;max.q <- 3;
all.AIC <- NULL
for(p in 0:max.p){
  for(q in 0:max.q){
    res <- arima(y,order=c(p,0,q))
    if(res$aic<min(all.AIC)){best.p<-p;best.q<-q}
    all.AIC <- c(all.AIC,res$aic)
  }
}
print(c(best.p,best.q))

## [1] 1 0
```

The best specification is an AR(1) model.

6.3 Multivariate models

This section presents Vector Auto-Regressive Moving-Average (SVARMA) models. These models are widely used in macroeconomic analysis. While simple and easy to estimate, they make it possible to conveniently capture the dynamics of complex multivariate systems. VAR popularity is notably due to Sims (1980)'s influential work. A nice survey if proposed by Stock and Watson (2016).

In economics, VAR models are often employed in order to identify *structural* shocks, that are independent primitive exogenous forces that drive economic variables (Ramey (2016)). They are often given a specific economic meaning (e.g., demand and supply shocks).

Working with these models (VAR and VARMA models) often is often based on two steps: in a first step, the **reduced-form** version of the model is estimated; in a second step, structural shocks are identified and IRFs are produced.

Kilian (1998) See this page

Sign restrictions: package, Danne (2015).

Pfaff (2008)

Korobilis (2013): R package to estimate Bayesian VAR models.

6.3.1 Definition of VARs (and SVARMA) models

Definition 6.22 ((S)VAR model). Let y_t denote a $n \times 1$ vector of random variables. Process y_t follows a p^{th} -order VAR if, for all t , we have

$$\begin{aligned} VAR: \quad y_t &= c + \Phi_1 y_{t-1} + \cdots + \Phi_p y_{t-p} + \varepsilon_t, \\ SVAR: \quad y_t &= c + \Phi_1 y_{t-1} + \cdots + \Phi_p y_{t-p} + B\eta_t, \end{aligned} \quad (6.35)$$

with $\varepsilon_t = B\eta_t$. We assume that $\{\eta_t\}$ is a white noise sequence whose components are mutually and serially independent.

The first line of Eq. (6.35) corresponds to the **reduced-form** of the VAR model (**structural form** for the second line).

While the **structural shocks** (the components of η_t) are mutually uncorrelated, this is not the case of the **innovations**, that are the components of ε_t . However, in boths cases, vectors η_t and ε_t are serially correlated (through time).

As was the case for univariate models, VARs can be extended with MA terms in η_t :

Definition 6.23 ((S)VARMA model). Let y_t denote a $n \times 1$ vector of random variables. Process y_t follows a VARMA model of order (p,q) if, for all t , we have

$$\begin{aligned} VARMA: \quad y_t &= c + \Phi_1 y_{t-1} + \cdots + \Phi_p y_{t-p} + \varepsilon_t + \Theta_1 \varepsilon_{t-1} + \cdots + \Theta_q, \\ SVARMA: \quad y_t &= c + \Phi_1 y_{t-1} + \cdots + \Phi_p y_{t-p} + B_0 \eta_t + B_1 \eta_{t-1} + \cdots + B_q \eta_{t-q}, \end{aligned} \quad (6.36)$$

with $\varepsilon_t = B_0 \eta_t$ (and $B_j = \Theta_j B_0$, for $j \geq 0$). We assume that $\{\eta_t\}$ is a white noise sequence whose components are are mutually and serially independent.

6.3.2 IRFs in SVARMA

One of the main objectives of macro-econometrics is to derive IRFs, that represent the dynamic effects of structural shocks (components of η_t) though the system of variables y_t .

Formally, an IRF is a difference in conditional expectations:

$$\boxed{\Psi_{i,j,h} = \mathbb{E}(y_{i,t+h} | \eta_{j,t} = 1) - \mathbb{E}(y_{i,t+h})}$$

(effect on $y_{i,t+h}$ of a one-unit shock on $\eta_{j,t}$).

If the dynamics of process y_t can be described as a VARMA model, and if y_t is covariance stationary (see Def. 6.4), then y_t admits the following infinite MA representation (MA(∞)):

$$y_t = \mu + \sum_{h=0}^{\infty} \Psi_h \eta_{t-h}. \quad (6.37)$$

This is also the Wold decomposition of process $\{y_t\}$ (see Theorem 6.3).

Estimating IRFs amounts to estimating the Ψ_h 's. In general, there exist three main approaches for that:

- Calibrate and solve a (purely structural) Dynamic Stochastic General Equilibrium (DSGE) model at the first order (linearization). The solution takes the form of Eq. (6.37).
- Directly estimate the Ψ_h based on **projection approaches** (see Section ??).
- Approximate the infinite MA representation by estimating a parsimonious type of model, e.g. **VAR(MA) models** (see Section 6.3.4). Once a (Structural) VARMA representation is obtained, Eq. (6.37) is easily deduced. For that, one can use the same recursive algorithm as for univariate processes (see Prop. 6.8).

Typically, consider the AR(2) case. The first steps of the algorithm are as follows:

$$\begin{aligned}
 y_t &= \Phi_1 \underbrace{y_{t-1}}_{=\Psi_0} + \Phi_2 y_{t-2} + B \eta_t \\
 &= \Phi_1 (\underbrace{\Phi_1 y_{t-2} + \Phi_2 y_{t-3} + B \eta_{t-1}}_{=\Psi_1}) + \Phi_2 y_{t-2} + B \eta_t \\
 &= B \eta_t + \Phi_1 B \eta_{t-1} + (\Phi_2 + \Phi_1^2) \underbrace{y_{t-2}}_{=\Psi_2} + \Phi_1 \Phi_2 y_{t-3} \\
 &= B \eta_t + \Phi_1 B \eta_{t-1} + (\Phi_2 + \Phi_1^2) (\underbrace{\Phi_1 y_{t-3} + \Phi_2 y_{t-4} + B \eta_{t-2}}_{f(y_{t-3}, y_{t-4})}) + \Phi_1 \Phi_2 y_{t-3} \\
 &= \underbrace{B \eta_t}_{=\Psi_0} + \underbrace{\Phi_1 B \eta_{t-1}}_{=\Psi_1} + \underbrace{(\Phi_2 + \Phi_1^2) B \eta_{t-2}}_{=\Psi_2} + f(y_{t-3}, y_{t-4}).
 \end{aligned}$$

In particular, we have $B = \Psi_0$. Matrix B indeed captures the contemporaneous impact of η_t on y_t .

Let us consider the following VARMA(1,1) model:

$$y_t = \underbrace{\begin{bmatrix} 0.5 & 0.3 \\ -0.4 & 0.7 \end{bmatrix}}_{\Phi_1} y_{t-1} + \underbrace{\begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}}_B \eta_t + \underbrace{\begin{bmatrix} 2 & 0 \\ 1 & 0.5 \end{bmatrix}}_{\Theta_1} \underbrace{\begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}}_B (\text{Eq. 38})$$

We can use function `simul.VARMA` of package `AEC` to produce IRFs (using `indic.IRF=1` in the list of arguments):

```

library(AEC)
distri <- list(type=c("gaussian","gaussian"),df=c(4,4))
n <- length(distri$type) # dimension of y_t
nb.sim <- 30
eps <- simul.distri(distri,nb.sim)
Phi <- array(NaN,c(n,n,1))
Phi[,,1] <- matrix(c(.5,-.4,.3,.7),2,2)

```

```

p <- dim(Phi)[3]
Theta <- array(NaN,c(n,n,1))
Theta[, ,1] <- -matrix(c(2,1,0,.5),2,2)
q <- dim(Theta)[3]
Mu <- rep(0,n)
C <- matrix(c(1,-1,2,1),2,2)
Model <- list(
  Mu = Mu, Phi = Phi, Theta = Theta, C = C, distri = distri
)
Y0 <- rep(0,n)
eta0 <- c(1,0)
res.sim.1 <- simul.VARMA(Model,nb.sim,Y0,eta0,indic.IRF=1)
eta0 <- c(0,1)
res.sim.2 <- simul.VARMA(Model,nb.sim,Y0,eta0,indic.IRF=1)

par(plt=c(.1,.95,.25,.8))
par(mfrow=c(2,2))
plot(res.sim.1$Y[1,],las=1,
     type="l",lwd=3,xlab="",ylab="",
     main=expression(paste("Response of ",y[1],"*,*",t)," to a one-unit increase in ",e))
abline(h=0,col="grey",lty=3)
plot(res.sim.2$Y[1,],las=1,
     type="l",lwd=3,xlab="",ylab="",
     main=expression(paste("Response of ",y[1],"*,*",t)," to a one-unit increase in ",e))
abline(h=0,col="grey",lty=3)
plot(res.sim.1$Y[2,],las=1,
     type="l",lwd=3,xlab="",ylab="",
     main=expression(paste("Response of ",y[2],"*,*",t)," to a one-unit increase in ",e))
abline(h=0,col="grey",lty=3)
plot(res.sim.2$Y[2,],las=1,
     type="l",lwd=3,xlab="",ylab="",
     main=expression(paste("Response of ",y[2],"*,*",t)," to a one-unit increase in ",e))
abline(h=0,col="grey",lty=3)

```

6.3.3 Covariance-stationary VARMA models

Let's come back to the infinite MA case (Eq. (6.37)):

$$y_t = \mu + \sum_{h=0}^{\infty} \Psi_h \eta_{t-h}.$$

For y_t to be covariance-stationary (and ergodic for the mean), it has to be the case that

$$\sum_{i=0}^{\infty} \|\Psi_i\| < \infty, \quad (6.39)$$




Figure 6.14: Impulse response functions

where $\|A\|$ denotes a norm of the matrix A (e.g. $\|A\| = \sqrt{\text{tr}(AA')}$).

This notably implies that if y_t is stationary (and ergodic for the mean), then $\|\Psi_h\| \rightarrow 0$ when h gets large.

What should be satisfied by Φ_k 's and Θ_k 's for a VARMA-based process (Eq. @ref{eq:VARMAstd}) to be stationary? The conditions will be similar to that we had in the univariate case (see Prop. 6.6). Let us introduce the following notations:

$$\begin{aligned}
 y_t &= c + \underbrace{\Phi_1 y_{t-1} + \cdots + \Phi_p y_{t-p}}_{\text{AR component}} + \\
 &\quad \underbrace{B \eta_t + \Theta_1 B \eta_{t-1} + \cdots + \Theta_q B \eta_{t-q}}_{\text{MA component}} \\
 &\Leftrightarrow \underbrace{(I - \Phi_1 L - \cdots - \Phi_p L^p)}_{=\Phi(L)} y_t = c + \underbrace{(I - \Theta_1 L - \cdots - \Theta_q L^q)}_{=\Theta(L)} B \eta_t.
 \end{aligned} \tag{6.40}$$

Process y_t is stationary iff the roots of $\det(\Phi(z)) = 0$ are strictly outside the unit circle or, equivalently, iff the eigenvalues of

$$\Phi = \begin{bmatrix} \Phi_1 & \Phi_2 & \cdots & \Phi_p \\ I & 0 & \cdots & 0 \\ 0 & \ddots & 0 & 0 \\ 0 & 0 & I & 0 \end{bmatrix} \tag{6.41}$$

lie strictly within the unit circle.

Hence, as was the case for univariate processes, the covariance-stationarity of a VARMA model depends only on the specification of its AR part.

Let's derive the first two unconditional moments of a (covariance-stationary) VARMA process.

Based on Eq. (6.40), we have $\mathbb{E}(\Phi(L)y_t) = c$, which gives $\Phi(1)\mathbb{E}(y_t) = c$, or::

$$\mathbb{E}(y_t) = (I - \Phi_1 - \cdots - \Phi_p)^{-1}c.$$

The autocovariances of y_t can be deduced from the infinite MA representation (Eq. (6.37)). We have:

$$\gamma_j \equiv \mathbb{C}ov(y_t, y_{t-j}) = \sum_{i=j}^{\infty} \Psi_i \Psi'_{i-j}.$$

(Note that this infinite sum exists as soon as Eq. (6.39) is satisfied.)

Conditional means and autocovariances can also be deduced from Eq. (6.37). For $0 \leq h$ and $0 \leq h_1 \leq h_2$:

$$\begin{aligned} \mathbb{E}_t(y_{t+h}) &= \mu + \sum_{k=0}^{\infty} \Psi_{k+h} \eta_{t-k} \\ \mathbb{C}ov_t(y_{t+1+h_1}, y_{t+1+h_2}) &= \sum_{k=0}^{h_1} \Psi_k \Psi'_{k+h_2-h_1}. \end{aligned}$$

The previous formula implies in particular that the forecasting error $y_{t+h} - \mathbb{E}_t(y_{t+h})$ has a variance equal to:

$$\mathbb{V}ar_t(y_{t+h}) = \sum_{k=1}^h \Psi_k \Psi'_k.$$

Because the η_t are mutually and serially independent (and therefore uncorrelated), we have:

$$\mathbb{V}ar(\Psi_k \eta_{t-k}) = \mathbb{V}ar\left(\sum_{i=1}^n \psi_{k,i} \eta_{i,t-k}\right) = \sum_{i=1}^n \psi_{k,i} \psi'_{k,i},$$

where $\psi_{k,i}$ denotes the i^{th} column of Ψ_k .

This suggests the following decomposition of the variance of the forecast error (called **variance decomposition**):

$$\mathbb{V}ar_t(y_{t+h}) = \sum_{i=1}^n \underbrace{\sum_{k=1}^h \psi_{k,i} \psi'_{k,i}}_{\text{Contribution of } \eta_{i,t}}.$$

Let us now turn to the estimation of VAR(MA) models.

If there is a MA component, OLS regressions yield biased estimates (even for asymptotically large samples).

Assume y_t follows a VARMA(1,1) model. We have:

$$y_{i,t} = \phi_i y_{t-1} + \varepsilon_{i,t},$$

where ϕ_i is the i^{th} row of Φ_1 , and where $\varepsilon_{i,t}$ is a linear combination of η_t and η_{t-1} .

Since y_{t-1} (the regressor) is correlated to η_{t-1} , it is also correlated to $\varepsilon_{i,t}$.

The OLS regression of $y_{i,t}$ on y_{t-1} yields a biased estimator of ϕ_i .

Estimation methods of VARMA models will be presented in Section ??.

6.3.4 VAR estimation

This section discusses the estimation of VAR models. (The estimation of SVARMA models is more challenging, see, e.g., Gouriéroux et al. (2020).) Eq. (6.35) can be written:

$$y_t = c + \Phi(L)y_{t-1} + \varepsilon_t,$$

with $\Phi(L) = \Phi_1 + \Phi_2 L + \dots + \Phi_p L^{p-1}$.

Consequently:

$$y_t \mid y_{t-1}, y_{t-2}, \dots, y_{-p+1} \sim \mathcal{N}(c + \Phi_1 y_{t-1} + \dots + \Phi_p y_{t-p}, \Omega).$$

Using Hamilton (1994)'s notations, denote with Π the matrix $[c \quad \Phi_1 \quad \Phi_2 \quad \dots \quad \Phi_p]'$ and with x_t the vector $[1 \quad y'_{t-1} \quad y'_{t-2} \quad \dots \quad y'_{t-p}]'$, we have:

$$y_t = \Pi' x_t + \varepsilon_t. \tag{6.42}$$

The previous representation is convenient to discuss the estimation of the VAR model, as parameters are gathered in two matrices only: Π and Ω .

Let us start with the case where the shocks are Gaussian.

Proposition 6.12 (MLE of a Gaussian VAR). *If y_t follows a VAR(p) (see Definition 6.22), and if $\varepsilon_t \sim i.i.d. \mathcal{N}(0, \Omega)$, then the ML estimate of Π , denoted by $\hat{\Pi}$ (see Eq. (6.42)), is given by*

$$\hat{\Pi} = \left[\sum_{t=1}^T x_t x_t' \right]^{-1} \left[\sum_{t=1}^T y_t' x_t \right] = (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{y}, \tag{6.43}$$

where \mathbf{X} is the $T \times (np)$ matrix whose t^{th} row is x_t and where \mathbf{y} is the $T \times n$ matrix whose t^{th} row is y_t' .

That is, the i^{th} column of $\hat{\Pi}$ (b_i , say) is the OLS estimate of β_i , where:

$$y_{i,t} = \beta'_i x_t + \varepsilon_{i,t}, \quad (6.44)$$

(i.e., $\beta'_i = [c_i, \phi'_{i,1}, \dots, \phi'_{i,p}]'$).

The ML estimate of Ω , denoted by $\hat{\Omega}$, coincides with the sample covariance matrix of the n series of the OLS residuals in Eq. (6.44), i.e.:

$$\hat{\Omega} = \frac{1}{T} \sum_{i=1}^T \hat{\varepsilon}_t \hat{\varepsilon}'_t, \quad \text{with } \hat{\varepsilon}_t = y_t - \hat{\Pi}' x_t. \quad (6.45)$$

The asymptotic distributions of these estimators are the ones resulting from standard OLS formula.

Proof. See Appendix 7.5. □

As stated by Proposition 6.13, when the shocks are not Gaussian, then the OLS regressions still provide consistent estimates of the model parameters. However, since x_t correlates to ε_s for $s < t$, the OLS estimator \mathbf{b}_i of β_i is biased in small sample. (That is also the case for the ML estimator.)

Indeed, denoting by ε_i the $T \times 1$ vector of $\varepsilon_{i,t}$'s, and using the notations of b_i and β_i introduced in Proposition 6.12, we have:

$$\mathbf{b}_i = \beta_i + (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \varepsilon_i. \quad (6.46)$$

We have non-zero correlation between x_t and $\varepsilon_{i,s}$ for $s < t$ and, therefore, $\mathbb{E}[(\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \varepsilon_i] \neq 0$.

However, when y_t is covariance stationary, then $\frac{1}{n} \mathbf{X}' \mathbf{X}$ converges to a positive definite matrix \mathbf{Q} , and $\frac{1}{n} \mathbf{X}' \varepsilon_i$ converges to 0. Hence $\mathbf{b}_i \xrightarrow{p} \beta_i$. More precisely:

Proposition 6.13 (Asymptotic distribution of the OLS estimate of β_i). *If y_t follows a VAR model, as defined in Definition 6.22, we have:*

$$\sqrt{T}(\mathbf{b}_i - \beta_i) = \underbrace{\left[\frac{1}{T} \sum_{t=p}^T x_t x_t' \right]^{-1}}_{\xrightarrow{p} \mathbf{Q}^{-1}} \underbrace{\sqrt{T} \left[\frac{1}{T} \sum_{t=1}^T x_t \varepsilon_{i,t} \right]}_{\xrightarrow{d} \mathcal{N}(0, \sigma_i^2 \mathbf{Q})},$$

where $\sigma_i = \text{Var}(\varepsilon_{i,t})$ and where $\mathbf{Q} = \text{plim} \frac{1}{T} \sum_{t=p}^T x_t x_t'$ is given by:

$$\mathbf{Q} = \begin{bmatrix} 1 & \mu' & \mu' & \dots & \mu' \\ \mu & \gamma_0 + \mu\mu' & \gamma_1 + \mu\mu' & \dots & \gamma_{p-1} + \mu\mu' \\ \mu & \gamma_1 + \mu\mu' & \gamma_0 + \mu\mu' & \dots & \gamma_{p-2} + \mu\mu' \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \mu & \gamma_{p-1} + \mu\mu' & \gamma_{p-2} + \mu\mu' & \dots & \gamma_0 + \mu\mu' \end{bmatrix}. \quad (6.47)$$

Proof. See Appendix 7.5. \square

The following proposition extends the previous proposition and includes covariances between different β_i 's as well as the asymptotic distribution of the ML estimates of Ω .

Proposition 6.14 (Asymptotic distribution of the OLS estimates). *If y_t follows a VAR model, as defined in Definition 6.22, we have:*

$$\sqrt{T} \begin{bmatrix} \text{vec}(\hat{\Pi} - \Pi) \\ \text{vec}(\hat{\Omega} - \Omega) \end{bmatrix} \sim \mathcal{N} \left(0, \begin{bmatrix} \Omega \otimes \mathbf{Q}^{-1} & 0 \\ 0 & \Sigma_{22} \end{bmatrix} \right), \quad (6.48)$$

where the component of Σ_{22} corresponding to the covariance between $\hat{\sigma}_{i,j}$ and $\hat{\sigma}_{k,l}$ (for $i, j, l, m \in \{1, \dots, n\}^4$) is equal to $\sigma_{i,l}\sigma_{j,m} + \sigma_{i,m}\sigma_{j,l}$.

Proof. See Hamilton (1994), Appendix of Chapter 11. \square

Naturally, in practice, Ω is replaced by $\hat{\Omega}$, Q is replaced with $\hat{\mathbf{Q}} = \frac{1}{T} \sum_{t=p}^T x_t x_t'$ and Σ with the matrix whose components are of the form $\hat{\sigma}_{i,l}\hat{\sigma}_{j,m} + \hat{\sigma}_{i,m}\hat{\sigma}_{j,l}$, where the $\hat{\sigma}_{i,l}$'s are the components of $\hat{\Omega}$.

The simplicity of the VAR framework and the tractability of its MLE open the way to convenient econometric testing. Let's illustrate this with the likelihood ratio test. The maximum value achieved by the MLE is

$$\log \mathcal{L}(Y_T; \hat{\Pi}, \hat{\Omega}) = -\frac{Tn}{2} \log(2\pi) + \frac{T}{2} \log |\hat{\Omega}^{-1}| - \frac{1}{2} \sum_{t=1}^T [\hat{\varepsilon}'_t \hat{\Omega}^{-1} \hat{\varepsilon}_t].$$

The last term is:

$$\begin{aligned} \sum_{t=1}^T \hat{\varepsilon}'_t \hat{\Omega}^{-1} \hat{\varepsilon}_t &= \text{Tr} \left[\sum_{t=1}^T \hat{\varepsilon}'_t \hat{\Omega}^{-1} \hat{\varepsilon}_t \right] = \text{Tr} \left[\sum_{t=1}^T \hat{\Omega}^{-1} \hat{\varepsilon}_t \hat{\varepsilon}'_t \right] \\ &= \text{Tr} \left[\hat{\Omega}^{-1} \sum_{t=1}^T \hat{\varepsilon}_t \hat{\varepsilon}'_t \right] = \text{Tr} [\hat{\Omega}^{-1} (T\hat{\Omega})] = Tn. \end{aligned}$$

Therefore, the optimized log-likelihood is simply obtained by:

$$\log \mathcal{L}(Y_T; \hat{\Pi}, \hat{\Omega}) = -(Tn/2) \log(2\pi) + (T/2) \log |\hat{\Omega}^{-1}| - Tn/2. \quad (6.49)$$

Assume that we want to test the null hypothesis that a set of variables follows a VAR(p_0) against the alternative specification of $p_1 (> p_0)$.

Let us denote by \hat{L}_0 and \hat{L}_1 the maximum log-likelihoods obtained with p_0 and p_1 lags, respectively.

Under the null hypothesis ($H_0: p = p_0$), we have:

$$2(\hat{L}_1 - \hat{L}_0) = T(\log |\hat{\Omega}_1^{-1}| - \log |\hat{\Omega}_0^{-1}|) \sim \chi^2(n^2(p_1 - p_0)).$$

What precedes can be used to help determine the appropriate number of lags to use in the specification. In a VAR, using too many lags consumes numerous degrees of freedom: with p lags, each of the n equations in the VAR contains $n \times p$ coefficients plus the intercept term. Adding lags improve in-sample fit, but is likely to result in over-parameterization and affect the **out-of-sample** prediction performance.

To select appropriate lag length, **selection criteria** can be used (see Definition 6.21). In the context of VAR models, using Eq. (6.49), we have:

$$\begin{aligned} AIC &= cst + \log |\hat{\Omega}| + \frac{2}{T} N \\ BIC &= cst + \log |\hat{\Omega}| + \frac{\log T}{T} N, \end{aligned}$$

where $N = p \times n^2$.

6.3.5 Block exogeneity and Granger causality

Block exogeneity

Let's decompose y_t into two subvectors $y_t^{(1)}$ ($n_1 \times 1$) and $y_t^{(2)}$ ($n_2 \times 1$), with $y_t' = [y_t^{(1)'}', y_t^{(2)'}']$ (and therefore $n = n_1 + n_2$), such that:

$$\begin{bmatrix} y_t^{(1)} \\ y_t^{(2)} \end{bmatrix} = \begin{bmatrix} \Phi^{(1,1)} & \Phi^{(1,2)} \\ \Phi^{(2,1)} & \Phi^{(2,2)} \end{bmatrix} \begin{bmatrix} y_{t-1}^{(1)} \\ y_{t-1}^{(2)} \end{bmatrix} + \varepsilon_t.$$

Using, e.g., a likelihood ratio test (see Def. 4.8), one can easily test for block exogeneity of $y_t^{(2)}$ (say). The null assumption can be expressed as $\Phi^{(2,1)} = 0$ and $\Sigma^{(2,1)} = 0$.

Granger Causality

Granger (1969) developed a method to explore **causal relationships** among variables. The approach consists in determining whether the past values of $y_{1,t}$ can help explain the current $y_{2,t}$ (beyond the information already included in the past values of $y_{2,t}$).

Formally, let us denote three information sets:

$$\begin{aligned} \mathcal{I}_{1,t} &= \{y_{1,t}, y_{1,t-1}, \dots\} \\ \mathcal{I}_{2,t} &= \{y_{2,t}, y_{2,t-1}, \dots\} \\ \mathcal{I}_t &= \{y_{1,t}, y_{1,t-1}, \dots, y_{2,t}, y_{2,t-1}, \dots\}. \end{aligned}$$

We say that $y_{1,t}$ Granger-causes $y_{2,t}$ if

$$\mathbb{E} [y_{2,t} | \mathcal{I}_{2,t-1}] \neq \mathbb{E} [y_{2,t} | \mathcal{I}_{t-1}].$$

To get the intuition behind the testing procedure, consider the following bivariate VAR(p) process:

$$\begin{aligned} y_{1,t} &= c_1 + \sum_{i=1}^p \Phi_i^{(11)} y_{1,t-i} + \sum_{i=1}^p \Phi_i^{(12)} y_{2,t-i} + \varepsilon_{1,t} \\ y_{2,t} &= c_2 + \sum_{i=1}^p \Phi_i^{(21)} y_{1,t-i} + \sum_{i=1}^p \Phi_i^{(22)} y_{2,t-i} + \varepsilon_{2,t}, \end{aligned}$$

where $\Phi_k^{(ij)}$ denotes the element (i, j) of Φ_k .

Then, $y_{1,t}$ is said not to Granger-cause $y_{2,t}$ if

$$\Phi_1^{(21)} = \Phi_2^{(21)} = \dots = \Phi_p^{(21)} = 0.$$

Therefore the hypothesis testing is

$$\begin{cases} H_0 : \Phi_1^{(21)} = \Phi_2^{(21)} = \dots = \Phi_p^{(21)} = 0 \\ H_1 : \Phi_1^{(21)} \neq 0 \text{ or } \Phi_2^{(21)} \neq 0 \text{ or } \dots \Phi_p^{(21)} \neq 0. \end{cases}$$

Loosely speaking, we reject H_0 if some of the coefficients on the lagged $y_{1,t}$'s are statistically significant. Formally, this can be tested using the F -test or asymptotic chi-square test. The F -statistic is

$$F = \frac{(RSS - USS)/p}{USS/(T - 2p - 1)},$$

where RSS is the Restricted sum of squared residuals and USS is the Unrestricted sum of squared residuals. Under H_0 , the F -statistic is distributed as $\mathcal{F}(p, T - 2p - 1)$.

Note that $pF \xrightarrow{T \rightarrow \infty} \chi^2(p)$. Therefore, for large samples and under H_0 :

$$F \sim \chi^2(p)/p.$$

6.3.6 Identification problem and standard identification techniques

The Identification Issue

In Section 6.3.4, we have seen how to estimate $\text{Var}(\varepsilon_t) = \Omega$ and the Φ_k matrices in the context of a VAR model. But the IRFs are functions of B and the Φ_k 's, not of Ω the Φ_k 's (see Section 6.3.2). We have $\Omega = BB'$, but this is not sufficient to recover B .

Indeed, seen a system of equations whose unknowns are the $b_{i,j}$'s (components of B), the system $\Omega = BB'$ contains only $n(n + 1)/2$ linearly independent equations. For instance, for $n = 2$:

$$\begin{aligned} & \begin{bmatrix} \omega_{11} & \omega_{12} \\ \omega_{12} & \omega_{22} \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{21} \\ b_{12} & b_{22} \end{bmatrix} \\ \Leftrightarrow & \begin{bmatrix} \omega_{11} & \omega_{12} \\ \omega_{12} & \omega_{22} \end{bmatrix} = \begin{bmatrix} b_{11}^2 + b_{12}^2 & b_{11}b_{21} + b_{12}b_{22} \\ b_{11}b_{21} + b_{12}b_{22} & b_{22}^2 + b_{21}^2 \end{bmatrix}. \end{aligned}$$

We then have 3 linearly independent equations but 4 unknowns. Therefore, B is not identified based on second-order moments. Additional restrictions are required to identify B .

This section covers two standard identification schemes: **short-run** and **long-run** restrictions:

1. a **short-run restriction (SRR)** prevents a structural shock from affecting an endogenous variable contemporaneously.
 - Easy to implement: the appropriate entries of B are set to 0.
 - Particular case: **Cholesky, or recursive approach.**
 - Examples: Bernanke (1986), Sims (1986), Galí (1992), Rubio-Ramírez et al. (2010).
2. a **long-run restriction (LRR)** prevents a structural shock from having a cumulative impact on one of the endogenous variables.
 - Additional computations are required to implement this. One needs to compute the cumulative effect of one of the structural shocks u_t on one of the endogenous variable.
 - Examples: Blanchard and Quah (1989), Faust and Leeper (1997), Galí (1999), Erceg et al. (2005), Christiano et al. (2007).

The two approaches can be combined (see, e.g., Gerlach and Smets (1995)).

Let us consider a simple example that could motivate short-run restrictions. Consider the following stylized macro model:

$$\begin{aligned} g_t &= \bar{g} - \lambda(i_{t-1} - \mathbb{E}_{t-1}\pi_t) + \underbrace{\sigma_d \eta_{d,t}}_{\text{demand shock}} && (\text{IS curve}) \\ \Delta\pi_t &= \beta(g_t - \bar{g}) + \underbrace{\sigma_\pi \eta_{\pi,t}}_{\substack{\text{cost push shock}}} && (\text{Phillips curve}) \\ i_t &= \rho i_{t-1} + [\gamma_\pi \mathbb{E}_t \pi_{t+1} + \gamma_g(g_t - \bar{g})] \\ &\quad + \underbrace{\sigma_{mp} \eta_{mp,t}}_{\text{Mon. Pol. shock}} && (\text{Taylor rule}), \end{aligned} \tag{6.50}$$

where:

$$\eta_t = \begin{bmatrix} \eta_{\pi,t} \\ \eta_{d,t} \\ \eta_{mp,t} \end{bmatrix} \sim i.i.d. \mathcal{N}(0, I). \quad (6.51)$$

Vector η_t is assumed to be a vector of structural shocks, mutually and serially independent. On date t :

- g_t is contemporaneously affected by $\eta_{d,t}$ only;
- π_t is contemporaneously affected by $\eta_{\pi,t}$ and $\eta_{d,t}$;
- i_t is contemporaneously affected by $\eta_{mp,t}$, $\eta_{\pi,t}$ and $\eta_{d,t}$.

System (6.50) could be rewritten in the form:

$$\begin{bmatrix} d_t \\ \pi_t \\ i_t \end{bmatrix} = \Phi(L) \begin{bmatrix} d_{t-1} \\ \pi_{t-1} \\ i_{t-1} \end{bmatrix} + \underbrace{\begin{bmatrix} 0 & \bullet & 0 \\ \bullet & \bullet & 0 \\ \bullet & \bullet & \bullet \end{bmatrix}}_{=B} \eta_t + \underbrace{\varepsilon_t}_{=B\varepsilon_t} \quad (6.52)$$

This is the **reduced-form** of the model. This representation suggests three additional restrictions on the entries of B ; the latter matrix is therefore identified (up to the signs of its columns) as soon as $\Omega = BB'$ is known.

There are particular cases in which some well-known matrix decomposition of $\Omega = \text{Var}(\varepsilon_t)$ can be used to easily estimate some specific SVAR.

Consider the following context:

- A first shock (say, $\eta_{n_1,t}$) can affect instantaneously (i.e., on date t) only one of the endogenous variable (say, $y_{n_1,t}$);
- A second shock (say, $\eta_{n_2,t}$) can affect instantaneously (i.e., on date t) the first two endogenous variables (say, $y_{n_1,t}$ and $y_{n_2,t}$);
- ...

This implies (1) that column n_1 of B has only 1 non-zero entry (this is the n_1^{th} entry), (2) that column n_2 of B has 2 non-zero entries (the n_1^{th} and the n_2^{th} ones), etc. Without loss of generality, we can set $n_1 = n$, $n_2 = n - 1$, etc. In this context, matrix B is lower triangular.

The Cholesky decomposition of Ω_ε then provides an appropriate estimate of B , that is a lower triangular matrix B that is such that:

$$\Omega_\varepsilon = BB'.$$

For instance, Dedola and Lippi (2005) estimate 5 structural VAR models for the US, the UK, Germany, France and Italy to analyse the monetary-policy transmission mechanisms. They estimate SVAR(5) models over the period 1975-1997.

The shock-identification scheme is based on Cholesky decompositions, the ordering of the endogenous variables being: the industrial production, the consumer price index, a commodity price index, the short-term rate, monetary aggregate and the effective exchange rate (except for the US). This ordering implies that monetary policy reacts to the shocks affecting the first three variables but that the latter react to monetary policy shocks with a one-period lag only.

Importantly, the Cholesky approach can be useful when one is interested in one specific structural shock. This was the case, e.g., of Christiano et al. (1996). Their identification is based on the following relationship between ε_t and η_t :

$$\begin{bmatrix} \varepsilon_{S,t} \\ \varepsilon_{r,t} \\ \varepsilon_{F,t} \end{bmatrix} = \begin{bmatrix} B_{SS} & 0 & 0 \\ B_{rS} & B_{rr} & 0 \\ B_{FS} & B_{Fr} & B_{FF} \end{bmatrix} \begin{bmatrix} \eta_{S,t} \\ \eta_{r,t} \\ \eta_{F,t} \end{bmatrix},$$

where S , r and F respectively correspond to *slow-moving variables*, the policy variable (short-term rate) and *fast-moving variables*. While $\eta_{r,t}$ is scalar, $\eta_{S,t}$ and $\eta_{F,t}$ may be vectors. The space spanned by $\varepsilon_{S,t}$ is the same as that spanned by $\eta_{S,t}$. As a result, because $\varepsilon_{r,t}$ is a linear combination of $\eta_{r,t}$ and $\eta_{S,t}$ (which are \perp), it comes that the $B_{rr}\eta_{r,t}$'s are the (population) residuals in the regression of $\varepsilon_{r,t}$ on $\varepsilon_{S,t}$. Because $\text{Var}(\eta_{r,t}) = 1$, B_{rr} is given by the square root of the variance of $B_{rr}\eta_{r,t}$. $B_{F,r}$ is finally obtained by regressing the components of $\varepsilon_{F,t}$ on $\eta_{r,t}$.

An equivalent approach consists in computing the Cholesky decomposition of BB' and the contemporaneous impacts of the monetary policy shock (on the n endogenous variables) are the components of the column of B corresponding to the policy variable.

```
library(AEC)
library(vars)

## Loading required package: strucchange

## Loading required package: urca

data("USmonthly")
# Select sample period:
First.date <- "1965-01-01"
Last.date <- "1995-06-01"
indic.first <- which(USmonthly$DATES==First.date)
indic.last <- which(USmonthly$DATES==Last.date)
USmonthly <- USmonthly[indic.first:indic.last,]
considered.variables <- c("LIP", "UNEMP", "LCPI", "LPCOM", "FFR", "NBR", "TTR", "M1")
y <- as.matrix(USmonthly[considered.variables])
```

```

res.svar.ordering <- svar.ordering(y,p=3,
                                     posit.of.shock = 5,
                                     nb.periods.IRF = 20,
                                     nb.bootstrap.replications = 100, # This is used in the parameter
                                     confidence.interval = 0.90, # expressed in pp.
                                     indic.plot = 1 # Plots are displayed if = 1.
)

```




Figure 6.15: Response to a monetary-policy shock. Identification approach of Christiano, Eichenbaum and Evans (1996).

Let us now turn to **Long-run restrictions**. Such a restriction concerns the long-run influence of a shock on an endogenous variable. Let us consider for instance a structural shock that is assumed to have no “long-run influence” on GDP. How to express this? The long-run change in GDP can be expressed as $GDP_{t+h} - GDP_t$, with h large. Note further that:

$$GDP_{t+h} - GDP_t = \Delta GDP_{t+h} + \Delta GDP_{t+h-1} + \cdots + \Delta GDP_{t+1}.$$

Hence, the fact that a given structural shock ($\eta_{i,t}$, say) has no long-run influence on GDP means that

$$\lim_{h \rightarrow \infty} \frac{\partial GDP_{t+h}}{\partial \eta_{i,t}} = \lim_{h \rightarrow \infty} \frac{\partial}{\partial \eta_{i,t}} \left(\sum_{k=1}^h \Delta GDP_{t+k} \right) = 0.$$

This can be easily formulated as a function of B and of the matrices Φ_i when y_t (including ΔGDP_t) follows a VAR process.

Without loss of generality, we will only consider the VAR(1) case. Indeed, one can always write a VAR(p) as a VAR(1). To see that, stack the last p values of vector y_t in vector $y_t^* = [y_t', \dots, y_{t-p+1}']'$; Eq. (6.35) can then be rewritten in its companion form:

$$y_t^* = \underbrace{\begin{bmatrix} c \\ 0 \\ \vdots \\ 0 \end{bmatrix}}_{=c^*} + \underbrace{\begin{bmatrix} \Phi_1 & \Phi_2 & \cdots & \Phi_p \\ I & 0 & \cdots & 0 \\ 0 & \ddots & 0 & 0 \\ 0 & 0 & I & 0 \end{bmatrix}}_{=\Phi} y_{t-1}^* + \underbrace{\begin{bmatrix} \varepsilon_t \\ 0 \\ \vdots \\ 0 \end{bmatrix}}_{=\varepsilon_t^*}, \quad (6.53)$$

where matrices Φ and $\Omega^* = \text{Var}(\varepsilon_t^*)$ are of dimension $np \times np$; Ω^* is filled with zeros, except the $n \times n$ upper-left block that is equal to $\Omega = \text{Var}(\varepsilon_t)$. (Matrix Φ had been introduced in Eq. (6.41).)

Focusing on the VAR(1) case:

$$y_t = c + \Phi y_{t-1} + \varepsilon_t \quad (6.54)$$

$$= c + \varepsilon_t + \Phi(c + \varepsilon_{t-1}) + \dots + \Phi^k(c + \varepsilon_{t-k}) + \dots \quad (6.55)$$

$$= \mu + \varepsilon_t + \Phi \varepsilon_{t-1} + \dots + \Phi^k \varepsilon_{t-k} + \dots \quad (6.56)$$

$$= \mu + B \eta_t + \Phi B \eta_{t-1} + \dots + \Phi^k B \eta_{t-k} + \dots, \quad (6.56)$$

The sequence of shocks $\{\eta_t\}$ determines the sequence $\{y_t\}$. What if $\{\eta_t\}$ is replaced by $\{\tilde{\eta}_t\}$, where $\tilde{\eta}_t = \eta_t$ if $t \neq s$ and $\tilde{\eta}_s = \eta_s + \gamma$?

Assume $\{\tilde{y}_t\}$ is the associated “perturbated” sequence. We have $\tilde{y}_t = y_t$ if $t < s$. For $t \geq s$, the Wold decomposition of $\{\tilde{y}_t\}$ implies:

$$\tilde{y}_t = y_t + \Phi^{t-s} B \gamma.$$

Therefore, the cumulative impact of γ on \tilde{y}_t will be (for $t \geq s$):

$$\begin{aligned} (\tilde{y}_t - y_t) + (\tilde{y}_{t-1} - y_{t-1}) + \dots + (\tilde{y}_s - y_s) &= \\ (Id + \Phi + \Phi^2 + \dots + \Phi^{t-s}) B \gamma. & \end{aligned} \quad (6.57)$$

Consider a shock on $\eta_{1,t}$, with a magnitude of 1. This shock corresponds to $\gamma = [1, 0, \dots, 0]'$. Given Eq. (6.57), the long-run cumulative effect of this shock on the endogenous variables is given by:

$$\underbrace{(Id + \Phi + \dots + \Phi^k + \dots)}_{=(Id - \Phi)^{-1}} B \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix},$$

that is the first column of $\Theta \equiv (Id - \Phi)^{-1} B$.

In this context, consider the following long-run restriction: “ j^{th} structural shock has no cumulative impact on the i^{th} endogenous variable”. It is equivalent to


$$\Theta_{ij} = 0,$$

where Θ_{ij} is the element (i, j) of Θ .

Blanchard and Quah (1989) implement long-run restrictions in a small-scale VAR. Two variables are considered: GDP and unemployment. Consequently, the VAR is affected by two types of shocks. The authors want to identify **supply shocks** (that can have a permanent effect on output) and **demand shocks** (that cannot have a permanent effect on output).²

Blanchard and Quah (1989)'s dataset is quarterly, spanning the period from 1950:2 to 1987:4. Their VAR features 8 lags. Here are the data they use:

```
library(AEC)
data(BQ)
par(mfrow=c(1, 2))
plot(BQ$Date, BQ$Dgdp, type="l", main="GDP quarterly growth rate", xlab="", ylab="", lwd=2)
plot(BQ$Date, BQ$unemp, type="l", ylim=c(-3, 6), main="Unemployment rate (gap)", xlab="", ylab="", lwd=2)
```



Estimate a reduced-form VAR(8) model:

```
library(vars)
y <- BQ[, 2:3]
est.VAR <- VAR(y, p=8)
Omega <- var(residuals(est.VAR))
```

²The motivation of the authors regarding their long-run restrictions can be obtained from a traditional Keynesian view of fluctuations. The authors propose a variant of a model from Fischer (1977).

Now, let us define a loss function (`loss`) that is equal to zero if (a) $BB' = \Omega$ and (b) the element (1,1) of ΘB is equal to zero:

```
# Compute (Id - Phi)^{-1}:
Phi <- Acoef(est.VAR)
PHI <- make.PHI(Phi)
sum.PHI.k <- solve(diag(dim(PHI)[1]) - PHI)[1:2,1:2]
loss <- function(param){
  B <- matrix(param,2,2)
  X <- Omega - B %*% t(B)
  Theta <- sum.PHI.k[1:2,1:2] %*% B
  loss <- 10000 * ( X[1,1]^2 + X[2,1]^2 + X[2,2]^2 + Theta[1,1]^2 )
  return(loss)
}
res.opt <- optim(c(1,0,0,1),loss,method="BFGS",hessian=FALSE)
print(res.opt$par)

## [1] 0.8570358 -0.2396345 0.1541395 0.1921221
```

(Note: one can use that type of approach, based on a loss function, to mix short- and long-run restrictions.)

Figure 6.16 displays the resulting IRFs. Note that, for GDP, we cumulate the GDP growth IRF, so as to have the response of the GDP level.

```
B.hat <- matrix(res.opt$par,2,2)
print(cbind(Omega,B.hat %*% t(B.hat)))

##          Dgdp      unemp
## Dgdp  0.7582704 -0.17576173  0.7582694 -0.17576173
## unemp -0.1757617  0.09433658 -0.1757617  0.09433558

nb.sim <- 40
par(mfrow=c(2,2))
Y <- simul.VAR(c=matrix(0,2,1),Phi,B.hat,nb.sim,y0.star=rep(0,2*8),indic.IRF = 1,u.sho
plot(cumsum(Y[,1]),type="l",lwd=2,xlab="",ylab="",main="Demand shock on GDP")
plot(Y[,2],type="l",lwd=2,xlab="",ylab="",main="Demand shock on UNEMP")
Y <- simul.VAR(c=matrix(0,2,1),Phi,B.hat,nb.sim,y0.star=rep(0,2*8),indic.IRF = 1,u.sho
plot(cumsum(Y[,1]),type="l",lwd=2,xlab="",ylab="",main="Supply shock on GDP")
plot(Y[,2],type="l",lwd=2,xlab="",ylab="",main="Supply shock on UNEMP")
```

6.3.7 Sign restrictions

To identify the structural shocks, we need to find a matrix B that satisfies $\Omega_\varepsilon = BB'$ (with $\Omega = \text{Var}(\varepsilon_t)$) and other restrictions. Indeed, $\Omega_\varepsilon = BB'$ is not




Figure 6.16: IRF of GDP and unemployment to demand and supply shocks.

sufficient to identify B since, if we take any orthogonal matrix Q (see Def. 6.24), then $\mathcal{P} = BQ$ also satisfies $\Omega = \mathcal{P}\mathcal{P}'$.

Definition 6.24 (Orthogonal matrix). An orthogonal matrix Q is a matrix such that $QQ' = I$, i.e., all columns (rows) of Q are orthogonal and unit vectors:

$$q_i' q_j = 0 \text{ if } i \neq j \text{ and } q_i' q_j = 1 \text{ if } i = j,$$

where q_i is the i^{th} column of Q .

The idea behind the sign-restriction approach is to “draw” random matrices \mathcal{P} that satisfy $\Omega = \mathcal{P}\mathcal{P}'$, and then to constitute a set of admissible matrices, keeping in this set only the simulated \mathcal{P} matrices that satisfy some predefined sign-based restriction. An example of restriction is “*after one year, a contractionary monetary-policy shocks has a negative impact on inflation*”.

As suggested above, if B is any matrix that satisfies $\Omega = BB'$ (for instance, B can be based on the Cholesky decomposition of Ω), then we also have $\Omega = \mathcal{P}\mathcal{P}'$ as soon as $\mathcal{P} = BQ$, where Q is an orthogonal matrix. Therefore, to draw \mathcal{P} matrices, it suffices to draw in the set of orthogonal matrices.

To fix ideas, consider dimension 2. In that case, the orthogonal matrices are rotation matrices, and the set of orthogonal matrices can be parameterized by the angle x , with:

$$Q_x = \begin{pmatrix} \cos(x) & \cos(x + \frac{\pi}{2}) \\ \sin(x) & \sin(x + \frac{\pi}{2}) \end{pmatrix} = \begin{pmatrix} \cos(x) & -\sin(x) \\ \sin(x) & \cos(x) \end{pmatrix}.$$

(This is an angle- x counter-clockwise rotation.) Hence, in that case, by drawing x randomly from $[0, 2\pi]$, we draw randomly from the set of 2×2 rotation matrices. For high-dimensional VAR, we lose this simple geometrical representation, though. It is not always possible to parametrize a rotation matrix (high-dimensional VARs).

How to proceed, then? Arias et al. (2018) provide a procedure. Their approach is based on the so-called QR decomposition: any square matrix X may be decomposed as $X = QR$ where Q is an orthogonal matrix and R is an upper diagonal matrix. With this in mind, they propose a two-step approach:

- i. Draw a random matrix X by drawing each element from independent standard normal distribution.
- ii. Let $X = QR$ be the QR decomposition of X with the diagonal of R normalized to be positive. The random matrix Q is orthogonal and is a draw from the uniform distribution over the set of orthogonal matrices.

Equipped with this procedure, the sign-restriction is based on the following algorithm:

1. Draw a random orthogonal matrix Q (using step i. and ii. described above).
2. Compute $B = PQ$ where P is the Cholesky decomposition of the reduced form residuals Ω_ε .
3. Compute the impulse response associated with B $y_{t,t+k} = \Phi^k B$ or the cumulated response $\bar{y}_{t,t+k} = \sum_{j=0}^k \Phi^j B$.
4. Are the sign restrictions satisfied?
 - a. **Yes.** Store the impulse response in the set of admissible response.
 - b. **No.** Discard the impulse response.
5. Perform N replications and report the median impulse response (and its “confidence” intervals).

Note: to take into account the uncertainty in B and Φ , you can draw B and Φ in Steps 2 and 3 using an inference method.

This method has the advantage of being relatively agnostic. Moreover, it is fairly flexible, as one can impose sign restrictions on any variable, at any horizon.

A prominent example is Uhlig (2005). Using US monthly data from 1965.I to 2003.XII, he employs sign restrictions to estimate the effect of monetary policy shocks.

According to conventional wisdom, monetary contractions should:³

- Raise the federal funds rate,
- Lower prices,
- Decrease non-borrowed reserves,
- Reduce real output.

The restrictions considered by Uhlig (2005) are as follows: an expansionary monetary policy shock leads to:

- Increases in prices
- Increase in nonborrowed reserves
- Decreases in the federal funds rate

What about output? Since is the response of interest, we leave it un-restricted.

```
library(AEC);library(vars);library(Matrix)
data("USmonthly")
First.date <- "1965-01-01"
Last.date <- "1995-06-01"
indic.first <- which(USmonthly$DATES==First.date)
indic.last <- which(USmonthly$DATES==Last.date)
USmonthly <- USmonthly[indic.first:indic.last,]
considered.variables <- c("LIP", "UNEMP", "LCPI", "LPCOM", "FFR", "NBR", "TTR", "M1")
n <- length(considered.variables)
y <- as.matrix(USmonthly[considered.variables])
sign.restrictions <- list()
horizon <- list()
#Define sign restrictions and horizon for restrictions
for(i in 1:n){
  sign.restrictions[[i]] <- matrix(0,n,n)
  horizon[[i]] <- 1
}
sign.restrictions[[1]][1,3] <- 1
sign.restrictions[[1]][2,5] <- -1
sign.restrictions[[1]][3,6] <- 1
horizon[[1]] <- 1:5
res.svar.signs <- svar.signs(y,p=3,
                                nb.shocks = 1, #number of identified shocks
                                nb.periods.IRF = 20,
```

³Standard identification schemes often fail to achieve these 4 points. Two puzzles regularly arise: *liquidity puzzle*: when identifying monetary policy shocks as surprise increases in the stock of money, interest rates tend to go up, not down; *price puzzle*: after a contractionary monetary policy shock, even with interest rates going up and money supply going down, inflation goes up rather than down.

```

bootstrap.replications = 1, # = 0 if no bootstrap
confidence.interval = 0.80, # expressed in pp.
indic.plot = 1, # Plots are displayed if = 1.
nb.draws = 10000, # number of draws
sign.restrictions,
horizon,
recursive =1 # =0 <- draw Q directly, =1 <- draw q recursive
)

```




Figure 6.17: IRF associated with a monetary policy shock; sign-restriction approach.

```

nb.rotations <- res.svar.signs$xx

```

A remarkable characteristic of sign restrictions is that it does not lead to a unique IRF. In that sense, we say that it is set-identified, not point-identified. By drawing the rotation matrices (Q) from a distribution and taking the median, we adopt an agnostic approach: “pure sign restriction approach”.

An alternative approach is the so-called **penalty-function approach** (PFA, Uhlig (2005), present in Danne (2015)’s package). This approach relies on a *penalty function*:

$$f(x) = \begin{cases} x & \text{if } x \leq 0 \\ 100.x & \text{if } x > 0 \end{cases}$$

which penalizes positive responses and rewards negative responses.

Let $\psi_k^j(q)$ be the impulse response of variable j . The $\psi_k^j(q)$'s are the elements of $\psi_k(q) = \Psi_k q$.

Let σ_j be the standard deviation of variable j . Let $\iota_{j,k} = 1$ if we restrict the response of variable j at the $k^t h$ horizon to be negative, $\iota_{j,k} = -1$ if we restrict it to be positive, and $\iota_{j,k} = 0$ if there is no restriction. The total penalty is given by

$$\mathbf{P}(q) = \sum_{j=1}^m \sum_{k=0}^K f \left(\iota_{j,k} \frac{\psi_k^j(q)}{\sigma_j} \right).$$

We are looking for a solution to

$$\min_q \mathbf{P}(q)$$

$$\text{s.t. } q' q = 1.$$

The problem is solved numerically.

6.3.8 Forecast error variance maximization

The approach presented in this section exploits the derivations of Uhlig (2004). Barsky and Sims (2011) exploit this approach to identify a TFP news shock, that they define as the shock (a) that is orthogonal to the innovation in current utilization-adjusted TFP and (b) that best explains variation in future TFP.

Consider a process $\{y_t\}$ that admits the infinite MA representation of Eq. (6.37). Let Q be an orthogonal matrix, an alternative decomposition is:

$$y_t = \sum_{h=0}^{+\infty} \underbrace{\Psi_h}_{Q \tilde{\eta}_{t-h}} \eta_{t-h} = \sum_{h=0}^{+\infty} \underbrace{\Psi_h Q}_{\tilde{\Psi}_h} \tilde{\eta}_{t-h} = \sum_{h=0}^{+\infty} \tilde{\Psi}_h \tilde{\eta}_{t-h}, \quad (6.58)$$

where $\tilde{\eta}_{t-h} = Q' \eta_{t-h}$ are the white-noise shocks associated with the new MA representation. (They also satisfy $\text{Var}(\tilde{\eta}_t) = Id$.)

The h -step ahead prediction error of y_{t+h} , given all the data up to and including $t-1$ is given by

$$e_{t+h}(h) = y_{t+h} - \mathbb{E}_{t-1}(y_{t+h}) = \sum_{j=0}^h \tilde{\Psi}_h \tilde{\eta}_{t+h-j}.$$

The variance-covariance matrix of $e_{t+h}(h)$ is

$$\Omega(h) = \sum_{j=0}^h \tilde{\Psi}_j \tilde{\Psi}'_j = \sum_{j=0}^h \Psi_j \Psi'_j.$$

We can decompose $\Omega(h)$ into the contribution of each shock l (l^{th} component of $\tilde{\eta}_t$):

$$\Omega^{(h)} = \sum_{l=1}^n \Omega_l^{(h)}(Q)$$

with

$$\Omega_l^{(h)}(Q) = \sum_{j=0}^h (\Psi_j q_l)(\Psi_j q_l)',$$

where q_l is the l^{th} column of Q .

This decomposition can be used with the objective of finding the **impulse vector** b that is s.t. that it explains as much as possible of the sum of the h -step ahead prediction error variance of some variable i , say, for prediction horizons $h \in [\underline{h}, \bar{h}]$.

Formally, the task is to explain as much as possible of the variance

$$\sigma^2(\underline{h}, \bar{h}, q_1) = \sum_{h=\underline{h}}^{\bar{h}} \sum_{j=0}^h [(\Psi_j q_1)(\Psi_j q_1)']_{i,i}$$

with a single impulse vector q_1 .

Denote by E_{ii} the matrix that is filled with zeros, except for its (i, i) entry, set to 1. We have:

$$\begin{aligned} \sigma^2(\underline{h}, \bar{h}, q_1) &= \sum_{h=\underline{h}}^{\bar{h}} \sum_{j=0}^h [(\Psi_j q_1)(\Psi_j q_1)']_{i,i} = \sum_{h=\underline{h}}^{\bar{h}} \sum_{j=0}^h \text{Tr}[E_{ii}(\Psi_j q_1)(\Psi_j q_1)'] \\ &= \sum_{h=\underline{h}}^{\bar{h}} \sum_{j=0}^h \text{Tr}[q_1' \Psi_j' E_{ii} \Psi_j q_1] \\ &= q_1' S q_1, \end{aligned}$$

where

$$\begin{aligned} S &= \sum_{h=\underline{h}}^{\bar{h}} \sum_{j=0}^h \Psi_j' E_{ii} \Psi_j \\ &= \sum_{j=0}^{\bar{h}} (\bar{h} + 1 - \max(\underline{h}, j)) \Psi_j' E_{ii} \Psi_j \\ &= \sum_{j=0}^{\bar{h}} (\bar{h} + 1 - \max(\underline{h}, j)) \Psi_{j,i}' \Psi_{j,i} \end{aligned}$$

where $\Psi_{j,i}$ denotes row i of Ψ_j , i.e., the response of variable i at horizon j (when $Q = \text{Id}$).

The maximization problem subject to the side constraint $q_1' q_1 = 1$ can be written as a Lagrangian:

$$L = q_1' S q_1 - \lambda(q_1' q_1 - 1),$$

with the first-order condition $S q_1 = \lambda q_1$ (the side constraint is $q_1' q_1 = 1$). From this equation, we see that the solution q_1 is an eigenvector of S , the

one associated with eigenvalue λ . We also see that $\sigma^2(\underline{h}, \bar{h}, q_1) = \lambda$. Thus, to maximize this variance, we need to find the eigenvector of S that is associated with the maximal eigenvalue λ . That defines the first principal component (see XXXXXX). That is, if S admits the following spectral decomposition:

$$S = \mathcal{P}D\mathcal{P}',$$

where D is diagonal matrix whose entries are the (ordered) eigenvalues: $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$, then $\sigma^2(\underline{h}, \bar{h}, q_1)$ is maximized for $q_1 = p_1$, where p_1 is the first column of \mathcal{P} .

6.3.9 Identification based on non-normality of the shocks

In this section, we show that the non-identification of the structural shocks (η_t) is specific to the Gaussian case. We propose consistent estimation approaches for SVAR in the context of non-Gaussian shocks.

We have seen in what precedes that we cannot identify B based on first and second moments only. Since a Gaussian distribution is perfectly determined by the first two moments, it comes that one cannot achieve identification when the structural shocks are Gaussian. That is, even if we observe an infinite number of i.i.d. $B\eta_t$, we cannot recover B is the η_t 's are Gaussian.

Indeed, if $\eta_t \sim \mathcal{N}(0, Id)$, then the distribution of $\varepsilon_t \equiv B\eta_t$ is $\mathcal{N}(0, BB')$. Hence $\Omega = BB'$ is observed (in the population), but for any orthogonal matrix Q (i.e. $QQ' = Id$), we also have $BQ\eta_t \sim \mathcal{N}(0, \Omega)$.

To illustrate, consider the following bivariate Gaussian situations, with $\Theta_1 = 0$:

$$\begin{bmatrix} \eta_{1,t} \\ \eta_{2,t} \end{bmatrix} \sim \mathcal{N}(0, Id), \text{ with } B = \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \text{ and } Q = \begin{bmatrix} \cos(\pi/3) & -\sin(\pi/3) \\ \sin(\pi/3) & \cos(\pi/3) \end{bmatrix} \text{ (rotation).}$$

Figure 6.18 shows that the distributions of $B\eta_t$ and of $BQ\eta_t$ are identical. However, the impulse response functions associated with one of the other impulse matrix (B or BQ) are different. This is illustrated by Figure 6.19, that shows the IRFs associated with two identical models (defined by Eq. (6.38)), the only difference being the impulse matrix (B or BQ).

Hence, in the Gaussian case, external restrictions (economic hypotheses) are needed to identify B (see previous sections). But such restrictions may not be necessary if the structural shocks are not Gaussian. That is, the identification problem is very specific to normally-distributed η_t 's (Rigobon (2003), Normandin and Phaneuf (2004), Lanne and Lütkepohl (2008)).

To better see why this can be the case, consider again a bivariate vector of independent structural shocks ($\eta_{1,t}$ and $\eta_{2,t}$) but, now, assume that one of them is not Gaussian any more. Specifically, assume that $\eta_{2,t}$ is drawn from




Figure 6.18: This figure compares the distributions of two Gaussian bivariate vectors, $B\eta_t$ and $BQ\eta_t$, where $\eta_t \sim \mathcal{N}(0, Id)$ (therefore $\eta_{1,t}$ and $\eta_{2,t}$ are independent), and Q is an orthogonal matrix.




Figure 6.19: This figure shows that the impulse response functions associated with an impulse matrix equal to B (black line) or BQ (red line) are different (even if $BB' = BQ(BQ)'$).

a Student distribution with 5 degrees of freedom: $\eta_{1,t} \sim \mathcal{N}(0, 1)$, $\eta_{2,t} \sim t(5)$, $B = \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}$ and $Q = \begin{bmatrix} \cos(\pi/3) & -\sin(\pi/3) \\ \sin(\pi/3) & \cos(\pi/3) \end{bmatrix}$.

Figure 6.20 shows that, in this case, $B\eta_t$ and $BQ\eta_t$ do not have the same distribution any more (in spite of the fact that, in both cases, we have $\text{Var}(\varepsilon_t) = BB'$). This opens the door to the identification of the impulse matrix (BQ) in the non-Gaussian case.




Figure 6.20: This figure compares the distributions of two Gaussian bivariate vectors, $B\eta_t$ and $BQ\eta_t$, where $\eta_{1,t} \sim \mathcal{N}(0, 1)$, $\eta_{2,t} \sim t(5)$, and Q is an orthogonal matrix.

The exercise that consists in identifying non-Gaussian independent shocks out of linear combinations of these shocks is a well-known problem of the signal-processing literature, called **independent component analysis (ICA)**. Without loss of generality, we can assume that $BB' = Id$ (i.e. B is orthogonal). (If this is not the case, i.e. if $\text{Var}(\varepsilon_t) = \Omega \neq Id$, then one can pre-multiply the data by $\Omega^{-1/2}$.) The classical ICA problem is as follows: Find B such that $\varepsilon_t = B\eta_t$ (or $\$ _t = B' _t \$$) given that

- i. We observe the ε_t 's,
- ii. The components of η_t are independent,
- iii. $BB' = Id$ (i.e., B is orthogonal).

Figure 6.21 represents again some bivariate distributions. The black (red) lines correspond to the distributions of η_t ($B\eta_t$). It is important to note that the two components of vector $B\eta_t$ are not independent (contrary to those of η_t).

In all cases, we have $\text{Var}(\varepsilon_t) = \text{Var}(\eta_t) = Id$. But the two components of ε_t are not independent. For instance: We have $\mathbb{E}(\varepsilon_{2,t} | \varepsilon_{1,t} > 4) < 0$ (whereas




Figure 6.21: The three plots represent the bivariate distributions of η_t (black) and of $B\eta_t$ (red), where the two components of η_t are independent, of unit variance, and B is orthogonal. Hence, for each of the three plots, $\text{Var}(B\eta_t) = \text{Id}$.

$\mathbb{E}(\eta_{2,t}|\eta_{1,t} > 4) = 0$). The objective of ICA is to rotate ε_t to retrieve independent components (η_t).

Hypothesis 6.1. Process η_t satisfies:

- i. The η_t 's are i.i.d. (across time) with $\mathbb{E}(\eta_t) = 0$ and $\text{Var}(\eta_t) = \text{Id}$.
- ii. The components $\eta_{1,t}, \dots, \eta_{n,t}$ are mutually independent. iii We have

$$\boxed{\varepsilon_t = B_0 \eta_t,}$$

with $\text{Var}(\varepsilon_t) = \text{Id}$ (i.e. B_0 is orthogonal).

Theorem 6.4 (Eriksson, Koivunen (2004)). *If Hypothesis 6.1 is satisfied and if at most one of the components of η is Gaussian, then matrix B_0 is identifiable up to the post multiplication by DP , where P is a permutation matrix and D is a diagonal matrix whose diagonal entries are 1 or -1 .}*

Hence, the structural shocks are identifiable. But how to estimate them based on observations of the ε_t 's? Gouriéroux et al. (2017) have proposed a **Pseudo-Maximum Likelihood (PML)** approach. This approach consists in maximizing a so-called **pseudo log-likelihood function**, based on a set of p.d.f. $g_i(\eta_i)$, $i = 1, \dots, n$ (that may be different from the true p.d.f. of the $\eta_{i,t}$'s):

$$\log \mathcal{L}_T(B) = \sum_{t=1}^T \sum_{i=1}^n \log g_i(b'_i Y_t), \quad (6.59)$$

where b_i is the i^{th} column of matrix B (or b'_i is the i^{th} row of B^{-1} since $B^{-1} = B'$).

The log-likelihood function (6.59) is computed as if the errors $\eta_{i,t}$ had the p.d.f. $g_i(\eta_i)$. The PML estimator of matrix B maximizes the pseudo log-likelihood

function:

$$\widehat{B}_T = \arg \max_B \sum_{t=1}^T \sum_{i=1}^n \log g_i(b'_i \varepsilon_t), \quad (6.60)$$

s.t. $B'B = Id$.

The restrictions $B'B = Id$ can be eliminated by parameterizing B in such a way that, whatever the consider parameters, B is orthogonal. Gouriéroux et al. (2017) propose to use, for that, the Cayley's representation: any orthogonal matrix with no eigenvalue equal to -1 can be written as

$$B(A) = (Id + A)(Id - A)^{-1}, \quad (6.61)$$

where A is a skew symmetric (or antisymmetric) matrix, such that $A' = -A$. There is a one-to-one relationship with A , since:

$$A = (B(A) + Id)^{-1}(B(A) - Id). \quad (6.62)$$

Hence, the PML estimator of matrix B is obtained as $\widehat{B}_T = B(\widehat{A}_T)$, where:

$$\widehat{A}_T = \arg \max_{a_{i,j}, i > j} \sum_{t=1}^T \sum_{i=1}^n \log g_i[b_i(A)' \varepsilon_t]. \quad (6.63)$$

Table 6.1: This table reports usual p.d.f. and their derivatives.

$\log g(x)$	$\frac{d \log g(x)}{dx}$	$\frac{d^2 \log g(x)}{dx^2}$
Gaussian	$-x^2/2$	-1
Student	$-\frac{1-\nu}{2} \log \left(1 + \frac{x^2}{\nu-2}\right) - \frac{x(1+\nu)}{\nu-2+x^2}$	$-(1+\nu) \frac{\nu-2-x^2}{\nu-2+x^2}$
$t(\nu > 4)$		
Hyperbolist	$-\log \left(\cosh \left\{\frac{\pi}{2}x\right\}\right)$	$-\left(\frac{\pi}{2} \frac{1}{\cosh \left(\frac{\pi}{2}x\right)}\right)^2$
se-		
cant		
Subgaussian	$\pi x^2 + \log \left(\cosh \left\{\frac{\pi}{2}x\right\}\right)$	$2\pi + \left(\frac{\pi}{2} \frac{1}{\cosh \left(\frac{\pi}{2}x\right)}\right)^2$

Under assumptions on the g_i functions (excluding the Gaussian distributions), Gouriéroux et al. (2017) derive the asymptotic properties of the PML estimator. Specifically, the PML estimator \widehat{B}_T of B_0 is consistent (in \mathcal{P}_0 , the set of

matrices obtained by permutation and sign change of the columns of B_0) and asymptotically normal, with speed of convergence $1/\sqrt{T}$.

The asymptotic variance-covariance matrix of $\text{vec}\sqrt{T}(\widehat{B_T} - B_0)$ is $A^{-1} \begin{bmatrix} \Gamma & 0 \\ 0 & 0 \end{bmatrix} (A')^{-1}$, where matrices A and Γ are detailed in Gouriéroux et al. (2017).

Note that the potential misspecification of pseudo-distributions g_i has no effect on the consistency of these specific PML estimators.

Let us apply the PML-ICA approach on U.S. data covering the period 1959:IV to 2015:I at the quarterly frequency ($T = 224$). We consider three dependent variables: inflation (π_t), economic activity (z_t , the output gap) and the nominal short-term interest rate (r_t). Changes in the log of oil prices added as an exogenous variable (x_t).

```
library(AEC)
First.date <- "1959-04-01"
Last.date <- "2015-01-01"
data <- US3var
data <- data[(data$Date>=First.date)&(data$Date<=Last.date),]
Y <- as.matrix(data[c("infl","y.gdp.gap","r")])
names.var <- c("inflation","real activity","short-term rate")
T <- dim(Y)[1]
n <- dim(Y)[2]
```

Let us denote by W_t the set of information made of the past values of $y_t = [\pi_t, z_t, r_t]$, that is $\{y_{t-1}, y_{t-2}, \dots\}$, and of exogenous variables $\{x_t, x_{t-1}, \dots\}$. The reduced-form VAR model reads:

$$y_t = \mu + \underbrace{\sum_{i=1}^p \Phi_i y_{t-i} + \Theta x_t + u_t}_{a(W_t; \theta)}$$

where the u_t 's are assumed to be serially independent, with zero mean and variance-covariance matrix Σ .

Matrices μ , Φ_i , Θ and Σ are consistently estimated by OLS. Jarque-Bera tests support the hypothesis of non-normality for all residuals.

```
nb.lags <- 6 # number of lags used in the VAR model
X <- NULL
for(i in 1:nb.lags){
  lagged.Y <- rbind(matrix(NaN,i,n),Y[1:(T-i),])
  X <- cbind(X,lagged.Y)}
X <- cbind(X,data$commo) # add exogenous variables
Phi <- matrix(0,n,n*nb.lags);mu <- rep(0,n)
```

```

effect.commo <- rep(0,n)
U <- NULL # Eta is the matrix of OLS residuals
for(i in 1:n){
  eq <- lm(Y[,i] ~ X)
  Phi[i,] <- eq$coef[2:(dim(Phi)[2]+1)]
  mu[i] <- eq$coef[1]
  U <- cbind(U,eq$residuals)
  effect.commo[i] <- eq$coef[length(eq$coef)]}
}
Omega <- var(U) # Omega_1 is the covariance matrix of the OLS residuals.
B <- t(chol(Omega)) # B_1 is the Cholesky matrix associated with Omega_1 (lower triangular)
Eps <- U %*% t(solve(B)) # Recover associated structural shocks

```

We want to estimate the orthogonal matrix B such that $u_t = SB\eta_t$, where

- S results from the Cholesky decomposition of Σ and
- the components of η_t are independent, zero-mean with unit variance.

The PML approach is applied on standardized VAR residuals given by:

$$\hat{\varepsilon}_t = \hat{S}_T^{-1} \underbrace{[y_t - a(W_t; \hat{\theta}_T)]}_{\text{VAR residuals}}.$$

By construction of \hat{S}_T^{-1} , it comes that the covariance matrix of these residuals is Id .

The pseudo density functions are distinct and asymmetric mixtures of Gaussian distributions.

```

distri <- list(
  type=c("mixt.gaussian","mixt.gaussian","mixt.gaussian"),
  df=c(NaN,NaN,NaN),
  p=c(0.5,.5,.5),mu=c(.1,.1,.1),sigma=c(.5,.7,1.3))
AA.0 <- c(0,0,0)
res.optim <- optim(AA.0,func.2.minimize,
  Y = Eps, distri = distri,
  gr = d.func.2.minimize,
  method="Nelder-Mead",
  control=list(trace=FALSE,maxit=1000))
AA.0 <- res.optim$par
res.optim <- optim(AA.0,func.2.minimize,d.func.2.minimize,
  Y = Eps, distri = distri,
  method="BFGS",
  control=list(trace=FALSE))
AA.est <- res.optim$par

```

```

n <- ncol(Y)
M <- make.M(n)
A.est <- matrix(M %*% AA.est,n,n)
C.PML <- (diag(n) + A.est) %*% solve(diag(n) - A.est)

eta.PML <- Eps %*% C.PML # eta.PML are the ICA-estimated structural shocks

A <- make.A.matrix(eta.PML,distri,C.PML)
Omega <- make.Omega(eta.PML,distri)
# Compute asymptotic covariance matrix of C.PML:
V <- make.Asympt.Cov.delta(eta.PML,distri,C.PML)
param <- c(C.PML)
st.dev <- sqrt(diag(V))
t.stat <- c(C.PML)/sqrt(diag(V))
cbind(param,st.dev,t.stat) # print results of PML estimation

##           param      st.dev      t.stat
## [1,]  0.94417705  0.040848382 23.1141845
## [2,] -0.32711569  0.118802653 -2.7534376
## [3,]  0.03905164  0.074172945  0.5264944
## [4,]  0.32070293  0.119270893  2.6888616
## [5,]  0.93977707  0.041629110 22.5749976
## [6,]  0.11818924  0.060821400  1.9432179
## [7,] -0.07536139  0.071980455 -1.0469702
## [8,] -0.09906759  0.062185577 -1.5930959
## [9,]  0.99222290  0.007785691 127.4418551

```

(Note: it is always useful to combine two optimization algorithms, such as Nelder-Mead and BFGS.)

We would obtain close results by neglecting commodity prices. In that case, one can simply use the function `estim.SVAR.ICA` of the `AEC` package. Let us compare the C matrix obtained in the two cases (with or without commodity prices):

```

ICA.res.no.commo <- estim.SVAR.ICA(Y,distri = distri,p=6)
round(cbind(ICA.res.no.commo$C.PML,NaN,C.PML),3)

```

```

##      [,1]  [,2]  [,3]  [,4]  [,5]  [,6]  [,7]
## [1,]  0.956  0.287 -0.059  NaN   0.944  0.321 -0.075
## [2,] -0.292  0.950 -0.108  NaN  -0.327  0.940 -0.099
## [3,]  0.025  0.121  0.992  NaN   0.039  0.118  0.992

```

Once B has been estimated, it remains to label the resulting structural shocks (components of η_t). Postulated shocks are monetary-policy, supply, and demand shocks. This labelling can be based on the following considerations:

- Contractionary **monetary-policy shocks** have a negative impact on real activity and on inflation.
- **Supply shock** have influences of opposite signs on economic activity and on inflation.
- **Demand shock** have influences of same signs on economic activity and on inflation.

Let us compute the IRFs associated with the three structural shocks. (For the sake of comparison, the first line of plots shows the IRFs to a monetary-policy shock obtained from a Cholesky-based approach where the short-term rate is ordered last.)

```
IRF.Chol <- array(NaN,c(n,41,n))
IRF.ICA <- array(NaN,c(n,41,n))
PHI <- list();for(i in 1:nb.lags){PHI[[i]]<-array(Phi,c(3,3,nb.lags))[,i]}
for(jjjj in 1:n){
  u.shock <- rep(0,n)
  u.shock[jjjj] <- 1
  IRF.Chol[,,jjjj] <- t(simul.VAR(c=rep(0,3),Phi=PHI,B=B,nb.sim=41,
                                y0.star=rep(0,3*nb.lags),indic.IRF = 1,u.shock = u.shock))
  IRF.ICA[,,jjjj] <- t(simul.VAR(c=rep(0,3),Phi=PHI,B=B*%C.PML,nb.sim=41,
                                y0.star=rep(0,3*nb.lags),indic.IRF = 1,u.shock = u.shock))
}
```

According to Figure 6.22, Shock 1 is a supply shock, Shock 2 is a demand shock, and Shock 3 is a monetary-policy shock. Note that Shock 3 is close to the one resulting from the Cholesky approach.

Relation with the Heteroskedasticity Identification

In some cases, where the ε_t 's are heteroskedastic, the B matrix can be identified (Rigobon (2003), Lanne et al. (2010)).

Consider the case where we still have $\varepsilon_t = B\eta_t$ but where η_t 's variance conditionally depends on a regime $s_t \in \{1, \dots, M\}$. That is:

$$\text{Var}(\eta_{k,t}|s_t) = \lambda_{s_t,k} \quad \text{for } k \in \{1, \dots, n\}$$

Denoting by Λ_i the diagonal matrix whose diagonal entries are the $\lambda_{i,k}$'s, it comes that:

$$\text{Var}(\eta_t|s_t) = \Lambda_{s_t}, \quad \text{and} \quad \text{Var}(\varepsilon_t|s_t) = B\Lambda_{s_t}B'.$$

Without loss of generality, it can be assumed that $\Lambda_1 = Id$.

In this context, B is identified, apart from sign reversal of its columns if for all $k \neq j \in \{1, \dots, n\}$, there is a regime i s.t. $\lambda_{i,k} \neq \lambda_{i,j}$. (Prop.1 in @Lanne et al. (2010)).




Figure 6.22: The first row of plots shows the responses of the three endogenous variables to the monetary policy shock in the context of a Cholesky-identified SVAR (ordering: inflation, output gap, interest rate). The next three rows of plots show the responses of the endogenous variables to the three structural shocks identified by ICA. The last one (Shock 3) is close to the Cholesky-identified monetary policy shock.

Bivariate regime case ($M = 2$): B identified if the $\lambda_{2,k}$'s are all different. That is, identification is ensured if “there is sufficient heterogeneity in the volatility changes” (Lütkepohl and Netšunajev (2017)).

If the regimes s_t are exogenous and serially independent, then this situation is consistent with the “non-Gaussian” situation described above.

6.3.10 Factor-Augmented VAR (FAVAR)

VAR models are subject to the curse of dimensionality: If n , is large, then the number of parameters (in n^2) explodes.

In the case where one suspects that the $y_{i,t}$'s are mainly driven by a small number of random sources, a factor structure may be imposed, and **principal component analysis** (PCA, see Appendix 7.1) can be employed to estimate the relevant factors (Bernanke et al. (2005)).

Let us denote by F_t a k -dimensional vector of latent factors accounting for important shares of the variances of the $y_{i,t}$'s (with $K \ll n$) and by x_t is a small M -dimensional subset of y_t (with $M \ll n$). The following factor structure is posited:

$$y_t = \Lambda^f F_t + \Lambda^x x_t + e_t,$$

where the e_t are “small” serially and mutually i.i.d. error terms. That is F_t and x_t are supposed to drive most of the fluctuations of y_t 's components.

The model is complemented by positing a VAR dynamics for $[F'_t, x'_t]'$:

$$\begin{bmatrix} F_t \\ x_t \end{bmatrix} = \Phi(L) \begin{bmatrix} F_{t-1} \\ x_{t-1} \end{bmatrix} + v_t. \quad (6.64)$$

Standard identification techniques of structural shocks can be employed in Eq. (6.64): Cholesky approach can be used for instance if the last component of x_t is the short-term interest rate and if it is assumed that a MP shock has no contemporaneous impact on other macro-variables (in x_t).

In their identification procedure, Bernanke et al. (2005) exploit the fact that macro-finance variables can be decomposed in two sets —fast-moving and slow-moving variables— and that only the former reacts contemporaneously to monetary-policy shocks. Now, how to estimate the (unobserved) factors F_t ? Bernanke et al. (2005) note that the first $K + M$ PCA of the whole dataset (y_t), that they denote by $\hat{C}(F_t, x_t)$ should span the same space as F_t and x_t . To get an estimate of F_t , the dependence of $\hat{C}(F_t, x_t)$ in x_t has to be removed. This is done by regressing, by OLS, $\hat{C}(F_t, x_t)$ on x_t and on $\hat{C}^*(F_t)$, where the latter is an estimate of the common components other than x_t . To proxy for $\hat{C}^*(F_t)$, Bernanke et al. (2005) take principal components from the set of slow-moving variables, that are not comtemporaneously correlated to x_t . Vector \hat{F}_t is then

computed as $\hat{C}(F_t, x_t) - b_x x_t$, where b_x are the coefficients coming from the previous OLS regressions.

Note that this approach implies that the vectorial space spanned by (\hat{F}_t, x_t) is the same as that spanned by $\hat{C}(F_t, x_t)$.

Below, we employ this method on the dataset built by McCracken and Ng (2016)—the FRED:MD database—that includes 119 time series.




Figure 6.23: Responses of a monetary-policy shock. FAVAR approach of Bernanke, Boivin, and Eliasz (2005). FRED-MD dataset.

6.3.11 Projections Methods

Consider the infinite MA representation of y_t (Eq. (6.37)):

$$y_t = \mu + \sum_{h=0}^{\infty} \Psi_h \eta_{t-h}.$$

As seen in Section 6.3.2, the entries (i, j) of the sequence of the Ψ_h matrices define the IRF of $\eta_{j,t}$ on $y_{i,t}$.

Assume that you observe $\eta_{j,t}$, then a consistent estimate of $\Psi_{i,j,h}$ is simply obtained by the OLS regression of $y_{i,t+h}$ on $\eta_{j,t}$:

$$y_{i,t+h} = \mu_i + \Psi_{i,j,h} \eta_{j,t} + u_{i,j,t+h}. \quad (6.65)$$

Because the residuals $u_{i,j,t+h}$ are autocorrelated (for $h > 0$), estimates of the covariance of the OLS estimators of the $\Psi_{i,j,h}$ then have to be based on robust estimators (e.g. Newey-West, see Eq. (2.39)). This is the core idea of the **local projection approach** proposed by Jordà (2005).

Now, how to proceed in the (usual) case where $\eta_{j,t}$ is not observed? We consider two situations.

Situation A: Without IV

This corresponds to the original Jordà (2005)'s approach.

Assume that the structural shock of interest ($\eta_{1,t}$, say) can be consistently obtained as the residual of a regression of a variable x_t on a set of control variables w_t independent from $\eta_{1,t}$:

$$\eta_{1,t} = x_t - \mathbb{E}(x_t|w_t), \quad (6.66)$$

where $\mathbb{E}(x_t|w_t)$ is affine in w_t and where w_t is an affine transformation of $\eta_{2:n,t}$ and of past shocks $\eta_{t-1}, \eta_{t-2}, \dots$

Eq. (6.66) implies that, conditional on w_t , the additional knowledge of x_t is useful only when it comes to forecast something that depends on $\eta_{1,t}$. Hence, given that $u_{i,1,t+h}$ (see Eq. (6.65)) is independent from $\eta_{1,t}$ (it depends on $\eta_{t+h}, \dots, \eta_{t+1}, \eta_{2:n,t}, \eta_{t-1}, \eta_{t-2}, \dots$), it comes that

$$\mathbb{E}(u_{i,1,t+h}|x_t, w_t) = \mathbb{E}(u_{i,1,t+h}|w_t).$$

This is the *conditional mean independence* case.

Let's rewrite Eq. (6.65) as follows:

$$\begin{aligned} y_{i,t+h} &= \mu_i + \Psi_{i,1,h}\eta_{1,t} + u_{i,1,t+h} \\ &= \mu_i + \Psi_{i,1,h}x_t - \Psi_{i,1,h}\mathbb{E}(x_t|w_t) + u_{i,1,t+h}, \end{aligned}$$

What precedes implies that the expectation of the blue term, conditional on x_t and w_t , is linear in w_t . Standard results in the conditional mean independence case imply that the regression of $y_{i,t+h}$ on x_t , controlling for w_t , provides a consistent estimate of $\Psi_{i,1,h}$:

$$y_{i,t+h} = \alpha_i + \Psi_{i,1,h}x_t + \beta'w_t + v_{i,t+h}. \quad (6.67)$$

This is for instance consistent with the case where $[\Delta GDP_t, \pi_t, i_t]'$ follows a VAR(1) and the monetary-policy shock do not contemporaneously affect ΔGDP_t and π_t .

The IRFs can be estimated by LP, taking $x_t = i_t$ and $w_t = [\Delta GDP_t, \pi_t, \Delta GDP_{t-1}, \pi_{t-1}, i_{t-1}]'$.

This approach closely relates to the SVAR Cholesky-based identification approach. Specifically, if $w_t = [y_{1,t}, \dots, y_{k-1,t}, y'_{t-1}, \dots, y'_{t-p}]'$, with $k \leq n$, and

$x_t = y_{k,t}$, then this approach corresponds, for $h = 0$, to the SVAR(p) Cholesky-based IRF (focusing on the responses to the k^{th} structural shock). However, the two approaches differ for $h > 0$, because the LP methodology does not assumes a VAR dynamics for y_t .⁴

Situation B: IV approach

Consider now that we have a valid instrument z_t for $\eta_{1,t}$ (with $\mathbb{E}(z_t) = 0$). That is:

$$\begin{cases} (IV.i) \quad \mathbb{E}(z_t \eta_{1,t}) \neq 0 & \text{(relevance condition)} \\ (IV.ii) \quad \mathbb{E}(z_t \eta_{j,t}) = 0 \quad \text{for } j > 1 & \text{(exogeneity condition)} \end{cases} \quad (6.68)$$

The instrument z_t can be used to identify the structural shock. Eq. (6.68) implies that there exist $\rho \neq 0$ and a mean-zero variable ξ_t such that:

$$\eta_{1,t} = \rho z_t + \xi_t,$$

where ξ_t is correlated neither to z_t , nor to $\eta_{j,t}$, $j \geq 2$.

Proof. Define $\rho = \frac{\mathbb{E}(\eta_{1,t} z_t)}{\text{Var}(z_t)}$ and $\xi_t = \eta_{1,t} - \rho z_t$. It is easily seen that ξ_t satisfies the moment restrictions given above. \square

Ramey (2016) reviews the different approaches employed to construct monetary policy-shocks (the two main approaches are presented in 6.5 and 6.6 below). She has also collected time series of such shocks, see her website.

Example 6.5 (Identification of Monetary-Policy Shocks Based on High-Frequency Data). Instruments for monetary-policy shocks can be extracted from high-frequency market data associated with interest-rate products.

The quotes of all interest-rate-related financial products are sensitive to monetary-policy announcements. That is because these quotes mainly depends on investors' expectations regarding future short-term rates: $\mathbb{E}_t(i_{t+s})$. Typically, if agents were risk-neutral, the maturity- h interest rate would approximatively be given by:

$$i_{t,h} \approx \mathbb{E}_t \left(\frac{1}{h} \int_0^h i_{t+s} ds \right) = \frac{1}{h} \int_0^h \mathbb{E}_t(i_{t+s}) ds.$$

In general, changes in $\mathbb{E}_t(i_{t+s})$, for $s > 0$, can be affected by all types of shocks that may trigger a reaction by the central bank.

However, if a MP announcement takes place between t and $t + \epsilon$, then most of $\mathbb{E}_{t+\epsilon}(i_{t+s}) - \mathbb{E}_t(i_{t+s})$ is to be attributed to the MP shock (see Figure 6.24, from

⁴This is reminiscent of the distinction between direct forecasting –based on regressions of y_{t+h} on $\{y_t, y_{t-1}, \dots\}$ – and iterated forecasting –based on a recursive model where $y_{t+1} = g(y_t, y_{t-1}, \dots) + \varepsilon_{t+1}$, see Marcellino et al. (2006).

Gürkaynak et al. (2005)). Hence, a monthly time series of MP shocks can be obtained by summing, over each month, the changes $i_{t+\epsilon,h} - i_{t,h}$ associated with a given interest rate (T-bills, futures, swaps) and a given maturity h .

See among others: Kuttner (2001), Cochrane and Piazzesi (2002), Gürkaynak et al. (2005), Piazzesi and Swanson (2008), Gertler and Karadi (2015).

Figure 1. Intraday Trading in Federal Funds Futures Contracts




Figure 6.24: Source: Gurkaynak, Sack and Swanson (2005). Transaction rates of Federal funds futures on June 25, 2003, day on which a regularly scheduled FOMC meeting was scheduled. At 2:15 p.m., the FOMC announced that it was lowering its target for the federal funds rate from 1.25% to 1%, while many market participants were expecting a 50 bp cut. This shows that (i) financial markets seem to fully adjust to the policy action within just a few minutes and (ii) the federal funds rate surprise is not necessarily in the same direction as the federal funds rate action itself.

Example 6.6 (Identification of Monetary-Policy Shocks Based on the Narrative Approach). Romer and Romer (2004) propose a two-step approach:

- derive a series for Federal Reserve intentions for the federal funds rate (the explicit target of the Fed) around FOMC meetings,
- control for Federal Reserve forecasts.

This gives a measure of intended monetary policy actions not driven by information about future economic developments. a. “intentions” are measured as a combination of narrative and quantitative evidence. Sources: (among others) Minutes of FOMC and “Blue Books”. b. Controls = variables spanning the information the Federal Reserve has about future developments. Data: Federal Reserve’s internal forecasts (inflation, real output and unemployment), “Green-book’s forecasts” – usually issued 6 days before the FOMC meeting.

The shock measure is the residual series in the linear regression of (a) on (b).

There are two main IV approaches to estimate IRFs see Stock and Watson (2018):

- a. The LP-IV approach, where y_t 's DGP is left unspecified,
- b. The SVAR-IV approach.

The LP-IV approach is based on a set of IV regressions (for each variable of interest, one for each forecast horizon). The SVAR-IV approach is based on IV regressions of VAR innovations only (one for each series of VAR innovations).

If the VAR adequately captures the DGP, then the IV-SVAR is optimal for all horizons. However, if the VAR is misspecified, then specification errors are compounded at each horizon and a local projection method would lead to better results.

Situation B.1: SVAR-IV approach

Assume you have consistent estimates of $\varepsilon_t = B\eta_t$, these estimates ($\hat{\varepsilon}_t$) coming from the estimation of a VAR model. You have, for $i \in \{1, \dots, n\}$:

$$\begin{aligned}\varepsilon_{i,t} &= b_{i,1}\eta_{1,t} + u_{i,t} (\#eq : eps_rho) \\ &= b_{i,1}\rho z_t + \underbrace{b_{i,1}\xi_t + u_{i,t}}_{\perp z_t}.\end{aligned}\tag{6.69}$$

($u_{i,t}$ is a linear combination of the $\eta_{j,t}$'s, $j \geq 2$).

Hence, up to a multiplicative factor (ρ), the (OLS) regressions of the $\hat{\varepsilon}_{i,t}$'s on z_t provide consistent estimates of the $b_{i,1}$'s.

Combined with the estimated VAR (the Φ_k matrices), this provides consistent estimates of the IRFs of $\eta_{1,t}$ on y_t , though up to a multiplicative factor. This scale ambiguity can be solved by rescaling the structural shock (“unit-effect normalisation”, see Stock and Watson (2018)). Let us consider $\tilde{\eta}_{1,t} = b_{1,1}\eta_{1,t}$; by construction, $\tilde{\eta}_{1,t}$ has a one-unit contemporaneous effect on $y_{1,t}$. Denoting by $\tilde{B}_{i,1}$ the contemporaneous impact of $\tilde{\eta}_{1,t}$, we get:

$$\tilde{B}_1 = \frac{1}{b_{1,1}}B_1,$$

where B_1 denotes the 1st column of B and $\tilde{B}_1 = [1, \tilde{B}_{2,1}, \dots, \tilde{B}_{n,1}]'$.

Eq. @ref(eq:eps_rho) gives:

$$\begin{aligned}\varepsilon_{1,t} &= \tilde{\eta}_{1,t} + u_{1,t} \\ \varepsilon_{i,t} &= \tilde{B}_{i,1}\tilde{\eta}_{1,t} + u_{i,t}.\end{aligned}$$

This suggests that $\tilde{B}_{i,1}$ can be estimated by regressing $\varepsilon_{i,t}$ on $\varepsilon_{1,t}$, using z_t as an instrument.

What about inference? Once cannot use the usual TSLS standard deviations because the $\varepsilon_{i,t}$'s are not directly observed. Bootstrap procedures can be resorted

to. Stock and Watson (2018) propose, in particular, a Gaussian parametric bootstrap:

Assume you have estimated $\{\widehat{\Phi}_1, \dots, \widehat{\Phi}_p, \widehat{B}_1\}$ using the SVAR-IV approach based on a size- T sample. Generate N (where N is large) size- T samples from the following VAR:

$$\begin{bmatrix} \widehat{\Phi}(L) & 0 \\ 0 & \widehat{\rho}(L) \end{bmatrix} \begin{bmatrix} y_t \\ z_t \end{bmatrix} = \begin{bmatrix} \varepsilon_t \\ e_t \end{bmatrix},$$

where $\begin{bmatrix} \varepsilon_t \\ e_t \end{bmatrix} \sim i.i.d. \mathcal{N} \left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \Omega & S'_{\varepsilon,e} \\ S_{\varepsilon,e} & \sigma_e^2 \end{bmatrix} \right)$,

where $\widehat{\rho}(L)$ and σ_e^2 result from the estimation of an AR process for z_t , and where Ω and $S_{\varepsilon,e}$ are sample covariances for the VAR/AR residuals.

For each simulated sample (of \tilde{y}_t and \tilde{z}_t , say), estimate $\{\widetilde{\Phi}_1, \dots, \widetilde{\Phi}_p, \widetilde{B}_1\}$ and associated $\widetilde{\Psi}_{i,1,h}$. This provides e.g. a sequence of N estimates of $\Psi_{i,1,h}$, from which quantiles and conf. intervals can be deduced.

```
# Load vars package:
library(vars)
library(AEC)
data("USmonthly")
First.date <- "1990-05-01"
Last.date <- "2012-6-01"
indic.first <- which(USmonthly$DATES==First.date)
indic.last <- which(USmonthly$DATES==Last.date)
USmonthly <- USmonthly[indic.first:indic.last,]
shock.name <- "FF4_TC" # "FF4_TC", "ED2_TC", "ff1_vr", "rrshock83b"
indic.shock.name <- which(names(USmonthly)==shock.name)
Z <- matrix(USmonthly[,indic.shock.name],ncol=1)
plot(USmonthly$DATES,Z,type="l",xlab="",ylab="")

considered.variables <- c("GS1","LIP","LCPI","EBP")
y <- as.matrix(USmonthly[,considered.variables])
n <- length(considered.variables)
colnames(y) <- considered.variables
res.svar.iv <- svar.iv(y,Z,p = 4,
                        names.of.variables=considered.variables,
                        nb.periods.IRF = 20,
                        z.AR.order=1, # This is used in the parametric bootstrap only
                        nb.bootstrap.replications = 100, # This is used in the parametric bootstrap
                        confidence.interval = 0.90, # This is used in the parametric bootstrap only
                        indic.plot=1)
```

Situation B.2: LP-IV




Figure 6.25: Gertler-Karadi monthly shocks, fed funds futures 3 months.




Figure 6.26: Reponses to a monetary-policy shock, SVAR-IV approach.

If you do not want to posit a VAR-type dynamics for y_t –e.g. because you suspect that the true generating model may be a non-invertible VARMA model– you can directly proceed by IV-projection methods to obtain the $\tilde{\Psi}_{i,1,h} \equiv \Psi_{i,1,h}/b_{1,1}$ (that are the IRFs of $\tilde{\eta}_{1,t}$ on $y_{i,t}$).

However, Assumptions (IV.i) and (IV.ii) (Eq. (6.68)) have to be complemented with (IV.iii):

$$(IV.iii) \quad \mathbb{E}(z_t \eta_{j,t+h}) = 0 \text{ for } h \neq 0 \quad (\text{lead-lag exogeneity})$$

When (IV.i), (IV.ii) and (IV.iii) are satisfied, $\tilde{\Psi}_{i,1,h}$ can be estimated by regressing $y_{i,t+h}$ on $y_{1,t}$, using z_t as an instrument, i.e. by considering the TSLS estimation of:

$$y_{i,t+h} = \alpha_i + \tilde{\Psi}_{i,1,h} y_{1,t} + \nu_{i,t+h}, \quad (6.70)$$

where $\nu_{i,t+h}$ is correlated to $y_{1,t}$, but not to z_t .

We have indeed:

$$\begin{aligned} y_{1,t} &= \alpha_1 + \tilde{\eta}_{1,t} + v_{1,t} \\ y_{i,t+h} &= \alpha_i + \tilde{\Psi}_{i,1,h} \tilde{\eta}_{1,t} + v_{i,t+h}, \end{aligned}$$

where the $v_{i,t+h}$'s are uncorrelated to z_t under (IV.i), (IV.ii) and (IV.iii).

Note again that, for $h > 0$, the $v_{i,t+h}$ (and $\nu_{i,t+h}$) are auto-correlated. Newey-West corrections therefore have to be used to compute std errors of the $\tilde{\Psi}_{i,1,h}$'s estimates.

Consider the linear regression:

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon,$$

where $\mathbb{E}(\varepsilon) = 0$, but where the explicative variables \mathbf{X} are supposed to be correlated to the residuals ε .

Moreover, the ε are supposed to be possibly heteroskedastic and auto-correlated.

We consider the instruments \mathbf{Z} , with $\mathbb{E}(\mathbf{X}'\mathbf{Z}) \neq 0$ but $\mathbb{E}(\varepsilon'\mathbf{Z}) = 0$.

The IV estimator of β is obtained by regressing $\hat{\mathbf{Y}}$ on $\hat{\mathbf{X}}$, where $\hat{\mathbf{Y}}$ and $\hat{\mathbf{X}}$ are the respective residuals of the regressions of \mathbf{Y} and \mathbf{X} on \mathbf{Z} .

$$\begin{aligned} \mathbf{b}_{iv} &= [\mathbf{X}'\mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{X}]^{-1}\mathbf{X}'\mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{Y} \\ \mathbf{b}_{iv} &= \beta + \frac{1}{\sqrt{T}} \underbrace{T[\mathbf{X}'\mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{X}]^{-1}\mathbf{X}'\mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}}_{=Q(\mathbf{X}, \mathbf{Z}) \xrightarrow{p} \mathbf{Q}_{xz}} \underbrace{\sqrt{T} \left(\frac{1}{T} \mathbf{Z}'\varepsilon \right)}_{\xrightarrow{d} \mathcal{N}(0, S)}, \end{aligned}$$

where \mathbf{S} is the long-run variance of $\mathbf{z}_t\varepsilon_t$ (see next slide).

The asymptotic covariance matrix of $\sqrt{T}\mathbf{b}_{iv}$ is $\mathbf{Q}_{xz}\mathbf{S}\mathbf{Q}'_{xz}$.

The covariance matrix of \mathbf{b}_{iv} can be approximated by $\frac{1}{T}Q(\mathbf{X}, \mathbf{Z})\hat{\mathbf{S}}Q(\mathbf{X}, \mathbf{Z})'$ where $\hat{\mathbf{S}}$ is the Newey-West estimator of \mathbf{S} (see Eq. (2.39))

(IV.iii) is usually not restrictive for $h > 0$ (z_t is usually not affected by future shocks). By contrast, it may be restrictive for $h < 0$. This can be solved by adding controls in Regression (6.70). These controls should span the space of $\{\eta_{t-1}, \eta_{t-2}, \dots\}$.

If z_t is suspected to be correlated to past values of $\eta_{1,t}$ but not to the $\eta_{j,t}$'s, $j > 1$, then one can add lags of z_t as controls (method e.g. advocated by Ramey, 2016, p.108, considering the instrument by Gertler and Karadi (2015)).

In the general case, one can use lags of y_t as controls. Note that, even if (IV.iii) holds, adding controls may reduce the variance of the regression error.

As noted by Stock and Watson (2018), the relevant variance is the long-run variance of the instrument-times-error term. They also recommend (p.926) using leads and lags of z_t to improve efficiency.

```
res.LP.IV <- make.LPIV.irf(y,Z,
                           nb.periods.IRF = 20,
                           nb.lags.Y.4.control=4,
                           nb.lags.Z.4.control=4,
                           indic.plot = 1, # Plots are displayed if = 1.
                           confidence.interval = 0.90)
```

6.3.12 Inference

Consider the following SVAR model:

$$y_t = \Phi_1 y_{t-1} + \dots + \Phi_p y_{t-p} + \varepsilon_t$$

with $\varepsilon_t = B\eta_t$, $\Omega_\varepsilon = BB'$.

The corresponding infinite MA representation (Eq. (6.37), or Wold theorem, Theorem 6.3) is:

$$y_t = \sum_{h=0}^{\infty} \Psi_h \eta_{t-h},$$

where $\Psi_0 = B$ and for $h = 1, 2, \dots$:

$$\Psi_h = \sum_{j=1}^h \Psi_{h-j} \Phi_j,$$

with $\Phi_j = 0$ for $j > p$ (see Prop. 6.8 for this recursive computation of the Ψ_j 's).

Inference on the VAR coefficients $\{\Phi_j\}_{j=1,\dots,p}$ is straightforward (standard OLS inference). But inference is more complicated regarding IRF. Indeed, as shown

by the previous equation, the (infinite) MA coefficients $\{\Psi_j\}_{j=1,\dots}$ are non-linear functions of the $\{\Phi_j\}_{j=1,\dots,p}$ and of Ω_ε . An other issue pertain to small sample bias: typically, for persistent process, auto-regressive parameters are known to be downward biased.

The main inference methods are the following:

- Monte Carlo method (Hamilton (1994))
- Asymptotic normal approximation (Lütkepohl (1990)), or Delta method
- Bootstrap method (Kilian_1998)

Monte Carlo method

We use Monte Carlo when we need to approximate the distribution of a variable whose distribution is unknown (here: the Ψ_j 's) but which is a function of another variable whose distribution is known (here, the Φ_j 's).

For instance, suppose we know the distribution of a random variable X , which takes values in \mathbb{R} , with density function p . Assume we want to compute the mean of $\varphi(X)$. We have:

$$\mathbb{E}(\varphi(X)) = \int_{-\infty}^{+\infty} \varphi(x)p(x)dx$$

Suppose that the above integral does not have a simple expression. We cannot compute $\mathbb{E}(\varphi(X))$ but, by virtue of the law of large numbers (Theorem ??), we can approximate it as follows:

$$\mathbb{E}(\varphi(X)) \approx \frac{1}{N} \sum_{i=1}^N \varphi(X^{(i)}),$$

where $\{X^{(i)}\}_{i=1,\dots,N}$ are N independent draws of X . More generally, the distribution of $\varphi(X)$ can be approximated by the empirical distribution of the $\varphi(X^{(i)})$'s. Typically, if 10'000 values of $\varphi(X^{(i)})$ are drawn, the 5th percentile of the p.d.f. of $\varphi(X)$ can be approximated by the 500th value of the 10'000 draws of $\varphi(X^{(i)})$ (after arranging these values in ascending order).

As regards the computation of confidence intervals around IRFs, one has to think of $\{\widehat{\Phi}_j\}_{j=1,\dots,p}$, and of $\widehat{\Omega}$ as X and $\{\widehat{\Psi}_j\}_{j=1,\dots}$ as $\varphi(X)$. (Proposition 6.14 provides us with the asymptotic distribution of the “ X .”)

To summarize, here are the steps one can implement to derive confidence intervals for the IRFs using the Monte-Carlo approach:

For each iteration k :

1. Draw $\{\widehat{\Phi}_j^{(k)}\}_{j=1,\dots,p}$ and $\widehat{\Omega}^{(k)}$ from their asymptotic distribution (using Proposition 6.14).

2. Compute the matrix $B^{(k)}$ so that $\widehat{\Omega}^{(k)} = B^{(k)}B^{(k)'}_t$, according to your identification strategy.
3. Compute the associated IRFs $\{\widehat{\Psi}_j\}^{(k)}$.

Perform N replications and report the median impulse response (and its confidence intervals).

Delta method

Suppose β is a vector of parameters and $\hat{\beta}$ is an estimator such that

$$\sqrt{T}(\hat{\beta} - \beta) \xrightarrow{d} \mathcal{N}(0, \Sigma_\beta),$$

where d denotes convergence in distribution, $\mathcal{N}(0, \Sigma_\beta)$ denotes the multivariate normal distribution with mean vector 0 and covariance matrix Σ_β and T is the sample size used for estimation.

Let $g(\beta) = (g_1(\beta), \dots, g_m(\beta))'$ be a continuously differentiable function with values in \mathbb{R}^m , and assume that $\partial g_i / \partial \beta' = (\partial g_i / \partial \beta_j)$ is nonzero at β for $i = 1, \dots, m$. Then

$$\sqrt{T}(g(\hat{\beta}) - g(\beta)) \xrightarrow{d} \mathcal{N}\left(0, \frac{\partial g}{\partial \beta'} \Sigma_\beta \frac{\partial g'}{\partial \beta}\right).$$

(This formula underlies the Delta method, see Eq. (4.15).)

Using this property, Lütkepohl (1990) provides the asymptotic distributions of the Ψ_j 's.

A limit of the last two approaches (Monte Carlo and the Delta method) is that they critically rely on asymptotic results. Bootstrapping approaches are more robust in small-sample situations.

Bootstrap

IRFs' confidence intervals are intervals where 90% (or 95%, 75%, ...) of the IRFs would lie, if we were to repeat the estimation a large number of times in similar conditions (T observations). We obviously cannot do this, because we have only one sample: $\{y_t\}_{t=1,\dots,T}$. But we can try to *construct* such samples.

Bootstrapping consists in:

- re-sampling N times, i.e., constructing N samples of T observations, using the estimated VAR coefficients and
 - a. a sample of residuals from the distribution $N(0, BB')$ (**parametric approach**), or
 - b. a sample of residuals drawn randomly from the set of the actual estimated residuals $\{\hat{\varepsilon}_t\}_{t=1,\dots,T}$. (**non-parametric approach**).
- re-estimating the SVAR N times.

Here is the algorithm:

1. Construct a sample

$$y_t^{(k)} = \widehat{\Phi}_1 y_{t-1}^{(k)} + \dots + \widehat{\Phi}_p y_{t-p}^{(k)} + \widehat{\varepsilon}_t^{(k)},$$

with $\widehat{\varepsilon}_t^{(k)} = \widehat{\varepsilon}_{s_t^{(k)}}$, where $\{s_1^{(k)}, \dots, s_T^{(k)}\}$ is a random set from $\{1, \dots, T\}^T$.

2. Re-estimate the SVAR and compute the IRFs $\{\widehat{\Psi}_j\}^{(k)}$.

Perform N replications and report the median impulse response (and its confidence intervals).

Bootstrap-after-bootstrap (Kilian (1998))

The previous simple bootstrapping procedure deals with non-normality and small sample distribution, since we use the actual residuals. However, it does not deal with the *small sample bias*, stemming, in particular, from small-sample bias associated with OLS coefficient estimates $\{\widehat{\Phi}_j\}_{j=1,\dots,p}$. The main idea of the bootstrap-after-bootstrap of Kilian (1998) is to run two consecutive bootstraps: the objective of the first is to compute the bias, which can further be used to correct the initial estimates of the Φ_i 's. Further, these corrected estimates are used—in the second bootstrap—to compute a set of IRFs (as in the standard bootstrap).

More formally, the algorithm is as follows:

1. Estimate the SVAR coefficients $\{\widehat{\Phi}_j\}_{j=1,\dots,p}$ and $\widehat{\Omega}$

2. **First bootstrap.** For each iteration k :

- a. Construct a sample

$$y_t^{(k)} = \widehat{\Phi}_1 y_{t-1}^{(k)} + \dots + \widehat{\Phi}_p y_{t-p}^{(k)} + \widehat{\varepsilon}_t^{(k)},$$

with $\widehat{\varepsilon}_t^{(k)} = \widehat{\varepsilon}_{s_t^{(k)}}$, where $\{s_1^{(k)}, \dots, s_T^{(k)}\}$ is a random set from $\{1, \dots, T\}^T$.

- b. Re-estimate the VAR and compute the coefficients $\{\widehat{\Phi}_j\}_{j=1,\dots,p}^{(k)}$.

3. Perform N replications and compute the median coefficients $\{\widehat{\Phi}_j\}_{j=1,\dots,p}^*$.

4. Approximate the bias terms by $\widehat{\Theta}_j = \widehat{\Phi}_j^* - \widehat{\Phi}_j$.

5. Construct the bias-corrected terms $\widetilde{\Phi}_j = \widehat{\Phi}_j - \widehat{\Theta}_j$.

6. **Second bootstrap.** For each iteration k :

- a. Construct a sample now from

$$y_t^{(k)} = \widetilde{\Phi}_1 y_{t-1}^{(k)} + \dots + \widetilde{\Phi}_p y_{t-p}^{(k)} + \widehat{\varepsilon}_t^{(k)}.$$

- b. Re-estimate the VAR and compute the coefficients $\{\widehat{\Phi}_j^*\}_{j=1,\dots,p}^{(k)}$.

- c. Construct the bias-corrected estimates $\tilde{\Phi}_j^{*(k)} = \widehat{\Phi}_j^{*(k)} - \widehat{\Theta}_j$.
 - d. Compute the associated IRFs $\{\tilde{\Psi}_j^{*(k)}\}_{j \geq 1}$.
7. Perform N replications and compute the median and the confidence interval of the set of IRFs.

It should be noted that correcting for the bias can generate non-stationary results ($\tilde{\Phi}$ with eigenvalue with modulus > 1). Solution (Kilian (1998)):

In step 5, check if the largest eigenvalue of $\tilde{\Phi}$ is of modulus < 1 . If not, shrink the bias: for all j s, set $\widehat{\Theta}_j^{(i+1)} = \delta_{i+1} \widehat{\Theta}_j^{(i)}$, with $\delta_{i+1} = \delta_i - 0.01$, starting with $\delta_1 = 1$ and $\widehat{\Theta}_j^{(1)} = \widehat{\Theta}_j$, and compute $\tilde{\Phi}_j^{(i+1)} = \widehat{\Phi}_j - \widehat{\Theta}_j^{(i+1)}$ until the largest eigenvalue of $\tilde{\Phi}^{(i+1)}$ has modulus < 1 .

Function `VAR.Boot` of package `VAR.ftp` (Kim (2022)) can be used to operate the bias-correction approach of Kilian (1998):

```
library(VAR.ftp)
library(vars) #standard VAR models
data(dat) # part of VAR.ftp package
corrected <- VAR.Boot(dat,p=2,nb=200,type="const")
noncorrec <- VAR(dat,p=2)
rbind(corrected$coef[,],
      (corrected$coef+corrected$Bias)[1,],
      noncorrec$varresult$inv$coefficients)

##           inv(-1)   inc(-1)   con(-1)   inv(-2)   inc(-2)   con(-2)       const
## [1,] -0.3380188 0.09704455 1.034321 -0.1513048 0.1205532 0.9948151 -0.01803010
## [2,] -0.3196310 0.14598883 0.961219 -0.1605511 0.1146050 0.9343938 -0.01672199
## [3,] -0.3196310 0.14598883 0.961219 -0.1605511 0.1146050 0.9343938 -0.01672199
```

6.4 Forecasting

Forecasting has always been an important part of the time series field (De Gooijer and Hyndman (2006)). Macroeconomic forecasts are done in many places: Public Administration (notably Treasuries), Central Banks, International Institutions (e.g. IMF, OECD), banks, big firms. These institutions are interested in the **point estimates** (\sim most likely value) of the variable of interest. They also sometimes need to measure the **uncertainty** (\sim dispersion of likely outcomes) associated to the point estimates.⁵

Forecasts produced by professional forecasters are available on these web pages:

⁵In its inflation report, the Bank of England displays charts showing the conditional distribution of future inflation, called fan charts. This fan charts show the uncertainty associated with future inflation. See this page.

- Philly Fed Survey of Professional Forecasters.
- ECB Survey of Professional Forecasters.
- IMF World Economic Outlook.
- OECD Global Economic Outlook.
- European Commission Economic Forecasts.

How to formalize the forecasting problem? Assume the current date is t . We want to forecast the value that variable y_t will take on date $t + 1$ (i.e., y_{t+1}) based on the observation of a set of variables gathered in vector x_t (x_t may contain lagged values of y_t).

The forecaster aims at minimizing (a function of) the forecast error. It is usual to consider the following (quadratic) loss function:

$$\underbrace{\mathbb{E}([y_{t+1} - y_{t+1}^*]^2)}_{\text{Mean square error (MSE)}}$$

where y_{t+1}^* is the forecast of y_{t+1} (function of x_t).

Proposition 6.15 (Smallest MSE). *The smallest MSE is obtained with MSE the expectation of y_{t+1} conditional on x_t .*

Proof. See Appendix 7.5. □

Proposition 6.16. *Among the class of linear forecasts, the smallest MSE is obtained with the linear projection of y_{t+1} on x_t . This projection, denoted by $\hat{P}(y_{t+1}|x_t) := \alpha' x_t$, satisfies:*

$$\mathbb{E}([y_{t+1} - \alpha' x_t] x_t) = \mathbf{0}. \quad (6.71)$$

Proof. Consider the function $f : \alpha \rightarrow \mathbb{E}([y_{t+1} - \alpha' x_t]^2)$. We have:

$$f(\alpha) = \mathbb{E}(y_{t+1}^2 - 2y_t x_t' \alpha + \alpha' x_t x_t' \alpha).$$

We have $\partial f(\alpha)/\partial \alpha = \mathbb{E}(-2y_t x_t + 2x_t x_t' \alpha)$. The function is minimised for $\partial f(\alpha)/\partial \alpha = 0$. □

Eq. (6.71) implies that $\mathbb{E}(y_{t+1} x_t) = \mathbb{E}(x_t x_t' \alpha)$. (Note that $x_t x_t' \alpha = x_t (x_t' \alpha) = (\alpha' x_t) x_t'$.)

Hence, if $\mathbb{E}(x_t x_t')$ is nonsingular,

$$\alpha = [\mathbb{E}(x_t x_t')]^{-1} \mathbb{E}(y_{t+1} x_t). \quad (6.72)$$

The MSE then is:

$$\mathbb{E}([y_{t+1} - \alpha' x_t]^2) = \mathbb{E}(y_{t+1}^2) - \mathbb{E}(y_{t+1} x_t) [\mathbb{E}(x_t x_t')]^{-1} \mathbb{E}(x_t y_{t+1}).$$

Consider the regression $y_{t+1} = \beta' \mathbf{x}_t + \varepsilon_{t+1}$. The OLS estimate is:

$$\mathbf{b} = \left[\underbrace{\frac{1}{T} \sum_{i=1}^T \mathbf{x}_t \mathbf{x}'_t}_{\mathbf{m}_1} \right]^{-1} \left[\underbrace{\frac{1}{T} \sum_{i=1}^T \mathbf{x}'_t y_{t+1}}_{\mathbf{m}_2} \right].$$

If $\{x_t, y_t\}$ is covariance-stationary and ergodic for the second moments then the sample moments (\mathbf{m}_1 and \mathbf{m}_2) converges in probability to the associated population moments and $\mathbf{b} \xrightarrow{p} \alpha$ (where α is defined in Eq. (6.72)).

Example 6.7 (Forecasting an MA(q) process). Consider the MA(q) process:

$$y_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q},$$

where $\{\varepsilon_t\}$ is a white noise sequence (Def. 6.1).

We have:

$$\begin{aligned} \mathbb{E}(y_{t+h} | \varepsilon_t, \varepsilon_{t-1}, \dots) &= \\ \begin{cases} \mu + \theta_h \varepsilon_t + \cdots + \theta_q \varepsilon_{t-q+h} & \text{for } h \in [1, q] \\ \mu & \text{for } h > q \end{cases} \end{aligned}$$

and

$$\begin{aligned} \text{Var}(y_{t+h} | \varepsilon_t, \varepsilon_{t-1}, \dots) &= \mathbb{E}([y_{t+h} - \mathbb{E}(y_{t+h} | \varepsilon_t, \varepsilon_{t-1}, \dots)]^2) = \\ \begin{cases} \sigma^2(1 + \theta_1^2 + \cdots + \theta_{h-1}^2) & \text{for } h \in [1, q] \\ \sigma^2(1 + \theta_1^2 + \cdots + \theta_q^2) & \text{for } h > q. \end{cases} \end{aligned}$$

Remark: The previous reasoning relies on the assumption that the ε_t s are observed. But this is generally not the case in practice. Note that consistent estimates are available if the MA process is invertible (see Eq. (6.32)).

Example 6.8 (Forecasting an AR(p) process). (See this web interface.) Consider the AR(p) process:

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + \varepsilon_t,$$

where $\{\varepsilon_t\}$ is a white noise sequence (Def. 6.1).

Using the notation of Eq. (6.9), we have:

$$\mathbf{y}_t - \mu = F(\mathbf{y}_{t-1} - \mu) + \xi_t,$$

with $\mu = [\mu, \dots, \mu]'$ (μ is defined in Eq. (6.13)). Hence:

$$\mathbf{y}_{t+h} - \mu = \xi_{t+h} + F\xi_{t+h-1} + \cdots + F^{h-1}\xi_{t+1} + F^h(\mathbf{y}_t - \mu).$$

Therefore:

$$\begin{aligned}\mathbb{E}(\mathbf{y}_{t+h}|y_t, y_{t-1}, \dots) &= \mu + F^h(\mathbf{y}_t - \mu) \\ \text{Var}([\mathbf{y}_{t+h} - \mathbb{E}(\mathbf{y}_{t+h}|y_t, y_{t-1}, \dots)]) &= \Sigma + F\Sigma F' + \dots + F^{h-1}\Sigma(F^{h-1})',\end{aligned}$$

where:

$$\Sigma = \begin{bmatrix} \sigma^2 & 0 & \dots \\ 0 & 0 & \ddots \\ \vdots & & \ddots \end{bmatrix}.$$

Alternative approach: Taking the (conditional) expectations of both sides of

$$y_{t+h} - \mu = \phi_1(y_{t+h-1} - \mu) + \phi_2(y_{t+h-2} - \mu) + \dots + \phi_p(y_{t-p} - \mu) + \varepsilon_{t+h},$$

we obtain:

$$\begin{aligned}\mathbb{E}(y_{t+h}|y_t, y_{t-1}, \dots) &= \mu + \phi_1(\mathbb{E}[y_{t+h-1}|y_t, y_{t-1}, \dots] - \mu) + \\ &\quad \phi_2(\mathbb{E}[y_{t+h-2}|y_t, y_{t-1}, \dots] - \mu) + \dots + \\ &\quad \phi_p(\mathbb{E}[y_{t+h-p}|y_t, y_{t-1}, \dots] - \mu),\end{aligned}$$

which can be exploited recursively.

The recursion begins with $\mathbb{E}(y_{t-k}|y_t, y_{t-1}, \dots) = y_{t-k}$ (for any $k \geq 0$).

Example 6.9 (Forecasting an ARMA(p,q) process). Consider the process:

$$y_t = c + \phi_1 y_{t-1} + \dots + \phi_p y_{t-p} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q}, \quad (6.73)$$

where $\{\varepsilon_t\}$ is a white noise sequence (Def. 6.1). We assume that the MA part of the process is invertible (see Eq. (6.32)), which implies that the information contained in $\{y_t, y_{t-1}, y_{t-2}, \dots\}$ is identical to that in $\{\varepsilon_t, \varepsilon_{t-1}, \varepsilon_{t-2}, \dots\}$.

While one could use a recursive algorithm to compute the conditional mean (as in Example 6.8), it is convenient to employ the Wold decomposition of this process (see Theorem 6.3 and Prop. 6.8 for the computation of the ψ_i 's in the context of ARMA processes):

$$y_t = \mu + \sum_{i=0}^{+\infty} \psi_i \varepsilon_{t-i}.$$

This implies:

$$\begin{aligned}y_{t+h} &= \mu + \sum_{i=0}^{h-1} \psi_i \varepsilon_{t+h-i} + \sum_{i=h}^{+\infty} \psi_i \varepsilon_{t+h-i} \\ &= \mu + \sum_{i=0}^{h-1} \psi_i \varepsilon_{t+h-i} + \sum_{i=0}^{+\infty} \psi_{i+h} \varepsilon_{t-i}.\end{aligned}$$

Since $\mathbb{E}(y_{t+h}|y_t, y_{t-1}, \dots) = \mu + \sum_{i=0}^{+\infty} \psi_{i+h} \varepsilon_{t-i}$, we get:

$$\text{Var}(y_{t+h}|y_t, y_{t-1}, \dots) = \text{Var}\left(\sum_{i=0}^{h-1} \psi_i \varepsilon_{t+h-i}\right) = \sigma^2 \sum_{i=0}^{h-1} \psi_i^2.$$

How to use the previous formulas in practice?

One has first to select a specification and to estimate the model. Two methods to determine relevant specifications:

- a. Information criteria (see Definition 6.21).
- b. Box-Jenkins approach.

Box and Jenkins (1976) have proposed an approach that is now widely used.

1. Data transformation. The data should be transformed to “make them stationary”. To do so, one can e.g. take logarithms, take changes in the considered series, remove (deterministic) trends.
2. Select p and q . This can be based on the PACF approach (see Section 6.2.4), or on selection criteria (see Definition 6.21).
3. Estimate the model parameters. See Section 6.2.8.
4. Check that the estimated model is consistent with the data. See below.

Assessing the performances of a forecasting model

Once one has fitted a model on a given dataset (of length T , say), one compute MSE (mean square errors) to evaluate the performance of the model. But this MSE is the **in-sample** one. It is easy to reduce in-sample MSE. Typically, if the model is estimated by OLS, adding covariates mechanically reduces the MSE (see Props. 2.4 and 2.5). That is, even if additional data are irrelevant, the R^2 of the regression increases. Adding irrelevant variables increases the (in-sample) R^2 but is bound to increase the **out-of-sample** MSE.

Therefore, it is important to analyse **out-of-sample** performances of the forecasting model:

- a. Estimate a model on a sample of reduced size ($1, \dots, T^*$, with $T^* < T$)
- b. Use the remaining available periods ($T^* + 1, \dots, T$) to compute **out-of-sample** forecasting errors (and compute their MSE). In an out-of-sample exercise, it is important to make sure that the data used to produce a forecasts (as of date T^*) where indeed available on date T^* .

Diebold-Mariano test

How to compare different forecasting approaches? Diebold and Mariano (1995) have proposed a simple test to address this question.

Assume that you want to compare approaches A and B. You have historical data sets and you have implemented both approaches in the past, providing you with two sets of forecasting errors: $\{e_t^A\}_{t=1,\dots,T}$ and $\{e_t^B\}_{t=1,\dots,T}$.

It may be the case that your forecasts serve a specific purpose and that, for instance, you dislike positive forecasting errors and you care less about negative

errors. We assume you are able to formalise this by means of a **loss function** $L(e)$. For instance:

- If you dislike large positive errors, you may set $L(e) = \exp(e)$.
- If you are concerned about both positive and negative errors (indifferently), you may set $L(e) = e^2$ (standard approach).

Let us define the sequence $\{d_t\}_{t=1,\dots,T} \equiv \{L(e_t^A) - L(e_t^B)\}_{t=1,\dots,T}$ and assume that this sequence is covariance stationary. We consider the following null hypothesis: $H_0 : \bar{d} = 0$, where \bar{d} denotes the population mean of the d_t s. Under H_0 and under the assumption of covariance-stationarity of d_t , we have (Theorem @ref{(hm:CLTcovstat)}):

$$\sqrt{T}\bar{d}_T \xrightarrow{d} \mathcal{N}\left(0, \sum_{j=-\infty}^{+\infty} \gamma_j\right),$$

where the γ_j s are the autocovariances of d_t .

Hence, assuming that $\hat{\sigma}^2$ is a consistent estimate of $\sum_{j=-\infty}^{+\infty} \gamma_j$ (for instance the one given by the Newey-West formula, see Def. ??), we have, under H_0 :

$$DM_T := \sqrt{T} \frac{\bar{d}_T}{\sqrt{\hat{\sigma}^2}} \xrightarrow{d} \mathcal{N}(0, 1).$$

DM_T is the test statistics. For a test of size α , the critical region is:⁶

$$] -\infty, -\Phi^{-1}(1 - \alpha/2)] \cup [\Phi^{-1}(1 - \alpha/2), +\infty[,$$

where Φ is the c.d.f. of the standard normal distribution.

Example 6.10 (Forecasting Swiss GDP growth). We use a long historical time series of the Swiss GDP growth taken from the Jordà et al. (2017) dataset (see Figure 6.3, and Example 6.4).

We want to forecast this GDP growth. We envision two specifications : an AR(1) specification (the one advocated by the AIC criteria, see Example 6.4), and an ARMA(2,2) specification. We are interested in 2-year-ahead forecasts (i.e., $h = 2$ since the data are yearly).

```
library(AEC)
library(forecast)

## Registered S3 method overwritten by 'quantmod':
##   method           from
##   as.zoo.data.frame zoo
```

⁶This ShinyApp application illustrates the notion of statistical test (illustrating the p-value and the critical region, in particular).

```

data <- subset(JST, iso=="CHE")
T <- dim(data)[1]
y <- c(NaN, log(data$gdp[2:T]/data$gdp[1:(T-1)]))
first.date <- T-50
e1 <- NULL; e2 <- NULL; h<-2
for(T.star in first.date:(T-h)){
  estim.model.1 <- arima(y[1:T.star], order=c(1,0,0))
  estim.model.2 <- arima(y[1:T.star], order=c(2,0,2))
  e1 <- c(e1, y[T.star+h] - predict(estim.model.1, n.ahead=h)$pred[h])
  e2 <- c(e2, y[T.star+h] - predict(estim.model.2, n.ahead=h)$pred[h])
}
res.DM <- dm.test(e1, e2, h = h, alternative = "greater")
res.DM

##
## Diebold-Mariano Test
##
## data: e1e2
## DM = -1.2379, Forecast horizon = 2, Loss function power = 2, p-value =
## 0.8891
## alternative hypothesis: greater

```

With `alternative = "greater"` The alternative hypothesis is that method 2 is more accurate than method 1. Since we do not reject the null (the p-value being of 0.889), we are not led to use the more sophisticated model (ARMA(2,2)) and we keep the simple AR(1) model.

Assume now that we want to compare the AR(1) process to a VAR model (see Def. 6.22). We consider a bivariate VAR, where GDP growth is complemented with CPI-based inflation rate.

```

library(vars)
infl <- c(NaN, log(data$cpi[2:T]/data$cpi[1:(T-1)]))
y_var <- cbind(y, infl)
e3 <- NULL
for(T.star in first.date:(T-h)){
  estim.model.3 <- VAR(y_var[2:T.star,], p=1)
  e3 <- c(e3, y[T.star+h] - predict(estim.model.3, n.ahead=h)$fcst$y[h, 1])
}
res.DM <- dm.test(e1, e2, h = h, alternative = "greater")
res.DM

##
## Diebold-Mariano Test
##

```

```
## data: e1e2
## DM = -1.2379, Forecast horizon = 2, Loss function power = 2, p-value =
## 0.8891
## alternative hypothesis: greater
```

Again, we do not find that the alternative model (here the VAR(1) model) is better than the AR(1) model to forecast GDP growth.

Chapter 7

Appendix

7.1 Principal component analysis (PCA)

Principal component analysis (PCA) is a classical and easy-to-use statistical method to reduce the dimension of large datasets containing variables that are linearly driven by a relatively small number of factors. This approach is widely used in data analysis and image compression.

Suppose that we have T observations of a n -dimensional random vector x , denoted by x_1, x_2, \dots, x_T . We suppose that each component of x is of mean zero.

Let denote with X the matrix given by $[x_1 \ x_2 \ \dots \ x_T]'$. Denote the j^{th} column of X by X_j .

We want to find the linear combination of the x_i 's ($x.u$), with $\|u\| = 1$, with “maximum variance.” That is, we want to solve:

$$\begin{array}{ll} \arg \max_u & u' X' X u. \\ \text{s.t.} & |u| = 1 \end{array} \quad (7.1)$$

Since $X'X$ is a positive definite matrix, it admits the following decomposition:

$$\begin{aligned} X'X &= PDP' \\ &= P \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} P', \end{aligned}$$

where P is an orthogonal matrix whose columns are the eigenvectors of $X'X$.

We can order the eigenvalues such that $\lambda_1 \geq \dots \geq \lambda_n$. (Since $X'X$ is positive definite, all these eigenvalues are positive.)

Since P is orthogonal, we have $u'X'Xu = u'PDP'u = y'Dy$ where $\|y\| = 1$. Therefore, we have $y_i^2 \leq 1$ for any $i \leq n$.

As a consequence:

$$y'Dy = \sum_{i=1}^n y_i^2 \lambda_i \leq \lambda_1 \sum_{i=1}^n y_i^2 = \lambda_1.$$

It is easily seen that the maximum is reached for $y = [1, 0, \dots, 0]'$. Therefore, the maximum of the optimization program (Eq. (7.1)) is obtained for $u = P[1, 0, \dots, 0]'$. That is, u is the eigenvector of $X'X$ that is associated with its larger eigenvalue (first column of P).

Let us denote with F the vector that is given by the matrix product XP (note that its last column is equal to Xu). The columns of F , denoted by F_j , are called **factors**. We have:

$$F'F = P'X'XP = D.$$

Therefore, in particular, the F_j 's are orthogonal.

Since $X = FP'$, the X_j 's are linear combinations of the factors. Let us then denote with $\hat{X}_{i,j}$ the part of X_i that is explained by factor F_j , we have:

$$\begin{aligned}\hat{X}_{i,j} &= p_{ij}F_j \\ X_i &= \sum_j \hat{X}_{i,j} = \sum_j p_{ij}F_j.\end{aligned}$$

Consider the share of variance that is explained –through the n variables (X_1, \dots, X_n) – by the first factor F_1 :

$$\frac{\sum_i \hat{X}_{i,1} \hat{X}'_{i,1}}{\sum_i X_i X'_i} = \frac{\sum_i p_{i1} F_1 F'_1 p_{i1}}{\text{tr}(X'X)} = \frac{\sum_i p_{i1}^2 \lambda_1}{\text{tr}(X'X)} = \frac{\lambda_1}{\sum_i \lambda_i}.$$

Intuitively, if the first eigenvalue is large, it means that the first factor embed a large share of the fluctuations of the n X_i 's.

Let us illustrate PCA on the term structure of yields. The term structure of yields (or yield curve) is known to be driven by only a small number of factors (e.g., Litterman and Scheinkman (1991)). One can typically employ PCA to recover such factors. The data used in the example below are taken from the Fred database (tickers: “DGS6MO”, “DGS1”, ...). The second plot shows the factor loadings, that indicate that the first factor is a level factor (loadings = black line), the second factor is a slope factor (loadings = blue line), the third factor is a curvature factor (loadings = red line).

To run a PCA, one simply has to apply function `prcomp` to a matrix of data:

```

library(AEC)
USyields <- USyields[complete.cases(USyields),]
yds <- USyields[c("Y1","Y2","Y3","Y5","Y7","Y10","Y20","Y30")]
PCA.yds <- prcomp(yds,center=TRUE,scale. = TRUE)

```

Let us know visualize some results. The first plot of Figure 7.1 shows the share of total variance explained by the different principal components (PCs). The second plot shows the facotr loadings. The two bottom plots show how yields (in black) are fitted by linear combinations of the first two PCs only.

```

par(mfrow=c(2,2))
par(plt=c(.1,.95,.2,.8))
barplot(PCA.yds$sdev^2/sum(PCA.yds$sdev^2),
        main="Share of variance expl. by PC's")
axis(1, at=1:dim(yds)[2], labels=colnames(PCA.yds$x))
nb.PC <- 2
plot(-PCA.yds$rotation[,1],type="l",lwd=2,ylim=c(-1,1),
      main="Factor loadings (1st 3 PCs)",xaxt="n",xlab="")
axis(1, at=1:dim(yds)[2], labels=colnames(yds))
lines(PCA.yds$rotation[,2],type="l",lwd=2,col="blue")
lines(PCA.yds$rotation[,3],type="l",lwd=2,col="red")
Y1.hat <- PCA.yds$x[,1:nb.PC] %*% PCA.yds$rotation["Y1",1:2]
Y1.hat <- mean(USyields$Y1) + sd(USyields$Y1) * Y1.hat
plot(USyields$date,USyields$Y1,type="l",lwd=2,
      main="Fit of 1-year yields (2 PCs)",
      ylab="Obs (black) / Fitted by 2PCs (dashed blue)")
lines(USyields$date,Y1.hat,col="blue",lty=2,lwd=2)
Y10.hat <- PCA.yds$x[,1:nb.PC] %*% PCA.yds$rotation["Y10",1:2]
Y10.hat <- mean(USyields$Y10) + sd(USyields$Y10) * Y10.hat
plot(USyields$date,USyields$Y10,type="l",lwd=2,
      main="Fit of 10-year yields (2 PCs)",
      ylab="Obs (black) / Fitted by 2PCs (dashed blue)")
lines(USyields$date,Y10.hat,col="blue",lty=2,lwd=2)

```

7.2 Linear algebra: definitions and results

Definition 7.1 (Eigenvalues). The eigenvalues of a matrix M are the numbers λ for which:

$$|M - \lambda I| = 0,$$

where $|\bullet|$ is the determinant operator.

Proposition 7.1 (Properties of the determinant). *We have:*




Figure 7.1: Some PCA results. The dataset contains 8 time series of U.S. interest rates of different maturities.

- $|MN| = |M| \times |N|$.
- $|M^{-1}| = |M|^{-1}$.
- If M admits the diagonal representation $M = TDT^{-1}$, where D is a diagonal matrix whose diagonal entries are $\{\lambda_i\}_{i=1,\dots,n}$, then:

$$|M - \lambda I| = \prod_{i=1}^n (\lambda_i - \lambda).$$

Definition 7.2 (Moore-Penrose inverse). If $M \in \mathbb{R}^{m \times n}$, then its Moore-Penrose pseudo inverse (exists and) is the unique matrix $M^* \in \mathbb{R}^{n \times m}$ that satisfies:

- i. $MM^*M = M$
- ii. $M^*MM^* = M^*$
- iii. $(MM^*)' = MM^*$.iv $(M^*M)' = M^*M$.

- Proposition 7.2** (Properties of the Moore-Penrose inverse). • If M is invertible then $M^* = M^{-1}$.
- The pseudo-inverse of a zero matrix is its transpose. *
 - *
 - The pseudo-inverse of the pseudo-inverse is the original matrix.

Definition 7.3 (Idempotent matrix). Matrix M is idempotent if $M^2 = M$.

If M is a symmetric idempotent matrix, then $M'M = M$.

Proposition 7.3 (Roots of an idempotent matrix). *The eigenvalues of an idempotent matrix are either 1 or 0.*

Proof. If λ is an eigenvalue of an idempotent matrix M then $\exists x \neq 0$ s.t. $Mx = \lambda x$. Hence $M^2x = \lambda Mx \Rightarrow (1 - \lambda)Mx = 0$. Either all element of Mx are zero, in which case $\lambda = 0$ or at least one element of Mx is nonzero, in which case $\lambda = 1$. \square

Proposition 7.4 (Idempotent matrix and chi-square distribution). *The rank of a symmetric idempotent matrix is equal to its trace.*

Proof. The result follows from Prop. 7.3, combined with the fact that the rank of a symmetric matrix is equal to the number of its nonzero eigenvalues. \square

Proposition 7.5 (Constrained least squares). *The solution of the following optimisation problem:*

$$\begin{aligned} \min_{\beta} \quad & \|y - X\beta\|^2 \\ \text{subject to } & R\beta = q \end{aligned}$$

is given by:

$$\beta^r = \beta_0 - (X'X)^{-1}R'\{R(X'X)^{-1}R'\}^{-1}(R\beta_0 - q),$$

where $\beta_0 = (X'X)^{-1}X'y$.

Proof. See for instance Jackman, 2007. \square

Proposition 7.6 (Inverse of a partitioned matrix). *We have:*

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}^{-1} = \begin{bmatrix} (A_{11} - A_{12}A_{22}^{-1}A_{21})^{-1} & -A_{11}^{-1}A_{12}(A_{22} - A_{21}A_{11}^{-1}A_{12})^{-1} \\ -(A_{22} - A_{21}A_{11}^{-1}A_{12})^{-1}A_{21}A_{11}^{-1} & (A_{22} - A_{21}A_{11}^{-1}A_{12})^{-1} \end{bmatrix}.$$

Definition 7.4 (Matrix derivatives). Consider a fonction $f : \mathbb{R}^K \rightarrow \mathbb{R}$. Its first-order derivative is:

$$\frac{\partial f}{\partial \mathbf{b}}(\mathbf{b}) = \begin{bmatrix} \frac{\partial f}{\partial b_1}(\mathbf{b}) \\ \vdots \\ \frac{\partial f}{\partial b_K}(\mathbf{b}) \end{bmatrix}.$$

We use the notation:

$$\frac{\partial f}{\partial \mathbf{b}'}(\mathbf{b}) = \left(\frac{\partial f}{\partial \mathbf{b}}(\mathbf{b}) \right)'.$$

Proposition 7.7. *We have:*

- If $f(\mathbf{b}) = A'\mathbf{b}$ where A is a $K \times 1$ vector then $\frac{\partial f}{\partial \mathbf{b}}(\mathbf{b}) = A$.
- If $f(\mathbf{b}) = \mathbf{b}'A\mathbf{b}$ where A is a $K \times K$ matrix, then $\frac{\partial f}{\partial \mathbf{b}}(\mathbf{b}) = 2A\mathbf{b}$.

Proposition 7.8 (Square and absolute summability). *We have:*

$$\underbrace{\sum_{i=0}^{\infty} |\theta_i| < +\infty}_{\text{Absolute summability}} \Rightarrow \underbrace{\sum_{i=0}^{\infty} \theta_i^2 < +\infty}_{\text{Square summability}} .$$

Proof. See Appendix 3.A in Hamilton. Idea: Absolute summability implies that there exist N such that, for $j > N$, $|\theta_j| < 1$ (deduced from Cauchy criterion, Theorem 7.2 and therefore $\theta_j^2 < |\theta_j|$). \square

7.3 Statistical analysis: definitions and results

7.3.1 Moments and statistics

Definition 7.5 (Partial correlation). The **partial correlation** between y and z , controlling for some variables \mathbf{X} is the sample correlation between y^* and z^* , where the latter two variables are the residuals in regressions of y on \mathbf{X} and of z on \mathbf{X} , respectively.

This correlation is denoted by $r_{yz}^{\mathbf{X}}$. By definition, we have:

$$r_{yz}^{\mathbf{X}} = \frac{\mathbf{z}^{*\prime} \mathbf{y}^*}{\sqrt{(\mathbf{z}^{*\prime} \mathbf{z}^*)(\mathbf{y}^{*\prime} \mathbf{y}^*)}}. \quad (7.2)$$

Definition 7.6 (Skewness and kurtosis). Let Y be a random variable whose fourth moment exists. The expectation of Y is denoted by μ .

- The skewness of Y is given by:

$$\frac{\mathbb{E}[(Y - \mu)^3]}{\{\mathbb{E}[(Y - \mu)^2]\}^{3/2}}.$$

- The kurtosis of Y is given by:

$$\frac{\mathbb{E}[(Y - \mu)^4]}{\{\mathbb{E}[(Y - \mu)^2]\}^2}.$$

Theorem 7.1 (Cauchy-Schwarz inequality). *We have:*

$$|\text{Cov}(X, Y)| \leq \sqrt{\text{Var}(X)\text{Var}(Y)}$$

and, if $X \neq 0$ and $Y \neq 0$, the equality holds iff X and Y are the same up to an affine transformation.

Proof. If $\text{Var}(X) = 0$, this is trivial. If this is not the case, then let's define Z as $Z = Y - \frac{\text{Cov}(X, Y)}{\text{Var}(X)}X$. It is easily seen that $\text{Cov}(X, Z) = 0$. Then, the variance of $Y = Z + \frac{\text{Cov}(X, Y)}{\text{Var}(X)}X$ is equal to the sum of the variance of Z and of the variance of $\frac{\text{Cov}(X, Y)}{\text{Var}(X)}X$, that is:

$$\text{Var}(Y) = \text{Var}(Z) + \left(\frac{\text{Cov}(X, Y)}{\text{Var}(X)} \right)^2 \text{Var}(X) \geq \left(\frac{\text{Cov}(X, Y)}{\text{Var}(X)} \right)^2 \text{Var}(X).$$

The equality holds iff $\text{Var}(Z) = 0$, i.e. iff $Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)}X + cst$. \square

Definition 7.7 (Asymptotic level). An asymptotic test with critical region Ω_n has an asymptotic level equal to α if:

$$\sup_{\theta \in \Theta} \lim_{n \rightarrow \infty} \mathbb{P}_\theta(S_n \in \Omega_n) = \alpha,$$

where S_n is the test statistic and Θ is such that the null hypothesis H_0 is equivalent to $\theta \in \Theta$.

Definition 7.8 (Asymptotically consistent test). An asymptotic test with critical region Ω_n is consistent if:

$$\forall \theta \in \Theta^c, \quad \mathbb{P}_\theta(S_n \in \Omega_n) \rightarrow 1,$$

where S_n is the test statistic and Θ^c is such that the null hypothesis H_0 is equivalent to $\theta \notin \Theta^c$.

Definition 7.9 (Kullback discrepancy). Given two p.d.f. f and f^* , the Kullback discrepancy is defined by:

$$I(f, f^*) = \mathbb{E}^* \left(\log \frac{f^*(Y)}{f(Y)} \right) = \int \log \frac{f^*(y)}{f(y)} f^*(y) dy.$$

Proposition 7.9 (Properties of the Kullback discrepancy). *We have:*

- i. $I(f, f^*) \geq 0$
- ii. $I(f, f^*) = 0$ iff $f \equiv f^*$.

Proof. $x \rightarrow -\log(x)$ is a convex function. Therefore $\mathbb{E}^*(-\log f(Y)/f^*(Y)) \geq -\log \mathbb{E}^*(f(Y)/f^*(Y)) = 0$ (proves (i)). Since $x \rightarrow -\log(x)$ is strictly convex, equality in (i) holds if and only if $f(Y)/f^*(Y)$ is constant (proves (ii)). \square

Definition 7.10 (Characteristic function). For any real-valued random variable X , the characteristic function is defined by:

$$\phi_X : u \rightarrow \mathbb{E}[\exp(iuX)].$$

7.3.2 Standard distributions

Definition 7.11 (F distribution). Consider $n = n_1 + n_2$ i.i.d. $\mathcal{N}(0, 1)$ r.v. X_i . If the r.v. F is defined by:

$$F = \frac{\sum_{i=1}^{n_1} X_i^2}{\sum_{j=n_1+1}^{n_1+n_2} X_j^2} \frac{n_2}{n_1}$$

then $F \sim \mathcal{F}(n_1, n_2)$. (See Table 7.4 for quantiles.)

Definition 7.12 (Student-t distribution). Z follows a Student-t (or t) distribution with ν degrees of freedom (d.f.) if:

$$Z = X_0 / \sqrt{\frac{\sum_{i=1}^{\nu} X_i^2}{\nu}}, \quad X_i \sim i.i.d. \mathcal{N}(0, 1).$$

We have $\mathbb{E}(Z) = 0$, and $\text{Var}(Z) = \frac{\nu}{\nu-2}$ if $\nu > 2$. (See Table 7.2 for quantiles.)

Definition 7.13 (Chi-square distribution). Z follows a χ^2 distribution with ν d.f. if $Z = \sum_{i=1}^{\nu} X_i^2$ where $X_i \sim i.i.d. \mathcal{N}(0, 1)$. We have $\mathbb{E}(Z) = \nu$. (See Table 7.3 for quantiles.)

Proposition 7.10 (Inner product of a multivariate Gaussian variable). Let X be a n -dimensional multivariate Gaussian variable: $X \sim \mathcal{N}(0, \Sigma)$. We have:

$$X' \Sigma^{-1} X \sim \chi^2(n).$$

Proof. Because Σ is a symmetrical definite positive matrix, it admits the spectral decomposition PDP' where P is an orthogonal matrix (i.e. $PP' = Id$) and D is a diagonal matrix with non-negative entries. Denoting by $\sqrt{D^{-1}}$ the diagonal matrix whose diagonal entries are the inverse of those of D , it is easily checked that the covariance matrix of $Y := \sqrt{D^{-1}}P'X$ is Id . Therefore Y is a vector of uncorrelated Gaussian variables. The properties of Gaussian variables imply that the components of Y are then also independent. Hence $Y'Y = \sum_i Y_i^2 \sim \chi^2(n)$.

It remains to note that $Y'Y = X'PD^{-1}P'X = X'\text{Var}(X)^{-1}X$ to conclude. \square

7.3.3 Stochastic convergences

Proposition 7.11 (Chebychev's inequality). If $\mathbb{E}(|X|^r)$ is finite for some $r > 0$ then:

$$\forall \varepsilon > 0, \quad \mathbb{P}(|X - c| > \varepsilon) \leq \frac{\mathbb{E}[|X - c|^r]}{\varepsilon^r}.$$

In particular, for $r = 2$:

$$\forall \varepsilon > 0, \quad \mathbb{P}(|X - c| > \varepsilon) \leq \frac{\mathbb{E}[(X - c)^2]}{\varepsilon^2}.$$

Proof. Remark that $\varepsilon^r \mathbb{I}_{\{|X| \geq \varepsilon\}} \leq |X|^r$ and take the expectation of both sides. \square

Definition 7.14 (Convergence in probability). The random variable sequence x_n converges in probability to a constant c if $\forall \varepsilon, \lim_{n \rightarrow \infty} \mathbb{P}(|x_n - c| > \varepsilon) = 0$.

It is denoted as: $\text{plim } x_n = c$.

Definition 7.15 (Convergence in the L^r norm). x_n converges in the r -th mean (or in the L^r -norm) towards x , if $\mathbb{E}(|x_n|^r)$ and $\mathbb{E}(|x|^r)$ exist and if

$$\lim_{n \rightarrow \infty} \mathbb{E}(|x_n - x|^r) = 0.$$

It is denoted as: $x_n \xrightarrow{L^r} c$.

For $r = 2$, this convergence is called **mean square convergence**.

Definition 7.16 (Almost sure convergence). The random variable sequence x_n converges almost surely to c if $\mathbb{P}(\lim_{n \rightarrow \infty} x_n = c) = 1$.

It is denoted as: $x_n \xrightarrow{a.s.} c$.

Definition 7.17 (Convergence in distribution). x_n is said to converge in distribution (or in law) to x if

$$\lim_{n \rightarrow \infty} F_{x_n}(s) = F_x(s)$$

for all s at which F_X –the cumulative distribution of X – is continuous.

It is denoted as: $x_n \xrightarrow{d} x$.

Proposition 7.12 (Rules for limiting distributions (Slutsky)). *We have:*

i. **Slutsky's theorem:** If $x_n \xrightarrow{d} x$ and $y_n \xrightarrow{p} c$ then

$$\begin{aligned} x_n y_n &\xrightarrow{d} xc \\ x_n + y_n &\xrightarrow{d} x + c \\ x_n / y_n &\xrightarrow{d} x/c \quad (\text{if } c \neq 0) \end{aligned}$$

ii. **Continuous mapping theorem:** If $x_n \xrightarrow{d} x$ and g is a continuous function then $g(x_n) \xrightarrow{d} g(x)$.

Proposition 7.13 (Implications of stochastic convergences). *We have:*

$$\begin{array}{ccc} \boxed{\xrightarrow{L^s}} & \xrightarrow[1 \leq r \leq s]{} & \boxed{\xrightarrow{L^r}} \\ & \Downarrow & \\ \boxed{\xrightarrow{a.s.}} & \Rightarrow & \boxed{\xrightarrow{p}} \Rightarrow \boxed{\xrightarrow{d}}. \end{array}$$

Proof. (of the fact that $\left(\xrightarrow{p}\right) \Rightarrow \left(\xrightarrow{d}\right)$). Assume that $X_n \xrightarrow{p} X$. Denoting by F and F_n the c.d.f. of X and X_n , respectively:

$$F_n(x) = \mathbb{P}(X_n \leq x, X \leq x+\varepsilon) + \mathbb{P}(X_n \leq x, X > x+\varepsilon) \leq F(x+\varepsilon) + \mathbb{P}(|X_n - X| > \varepsilon). \quad (7.3)$$

Besides,

$$F(x-\varepsilon) = \mathbb{P}(X \leq x-\varepsilon, X_n \leq x) + \mathbb{P}(X \leq x-\varepsilon, X_n > x) \leq F_n(x) + \mathbb{P}(|X_n - X| > \varepsilon),$$

which implies:

$$F(x-\varepsilon) - \mathbb{P}(|X_n - X| > \varepsilon) \leq F_n(x). \quad (7.4)$$

Eqs. (7.3) and (7.4) imply:

$$F(x-\varepsilon) - \mathbb{P}(|X_n - X| > \varepsilon) \leq F_n(x) \leq F(x+\varepsilon) + \mathbb{P}(|X_n - X| > \varepsilon).$$

Taking limits as $n \rightarrow \infty$ yields

$$F(x-\varepsilon) \leq \liminf_{n \rightarrow \infty} F_n(x) \leq \limsup_{n \rightarrow \infty} F_n(x) \leq F(x+\varepsilon).$$

The result is then obtained by taking limits as $\varepsilon \rightarrow 0$ (if F is continuous at x). \square

Proposition 7.14 (Convergence in distribution to a constant). *If X_n converges in distribution to a constant c , then X_n converges in probability to c .*

Proof. If $\varepsilon > 0$, we have $\mathbb{P}(X_n < c - \varepsilon) \xrightarrow{n \rightarrow \infty} 0$ i.e. $\mathbb{P}(X_n \geq c - \varepsilon) \xrightarrow{n \rightarrow \infty} 1$ and $\mathbb{P}(X_n < c + \varepsilon) \xrightarrow{n \rightarrow \infty} 1$. Therefore $\mathbb{P}(c - \varepsilon \leq X_n < c + \varepsilon) \xrightarrow{n \rightarrow \infty} 1$, which gives the result. \square

Example 7.1 (Convergence in probability but not L^r). Let $\{x_n\}_{n \in \mathbb{N}}$ be a series of random variables defined by:

$$x_n = nu_n,$$

where u_n are independent random variables s.t. $u_n \sim \mathcal{B}(1/n)$.

We have $x_n \xrightarrow{p} 0$ but $x_n \not\xrightarrow{L^r} 0$ because $\mathbb{E}(|X_n - 0|) = \mathbb{E}(X_n) = 1$.

Theorem 7.2 (Cauchy criterion (non-stochastic case)). *We have that $\sum_{i=0}^T a_i$ converges ($T \rightarrow \infty$) iff, for any $\eta > 0$, there exists an integer N such that, for all $M \geq N$,*

$$\left| \sum_{i=N+1}^M a_i \right| < \eta.$$

Theorem 7.3 (Cauchy criterion (stochastic case)). *We have that $\sum_{i=0}^T \theta_i \varepsilon_{t-i}$ converges in mean square ($T \rightarrow \infty$) to a random variable iff, for any $\eta > 0$, there exists an integer N such that, for all $M \geq N$,*

$$\mathbb{E} \left[\left(\sum_{i=N+1}^M \theta_i \varepsilon_{t-i} \right)^2 \right] < \eta.$$

7.3.4 Central limit theorem

Theorem 7.4 (Law of large numbers). *The sample mean is a consistent estimator of the population mean.*

Proof. Let's denote by ϕ_{X_i} the characteristic function of a r.v. X_i . If the mean of X_i is μ then the Taylor expansion of the characteristic function is:

$$\phi_{X_i}(u) = \mathbb{E}(\exp(iuX_i)) = 1 + iu\mu + o(u).$$

The properties of the characteristic function (see Def. 7.10) imply that:

$$\phi_{\frac{1}{n}(X_1 + \dots + X_n)}(u) = \prod_{i=1}^n \left(1 + i\frac{u}{n}\mu + o\left(\frac{u}{n}\right)\right) \rightarrow e^{iu\mu}.$$

The facts that (a) $e^{iu\mu}$ is the characteristic function of the constant μ and (b) that a characteristic function uniquely characterises a distribution imply that the sample mean converges in distribution to the constant μ , which further implies that it converges in probability to μ . \square

Theorem 7.5 (Lindberg-Levy Central limit theorem, CLT). *If x_n is an i.i.d. sequence of random variables with mean μ and variance σ^2 ($\in]0, +\infty[$), then:*

$$\boxed{\sqrt{n}(\bar{x}_n - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2), \quad \text{where} \quad \bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i.}$$

Proof. Let us introduce the r.v. $Y_n := \sqrt{n}(\bar{X}_n - \mu)$. We have $\phi_{Y_n}(u) = [\mathbb{E}(\exp(i\frac{1}{\sqrt{n}}u(X_1 - \mu)))]^n$. We have:

$$\begin{aligned} \left[\mathbb{E} \left(\exp \left(i \frac{1}{\sqrt{n}} u (X_1 - \mu) \right) \right) \right]^n &= \left[\mathbb{E} \left(1 + i \frac{1}{\sqrt{n}} u (X_1 - \mu) - \frac{1}{2n} u^2 (X_1 - \mu)^2 + o(u^2) \right) \right]^n \\ &= \left(1 - \frac{1}{2n} u^2 \sigma^2 + o(u^2) \right)^n. \end{aligned}$$

Therefore $\phi_{Y_n}(u) \xrightarrow{n \rightarrow \infty} \exp(-\frac{1}{2}u^2\sigma^2)$, which is the characteristic function of $\mathcal{N}(0, \sigma^2)$. \square

7.4 Some properties of Gaussian variables

Proposition 7.15. *If \mathbf{A} is idempotent and if \mathbf{x} is Gaussian, \mathbf{Lx} and $\mathbf{x}'\mathbf{Ax}$ are independent if $\mathbf{LA} = \mathbf{0}$.*

Proof. If $\mathbf{L}\mathbf{A} = \mathbf{0}$, then the two Gaussian vectors \mathbf{Lx} and \mathbf{Ax} are independent. This implies the independence of any function of \mathbf{Lx} and any function of \mathbf{Ax} . The results then follows from the observation that $\mathbf{x}'\mathbf{Ax} = (\mathbf{Ax})'(\mathbf{Ax})$, which is a function of \mathbf{Ax} . \square

Proposition 7.16 (Bayesian update in a vector of Gaussian variables). *If*

$$\begin{bmatrix} Y_1 \\ Y_2 \end{bmatrix} \sim \mathcal{N} \left(0, \begin{bmatrix} \Omega_{11} & \Omega_{12} \\ \Omega_{21} & \Omega_{22} \end{bmatrix} \right),$$

then

$$\begin{aligned} Y_2|Y_1 &\sim \mathcal{N}(\Omega_{21}\Omega_{11}^{-1}Y_1, \Omega_{22} - \Omega_{21}\Omega_{11}^{-1}\Omega_{12}), \\ Y_1|Y_2 &\sim \mathcal{N}(\Omega_{12}\Omega_{22}^{-1}Y_2, \Omega_{11} - \Omega_{12}\Omega_{22}^{-1}\Omega_{21}). \end{aligned}$$

Proposition 7.17 (Truncated distributions). *If X is a random variable distributed according to some p.d.f. f , with c.d.f. F , with infinite support. Then the p.d.f. of $X|a \leq X < b$ is*

$$g(x) = \frac{f(x)}{F(b) - F(a)} \mathbb{I}_{\{a \leq x < b\}},$$

for any $a < b$.

In particular, for a Gaussian variable $X \sim \mathcal{N}(\mu, \sigma^2)$, we have

$$f(X = x|a \leq X < b) = \frac{\frac{1}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right)}{Z},$$

with $Z = \Phi(\beta) - \Phi(\alpha)$, where $\alpha = \frac{a-\mu}{\sigma}$ and $\beta = \frac{b-\mu}{\sigma}$.

Moreover:

$$\mathbb{E}(X|a \leq X < b) = \mu - \frac{\phi(\beta) - \phi(\alpha)}{Z} \sigma. \quad (7.5)$$

We also have:

$$\begin{aligned} \text{Var}(X|a \leq X < b) &= \sigma^2 \left[1 - \frac{\beta\phi(\beta) - \alpha\phi(\alpha)}{Z} - \left(\frac{\phi(\beta) - \phi(\alpha)}{Z} \right)^2 \right] \end{aligned} \quad (7.6)$$

In particular, for $b \rightarrow \infty$, we get:

$$\text{Var}(X|a < X) = \sigma^2 [1 + \alpha\lambda(-\alpha) - \lambda(-\alpha)^2], \quad (7.7)$$

with $\lambda(x) = \frac{\phi(x)}{\Phi(x)}$ is called the **inverse Mills ratio**.




Figure 7.2: $\mathbb{E}(X|X < b)$ as a function of b when $X \sim \mathcal{N}(0, 1)$ (in black).

Consider the case where $a \rightarrow -\infty$ (i.e. the conditioning set is $X < b$) and $\mu = 0$, $\sigma = 1$. Then Eq. (7.5) gives $\mathbb{E}(X|X < b) = -\lambda(b) = -\frac{\phi(b)}{\Phi(b)}$, where λ is the function computing the inverse Mills ratio.

Proposition 7.18 (p.d.f. of a multivariate Gaussian variable). *If $Y \sim \mathcal{N}(\mu, \Omega)$ and if Y is a n -dimensional vector, then the density function of Y is:*

$$\frac{1}{(2\pi)^{n/2} |\Omega|^{1/2}} \exp \left[-\frac{1}{2} (Y - \mu)' \Omega^{-1} (Y - \mu) \right].$$

7.5 Proofs

Proof of Proposition 4.4

Proof. Assumptions (i) and (ii) (in the set of Assumptions 4.1) imply that θ_{MLE} exists ($= \operatorname{argmax}_{\theta} (1/n) \log \mathcal{L}(\theta; \mathbf{y})$).

$(1/n) \log \mathcal{L}(\theta; \mathbf{y})$ can be interpreted as the sample mean of the r.v. $\log f(Y_i; \theta)$ that are i.i.d. Therefore $(1/n) \log \mathcal{L}(\theta; \mathbf{y})$ converges to $\mathbb{E}_{\theta_0}(\log f(Y; \theta))$ – which exists (Assumption iv).

Because the latter convergence is uniform (Assumption v), the solution θ_{MLE} almost surely converges to the solution to the limit problem:

$$\operatorname{argmax}_{\theta} \mathbb{E}_{\theta_0}(\log f(Y; \theta)) = \operatorname{argmax}_{\theta} \int_Y \log f(y; \theta) f(y; \theta_0) dy.$$

Properties of the Kullback information measure (see Prop. 7.9), together with the identifiability assumption (ii) implies that the solution to the limit problem is unique and equal to θ_0 .

Consider a r.v. sequence θ that converges to θ_0 . The Taylor expansion of the score in a neighborhood of θ_0 yields to:

$$\frac{\partial \log \mathcal{L}(\theta; \mathbf{y})}{\partial \theta} = \frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} + \frac{\partial^2 \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta \partial \theta'} (\theta - \theta_0) + o_p(\theta - \theta_0)$$

θ_{MLE} converges to θ_0 and satisfies the likelihood equation $\frac{\partial \log \mathcal{L}(\theta; \mathbf{y})}{\partial \theta} = \mathbf{0}$. Therefore:

$$\frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} \approx -\frac{\partial^2 \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta \partial \theta'} (\theta_{MLE} - \theta_0),$$

or equivalently:

$$\frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} \approx \left(-\frac{1}{n} \sum_{i=1}^n \frac{\partial^2 \log f(y_i; \theta_0)}{\partial \theta \partial \theta'} \right) \sqrt{n} (\theta_{MLE} - \theta_0),$$

By the law of large numbers, we have: $\left(-\frac{1}{n} \sum_{i=1}^n \frac{\partial^2 \log f(y_i; \theta_0)}{\partial \theta \partial \theta'} \right) \rightarrow \frac{1}{n} \mathbf{I}(\theta_0) = \mathcal{J}_Y(\theta_0)$.

Besides, we have:

$$\begin{aligned} \frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} &= \sqrt{n} \left(\frac{1}{n} \sum_i \frac{\partial \log f(y_i; \theta_0)}{\partial \theta} \right) \\ &= \sqrt{n} \left(\frac{1}{n} \sum_i \left\{ \frac{\partial \log f(y_i; \theta_0)}{\partial \theta} - \mathbb{E}_{\theta_0} \frac{\partial \log f(Y_i; \theta_0)}{\partial \theta} \right\} \right) \end{aligned}$$

which converges to $\mathcal{N}(0, \mathcal{J}_Y(\theta_0))$ by the CLT.

Collecting the preceding results leads to (b). The fact that θ_{MLE} achieves the FDCR bound proves (c). \square

Proof of Proposition 4.5

Proof. We have $\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} \mathcal{N}(0, \mathcal{J}(\theta_0)^{-1})$ (Eq. ??eq:normMLE)). A Taylor expansion around θ_0 yields to:

$$\sqrt{n}(h(\hat{\theta}_n) - h(\theta_0)) \xrightarrow{d} \mathcal{N}\left(0, \frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{\partial h(\theta_0)'}{\partial \theta}\right). \quad (7.8)$$

Under H_0 , $h(\theta_0) = 0$ therefore:

$$\sqrt{n}h(\hat{\theta}_n) \xrightarrow{d} \mathcal{N}\left(0, \frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{I}(\theta_0)^{-1} \frac{\partial h(\theta_0)'}{\partial \theta}\right). \quad (7.9)$$

Hence

$$\sqrt{n} \left(\frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{I}(\theta_0)^{-1} \frac{\partial h(\theta_0)'}{\partial \theta} \right)^{-1/2} h(\hat{\theta}_n) \xrightarrow{d} \mathcal{N}(0, Id).$$

Taking the quadratic form, we obtain:

$$nh(\hat{\theta}_n)' \left(\frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{I}(\theta_0)^{-1} \frac{\partial h(\theta_0)'}{\partial \theta} \right)^{-1} h(\hat{\theta}_n) \xrightarrow{d} \chi^2(r).$$

The fact that the test has asymptotic level α directly stems from what precedes.

Consistency of the test: Consider $\theta_0 \in \Theta$. Because the MLE is consistent, $h(\hat{\theta}_n)$ converges to $h(\theta_0) \neq 0$. Eq. (7.8) is still valid. It implies that ξ_n^W converges to $+\infty$ and therefore that $\mathbb{P}_\theta(\xi_n^W \geq \chi_{1-\alpha}^2(r)) \rightarrow 1$. \square

Proof of Proposition 4.6

Proof. Notations: “ \approx ” means “equal up to a term that converges to 0 in probability”. We are under H_0 . $\hat{\theta}^0$ is the constrained ML estimator; $\hat{\theta}$ denotes the unconstrained one.

We combine the two Taylor expansion: $h(\hat{\theta}_n) \approx \frac{\partial h(\theta_0)}{\partial \theta'} (\hat{\theta}_n - \theta_0)$ and $h(\hat{\theta}_n^0) \approx \frac{\partial h(\theta_0)}{\partial \theta'} (\hat{\theta}_n^0 - \theta_0)$ and we use $h(\hat{\theta}_n^0) = 0$ (by definition) to get:

$$\sqrt{n}h(\hat{\theta}_n) \approx \frac{\partial h(\theta_0)}{\partial \theta'} \sqrt{n}(\hat{\theta}_n - \hat{\theta}_n^0). \quad (7.10)$$

Besides, we have (using the definition of the information matrix):

$$\frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \approx \frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} - \mathcal{I}(\theta_0) \sqrt{n}(\hat{\theta}_n^0 - \theta_0) \quad (7.11)$$

and:

$$0 = \frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\hat{\theta}_n; \mathbf{y})}{\partial \theta} \approx \frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} - \mathcal{I}(\theta_0) \sqrt{n}(\hat{\theta}_n - \theta_0). \quad (7.12)$$

Taking the difference and multiplying by $\mathcal{I}(\theta_0)^{-1}$:

$$\sqrt{n}(\hat{\theta}_n - \hat{\theta}_n^0) \approx \mathcal{I}(\theta_0)^{-1} \frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \mathcal{I}(\theta_0). \quad (7.13)$$

Eqs. (7.10) and (7.13) yield to:

$$\sqrt{n}h(\hat{\theta}_n) \approx \frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta}. \quad (7.14)$$

Recall that $\hat{\theta}_n^0$ is the MLE of θ_0 under the constraint $h(\theta) = 0$. The vector of Lagrange multipliers $\hat{\lambda}_n$ associated to this program satisfies:

$$\frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} + \frac{\partial h'(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \hat{\lambda}_n = 0. \quad (7.15)$$

Substituting the latter equation in Eq. (7.14) gives:

$$\sqrt{n}h(\hat{\theta}_n) \approx -\frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{\partial h'(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \frac{\hat{\lambda}_n}{\sqrt{n}} \approx -\frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{\partial h'(\theta_0; \mathbf{y})}{\partial \theta} \frac{\hat{\lambda}_n}{\sqrt{n}}$$

which yields:

$$\frac{\hat{\lambda}_n}{\sqrt{n}} \approx -\left(\frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{\partial h'(\theta_0; \mathbf{y})}{\partial \theta} \right)^{-1} \sqrt{n}h(\hat{\theta}_n). \quad (7.16)$$

It follows, from Eq. (7.9), that:

$$\frac{\hat{\lambda}_n}{\sqrt{n}} \xrightarrow{d} \mathcal{N}\left(0, \left(\frac{\partial h(\theta_0)}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{\partial h'(\theta_0; \mathbf{y})}{\partial \theta} \right)^{-1}\right).$$

Taking the quadratic form of the last equation gives:

$$\frac{1}{n} \hat{\lambda}'_n \frac{\partial h(\hat{\theta}_n^0)}{\partial \theta'} \mathcal{J}(\hat{\theta}_n^0)^{-1} \frac{\partial h'(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \hat{\lambda}_n \xrightarrow{d} \chi^2(r).$$

Using Eq. (7.15), it appears that the left-hand side term of the last equation is ξ^{LM} as defined in Eq. (4.18). Consistency: see Remark 17.3 in Gouriéroux and Monfort (1995). \square

Proof of Proposition 4.7

Proof. Let us first demonstrate the asymptotic equivalence of ξ^{LM} and ξ^{LR} .

The second-order taylor expansions of $\log \mathcal{L}(\hat{\theta}_n^0, \mathbf{y})$ and $\log \mathcal{L}(\hat{\theta}_n, \mathbf{y})$ are:

$$\begin{aligned} \log \mathcal{L}(\hat{\theta}_n, \mathbf{y}) &\approx \log \mathcal{L}(\theta_0, \mathbf{y}) + \frac{\partial \log \mathcal{L}(\theta_0, \mathbf{y})}{\partial \theta'} (\hat{\theta}_n - \theta_0) - \frac{n}{2} (\hat{\theta}_n - \theta_0)' \mathcal{J}(\theta_0) (\hat{\theta}_n - \theta_0) \\ \log \mathcal{L}(\hat{\theta}_n^0, \mathbf{y}) &\approx \log \mathcal{L}(\theta_0, \mathbf{y}) + \frac{\partial \log \mathcal{L}(\theta_0, \mathbf{y})}{\partial \theta'} (\hat{\theta}_n^0 - \theta_0) - \frac{n}{2} (\hat{\theta}_n^0 - \theta_0)' \mathcal{J}(\theta_0) (\hat{\theta}_n^0 - \theta_0). \end{aligned}$$

Taking the difference, we obtain:

$$\xi_n^{LR} \approx 2 \frac{\partial \log \mathcal{L}(\theta_0, \mathbf{y})}{\partial \theta'} (\hat{\theta}_n - \hat{\theta}_n^0) + n(\hat{\theta}_n^0 - \theta_0)' \mathcal{J}(\theta_0)(\hat{\theta}_n^0 - \theta_0) - n(\hat{\theta}_n - \theta_0)' \mathcal{J}(\theta_0)(\hat{\theta}_n - \theta_0).$$

Using $\frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\theta_0; \mathbf{y})}{\partial \theta} \approx \mathcal{J}(\theta_0) \sqrt{n}(\hat{\theta}_n - \theta_0)$ (Eq. (7.12)), we have:

$$\xi_n^{LR} \approx 2n(\hat{\theta}_n - \theta_0)' \mathcal{J}(\theta_0)(\hat{\theta}_n - \hat{\theta}_n^0) + n(\hat{\theta}_n^0 - \theta_0)' \mathcal{J}(\theta_0)(\hat{\theta}_n^0 - \theta_0) - n(\hat{\theta}_n - \theta_0)' \mathcal{J}(\theta_0)(\hat{\theta}_n - \theta_0).$$

In the second of the three terms in the sum, we replace $(\hat{\theta}_n^0 - \theta_0)$ by $(\hat{\theta}_n^0 - \hat{\theta}_n + \hat{\theta}_n - \theta_0)$ and we develop the associated product. This leads to:

$$\xi_n^{LR} \approx n(\hat{\theta}_n^0 - \hat{\theta}_n)' \mathcal{J}(\theta_0)^{-1} (\hat{\theta}_n^0 - \hat{\theta}_n). \quad (7.17)$$

The difference between Eqs. (7.11) and (7.12) implies:

$$\frac{1}{\sqrt{n}} \frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \approx \mathcal{J}(\theta_0) \sqrt{n}(\hat{\theta}_n - \hat{\theta}_n^0),$$

which, associated to Eq. @??eq:lr10), gives:

$$\xi_n^{LR} \approx \frac{1}{n} \frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta'} \mathcal{J}(\theta_0)^{-1} \frac{\partial \log \mathcal{L}(\hat{\theta}_n^0; \mathbf{y})}{\partial \theta} \approx \xi_n^{LM}.$$

Hence ξ_n^{LR} has the same asymptotic distribution as ξ_n^{LM} .

Let's show that the LR test is consistent. For this, note that:

$$\frac{\log \mathcal{L}(\hat{\theta}, \mathbf{y}) - \log \mathcal{L}(\hat{\theta}^0, \mathbf{y})}{n} = \frac{1}{n} \sum_{i=1}^n [\log f(y_i; \hat{\theta}_n) - \log f(y_i; \hat{\theta}_n^0)] \rightarrow \mathbb{E}_0[\log f(Y; \theta_0) - \log f(Y; \theta_\infty)],$$

where θ_∞ , the pseudo true value, is such that $h(\theta_\infty) \neq 0$ (by definition of H_1). From the Kullback inequality and the asymptotic identifiability of θ_0 , it follows that $\mathbb{E}_0[\log f(Y; \theta_0) - \log f(Y; \theta_\infty)] > 0$. Therefore $\xi_n^{LR} \rightarrow +\infty$ under H_1 .

Let us now demonstrate the equivalence of ξ^{LM} and ξ^W .

We have (using Eq. ??eq:multiplier)):

$$\xi_n^{LM} = \frac{1}{n} \hat{\lambda}_n' \frac{\partial h(\hat{\theta}_n)}{\partial \theta'} \mathcal{J}(\hat{\theta}_n)^{-1} \frac{\partial h'(\hat{\theta}_n; \mathbf{y})}{\partial \theta} \hat{\lambda}_n.$$

Since, under H_0 , $\hat{\theta}_n^0 \approx \hat{\theta}_n \approx \theta_0$, Eq. (7.16) therefore implies that:

$$\xi^{LM} \approx nh(\hat{\theta}_n)' \left(\frac{\partial h(\hat{\theta}_n)}{\partial \theta'} \mathcal{J}(\hat{\theta}_n)^{-1} \frac{\partial h'(\hat{\theta}_n; \mathbf{y})}{\partial \theta} \right)^{-1} h(\hat{\theta}_n) = \xi^W,$$

which gives the result. \square

Proof of Eq. (6.3)

Proof. We have:

$$\begin{aligned}
 & T\mathbb{E}[(\bar{y}_T - \mu)^2] \\
 = & T\mathbb{E}\left[\left(\frac{1}{T}\sum_{t=1}^T(y_t - \mu)\right)^2\right] = \frac{1}{T}\mathbb{E}\left[\sum_{t=1}^T(y_t - \mu)^2 + 2\sum_{s < t \leq T}(y_t - \mu)(y_s - \mu)\right] \\
 = & \gamma_0 + \frac{2}{T}\left(\sum_{t=2}^T\mathbb{E}[(y_t - \mu)(y_{t-1} - \mu)]\right) + \frac{2}{T}\left(\sum_{t=3}^T\mathbb{E}[(y_t - \mu)(y_{t-2} - \mu)]\right) + \dots \\
 & + \frac{2}{T}\left(\sum_{t=T-1}^T\mathbb{E}[(y_t - \mu)(y_{t-(T-2)} - \mu)]\right) + \frac{2}{T}\mathbb{E}[(y_T - \mu)(y_{T-(T-1)} - \mu)] \\
 = & \gamma_0 + 2\frac{T-1}{T}\gamma_1 + \dots + 2\frac{1}{T}\gamma_{T-1}.
 \end{aligned}$$

Therefore:

$$T\mathbb{E}[(\bar{y}_T - \mu)^2] - \sum_{j=-\infty}^{+\infty}\gamma_j = -2\frac{1}{T}\gamma_1 - 2\frac{2}{T}\gamma_2 - \dots - 2\frac{T-1}{T}\gamma_{T-1} - 2\gamma_T - 2\gamma_{T+1} + \dots$$

And then:

$$\left|T\mathbb{E}[(\bar{y}_T - \mu)^2] - \sum_{j=-\infty}^{+\infty}\gamma_j\right| \leq 2\frac{1}{T}|\gamma_1| + 2\frac{2}{T}|\gamma_2| + \dots + 2\frac{T-1}{T}|\gamma_{T-1}| + 2|\gamma_T| + 2|\gamma_{T+1}| + \dots$$

For any $q \leq T$, we have:

$$\begin{aligned}
 \left|T\mathbb{E}[(\bar{y}_T - \mu)^2] - \sum_{j=-\infty}^{+\infty}\gamma_j\right| &\leq 2\frac{1}{T}|\gamma_1| + 2\frac{2}{T}|\gamma_2| + \dots + 2\frac{q-1}{T}|\gamma_{q-1}| + 2\frac{q}{T}|\gamma_q| + \\
 &\quad 2\frac{q+1}{T}|\gamma_{q+1}| + \dots + 2\frac{T-1}{T}|\gamma_{T-1}| + 2|\gamma_T| + 2|\gamma_{T+1}| + \dots \\
 &\leq \frac{2}{T}(|\gamma_1| + 2|\gamma_2| + \dots + (q-1)|\gamma_{q-1}| + q|\gamma_q|) + \\
 &\quad 2|\gamma_{q+1}| + \dots + 2|\gamma_{T-1}| + 2|\gamma_T| + 2|\gamma_{T+1}| + \dots
 \end{aligned}$$

Consider $\varepsilon > 0$. The fact that the autocovariances are absolutely summable implies that there exists q_0 such that (Cauchy criterion, Theorem 7.2):

$$2|\gamma_{q_0+1}| + 2|\gamma_{q_0+2}| + 2|\gamma_{q_0+3}| + \dots < \varepsilon/2.$$

Then, if $T > q_0$, it comes that:

$$\left|T\mathbb{E}[(\bar{y}_T - \mu)^2] - \sum_{j=-\infty}^{+\infty}\gamma_j\right| \leq \frac{2}{T}(|\gamma_1| + 2|\gamma_2| + \dots + (q_0-1)|\gamma_{q_0-1}| + q_0|\gamma_{q_0}|) + \varepsilon/2.$$

If $T \geq 2 \left(|\gamma_1| + 2|\gamma_2| + \dots + (q_0 - 1)|\gamma_{q_0-1}| + q_0|\gamma_{q_0}| \right) / (\varepsilon/2)$ ($= f(q_0)$, say) then

$$\frac{2}{T} \left(|\gamma_1| + 2|\gamma_2| + \dots + (q_0 - 1)|\gamma_{q_0-1}| + q_0|\gamma_{q_0}| \right) \leq \varepsilon/2.$$

Then, if $T > f(q_0)$ and $T > q_0$, i.e. if $T > \max(f(q_0), q_0)$, we have:

$$\left| T \mathbb{E} [(\bar{y}_T - \mu)^2] - \sum_{j=-\infty}^{+\infty} \gamma_j \right| \leq \varepsilon.$$

□

Proof of Proposition 6.15

Proof. We have:

$$\begin{aligned} \mathbb{E}([y_{t+1} - y_{t+1}^*]^2) &= \mathbb{E}([\{y_{t+1} - \mathbb{E}(y_{t+1}|x_t)\} + \{\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*\}]^2) \\ &= \mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)]^2) + \mathbb{E}([\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*]^2) \\ &\quad + 2\mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)][\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*]). \end{aligned} \quad (7.18)$$

Let us focus on the last term. We have:

$$\begin{aligned} &\mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)][\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*]) \\ &= \mathbb{E}(\mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)][\underbrace{\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*}_{\text{function of } x_t}]|x_t)) \\ &= \mathbb{E}([\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*]\mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)]|x_t)) \\ &= \mathbb{E}([\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*]\underbrace{[\mathbb{E}(y_{t+1}|x_t) - \mathbb{E}(y_{t+1}|x_t)]}_{=0}) = 0. \end{aligned}$$

Therefore, Eq. (7.18) becomes:

$$\begin{aligned} &\mathbb{E}([y_{t+1} - y_{t+1}^*]^2) \\ &= \underbrace{\mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)]^2)}_{\geq 0 \text{ and does not depend on } y_{t+1}^*} + \underbrace{\mathbb{E}([\mathbb{E}(y_{t+1}|x_t) - y_{t+1}^*]^2)}_{\geq 0 \text{ and depends on } y_{t+1}^*}. \end{aligned}$$

This implies that $\mathbb{E}([y_{t+1} - y_{t+1}^*]^2)$ is always larger than $\mathbb{E}([y_{t+1} - \mathbb{E}(y_{t+1}|x_t)]^2)$, and is therefore minimized if the second term is equal to zero, that is if $\mathbb{E}(y_{t+1}|x_t) = y_{t+1}^*$. □

Proof of Proposition 6.12

Proof. Using Proposition ?? (in Appendix ??), we obtain that, conditionally on x_1 , the log-likelihood is given by

$$\begin{aligned}\log \mathcal{L}(Y_T; \theta) &= -(Tn/2) \log(2\pi) + (T/2) \log |\Omega^{-1}| \\ &\quad - \frac{1}{2} \sum_{t=1}^T [(y_t - \Pi' x_t)' \Omega^{-1} (y_t - \Pi' x_t)].\end{aligned}$$

Let's rewrite the last term of the log-likelihood:

$$\begin{aligned}\sum_{t=1}^T [(y_t - \Pi' x_t)' \Omega^{-1} (y_t - \Pi' x_t)] &= \\ \sum_{t=1}^T [(y_t - \hat{\Pi}' x_t + \hat{\Pi}' x_t - \Pi' x_t)' \Omega^{-1} (y_t - \hat{\Pi}' x_t + \hat{\Pi}' x_t - \Pi' x_t)] &= \\ \sum_{t=1}^T [(\hat{\varepsilon}_t + (\hat{\Pi} - \Pi)' x_t)' \Omega^{-1} (\hat{\varepsilon}_t + (\hat{\Pi} - \Pi)' x_t)],\end{aligned}$$

where the j^{th} element of the $(n \times 1)$ vector $\hat{\varepsilon}_t$ is the sample residual, for observation t , from an OLS regression of $y_{j,t}$ on x_t . Expanding the previous equation, we get:

$$\begin{aligned}\sum_{t=1}^T [(y_t - \Pi' x_t)' \Omega^{-1} (y_t - \Pi' x_t)] &= \sum_{t=1}^T \hat{\varepsilon}_t' \Omega^{-1} \hat{\varepsilon}_t \\ &\quad + 2 \sum_{t=1}^T \hat{\varepsilon}_t' \Omega^{-1} (\hat{\Pi} - \Pi)' x_t + \sum_{t=1}^T x_t' (\hat{\Pi} - \Pi) \Omega^{-1} (\hat{\Pi} - \Pi)' x_t.\end{aligned}$$

Let's apply the trace operator on the second term (that is a scalar):

$$\begin{aligned}\sum_{t=1}^T \hat{\varepsilon}_t' \Omega^{-1} (\hat{\Pi} - \Pi)' x_t &= Tr \left(\sum_{t=1}^T \hat{\varepsilon}_t' \Omega^{-1} (\hat{\Pi} - \Pi)' x_t \right) \\ &= Tr \left(\sum_{t=1}^T \Omega^{-1} (\hat{\Pi} - \Pi)' x_t \hat{\varepsilon}_t' \right) = Tr \left(\Omega^{-1} (\hat{\Pi} - \Pi)' \sum_{t=1}^T x_t \hat{\varepsilon}_t' \right).\end{aligned}$$

Given that, by construction (property of OLS estimates), the sample residuals are orthogonal to the explanatory variables, this term is zero. Introducing $\tilde{x}_t = (\hat{\Pi} - \Pi)' x_t$, we have

$$\sum_{t=1}^T [(y_t - \Pi' x_t)' \Omega^{-1} (y_t - \Pi' x_t)] = \sum_{t=1}^T \hat{\varepsilon}_t' \Omega^{-1} \hat{\varepsilon}_t + \sum_{t=1}^T \tilde{x}_t' \Omega^{-1} \tilde{x}_t.$$

Since Ω is a positive definite matrix, Ω^{-1} is as well. Consequently, the smallest value that the last term can take is obtained for $\tilde{x}_t = 0$, i.e. when $\Pi = \hat{\Pi}$.

The MLE of Ω is the matrix $\hat{\Omega}$ that maximizes $\Omega \xrightarrow{\ell} L(Y_T; \hat{\Pi}, \Omega)$. We have:

$$\log \mathcal{L}(Y_T; \hat{\Pi}, \Omega) = -(Tn/2) \log(2\pi) + (T/2) \log |\Omega^{-1}| - \frac{1}{2} \sum_{t=1}^T [\hat{\varepsilon}'_t \Omega^{-1} \hat{\varepsilon}_t].$$

Matrix $\hat{\Omega}$ is a symmetric positive definite. It is easily checked that the (unrestricted) matrix that maximizes the latter expression is symmetric positive definite matrix. Indeed:

$$\frac{\partial \log \mathcal{L}(Y_T; \hat{\Pi}, \Omega)}{\partial \Omega} = \frac{T}{2} \Omega' - \frac{1}{2} \sum_{t=1}^T \hat{\varepsilon}_t \hat{\varepsilon}'_t \Rightarrow \hat{\Omega}' = \frac{1}{T} \sum_{t=1}^T \hat{\varepsilon}_t \hat{\varepsilon}'_t,$$

which leads to the result. \square

Proof of Proposition 6.13

Proof. Let us drop the i subscript. Rearranging Eq. (6.46), we have:

$$\sqrt{T}(\mathbf{b} - \beta) = (X' X/T)^{-1} \sqrt{T}(X' \varepsilon/T).$$

Let us consider the autocovariances of $\mathbf{v}_t = x_t \varepsilon_t$, denoted by γ_j^v . Using the fact that x_t is a linear combination of past ε_t s and that ε_t is a white noise, we get that $\mathbb{E}(\varepsilon_t x_t) = 0$. Therefore

$$\gamma_j^v = \mathbb{E}(\varepsilon_t \varepsilon_{t-j} x_t x'_{t-j}).$$

If $j > 0$, we have $\mathbb{E}(\varepsilon_t \varepsilon_{t-j} x_t x'_{t-j}) = \mathbb{E}(\mathbb{E}[\varepsilon_t \varepsilon_{t-j} x_t x'_{t-j} | \varepsilon_{t-j}, x_t, x_{t-j}]) = \mathbb{E}(\varepsilon_{t-j} x_t x'_{t-j} \mathbb{E}[\varepsilon_t | \varepsilon_{t-j}, x_t, x_{t-j}]) = 0$. Note that we have $\mathbb{E}[\varepsilon_t | \varepsilon_{t-j}, x_t, x_{t-j}] = 0$ because $\{\varepsilon_t\}$ is an i.i.d. white noise sequence. If $j = 0$, we have:

$$\gamma_0^v = \mathbb{E}(\varepsilon_t^2 x_t x'_t) = \mathbb{E}(\varepsilon_t^2) \mathbb{E}(x_t x'_t) = \sigma^2 \mathbf{Q}.$$

The convergence in distribution of $\sqrt{T}(X' \varepsilon/T) = \sqrt{T} \frac{1}{T} \sum_{t=1}^T v_t$ results from the Central Limit Theorem for covariance-stationary processes, using the γ_j^v computed above. \square

7.6 Additional codes

7.6.1 Simulating GEV distributions

The following lines of code have been used to generate Figure 5.6.

```

n.sim <- 4000
par(mfrow=c(1,3),
    plt=c(.2,.95,.2,.85))
all.rhos <- c(.3,.6,.95)
for(j in 1:length(all.rhos)){
  theta <- 1/all.rhos[j]
  v1 <- runif(n.sim)
  v2 <- runif(n.sim)
  w <- rep(.000001,n.sim)
  # solve for f(w) = w*(1 - log(w)/theta) - v2 = 0
  for(i in 1:20){
    f.i <- w * (1 - log(w)/theta) - v2
    f.prime <- 1 - log(w)/theta - 1/theta
    w <- w - f.i/f.prime
  }
  u1 <- exp(v1^(1/theta) * log(w))
  u2 <- exp((1-v1)^(1/theta) * log(w))

  # Get eps1 and eps2 using the inverse of
  # the Gumbel distribution's cdf:
  eps1 <- -log(-log(u1))
  eps2 <- -log(-log(u2))
  cbind(cor(eps1,eps2),1-all.rhos[j]^2)
  plot(eps1,eps2,pch=19,col="#FF000044",
       main=paste("rho = ",toString(all.rhos[j]),sep=""),
       xlab=expression(epsilon[1]),
       ylab=expression(epsilon[2]),
       cex.lab=2,cex.main=1.5)
}

```

7.6.2 Computing the covariance matrix of IRF using the delta method

```

irf.function <- function(THETA){
  c <- THETA[1]
  phi <- THETA[2:(p+1)]
  if(q>0){
    theta <- c(1,THETA[(1+p+1):(1+p+q)])
  }else{
    theta <- 1
  }
  sigma <- THETA[1+p+q+1]
  r <- dim(Matrix.of.Exog)[2] - 1

```

```
beta <- THETA[(1+p+q+1):(1+p+q+1+(r+1))]

irf <- sim.arma(0,phi,beta,sigma=sd(Ramey$ED3_TC,na.rm=TRUE),T=60,
                y.0=rep(0,length(x$phi)),nb.sim=1,make.IRF=1,
                X=NaN,beta=NaN)
return(irf)
}

IRF.0 <- 100*irf.function(x$THETA)
eps <- .00000001
d.IRF <- NULL
for(i in 1:length(x$THETA)){
  THETA.i <- x$THETA
  THETA.i[i] <- THETA.i[i] + eps
  IRF.i <- 100*irf.function(THETA.i)
  d.IRF <- cbind(d.IRF,
                  (IRF.i - IRF.0)/eps
                  )
}
mat.var.cov.IRF <- d.IRF %*% x$I %*% t(d.IRF)
```

7.7 Statistical Tables

Table 7.1: Quantiles of the $\mathcal{N}(0, 1)$ distribution. If a and b are respectively the row and column number; then the corresponding cell gives $\mathbb{P}(0 < X \leq a + b)$, where $X \sim \mathcal{N}(0, 1)$.

	0	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0	0.5000	0.6179	0.7257	0.8159	0.8849	0.9332	0.9641	0.9821	0.9918	0.9965
0.1	0.5040	0.6217	0.7291	0.8186	0.8869	0.9345	0.9649	0.9826	0.9920	0.9966
0.2	0.5080	0.6255	0.7324	0.8212	0.8888	0.9357	0.9656	0.9830	0.9922	0.9967
0.3	0.5120	0.6293	0.7357	0.8238	0.8907	0.9370	0.9664	0.9834	0.9925	0.9968
0.4	0.5160	0.6331	0.7389	0.8264	0.8925	0.9382	0.9671	0.9838	0.9927	0.9969
0.5	0.5199	0.6368	0.7422	0.8289	0.8944	0.9394	0.9678	0.9842	0.9929	0.9970
0.6	0.5239	0.6406	0.7454	0.8315	0.8962	0.9406	0.9686	0.9846	0.9931	0.9971
0.7	0.5279	0.6443	0.7486	0.8340	0.8980	0.9418	0.9693	0.9850	0.9932	0.9972
0.8	0.5319	0.6480	0.7517	0.8365	0.8997	0.9429	0.9699	0.9854	0.9934	0.9973
0.9	0.5359	0.6517	0.7549	0.8389	0.9015	0.9441	0.9706	0.9857	0.9936	0.9974
1	0.5398	0.6554	0.7580	0.8413	0.9032	0.9452	0.9713	0.9861	0.9938	0.9974
1.1	0.5438	0.6591	0.7611	0.8438	0.9049	0.9463	0.9719	0.9864	0.9940	0.9975
1.2	0.5478	0.6628	0.7642	0.8461	0.9066	0.9474	0.9726	0.9868	0.9941	0.9976
1.3	0.5517	0.6664	0.7673	0.8485	0.9082	0.9484	0.9732	0.9871	0.9943	0.9977
1.4	0.5557	0.6700	0.7704	0.8508	0.9099	0.9495	0.9738	0.9875	0.9945	0.9977
1.5	0.5596	0.6736	0.7734	0.8531	0.9115	0.9505	0.9744	0.9878	0.9946	0.9978
1.6	0.5636	0.6772	0.7764	0.8554	0.9131	0.9515	0.9750	0.9881	0.9948	0.9979
1.7	0.5675	0.6808	0.7794	0.8577	0.9147	0.9525	0.9756	0.9884	0.9949	0.9979
1.8	0.5714	0.6844	0.7823	0.8599	0.9162	0.9535	0.9761	0.9887	0.9951	0.9980
1.9	0.5753	0.6879	0.7852	0.8621	0.9177	0.9545	0.9767	0.9890	0.9952	0.9981
2	0.5793	0.6915	0.7881	0.8643	0.9192	0.9554	0.9772	0.9893	0.9953	0.9981
2.1	0.5832	0.6950	0.7910	0.8665	0.9207	0.9564	0.9778	0.9896	0.9955	0.9982
2.2	0.5871	0.6985	0.7939	0.8686	0.9222	0.9573	0.9783	0.9898	0.9956	0.9982
2.3	0.5910	0.7019	0.7967	0.8708	0.9236	0.9582	0.9788	0.9901	0.9957	0.9983
2.4	0.5948	0.7054	0.7995	0.8729	0.9251	0.9591	0.9793	0.9904	0.9959	0.9984
2.5	0.5987	0.7088	0.8023	0.8749	0.9265	0.9599	0.9798	0.9906	0.9960	0.9984
2.6	0.6026	0.7123	0.8051	0.8770	0.9279	0.9608	0.9803	0.9909	0.9961	0.9985
2.7	0.6064	0.7157	0.8078	0.8790	0.9292	0.9616	0.9808	0.9911	0.9962	0.9985
2.8	0.6103	0.7190	0.8106	0.8810	0.9306	0.9625	0.9812	0.9913	0.9963	0.9986
2.9	0.6141	0.7224	0.8133	0.8830	0.9319	0.9633	0.9817	0.9916	0.9964	0.9986

Table 7.2: Quantiles of the Student- t distribution. The rows correspond to different degrees of freedom (ν , say); the columns correspond to different probabilities (z , say). The cell gives q that is s.t. $\mathbb{P}(-q < X < q) = z$, with $X \sim t(\nu)$.

	0.05	0.1	0.75	0.9	0.95	0.975	0.99	0.999
1	0.079	0.158	2.414	6.314	12.706	25.452	63.657	636.619
2	0.071	0.142	1.604	2.920	4.303	6.205	9.925	31.599
3	0.068	0.137	1.423	2.353	3.182	4.177	5.841	12.924
4	0.067	0.134	1.344	2.132	2.776	3.495	4.604	8.610
5	0.066	0.132	1.301	2.015	2.571	3.163	4.032	6.869
6	0.065	0.131	1.273	1.943	2.447	2.969	3.707	5.959
7	0.065	0.130	1.254	1.895	2.365	2.841	3.499	5.408
8	0.065	0.130	1.240	1.860	2.306	2.752	3.355	5.041
9	0.064	0.129	1.230	1.833	2.262	2.685	3.250	4.781
10	0.064	0.129	1.221	1.812	2.228	2.634	3.169	4.587
20	0.063	0.127	1.185	1.725	2.086	2.423	2.845	3.850
30	0.063	0.127	1.173	1.697	2.042	2.360	2.750	3.646
40	0.063	0.126	1.167	1.684	2.021	2.329	2.704	3.551
50	0.063	0.126	1.164	1.676	2.009	2.311	2.678	3.496
60	0.063	0.126	1.162	1.671	2.000	2.299	2.660	3.460
70	0.063	0.126	1.160	1.667	1.994	2.291	2.648	3.435
80	0.063	0.126	1.159	1.664	1.990	2.284	2.639	3.416
90	0.063	0.126	1.158	1.662	1.987	2.280	2.632	3.402
100	0.063	0.126	1.157	1.660	1.984	2.276	2.626	3.390
200	0.063	0.126	1.154	1.653	1.972	2.258	2.601	3.340
500	0.063	0.126	1.152	1.648	1.965	2.248	2.586	3.310

Table 7.3: Quantiles of the χ^2 distribution. The rows correspond to different degrees of freedom; the columns correspond to different probabilities.

	0.05	0.1	0.75	0.9	0.95	0.975	0.99	0.999
1	0.004	0.016	1.323	2.706	3.841	5.024	6.635	10.828
2	0.103	0.211	2.773	4.605	5.991	7.378	9.210	13.816
3	0.352	0.584	4.108	6.251	7.815	9.348	11.345	16.266
4	0.711	1.064	5.385	7.779	9.488	11.143	13.277	18.467
5	1.145	1.610	6.626	9.236	11.070	12.833	15.086	20.515
6	1.635	2.204	7.841	10.645	12.592	14.449	16.812	22.458
7	2.167	2.833	9.037	12.017	14.067	16.013	18.475	24.322
8	2.733	3.490	10.219	13.362	15.507	17.535	20.090	26.124
9	3.325	4.168	11.389	14.684	16.919	19.023	21.666	27.877
10	3.940	4.865	12.549	15.987	18.307	20.483	23.209	29.588
20	10.851	12.443	23.828	28.412	31.410	34.170	37.566	45.315
30	18.493	20.599	34.800	40.256	43.773	46.979	50.892	59.703
40	26.509	29.051	45.616	51.805	55.758	59.342	63.691	73.402
50	34.764	37.689	56.334	63.167	67.505	71.420	76.154	86.661
60	43.188	46.459	66.981	74.397	79.082	83.298	88.379	99.607
70	51.739	55.329	77.577	85.527	90.531	95.023	100.425	112.317
80	60.391	64.278	88.130	96.578	101.879	106.629	112.329	124.839
90	69.126	73.291	98.650	107.565	113.145	118.136	124.116	137.208
100	77.929	82.358	109.141	118.498	124.342	129.561	135.807	149.449
200	168.279	174.835	213.102	226.021	233.994	241.058	249.445	267.541
500	449.147	459.926	520.950	540.930	553.127	563.852	576.493	603.446

Table 7.4: Quantiles of the \mathcal{F} distribution. The columns and rows correspond to different degrees of freedom (resp. n_1 and n_2). The different panels correspond to different probabilities (α) The corresponding cell gives z that is s.t. $\mathbb{P}(X \leq z) = \alpha$, with $X \sim \mathcal{F}(n_1, n_2)$.

	1	2	3	4	5	6	7	8	9	10
alpha = 0.9										
5	4.060	3.780	3.619	3.520	3.453	3.405	3.368	3.339	3.316	3.297
10	3.285	2.924	2.728	2.605	2.522	2.461	2.414	2.377	2.347	2.323
15	3.073	2.695	2.490	2.361	2.273	2.208	2.158	2.119	2.086	2.059
20	2.975	2.589	2.380	2.249	2.158	2.091	2.040	1.999	1.965	1.937
50	2.809	2.412	2.197	2.061	1.966	1.895	1.840	1.796	1.760	1.729
100	2.756	2.356	2.139	2.002	1.906	1.834	1.778	1.732	1.695	1.663
500	2.716	2.313	2.095	1.956	1.859	1.786	1.729	1.683	1.644	1.612
alpha = 0.95										
5	6.608	5.786	5.409	5.192	5.050	4.950	4.876	4.818	4.772	4.735
10	4.965	4.103	3.708	3.478	3.326	3.217	3.135	3.072	3.020	2.978
15	4.543	3.682	3.287	3.056	2.901	2.790	2.707	2.641	2.588	2.544
20	4.351	3.493	3.098	2.866	2.711	2.599	2.514	2.447	2.393	2.348
50	4.034	3.183	2.790	2.557	2.400	2.286	2.199	2.130	2.073	2.026
100	3.936	3.087	2.696	2.463	2.305	2.191	2.103	2.032	1.975	1.927
500	3.860	3.014	2.623	2.390	2.232	2.117	2.028	1.957	1.899	1.850
alpha = 0.99										
5	16.258	13.274	12.060	11.392	10.967	10.672	10.456	10.289	10.158	10.051
10	10.044	7.559	6.552	5.994	5.636	5.386	5.200	5.057	4.942	4.849
15	8.683	6.359	5.417	4.893	4.556	4.318	4.142	4.004	3.895	3.805
20	8.096	5.849	4.938	4.431	4.103	3.871	3.699	3.564	3.457	3.368
50	7.171	5.057	4.199	3.720	3.408	3.186	3.020	2.890	2.785	2.698
100	6.895	4.824	3.984	3.513	3.206	2.988	2.823	2.694	2.590	2.503
500	6.686	4.648	3.821	3.357	3.054	2.838	2.675	2.547	2.443	2.356

Bibliography

- Abadie, A. and Cattaneo, M. D. (2018). Econometric Methods for Program Evaluation. *Annual Review of Economics*, 10(1):465–503.
- Anderson, T. (1971). *The Statistical Analysis of Time Series*. Wiley.
- Anderson, T. W. and Hsiao, C. (1982). Formulation and estimation of dynamic models using panel data. *Journal of Econometrics*, 18(1):47–82.
- Andrews, I., Stock, J. H., and Sun, L. (2019). Weak instruments in instrumental variables regression: Theory and practice. *Annual Review of Economics*, 11(1):727–753.
- Angrist, J. D. and Pischke, J.-S. (2008). *Mostly Harmless Econometrics: An Empiricist's Companion*. Princeton University Press.
- Arellano, M. and Bond, S. (1991). Some Tests of Specification for Panel Data: Monte Carlo Evidence and an Application to Employment Equations. *Review of Economic Studies*, 58(2):277–297.
- Arias, J. E., Rubio-Ramírez, J. F., and Waggoner, D. F. (2018). Inference based on structural vector autoregressions identified with sign and zero restrictions: Theory and applications. *Econometrica*, 86(2):685–720.
- Barsky, R. B. and Sims, E. R. (2011). News shocks and business cycles. *Journal of Monetary Economics*, 58(3):273–289.
- Bernanke, B. S. (1986). Alternative explanations of the money-income correlation. *Carnegie-Rochester Conference Series on Public Policy*, 25:49–99.
- Bernanke, B. S., Boivin, J., and Eliasz, P. (2005). Measuring the Effects of Monetary Policy: A Factor-Augmented Vector Autoregressive (FAVAR) Approach*. *The Quarterly Journal of Economics*, 120(1):387–422.
- Blanchard, O. J. and Quah, D. (1989). The Dynamic Effects of Aggregate Demand and Supply Disturbances. *American Economic Review*, 79(4):655–673.

- Box, G. and Jenkins, G. M. (1976). *Time Series Analysis: Forecasting and Control*. Holden-Day.
- Cameron, A. C. and Miller, D. L. (2014). A practitioner's guide to cluster-robust inference. *The Journal of Human Resources*, 50(2).
- Cameron, A. C. and Trivedi, P. K. (2005). *Microeconometrics: Methods and Applications*. Cambridge University Press.
- Christiano, L. J., Eichenbaum, M., and Evans, C. (1996). The effects of monetary policy shocks: Evidence from the flow of funds. *The Review of Economics and Statistics*, 78(1):16–34.
- Christiano, L. J., Eichenbaum, M., and Vigfusson, R. (2007). *Assessing Structural VARs*, pages 1–106. MIT Press.
- Cochrane, D. and Orcutt, G. H. (1949). Application of least squares regression to relationships containing auto-correlated error terms. *Journal of the American Statistical Association*, 44(245):32–61.
- Cochrane, J. H. and Piazzesi, M. (2002). The Fed and Interest Rates - A High-Frequency Identification. *American Economic Review*, 92(2):90–95.
- Danne, C. (2015). VARsignR: Estimating VARs using sign restrictions in R. MPRA Paper 68429, University Library of Munich, Germany.
- De Gooijer, J. G. and Hyndman, R. J. (2006). 25 years of time series forecasting. *International Journal of Forecasting*, 22(3):443–473. Twenty five years of forecasting.
- Dedola, L. and Lippi, F. (2005). The monetary transmission mechanism: Evidence from the industries of five oecd countries. *European Economic Review*, 49(6):1543–1569.
- Diebold, F. and Mariano, R. (1995). Comparing predictive accuracy. *Journal of Business & Economic Statistics*, 13(3):253–63.
- Duan, N., Manning, W. G., Morris, C. N., and Newhouse, J. P. (1983). A comparison of alternative models for the demand for medical care. *Journal of Business & Economic Statistics*, 1(2):115–126.
- Durbin, J. (1954). Errors in variables. *Revue de l'Institut International de Statistique / Review of the International Statistical Institute*, 22(1/3):23–32.
- Durbin, J. and Watson, G. S. (1950). Testing for Serial Correlation in Least Squares Regression. I. *Biometrika*, 37(3-4):409–428.
- Durbin, J. and Watson, G. S. (1951). Testing for Serial Correlation in Least Squares Regression. II. *Biometrika*, 38(1-2):159–178.

- Erceg, C. J., Guerrieri, L., and Gust, C. (2005). Can Long-Run Restrictions Identify Technology Shocks? *Journal of the European Economic Association*, 3(6):1237–1278.
- Faust, J. and Leeper, E. M. (1997). When do long-run identifying restrictions give reliable results? *Journal of Business & Economic Statistics*, 15(3):345–353.
- Fischer, S. (1977). Long-term contracts, rational expectations, and the optimal money supply rule. *Journal of Political Economy*, 85(1):191–205.
- Fritsch, M., Pua, A. A. Y., and Schnurbus, J. (2019). Pdynmc - An R-package for estimating linear dynamic panel data models based on linear and non-linear moment conditions. Passauer Diskussionspapiere, Betriebswirtschaftliche Reihe B-39-19, University of Passau, Faculty of Business and Economics.
- Galí, J. (1999). Technology, employment, and the business cycle: Do technology shocks explain aggregate fluctuations? *American Economic Review*, 89(1):249–271.
- Galí, J. (1992). How well does the is-lm model fit postwar u.s. data? *The Quarterly Journal of Economics*, 107(2):709–738.
- Gerlach, S. and Smets, F. (1995). The Monetary Transmission Mechanism: Evidence from the G-7 Countries. CEPR Discussion Papers 1219, C.E.P.R. Discussion Papers.
- Gertler, M. and Karadi, P. (2015). Monetary Policy Surprises, Credit Costs, and Economic Activity. *American Economic Journal: Macroeconomics*, 7(1):44–76.
- Gouriéroux, C. and Monfort, A. (1995). *Statistics and Econometric Models*, volume 1 of *Themes in Modern Econometrics*. Cambridge University Press.
- Gouriéroux, C., Monfort, A., and Renne, J.-P. (2020). Identification and Estimation in Non-Fundamental Structural VARMA Models. *Review of Economic Studies*, 87(4):1915–1953.
- Gouriéroux, C., Monfort, A., and Renne, J.-P. (2017). Statistical inference for independent component analysis: Application to structural var models. *Journal of Econometrics*, 196(1):111–126.
- Granger, C. W. J. (1969). Investigating causal relations by econometric models and cross-spectral methods. *Econometrica*, 37(3):424–438.
- Greene, W. H. (2003). *Econometric Analysis*. Pearson Education, fifth edition.
- Gürkaynak, R. S., Sack, B., and Swanson, E. (2005). Do Actions Speak Louder Than Words? The Response of Asset Prices to Monetary Policy Actions and Statements. *International Journal of Central Banking*, 1(1).

- Hamilton, J. D. (1994). *Time Series Analysis*. Princeton University Press.
- Hansen, L. P. (1982). Large sample properties of generalized method of moments estimators. *Econometrica*, 50(4):1029–1054.
- Hausman, J. A. (1978). Specification tests in econometrics. *Econometrica*, 46(6):1251–1271.
- Heiss, F. (2002). Structural choice analysis with nested logit models. *Stata Journal*, 2(3):227–252(26).
- Hensher, D. and Greene, W. (2002). Specification and estimation of the nested logit model: alternative normalisations. *Transportation Research Part B: Methodological*, 36(1):1–17.
- James, G., Witten, D., Hastie, T., and Tibshirani, R. (2013). *An Introduction to Statistical Learning: with Applications in R*. Springer.
- Jordà, O. (2005). Estimation and Inference of Impulse Responses by Local Projections. *American Economic Review*, 95(1):161–182.
- Jordà, O., Schularick, M., and Taylor, A. M. (2017). Macrofinancial History and the New Business Cycle Facts. *NBER Macroeconomics Annual*, 31(1):213–263.
- Kilian, L. (1998). Small-sample confidence intervals for impulse response functions. *The Review of Economics and Statistics*, 80(2):218–230.
- Kim, J. H. (2022). *VAR. etp: VAR Modelling: Estimation, Testing, and Prediction*. R package version 1.0.
- Korobilis, D. (2013). Var forecasting using bayesian variable selection. *Journal of Applied Econometrics*, 28(2):204–230.
- Kuttner, K. N. (2001). Monetary policy surprises and interest rates: Evidence from the fed funds futures market. *Journal of Monetary Economics*, 47(3):523–544.
- Lanne, M. and Lütkepohl, H. (2008). Identifying monetary policy shocks via changes in volatility. *Journal of Money, Credit and Banking*, 40(6):1131–1149.
- Lanne, M., Lütkepohl, H., and Maciejowska, K. (2010). Structural vector autoregressions with markov switching. *Journal of Economic Dynamics and Control*, 34(2):121–131.
- Litterman, R. and Scheinkman, J. (1991). Common Factors Affecting Bond Returns. *Journal of Fixed Income*, (1):54–61.
- Lütkepohl, H. (1990). Asymptotic distributions of impulse response functions and forecast error variance decompositions of vector autoregressive models. *The Review of Economics and Statistics*, 72(1):116–25.

- Lütkepohl, H. and Netšunajev, A. (2017). Structural vector autoregressions with heteroskedasticity: A review of different volatility models. *Econometrics and Statistics*, 1:2–18.
- MacKinnon, J. and White, H. (1985). Some heteroskedasticity-consistent covariance matrix estimators with improved finite sample properties. *Journal of Econometrics*, 29(3):305–325.
- MacKinnon, J. G., Ørregaard Nielsen, M., and Webb, M. D. (2022). Cluster-robust inference: A guide to empirical practice. *Journal of Econometrics*.
- Marcellino, M., Stock, J. H., and Watson, M. W. (2006). A comparison of direct and iterated multistep AR methods for forecasting macroeconomic time series. *Journal of Econometrics*, 135(1-2):499–526.
- McCracken, M. W. and Ng, S. (2016). FRED-MD: A Monthly Database for Macroeconomic Research. *Journal of Business & Economic Statistics*, 34(4):574–589.
- Meyer, B. D., Viscusi, W. K., and Durbin, D. L. (1995). Workers' Compensation and Injury Duration: Evidence from a Natural Experiment. *American Economic Review*, 85(3):322–340.
- Mroz, T. A. (1987). The sensitivity of an empirical model of married women's hours of work to economic and statistical assumptions. *Econometrica*, 55(4):765–799.
- Nakosteen, R. A. and Zimmer, M. (1980). Migration and income: The question of self-selection. *Southern Economic Journal*, 46(3):840–851.
- Newey, W. K. and West, K. D. (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, 55(3):703–708.
- Normandin, M. and Phaneuf, L. (2004). Monetary policy shocks:: Testing identification conditions under time-varying conditional volatility. *Journal of Monetary Economics*, 51(6):1217–1243.
- Pfaff, B. (2008). Var, svar and svec models: Implementation within R package vars. *Journal of Statistical Software*, 27(4).
- Piazzesi, M. and Swanson, E. T. (2008). Futures prices as risk-adjusted forecasts of monetary policy. *Journal of Monetary Economics*, 55(4):677–691.
- Ramey, V. A. (2016). Macroeconomic Shocks and Their Propagation. NBER Working Papers 21978, National Bureau of Economic Research, Inc.
- Rigobon, R. (2003). Identification Through Heteroskedasticity. *The Review of Economics and Statistics*, 85(4):777–792.

- Roberts, G. O., Gelman, A., and Gilks, W. R. (1997). Weak convergence and optimal scaling of random walk metropolis algorithms. *The Annals of Applied Probability*, 7(1):110–120.
- Romer, C. D. and Romer, D. H. (2004). A New Measure of Monetary Shocks: Derivation and Implications. *American Economic Review*, 94(4):1055–1084.
- Ruibio-Ramírez, J. F., Waggoner, D. F., and Zha, T. (2010). Structural vector autoregressions: Theory of identification and algorithms for inference. *The Review of Economic Studies*, 77(2):665–696.
- Sargan, J. D. (1958). The estimation of economic relationships using instrumental variables. *Econometrica*, 26(3):393–415.
- Sims, C. A. (1980). Macroeconomics and reality. *Econometrica*, 48(1):1–48.
- Sims, C. A. (1986). Are forecasting models usable for policy analysis? *Quarterly Review*, 10(Win):2–16.
- Stock, J. and Watson, M. (2016). Dynamic Factor Models, Factor-Augmented Vector Autoregressions, and Structural Vector Autoregressions in Macroeconomics. In Taylor, J. B. and Uhlig, H., editors, *Handbook of Macroeconomics*, volume 2 of *Handbook of Macroeconomics*, chapter 0, pages 415–525. Elsevier.
- Stock, J. and Watson, M. W. (2003). *Introduction to Econometrics*. Prentice Hall, New York.
- Stock, J. H. and Watson, M. W. (2018). Identification and estimation of dynamic causal effects in macroeconomics using external instruments. *The Economic Journal*, 128(610):917–948.
- Stock, J. H. and Yogo, M. (2005). *Testing for Weak Instruments in Linear IV Regression*, page 80–108. Cambridge University Press.
- Tibshirani, R. (2011). Regression shrinkage and selection via the lasso: a retrospective. *Journal of the Royal Statistical Society Series B*, 73(3):273–282.
- Tobin, J. (1956). Estimation of relationships for limited dependent variables. Cowles Foundation Discussion Papers 3R, Cowles Foundation for Research in Economics, Yale University.
- Uhlig, H. (2004). What moves GNP? Econometric Society 2004 North American Winter Meetings 636, Econometric Society.
- Uhlig, H. (2005). What are the effects of monetary policy on output? results from an agnostic identification procedure. *Journal of Monetary Economics*, 52(2):381–419.
- White, H. (1980). A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica*, 48(4):817–838.
- Wu, D.-M. (1973). Alternative tests of independence between stochastic regressors and disturbances. *Econometrica*, 41(4):733–750.