
An Introduction to Calculus

Rob Hayward

Friday, 20th September 2013

Usain Bolt



Average and Instantaneous Speed

Average and Instantaneous Speed

Slope is constant with average speed but varies over time.

■ Average Speed = $\frac{Distance}{Time}$

Average and Instantaneous Speed

- Average Speed = $\frac{Distance}{Time}$
- Average Speed = $\frac{100}{9.58}$ = 10.43 $\frac{m}{s}$

Average and Instantaneous Speed

- Average Speed = $\frac{Distance}{Time}$
- Average Speed = $\frac{100}{9.58}$ = 10.43 $\frac{m}{s}$
- Need to find the tangent to the line for Instantaneous Speed

Average and Instantaneous Speed

- Average Speed = $\frac{\textit{Distance}}{\textit{Time}}$
- Average Speed = $\frac{100}{9.58}$ = 10.43 $\frac{m}{s}$
- Need to find the tangent to the line for Instantaneous Speed
- Instantaneous Speed = $\frac{\Delta y}{\Delta x}$

Small change (h)

Instantaneous speed is the tangent to the curve. For f(x)

Small change (h)

Instantaneous speed is the tangent to the curve. For f(x)

■ Instantaneous speed = $\frac{f(x_0+h_0)-f(x_0)}{(x_0+h_0)-x_0}$

Small change (h)

Instantaneous speed is the tangent to the curve. For f(x)

- Instantaneous speed = $\frac{f(x_0+h_0)-f(x_0)}{(x_0+h_0)-x_0}$
- \blacksquare h_0 is small


Small change (h)

Instantaneous speed is the tangent to the curve. For f(x)

- Instantaneous speed = $\frac{f(x_0+h_0)-f(x_0)}{(x_0+h_0)-x_0}$
- \blacksquare h_0 is small
- Instantaneous speed = $\frac{f(x_0+h_0)-f(x_0)}{h_0}$

Example page 1

$$f(x) = y = x^2$$



Example page 2

Instantaneous change at (3, 9) when $f(x) = x^2$

Instantaneous speed = $\frac{f(x_0 + h_0) - f(x_0)}{h_0}$

h	x + h	f(x + h)	$\frac{f(x+h)-f(x)}{h}$
0.1	3.1	9.61	6.1
0.01	3.01	9.0601	6.01
0.001	3.001	9.0060	6.001

Example page 3

Calculation

Instantaneous speed
$$= \frac{f(x_0 + h_0) - f(x_0)}{h_0}$$

$$= \frac{(x_0 + h_0)^2 - x_0^2}{h_0}$$

$$= \frac{x_0^2 + 2x_0h_0 + h_0^2 - x_0^2}{h_0}$$

$$= \frac{h_0(2x_0 + h_0)}{h_0}$$

$$= 2x + h$$

Instantaneous rate of change

Instantaneous rate of change

$$f'(x_0) = \lim_{h \to 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

Instantaneous rate of change

$$f'(x_0) = \lim_{h \to 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

$$\blacksquare f'(x)$$
 or

Instantaneous rate of change

$$f'(x_0) = \lim_{h \to 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

- f'(x) or
- $\blacksquare \frac{\mathrm{d}y}{\mathrm{d}x}(x)$

Instantaneous rate of change

$$f'(x_0) = \lim_{h \to 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

$$f'(x)$$
 or

$$\blacksquare \frac{\mathrm{d}y}{\mathrm{d}x}(x)$$

Instantaneous rate of change

The instantaneous rate of change


$$f'(x_0) = \lim_{h \to 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

- f'(x) or

For any positive integer k, the derivative of $f(x) = x^k$ at x_0 is $f'(x) = kx^{k-1}$

Example 2 page 1

 $TPP = 100 + 32Q + 10Q^2 - Q^3$



Example 2 page 2

Differentiating TPP

$$TPP = 100 + 32Q + 10Q2 - Q^3$$

 $TPP' = 32 + 20Q - 3Q^2$

Gradient at maximum is zero, therefore

$$3Q^2 - 20Q - 32 = 0$$

 $(3Q + 4)(Q - 8) = 0$

So
$$Q=8$$
, or $Q=-1.33$

Quadratic Solution

For a quadratic equation of the form

$$ax^2 + bx + c$$

The solution or the roots can be found with

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$