

Tauchen Method*

November 2, 2020

We are going to discretize the continuous AR(1) process,

$$y_{t+1} = \rho y_t + \epsilon_{t+1}$$

where $\epsilon \sim \mathcal{N}(0, \sigma_\epsilon^2)$.

The simplest way to discretize this process and to put into a discrete Markov chain form is using Tauchen (1986) method.

Let \tilde{y} be the discrete-valued process that approximates y and let $\{y_1, y_2, \dots, y_N\}$ be the finite set of possible realizations of \tilde{y} .

Choice of Points Tauchen suggests to select a maximum value y_N as a multiple m (e.g. $m = 3$) of the unconditional standard deviation, i.e.

$$y_N = m \left(\frac{\sigma_\epsilon^2}{1 - \rho^2} \right)^{\frac{1}{2}}$$

and let $y_1 = -y_N$ (assuming G is symmetric), and $\{y_2, y_3, \dots, y_{N-1}\}$ be located in an evenly-spaced manner over the interval $[y_1, y_N]$. Denote with d the distance between successive points in the state space.

Transition Probabilities Let

$$\begin{aligned} \pi_{jk} &= \Pr \{ \tilde{y}_t = y_k | \tilde{y}_{t-1} = y_j \} = \Pr \left\{ y_k - \frac{d}{2} < \rho y_j + \epsilon < y_k + \frac{d}{2} \right\} \\ &= \Pr \left\{ y_k - \frac{d}{2} - \rho y_j < \epsilon < y_k + \frac{d}{2} - \rho y_j \right\} \end{aligned}$$

be the (generic) transition probability. Then, if $1 < k < N - 1$, for each j choose

$$\pi_{jk} = F \left(\frac{y_k + d/2 - \rho y_j}{\sigma_\epsilon} \right) - F \left(\frac{y_k - d/2 - \rho y_j}{\sigma_\epsilon} \right)$$

where F is the cumulative density function of the standard normal distribution. And for boundaries of the interval $k = 1$ and $k = N$,

$$\begin{aligned} \pi_{j1} &= F \left(\frac{y_1 + d/2 - \rho y_j}{\sigma_\epsilon} \right) \\ \pi_{jN} &= 1 - F \left(\frac{y_N - d/2 - \rho y_j}{\sigma_\epsilon} \right) \end{aligned}$$

*Note prepared for UBC ECON407, fall 2020