

EPI809

Testing in Multiple Regression Models

- Long & Short regressions, and various ways of formulating H_0 & H_a
- Type-I error
- Single-test p-values & type-I error rate
- Multiple testing
- Sequential testing:
 - F-test, if significant,
 - Examine t-test for individual predictors

Various ways of describing H0 and Ha

$$\left\{ \begin{array}{l} \text{Model: } y_i = \mu + x_{1i}\beta_1 + x_{2i}\beta_2 + x_{3i}\beta_3 + x_{4i}\beta_4 + \varepsilon_i \\ H_0: \beta_1 = \beta_2 = \beta_3 = \beta_4 = 0 \\ H_a: \text{At least one of the } \beta's \neq 0 \end{array} \right.$$

$$\left\{ \begin{array}{l} H_0: y_i = \mu + \varepsilon_i \\ H_a: y_i = \mu + x_{1i}\beta_1 + x_{2i}\beta_2 + x_{3i}\beta_3 + x_{4i}\beta_4 + \varepsilon_i \end{array} \right.$$

$$\left\{ \begin{array}{l} H_0: \beta_1 = \beta_2 = \beta_3 = \beta_4 = 0 \\ H_a: y_i = \mu + x_{1i}\beta_1 + x_{2i}\beta_2 + x_{3i}\beta_3 + x_{4i}\beta_4 + \varepsilon_i \end{array} \right.$$

F-test Vs t-test

- The preceding example is a 3 DF test.
- For 1-DF tests, the F-test and the t-test are equivalent:
 - The p-values are identical
 - the $t\text{-stat} = \sqrt{F\text{-stat}}$
(verify this with an example)
- However, for test involving more than 1 DF the two tests behave different.
- How do we proceed?
 - For more than 1DF tests we use the F-test
 - If the F-test leads to rejection, then, at least one of the individual effects is non-zero, and only then we evaluate individual effects using the t-test.
- Why don't we just focus on the t-test?
 - The problem is that when we carry multiple test, rejection based on individual p-values do not control type-I error rate accurately. This problem emerges from multiple-testing/

Hypothesis Testing & Type-I Error

State of nature	Do not reject H_0	Reject H_0
H_0 holds	✓	Type-I error
H_a holds	Type-II error	✓

- We use p-values to control type-I error rate (i.e., the probability of rejecting H_0 when H_0 holds).
- P-values are estimates of type-I error rate.
- We reject when p-values are small to keep the probability of type-I error low.
- (If the p-value is correct) and we conduct 1 test $p(\text{type-I error}) = \text{pvalue}$
- What if we do two tests? What is the probability of making at least one mistake?

Hypothesis Testing & Type-I Error

$$\begin{cases} H_0: y_i = \mu + \varepsilon_i \\ H_a: y_i = \mu + x_{1i}\beta_1 + x_{2i}\beta_2 + \varepsilon_i \end{cases}$$

$$pvalue_1 = P(\text{rejecting } H_{01}: \beta_1=0 \mid \beta_1=0, \mu \text{ and } \beta_2 \text{ free})$$

$$pvalue_2 = P(\text{rejecting } H_{02}: \beta_2=0 \mid \beta_2=0, \mu \text{ and } \beta_1 \text{ free})$$


What is the probability of making at least one mistake (FWER=Family-wise error rate)?

$$P[(\text{reject } H_{01} \mid H_{01}, H_{02}) \text{ or } (\text{reject } H_{02} \mid H_{01}, H_{02}) \text{ or } (\text{reject } H_{01} \& H_{02} \mid H_{01}, H_{02})]$$

$$p(A \text{ or } B) = p(A) + p(B) - p(A \& B)$$

Thus, if we reject t-tests at an α -level (significance, say 0.05), and we conduct multiple tests, the probability of making at least 1 type-I error can be larger than α (up to α times the number of tests)

To address this problem, if our test involves more than 1 DF, we use the F-test



Post-hoc analysis. What if the F-test is rejected?

- If the F-test is rejected, we can then proceed and evaluate individual t-tests, but we still need to account for multiple testing.
- For factors, when comparing the effects of various levels, two common approaches are Duncan's test and Tukey's Honest Significant Difference (HSD)
- Both tests are implemented in the package Agricolae (See examples)