

Essentials of Applied Data Analysis

IPSA-USP Summer School 2017

Leonardo Sangali Barone
leonardo.barone@usp.br

jan/17

Axioms and theorems of probability - summary

(Also know as “rules of probability”)

1. For every event A , $0 \leq P(A) \leq 1$
2. $P(S) = 1$
3. $P(\emptyset) = 0$
4. If A and B are mutually exclusive, then:

$$P(A \cup B) = P(A) + P(B)$$

5. If A_1, A_2, \dots is a sequence of mutually exclusive events, then:

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

6. $P(A^c) = 1 - P(A)$
7. $P(A \cup A^c) = P(A) + P(A^c) = 1$ (because they are mutually exclusive)
8. $P(A \cap B)$ is the the probability of A **AND** B happening at the same time.
9. $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
10. If A and B are independent events, then $P(A \cap B) = P(A) * P(B)$

11. If A and B are not independent events, then the conditional probability of A given B, $P(A|B)$, is defined as

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

12. If A and B are independent events, then $P(A|B) = P(A)$ and the formula

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

can be rewritten as (as previously seen)

$$P(A \cap B) = P(A) * P(B)$$

13. If A and B are mutually exclusive, then

$$P(A \cap B) = 0$$

hence

$$P(A|B) = P(B|A) = 0$$