

# Comparing Groups (Chapter 20)

Patrick Frenett and Nicholas Horton ([nhorton@amherst.edu](mailto:nhorton@amherst.edu))

June 16, 2016

## Introduction and background

This document is intended to help describe how to undertake analyses introduced as examples in the Fourth Edition of *Intro Stats* (2013) by De Veaux, Velleman, and Bock. More information about the book can be found at [http://wps.aw.com/aw\\_deveaux\\_stats\\_series](http://wps.aw.com/aw_deveaux_stats_series). This file as well as the associated R Markdown reproducible analysis source file used to create it can be found at <http://www.amherst.edu/~nhorton/sdm4>.

This work leverages initiatives undertaken by Project MOSAIC (<http://www.mosaic-web.org>), an NSF-funded effort to improve the teaching of statistics, calculus, science and computing in the undergraduate curriculum. In particular, we utilize the `mosaic` package, which was written to simplify the use of R for introductory statistics courses. A short summary of the R needed to teach introductory statistics can be found in the `mosaic` package vignettes (<http://cran.r-project.org/web/packages/mosaic>).

Note that some of the figures in this document may differ slightly from those in the IS4 book due to small differences in datasets. However in all cases the analysis and techniques in R are accurate.

## Chapter 20: Comparing Groups

### Section 20.1: The standard deviation of a difference

We can replicate the calculations in the example on the bottom of page 543.

```
n1 <- 248; p1 <- 0.57
n2 <- 256; p2 <- 0.70
sediff <- sqrt(p1*(1-p1)/n1 + p2*(1-p2)/n2); sediff
```

```
## [1] 0.04252786
```

### Section 20.3: Confidence interval for a difference

We can replicate the values from the example on page 546.

```
(p2 - p1) + c(-1.96, 1.96)*sediff
```

```
## [1] 0.04664539 0.21335461
```

### Section 20.4: Testing for a difference in proportions

We can replicate the values from the example on pages 550-551.

```
n1 <- 293; y1 <- 205
n2 <- 469; y2 <- 235
ppooled <- (y1+y2)/(n1+n2); ppooled
```

```
## [1] 0.5774278
```

```
sepoled <- sqrt(ppooled*(1-ppooled)/n1 + ppooled*(1-ppooled)/n2); sepoled
```

```
## [1] 0.0367838
```

```
z <- (y1/n1 - y2/n2)/sepoled; z
```

```
## [1] 5.398915
```

```
pval <- 2*pnorm(z, lower.tail = FALSE); pval
```

```
## [1] 6.704501e-08
```

## Section 20.6: Testing for a difference in means

```
n1 <- 8; n2 <- 7
ybar1 <- 281.88; ybar2 <- 211.43
s1 <- 18.31; s2 <- 46.43
sediff <- sqrt(s1^2/n1 + s2^2/n2); sediff
```

```
## [1] 18.70483
```

```
t <- (ybar1 - ybar2)/sediff; t
```

```
## [1] 3.766407
```

```
pval <- 2*pt(t, df=7.62); pval
```

```
## [1] 1.993996
```

```
prices <- read.csv("http://www.amherst.edu/~nhorton/sdm4/data/Camera_prices.csv")
prices
```

```
##   Buying.from.a.Friend Buying.from.a.Stranger
## 1                   275                   260
## 2                   300                   250
## 3                   260                   175
## 4                   300                   130
## 5                   255                   200
## 6                   275                   225
## 7                   290                   240
## 8                   300                    NA
```

```
with(prices, t.test(Buying.from.a.Friend, Buying.from.a.Stranger))
```

```
##
## Welch Two Sample t-test
##
## data: c(275L, 300L, 260L, 300L, 255L, 275L, 290L, 300L) and c(260L, 250L, 175L, 130L, 200L, 225L, 240L, NA)
## t = 3.766, df = 7.6229, p-value = 0.006003
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## 26.93688 113.95597
## sample estimates:
## mean of x mean of y
## 281.8750 211.4286
```

Let's turn this dataset in a lattice friendlier version.

```
ds <- with(prices,
  data.frame(price=c(Buying.from.a.Friend, Buying.from.a.Stranger),
    group=c(rep("Friend", nrow(prices)), rep("Stranger", nrow(prices)))))
ds
```

```
##   price  group
## 1   275  Friend
## 2   300  Friend
## 3   260  Friend
## 4   300  Friend
## 5   255  Friend
## 6   275  Friend
## 7   290  Friend
## 8   300  Friend
## 9   260 Stranger
## 10  250 Stranger
## 11  175 Stranger
## 12  130 Stranger
## 13  200 Stranger
## 14  225 Stranger
## 15  240 Stranger
## 16   NA Stranger
```

```
t.test(price ~ group, data=ds) # Unpooled
```

```
##
## Welch Two Sample t-test
##
## data: price by group
## t = 3.766, df = 7.6229, p-value = 0.006003
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## 26.93688 113.95597
## sample estimates:
## mean in group Friend mean in group Stranger
## 281.8750 211.4286
```

```
t.test(price ~ group, var.equal=TRUE, data=ds) # Pooled
```

```
##  
## Two Sample t-test  
##  
## data: price by group  
## t = 3.9699, df = 13, p-value = 0.0016  
## alternative hypothesis: true difference in means is not equal to 0  
## 95 percent confidence interval:  
## 32.11047 108.78238  
## sample estimates:  
## mean in group Friend mean in group Stranger  
## 281.8750 211.4286
```

```
bwplot(group ~ price, data=ds)
```

