

Problem Set 2

Prof. Conlon

Due: 3/1/19

Packages to Install

The packages used this week are

- estimatr (Tidyverse version of lm function)

Problem 1 (Analytical Exercise)

Consider the estimation of the individual effects model:

$$y_{it} = x'_{it}\beta + \alpha_i + \epsilon_{it}, \mathbb{E}[\epsilon_{it} | x_{it}, \alpha_i] = 0$$

where $i = \{1, \dots, n\}$ and $t = \{1, \dots, T\}$.

This exercises ask you to relate the (random effects) GLS estimator $\hat{\beta}_{GLS} = (X'_*X_*)^{-1}X'_*y_*$ to the “within” (fixed-effects) estimator $\hat{\beta}_{FE} = (\dot{X}'\dot{X})^{-1}\dot{X}'\dot{y}$ and the “between” estimator $\hat{\beta}_{BW} = (\bar{X}'\bar{X})^{-1}\bar{X}'\bar{y}$ where $w = \{x, y\}$:

$$\begin{aligned}\bar{w}_i &:= \frac{1}{T} \sum_{i=1}^T w_i \\ \dot{w}_i &:= w_{it} - \bar{w}_i \\ w_{it,*} &:= w_{it} - (1 - \lambda)\bar{w}_i \\ \lambda^2 &= \frac{Var(\epsilon)}{T Var(\alpha_i) + Var(\epsilon_{it})}\end{aligned}$$

1. Express the GLS estimator in terms of \bar{X} , \dot{X} , \bar{y} , \dot{y} , λ , and T .
2. Show that there is a matrix R depending on \bar{X} , \dot{X} , λ and T such that the GLS estimator is a weighted average of the "within" and "between" estimators:

$$\hat{\beta}_{GLS} = R\hat{\beta}_{FE} + (I - R)\hat{\beta}_W$$

3. What happens to the relative weights on the "within" and "between" estimators as we increase the sample size, i.e. $T \rightarrow \infty$?
4. Suppose that the random effects assumption $\mathbb{E}[\alpha_i | x_{i1}, \dots, x_{iT}] = 0$ does not hold. Characterize the bias of the estimators $\hat{\beta}_{FE}$, $\hat{\beta}_W$. (Note: An estimator $\hat{\beta}$ is unbiased if $\mathbb{E}[\hat{\beta}] = \beta$)
5. Use your result from (d) to give a formula for the bias of our random effects estimator $\hat{\beta}_{GLS}$. What happens to the bias as $T \rightarrow \infty$.

Problem 2 (Coding Exercise)

We observe N observations of the random variable X_i where each X_i is drawn from the Weibull distribution:

$$X_i \sim W(\gamma)$$

The probability density function for the Weibull is the following:

$$f(x; \gamma) = \gamma x^{\gamma-1} \exp(-(x^\gamma)) \quad ; x \geq 0, \gamma > 0$$

1. Assume our N observations are independent and identically distributed, what is the log-likelihood function?
2. Calculate the gradient (or first derivative) of your log-likelihood function.
3. Using the first order condition, what is the MLE estimator for γ ?
4. Verify that the second order condition guarantees a unique global solution.
5. In R, I want you to write a function called `mle_weibull` that takes two arguments (X, γ) , where X is a vector of data and γ is a scalar. The function returns the value of the log-likelihood function you derived in the last part.
6. Optimization routines can either be given a first derivative (or gradient) or the optimization routines calculate numerical derivatives. We will be using the R function *optim*, which accepts the first derivative as an argument *gr*.
 - a. We first want you to run *optim* without supplying a first derivative (leaving *gr* out of the function). Note, to run *optim* you will need to supply your data X as an additional parameter at the end of the function. We have provided you with simulated data in the file `'prob_4_simulation.rda'` located in the data folder.
 - b. We now want you to create a new function called *gradient*, which takes the same two arguments as your likelihood function. Now calculate the MLE using *optim* with the gradient.
 - c. Compare both the number of iterations until convergence and your estimated γ values from both runs.