

# Introduction to Bayesian Statistics with R

## Day 3

Oliver Lindemann

Erasmus University Rotterdam, NL

### Bayes rule


$$\underbrace{p(\theta|D)}_{\text{Posterior}} = \underbrace{p(D|\theta)}_{\text{Likelihood}} \underbrace{p(\theta)}_{\text{Prior}} / \underbrace{p(D)}_{\text{Evidence}}$$

Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 2/44

### Discrete Likelihood $p(D|\theta)$

Example from previous session: Discrete values



Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 3/44

### How to determine $p(D|\theta)$ ?

How to determine the posterior distribution of continuous parameters?

1. Analytically
2. Approximation (Grid approximation)
3. Simulation (Markov Chain Monte Carlo)

Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 4/44

### Likelihood $p(D|\theta)$

$$\underbrace{p(\theta|D)}_{\text{Posterior}} = \underbrace{p(D|\theta)}_{\text{Likelihood}} \underbrace{p(\theta)}_{\text{Prior}} / \underbrace{p(D)}_{\text{Evidence}}$$

Coin flip example:  $D = \{y_1, \dots, y_N\}$

- $N$ : number of flips
- $z$ : Number of 'heads',  $z = \sum_{i=1}^N y_i$

Bernoulli Distribution

$$p(y_i|\theta) = \theta^{y_i} (1 - \theta)^{(1-y_i)}$$

Multiple coin flips

$$p(D|\theta) = p(\{y_1, \dots, y_N\}|\theta) = \prod_i \theta^{y_i} (1 - \theta)^{(1-y_i)}$$

$$p(D|\theta) = p(z, N|\theta) = \theta^z (1 - \theta)^{N-z}$$




Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 5/44

### Continuous likelihood function: $p(D|\theta)$

$$\underbrace{p(\theta|D)}_{\text{Posterior}} = \underbrace{p(D|\theta)}_{\text{Likelihood}} \underbrace{p(\theta)}_{\text{Prior}} / \underbrace{p(D)}_{\text{Evidence}}$$

$$p(D|\theta) = p(z = 3, N = 12|\theta) = \theta^z (1 - \theta)^{N-z} = \theta^3 (1 - \theta)^9$$



Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 6/44

## Continuous likelihood function: $p(D|\theta)$

$$p(\theta|D) = \frac{p(D|\theta) p(\theta)}{p(D)}$$

Posterior Likelihood Prior Evidence

```
Theta = seq(0,1, by=0.01)
z = 3
N = 12
likelihood = Theta^z * (1-Theta)^(N-z) # Bernoulli likelihood

plot( Theta, likelihood, type="l", lwd=3,
      main="Likelihood",
      xlim=c(0,1),
      xlab=bquote(theta),
      ylim=c(0,1.1*max(likelihood)),
      ylab=bquote(p("D|",theta,"")),
      col="skyblue" )
```

## Continuous description of beliefs: Prior $p(\theta)$

$$p(\theta|D) = \frac{p(D|\theta) p(\theta)}{p(D)}$$

Posterior Likelihood Prior Evidence

### Problem

We need a mathematical formula that describes the prior beliefs.

Formula of  $p(\theta)$  should have certain characteristics

1. Bayes' nominator:  
 $p(D|\theta) \cdot p(\theta)$  should have the same form as  $p(\theta)$
2. Bayes' denominator:  
 $p(D) = \int d\theta p(D|\theta)p(\theta)$  should be solvable analytically

## Continuous description of beliefs: Prior $p(\theta)$

Think about it:

- Form of likelihood,  $p(D|\theta)$
- If the prior,  $p(\theta)$ , has the form
- Likelihood  $\times$  prior looks then like this

$$\theta^z (1-\theta)^{N-z}$$

$$\theta^a (1-\theta)^b$$

$$\theta^{z+a} (1-\theta)^{N-z+b}$$

## Continuous prior: Beta Distribution

Probability function of the form is called


### Beta Distribution

$$p(\theta|a, b) = \text{beta}(\theta|a, b) = \theta^{a-1} (1-\theta)^{b-1} / B(a, b) \quad (3.1)$$


$B(a, b)$  is a normalizing constants  $B(a, b) = \int d\theta \theta^{a-1} (1-\theta)^{b-1}$

In R


```
# beta distribution
dbeta(theta, a, b)
# normalizing constant
beta(a, b)
```



## Updating Continuous Prior & Highest Density Interval (HDI)




## Impact of Priors



Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 13/44


## Impact of Priors



Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 14/44


## Impact of Priors: Arbitrarily shaped Distributions



Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 15/44

## Examples of 95% highest density intervals (HDIs)



Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 16/44

## Calculating Posterior

Data  $D = z, N$

Likelihood  $p(\theta|z, N) = \theta^z (1 - \theta)^{(N-z)}$

Prior  $p(\theta) = \text{Beta}(\theta|a, b) = \theta^{(a-1)} (1 - \theta)^{(b-1)} / B(a, b)$

Evidence  $p(D) = p(z, N)$

Posterior (Bayes rule)

$$p(\theta|z, N) = \frac{p(z, N|\theta) p(\theta)}{p(z, N)}$$

Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 17/44

## Calculating Posterior

$$p(\theta|z, N) = \frac{p(z, N|\theta) p(\theta)}{p(z, N)}$$

$$p(\theta|z, N) \propto p(z, N|\theta) p(\theta)$$

$$\propto \theta^z (1 - \theta)^{(N-z)} \theta^{(a-1)} (1 - \theta)^{(b-1)}$$

$$\propto \theta^{(z+a-1)} (1 - \theta)^{(N-z+b-1)}$$

$$p(\theta|z, N) = \frac{\theta^{(z+a-1)} (1 - \theta)^{(N-z+b-1)}}{B(z+a, N-z+b)}$$

$$p(\theta|a, b) = \theta^{(a-1)} (1 - \theta)^{(b-1)} / B(a, b)$$

Finding appropriate parameter  $(a, b)$  for my Beta prior distribution?

Analytically Impact of Priors Grid Approximation MCMC simulation JAGS MCMC diagnostics

bayes18-3 18/44

## Approximating Posterior Distributions

## Bayes theorem for discrete & continuous values


### Bayes Rule (continuous values)

$$p(\theta|D) = \frac{p(D|\theta) \cdot p(\theta)}{\int d\theta p(D|\theta) p(\theta)} \quad (2.6)$$


### Bayes Rule (discrete values)

$$p(\theta|D) = \frac{p(D|\theta) \cdot p(\theta)}{\sum_{\theta} p(D|\theta) p(\theta)} \quad (2.5)$$


## Grid Approximation



## Discretizing



## Advantage: Modelling Arbitrarily shaped Distributions




## Problem with grid approximation

- not very handy method
- not appropriate for models with large parameter space
  - ▶ e.g. 6 parameter
  - ▶ each modeled 1000 values (grid density)
  - ▶  $1,000^6 = 1,000,000,000,000,000,000$

# Markov Chain Monte Carlo (MCMC) simulation

## Markov Chain Monte Carlo simulation

### Metaphor



- Politician lives on a long chain of islands
- Goal: Visit each island in proportion to its population
- Problem:
  - ▶ Unknown number of islands?
  - ▶ Unknown total population?
- But: Major of the current and the next island can be asked


## MCMC: Metropolis Algorithm

### Random walk:

1. choose randomly step to the left or right (*proposal distribution*)
2. if  $p(\text{proposed}) > p(\text{current})$ 
  - ▶ move to proposed
3. else
  - ▶ calculate ratio  $p_{\text{move}} = \frac{p(\text{proposed})}{p(\text{current})}$
  - ▶ move in with likelihood  $p_{\text{move}}$  to next place

## MCMC: Metropolis Algorithm

(Demonstration)



## MCMC: Metropolis Algorithm

### Random walk:

1. choose randomly step to the left or right (*proposal distribution*)
2. if  $p(\text{proposed}) > p(\text{current})$ 
  - ▶ move to proposed
3. else
  - ▶ calculate ratio  $p_{\text{move}} = \frac{p(\text{proposed})}{p(\text{current})}$
  - ▶ move in with likelihood  $p_{\text{move}}$  to next place

### Target Distribution

(step 2 & 3)

$$p_{\text{move}} = \min\left(\frac{p(\text{proposed})}{p(\text{current})}, 1\right)$$

## MCMC: Metropolis Algorithm

### Markov Chain Monte Carlo (MCMC)

We must be able to

1. create a random value from the **proposal distribution**
2. compute ratio to evaluate the **target distribution** at any position:  $p_{\text{move}} = p(\text{proposed})/p(\text{current})$
3. generate random value from uniform distribution (to move or not move according  $p_{\text{move}}$ )

Note: Target distribution does not have to be normalized, because merely need to know the ratio between two positions (e.g. slope).

## MCMC: Continuous data

Metropolis algorithm is applied to

- continuous data
- any number of dimensions
- more general proposal distributions

But, the essence of the procedure is always the same as in our discrete one-dimensional example of the islands.

We just need to be able to compute values from not normalized target distribution  $P(\theta)$ .

## MCMC: Applied to Bayes Theorem

$$p(\theta|D) = \frac{p(D|\theta) \cdot p(\theta)}{\int d\theta p(D|\theta) p(\theta)}$$

Sample from not normalized *target distribution*

$$p(\theta|D) \propto p(D|\theta) \cdot p(\theta)$$

## Random walk through continuous parameter space

Algorithm

1. start at arbitrary point
2. (randomly) propose a movement
  - ▶ proposal distribution can have any form
  - ▶ should be center round zero
  - ▶ typically normal distribution
3. Decide accept proposal or not
  - ▶ calculate  $p(\theta_{\text{move}})$
  - ▶ draw random number between  $[0, 1]$

**Good news!**

- This can be automatised
- The software packages like BUGS or JAGS will do the sampling for us.

**JAGS**

## JAGS: "Just Another Gibbs Sampler"

### JAGS

- using the model description language BUGS
- <http://mcmc-jags.sourceforge.net/>

### BUGS language (WinBUGS)

- probabilistic modelling language
- Markov chain Monte Carlo (MCMC) methods.
- Declarative Programming language (!)
  - ▶ In contrast to imperative language, a declarative language declares *what* needs to be done rather than *how* to do it.

Similar software: Stan, NIMBLE

## MCMC sampling in R and JAGS

Data

- $y_i = \{y_1, \dots, y_N\}$ : coin flips
- $N$ : number of flips


Model

- likelihood: flips are Bernoulli distributed  
 $p(y_i|\theta) = \theta^{y_i} (1 - \theta)^{(1-y_i)}$
- prior  $p(\theta)$ : uniform  
 $\text{unif}(0, 1) = \text{beta}(1, 1)$

```
# The JAGS model
model
{
  # Likelihood
  for (i in 1:N)
  {
    y[i] ~ dbern( theta )
  }

  # Prior of theta
  theta ~ dunif(0, 1)
}
```






Distribution	Density	Distribution	Quantile
Bernoulli	dbern	pbern	qbern
Beta	dbeta	pbeta	qbeta
Binomial	dbin	pbm	qbin
Chi-square	dchisqr	pchisqr	qchisqr
Double exponential	dexp	pexp	qexp
Exponential	dexp	pexp	qexp
F	df	pf	qf
Gamma	dgamma	pgamma	qgamma
Generalized gamma	dgen.gamma	pgen.gamma	qgen.gamma
Noncentral hypergeometric	dhyper	phyper	qhyper
Logistic	dlogis	plogis	qlogis
Log-normal	dlnorm	plnorm	qlnorm
Negative binomial	dnegbin	pnegbin	qnegbin
Noncentral Chi-square	dchisqr	pchisqr	qchisqr
Normal	dnorm	pnorm	qnorm
Pareto	dpar	ppar	qpar
Poisson	dpois	ppois	qpois
Student t	dt	pt	qt
Weibull	dweib	pweib	qweib

Table 5.3: Functions to calculate the probability density, probability function, and quantiles of some of the distributions provided by the bugs module.

MCMC Sample Diagnostic

```
# The JAGS model
model
{
  # Likelihood
  for (i in 1:N)
  {
    y[i] ~ dbern( theta )
  }

  # Prior of theta
  theta ~ dbeta(a, b)
  a = 1
  b = 1
}
```



- Samples should be
1. **representative** of the posterior distribution
  2. sufficient size so that estimates are **accurate** and **stable**
    - burn-in
    - thinning
  3. sample process should be **efficient**




Figure 7.10: Illustration of MCMC diagnostics. Three chains were generated by starting a Metropolis algorithm at different initial values, with proposal SD=0.02

Trace plots (later samples after burn in)




Figure 7.11: Illustration of MCMC diagnostics. Three chains were generated by

Two Coinflips

```
# two independent coin flips
model
{
  # Likelihood
  for (i in 1:N)
  {
    ya[i] ~ dbern( theta_a )
    yb[i] ~ dbern( theta_b )
  }

  theta_a ~ dbeta(a, b)
  theta_b ~ dbeta(a, b)
  a <- 1
  b <- 1
}
```