

The entire set of m tests is called a "family" is "any logical collection of inferences for which it is meaningful to take into account some combined measure of error" or a set of tests where you wish to prevent "data dredging" (e.g. the spurious correlations in 342) or to "ensure a correct 'overall' decision in the collection of tests".

We'll discuss two error properties/metrics for a family of tests.

The first is called "familywise error rate" (FWER) defined as:

$$\text{FWER} := P(V > 0) \leq \text{FWER}_0 \leftarrow \text{this is the level of control that I choose e.g. 5\%}$$

If you can show that $\text{FWER} \leq \text{FWER}_0$ for any $m_0 \leq m$ subset of the m tests, this is called "strong FWER control". We won't study it. If you can show that $\text{FWER} \leq \text{FWER}_0$ for $m_0 = m$ then this is called "weak FWER control" which we will study. If $m_0 = m$...

		Decision		
		Retain H_0	Reject H_0	
Truth	H_0	U	V	$m_0 \Rightarrow V = R \Rightarrow \text{FWER} = P(R > 0)$
	H_a	O	O	O
		F	R	m

Our goal is weak FWER control under the most general settings.

$R_1 = 1$ if H_{01} is rejected, $R_1 = 0$ if H_{01} is retained.

$R_2 = 1$ " H_{02} " " $R_2 = 0$ " H_{02} " "

$R_m = 1$ " H_{0m} " " $R_m = 0$ " H_{0m} " "

$$\text{FWER} = P(R > 0) = P(R_1 = 1 \cup R_2 = 1 \cup \dots \cup R_m = 1) \\ \leq \sum_{i=1}^m P(R_i = 1) = m\alpha.$$

recall from Math 241, $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
the principle of inclusion - exclusion.

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = \left(\sum P(A_i) - \sum P(A_i \cap A_j) + \sum P(A_i \cap A_j \cap A_k) \right. \\ \left. - \dots + \dots \right)$$

and from here you can prove "Boole's Inequality."

$$P(A_1 \cup A_2 \cup \dots \cup A_n) \leq \sum P(A_i)$$

$$\Rightarrow \text{FWER} \leq \text{FWER}_0 \Rightarrow m\alpha = \text{FWER}_0 \Rightarrow \alpha = \frac{\text{FWER}_0}{m}.$$

this is called the Bonferroni correction (1936)

\Rightarrow a pval for an individual test must be less than FWER_0 / m . Equivalently, you can multiple the p-values by m / FWER_0 and compare each to $\alpha = 5\%$.

$$\text{pval} \leq \frac{\text{FWER}_0}{m} = \alpha \Rightarrow \frac{m}{\text{FWER}_0} \text{pval} \leq \alpha.$$

Adjusted P-value

e.g. if $m = 30$, $\text{FWER}_0 = 5\% \Rightarrow \alpha = \text{FWER}_0/m = 0.00167$

The obvious problem with this correction is... it gives you really bad power! Because it is ultra-conservative.

We can do a bit better if we assume the tests are independent. Then, $R_1, R_2, \dots, R_m \stackrel{\text{iid}}{\sim} \text{Bern}(\alpha) \Rightarrow R \sim \text{Bin}(m, \alpha)$

$$\text{FWER} = P(R > 0) = 1 - P(R = 0) = 1 - (1 - \alpha)^m \leq \text{FWER}_0$$

$$\Rightarrow 1 - \text{FWER}_0 = (1 - \alpha)^m \Rightarrow 1 - \alpha = (1 - \text{FWER}_0)^{1/m}$$

$$\Rightarrow \alpha = 1 - (1 - \text{FWER}_0)^{1/m} \Rightarrow 1 - \alpha = (1 - \text{FWER}_0)^{1/m}$$

$$\Rightarrow \alpha = 1 - (1 - \text{FWER}_0)^{1/m} \quad \text{Dunn-Sidak Correction (1967)}$$

e.g. if $m = 30$, $\text{FWER}_0 = 5\%$

$$\Rightarrow \alpha = 1 - (95\%)^{1/30} = 0.00171 > 0.00167$$

(the Bonferroni)

Thus, you get slightly higher power with Sidak.

$$1 - (1 - x)^{1/c} \approx x/c \quad (1^{\text{st}} \text{ order Taylor series})$$

There are other methods e.g. the "Holm step-down" procedure (1979) but it is similar to the Simes procedure (1986) which we talk about now.

Bonferroni and Sidak never looked at the p-values and there's a lot of information there. Remember, Fisher created the p-val to gauge the "strength" of a rejection. Rejecting with a p-value of 0.00001 is much stronger than rejecting with a p-value of 0.01. Holm and Simes used this. For the m tests, you get p-values p_1, p_2, \dots, p_m but don't retain / reject anything yet!! Order them from smallest to largest.

$$p(1) \leq p(2) \leq \dots \leq p(m) \quad (\text{order statistics})$$

\uparrow min pval \uparrow max pval "linear step-up"

Then locate the following: $a_* := \max \{a : p_{(a)} \leq \frac{\alpha}{m} \text{FWER}_0\}$

$\nearrow \in \{\text{Bonferroni}, \dots, \text{Holm}\}$

or let $a_* = 0$ if max doesn't exist.

Then set $\alpha = \frac{a_*}{m} \text{FWER}_0$.

You can prove that this gives you weak FWER control. It is rare that this is not more powerful than Bonferroni / Sidak.

By construction you reject all tests up to the a_* th test (if the tests are in order of p-value). Then you retain all the other $m - a_*$ tests.

The problem with FWER in general is maybe it's too conservative. What if you want to trade some false rejections for more power? Let's consider another metric of familywise control (not FWER), called "False Discovery Rate" (FDR). First, define the "false Discovery Proportion" (FDP),

$$\text{FDP} := \begin{cases} V/R & \text{if } R > 0 \\ 0 & \text{if } R = 0 \end{cases}, \text{ the random proportion of rejections that are Type I errors}$$

$\text{FDR} := E[\text{FDP}]$, the expected proportion of rejections that are Type I errors.

Now we wish to control FDR so we want:

$\text{FDR} \leq \text{FDR}_0$, a constant you set. For example if $\text{FDR}_0 = 5\%$ and I run m tests and get 100 rejections, then I expect ≤ 5 of the rejections to be Type I errors and ≥ 95 of the rejections to be real discoveries.

Note: if $m = m_0$ then $\text{FWER} = \text{FDR}$. Proof

$$m = m_0 \Rightarrow V = R \Rightarrow \text{FDP} = \begin{cases} 1 & \text{if } R > 0 \\ 0 & \text{if } R = 0 \end{cases} = \text{Bern}(P(R > 0))$$

$$\Rightarrow \text{FDR} = E[\text{FDP}] = P(R > 0) = \text{FWER}$$

Not on test

Note: the FDR procedure is more powerful than the FWER procedure.

$$\mathbb{1}_{V \geq 1} \geq V/R \quad \text{if } V=0 \Rightarrow 0 \geq 0 \quad \checkmark$$

$$V=1 \Rightarrow 1 \geq 1 \text{ or } 1/2 \text{ or } 1/3 \dots V/R$$

$$V \geq 1 \Rightarrow 1 \geq 1 \text{ or } V/V_{+1} \text{ or } \dots V/R$$

$$\Rightarrow E[\mathbb{1}_{V \geq 1}] \geq E[V/R]$$

$$P(V \geq 1) \geq \text{FDR}$$

||

$$\text{FWER} \geq \text{FDR} \quad \checkmark$$

Benjamini and Hochberg (1995) proved the Simes procedure controls FDR for any m_0 subset of the m tests. In fact $\text{FDR} = m_0/m \text{FDR}_0 \leq \text{FDR}_0$, thus for a small m_0 (which don't observe), the FDR control is much better than FDR_0 .