

Lec 16

11/02/2020


$\hat{\theta}_{1.} = \frac{n_{1.}}{n}$ proportion of people with black hair.

$$\Rightarrow \hat{\theta}_{i.} = \frac{n_{i.}}{n}, \hat{\theta}_{.j} = \frac{n_{.j}}{n}$$

$$\Rightarrow \hat{\phi} = \sum_{i=1}^r \sum_{j=1}^c \frac{(O_{ij} - n \hat{\theta}_{i.} \hat{\theta}_{.j})^2}{n \hat{\theta}_{i.} \hat{\theta}_{.j}} \xrightarrow{d} \chi^2_{(r-1)(c-1)} \quad \text{a fact that}$$

could be proved in 368. if time permitted.

"Chi-squared test of independence"



$$\hat{\phi} = 41.28 \quad \alpha = 5\% \quad P_{\chi^2_9}(16.99) = 95\%$$

$\notin \text{RET}$



$\Rightarrow \text{Reject } H_0$

$$\text{RET} = [0, 16.99]$$

We conclude that hair color and eye color are dependent.

This class has focused mainly on the 3 goals of inference: estimation, testing and confidence sets. We will continue to study these 3 goals but we will also study some tangential "meta" concepts that are classic.

Here's one such classic "meta concept". Usually you are.

- given a dataset x_1, \dots, x_n , then you
- assume a DGP, then you
- define one (or many) inferential target parameters θ , then
- compute $\hat{\theta} = w(x_1, \dots, x_n)$ and then you
- make a CI / run a test at size α .

Let's examine (b). How do you just "assume a DGP"? Sometimes you really do know the DGP e.g. a coin flipped repeatedly is iid Bernoulli (θ), a die rolled repeatedly is iid uniform discrete. But what about "daily wind speeds at JFK airport" or "rat survival times" (like on midterm) or "daily percentage returns of the S&P 500"? The DGP's for the last three are very complicated and they're unknown.

What if we wanted to "guess" the DGP's? This is actually a really big part of what statisticians do. This is called "model fitting". DGP = model. Models you kinda make up & hope fully they're useful for whatever you're doing. Why don't we proceed as follows, let's guess M candidate models. DGPs $m = 1, 2, \dots, M$ and then

(1) Pick the "best" model out of my M guesses and maybe (2) provide a weighting score to each of the M guesses (low scores indicate bad guesses and high scores indicate good guesses). Goal (1) is famous and called "model selection". In 3h2, we do a little of this theoretically. Here we'll do it more theoretically.

Model selection is more fundamental than you realize. It's actually the entire problem of all of science. For example, let's say you have some astronomical data on movement of different planets, stars, etc. You want to fit a model (guess DGP) for the force on two celestial bodies with masses m_1, m_2 at a distance r from each other (i.e. "gravity"). Consider the following models:

Mod 1: $F = G \frac{M_1 M_2}{r^2}$ Newton's law

Mod 2: $F = G_1 \frac{M_1 M_2}{r^2} + G_2 \frac{M_1 M_2}{r^3}$ Newton's extension

Mod 3: $F = G_1 \frac{M_1 M_2}{r^2} e^{-G_2 r}$ Laplace's extension

Which model is the best? We know all these are wrong because Einstein came and disproved them with general relativity.

Let's talk about model selection techniques. Our data x_1, \dots, x_n comes from an unknown DGP. Here are M candidate models.

Mod 1: $f_1(x_1, \dots, x_n; \theta_{11}, \dots, \theta_{1k_1}) = \mathcal{L}_1(\theta_{11}, \dots, \theta_{1k_1}; x_1, \dots, x_n)$

Mod 2: $f_2(x_1, \dots, x_n; \theta_{21}, \dots, \theta_{2k_2}) = \mathcal{L}_2(\theta_{21}, \dots, \theta_{2k_2}; x_1, \dots, x_n)$

!

$$\text{Mod } m : f_m(x_1, \dots, x_n; \theta_{m1}, \dots, \theta_{mk_m}) = f_{k_m}(\theta_{m1}, \dots, \theta_{k_m}; x_1, \dots, x_n)$$

k_1 is the # of parameters in model 1, k_2 is the number of parameters in model 2, ..., k_M is the number of parameters in model M . Each k_m could be different.

Why don't we just select the model that has the highest likelihood?

$$m_k = \underset{m}{\operatorname{argmax}} \left[\ell(\theta_{m1}, \dots, \theta_{mk_m}; x_1, \dots, x_n) \right]$$

$$= \underset{m}{\operatorname{argmax}} \left[\ln(\theta_{m1}, \dots, \theta_{mk_m}; x_1, \dots, x_n) \right]$$

The problem with this is we don't know the values of θ for any of the models!

So let'sRichardize $k_1 + k_2 + \dots + k_M$ times! We'll estimate each of the parameters using MLE's and plug them all in and then...

$$m_k = \underset{m}{\operatorname{argmax}} \left\{ \ell(\hat{\theta}_{m1}^{\text{MLE}}, \dots, \hat{\theta}_{mk_m}^{\text{MLE}}; x_1, \dots, x_n) \right\}$$

You could do this. But... it will not give you the best model. Why?

$\ell(\hat{\theta}_{m1}^{\text{MLE}}, \dots, \hat{\theta}_{mk_m}^{\text{MLE}}; x_1, \dots, x_n)$ is an estimator for

$\ell(\theta_{m1}, \dots, \theta_{mk_m}; x_1, \dots, x_n)$ and it's biased...

with many assumptions, you can prove that.
 asymptotically ... meaning as n gets large ...
 Bias $[l(\hat{\theta}_{m_1}^{MLE}, \dots, \hat{\theta}_{m_{k_m}}^{MLE}; x_1, \dots, x_n)] = k_m > 0$

There is positive bias (meaning that log-likelihood would appear higher on average) and this bias increases you use more parameters in your candidate models. This was figured out by H. Akaike, a Japanese statistician, and he published it in 1974. Once you have the bias, you just use it to correct your estimate!

$$l(\theta_{m_1}, \dots, \theta_{m_{k_m}}) \approx l(\hat{\theta}_{m_1}^{MLE}, \dots, \hat{\theta}_{m_{k_m}}^{MLE}; x_1, \dots, x_n) - k_m$$

Recall that log likelihood is always negative for discrete DGP's and almost always negative for continuous DGP's. So let's flip its sign and multiply by two.

$$AIC_m := -2 l(\hat{\theta}_{m_1}^{MLE}, \dots, \hat{\theta}_{m_{k_m}}^{MLE}; x_1, \dots, x_n) + 2k_m$$

complexing penalty

(Akaike's Information Criterion)

The "best" log likelihood is the largest i.e. closest to zero so once negated, the "best" negative log likelihood is the smallest i.e. closest to zero.

$$m_k := \operatorname{argmin} [AIC_n]$$