


HW 2: Linear regression and prediction using MLB players

Stats and sports class

3. Make a correlation matrix - both a matrix of the variables, as well as a visualization – using hits, doubles, triples, home runs, RBI, and strikeouts.

Answer

```
library(corrplot)
library(tidyverse)
library(Lahman)
Batting_1 <- Batting %>%
  filter(yearID >= 2000) %>%
  select(playerID, yearID, AB:SO) %>%
  filter(AB >= 500)
var_cor <- Batting_1 %>%
  select(H, X2B, X3B, HR, RBI, SO)
cor_mat <- cor(var_cor)
corrplot(cor_mat, method = "number")
```




```
## Note -- any correlation plot would be acceptable
```

4. Make a scatter plot of runs batted in (RBI, the y-variable) and home runs (HR, the x-variable). Estimate and write the regression line using the `lm` command. Finally, interpret the slope and intercept of this line.

Answer

```
ggplot(data = Batting_1, aes(x = HR, y = RBI)) +
  geom_point()
```



```
fit_1 <- lm(RBI ~ HR, data = Batting_1)
summary(fit_1)
```

```
##
## Call:
## lm(formula = RBI ~ HR, data = Batting_1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -37.663  -8.697  -0.732   8.320  52.472
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  42.76593   0.60043   71.23  <2e-16 ***
## HR           1.79657   0.02516   71.41  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 12.59 on 2089 degrees of freedom
## Multiple R-squared:  0.7094, Adjusted R-squared:  0.7092
## F-statistic: 5099 on 1 and 2089 DF, p-value: < 2.2e-16
```

Slope of 1.79: For each additional home run, a batter is expected to have an additional 1.79 RBIs

Intercept of 42.8: A batter that hits 0 HRs would be expected to have 42.8 RBIs.

Estimated line: $\text{RBI-hat} = 42.8 + 1.79 \cdot \text{HR}$

5. Pete Alonso – currently with the New York Mets – hit 53 home runs and batted in 120 runs in 2019. Given his home runs, what is his residual? That is, how many more or fewer runs batted in has he hit than we'd expect given his home runs?

Expected RBIs: $42.8 + 1.79 \cdot 53 = 137.7$ We expect Alonso to have roughly 137.7 runs batted in. He actually has 120, which means he has fewer RBIs than we expect


Part II: Predictability of player metrics

Note: The code drops the last year of a players' career – there is no future variable to look at.


10. Use (i) scatter plots and (ii) correlation coefficients to assess the year-over-year repeatability of strikeout rate, walk rate (BB_rate), HR rate, and RBI rate. That is, compare each metric in a players' current year to the metric that he records in the following year. Which of these metrics is most repeatable? Which of these is least repeatable?

#i Scatter plots


```
ggplot(Batting_2, aes(K_rate, K_rate_next)) +  
  geom_point()
```




```
ggplot(Batting_2, aes(BB_rate, BB_rate_next)) +  
  geom_point()
```



```
ggplot(Batting_2, aes(HR_rate, HR_rate_next)) +  
  geom_point()
```



```
ggplot(Batting_2, aes(RBI_rate, RBI_rate_next)) +  
  geom_point()
```



```
# ii correlation coefficient  
Batting_2 %>%  
  summarise(cor_k = cor(K_rate, K_rate_next),  
            cor_bb = cor(BB_rate, BB_rate_next),  
            cor_hr = cor(HR_rate, HR_rate_next),  
            cor_rbi = cor(RBI_rate, RBI_rate_next))
```

```
## # A tibble: 1 x 4  
##   cor_k cor_bb cor_hr cor_rbi  
##   <dbl> <dbl> <dbl> <dbl>  
## 1 0.854 0.784 0.724 0.670
```

By both looking at the scatter plots and the correlation coefficients, strikeouts are the most repeatable from year to year, and rbi rate is the least repeatable.