

Bayesian estimation of bivariate regression models via MCMC

Applied Bayesian Statistics
Winter Term 2018

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Susumu Shikano
GSDS

Linear Regression Model

Two approaches to obtain posterior

- Conjugacy analysis
 - Conjugacy: The property that the prior and posterior have the same probability form depending on the form of the distribution used to calculate the likelihood.
 - Posterior can be obtained analytically.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Linear Regression Model

Two approaches to obtain posterior

- Conjugacy analysis
 - Conjugacy: The property that the prior and posterior have the same probability form depending on the form of the distribution used to calculate the likelihood.
 - Posterior can be obtained analytically.
- Deriving posterior per Gibbs Sampling
 - Conjugacy is not must.
 - Use of MCMC.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Today's session

- An alternative approach to obtain posterior
 - Derive conditional posterior
 - Run Gibb sampling
- Gibbs sampling and its alternatives
 - Gibbs sampling
 - Metropolis-Hasting algorithm
 - Slice Sampling

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for the slope parameter β_1

- Prior:

$$\begin{aligned}f(\beta_1) = f_N(\mu_1^0, V_1^0) &= \frac{1}{\sqrt{2\pi V_1^0}} \exp \left\{ -\frac{(\beta_1 - \mu_1^0)'(\beta_1 - \mu_1^0)}{2V_1^0} \right\} \\&= \frac{1}{\sqrt{2\pi V_1^0}} \exp \left\{ -\frac{(\beta_1 - \mu_1^0)^2}{2V_1^0} \right\}\end{aligned}$$

- Likelihood:

$$\begin{aligned}f(\mathbf{y}|\beta, \sigma^2) &= \prod_i^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{(y_i - \mathbf{X}\beta)'(y_i - \mathbf{X}\beta)}{2\sigma^2} \right\} \\&= \prod_i^n \frac{1}{\sqrt{\pi\sigma^2}} \exp \left\{ -\frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2} \right\}\end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for the slope parameter β_1

$$f(\beta_1 | y, \beta_0, \sigma^2) \propto f(\beta_1) f(y | \beta_0, \beta_1, \sigma^2)$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for the slope parameter β_1

$$\begin{aligned} f(\beta_1 | y, \beta_0, \sigma^2) &\propto f(\beta_1) f(y | \beta_0, \beta_1, \sigma^2) \\ &\propto \frac{1}{\sqrt{2\pi V_1^0}} \exp \left\{ -\frac{(\beta_1 - \mu_1^0)^2}{2V_1^0} \right\} \times \\ &\quad \prod_i^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2} \right\} \end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for the slope parameter β_1

$$\begin{aligned} f(\beta_1 | y, \beta_0, \sigma^2) &\propto f(\beta_1) f(y | \beta_0, \beta_1, \sigma^2) \\ &\propto \frac{1}{\sqrt{2\pi V_1^0}} \exp \left\{ -\frac{(\beta_1 - \mu_1^0)^2}{2V_1^0} \right\} \times \\ &\quad \prod_i^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2} \right\} \\ &\propto \exp \left\{ -\left(\frac{1}{2V_1^0} + \frac{\sum_i^n x_i^2}{2\sigma^2} \right) \beta_1^2 + \right. \\ &\quad \left. \left(\frac{\mu_1^0}{V_1^0} + \frac{\sum_i^n (y_i - \beta_0) x_i}{\sigma^2} \right) \beta_1 + \text{const} \right\} \end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

If a pdf can be expressed as...

$$f(\theta) \propto \exp(a\theta^2 + b\theta + \text{const})$$

The corresponding distribution is a normal distribution: $N\left(-\frac{b}{2a}, -\frac{1}{2a}\right)$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

If a pdf can be expressed as...

$$f(\theta) \propto \exp(a\theta^2 + b\theta + \text{const})$$

The corresponding distribution is a normal distribution: $N\left(-\frac{b}{2a}, -\frac{1}{2a}\right)$

Conditional posterior for β_1 : $N(\mu_1^*, V_1^*)$

$$\begin{aligned}\mu_1^* &= V_1^* \left(\frac{\mu_1^0}{V_1^0} + \frac{\sum_i^n (y_i - \beta_0)x_i}{\sigma^2} \right) \\ V_1^* &= \left(\frac{1}{V_1^0} + \frac{\sum_i^n x_i^2}{\sigma^2} \right)^{-1}\end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for the intercept $\beta_0: N(\mu_0^*, V_0^*)$

Analogously to the conditional posterior for β_1 :

$$\begin{aligned}\mu_0^* &= V_0^* \left(\frac{\mu_0^0}{V_0^0} + \frac{\sum_i^n (y_i - \beta_1 x_i)}{\sigma^2} \right) \\ V_0^* &= \left(\frac{1}{V_0^0} + \frac{n}{\sigma^2} \right)^{-1}\end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for β_0 and β_1

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for β_0 and β_1

- It is also possible to derive conditional posterior for β_0 and β_1 .

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for β_0 and β_1

- It is also possible to derive conditional posterior for β_0 and β_1 .
- Prior: multivariate normal.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for β_0 and β_1

- It is also possible to derive conditional posterior for β_0 and β_1 .
- Prior: multivariate normal.
- For more details see Shikano (2014) Bayesian estimation of regression models,...

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Conditional posterior for σ^2

$$\begin{aligned}f(\sigma^2|y, \beta_0, \beta_1) &\propto f_{\Gamma^{-1}}(\sigma^2)f(y|\beta_0, \beta_1, \sigma^2) \\&\propto \frac{a0^{d0}}{\Gamma(d0)} (\sigma^2)^{-d0-1} \exp\left(-\frac{a0}{\sigma^2}\right) \\&\quad \prod_i^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2}\right\} \\&\propto (\sigma^2)^{-d0-1-\frac{n}{2}} \\&\quad \exp\left\{-\frac{1}{\sigma^2} \left(a0 + \frac{1}{2} \sum_i^n (y_i - \beta_0 - \beta_1 x_i)^2\right)\right\}\end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

An alternative approach

Bayesian estimation of bivariate regression models via MCMC

Susumu Shikano

Conditional posterior for σ^2

$$\sigma^2 | y, \beta_0, \beta_1 \sim IG \left(a0 + \frac{1}{2} \sum_i^n (y_i - \beta_0 - \beta_1 x_i)^2, d0 + \frac{n}{2} \right)$$

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Conditional posterior

$$\beta_1 | \beta_0, \sigma^2$$



σ^2 is set to 253.11. Prior for β_1 is $N(0, 10000)$.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Conditional posterior

$$\beta_1 | \beta_0, \sigma^2$$



σ^2 is set to 253.11. Prior for β_1 is $N(0, 10000)$.

**Bayesian
estimation of
bivariate
regression
models via MCMC**

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Conditional posterior

$$\beta_1 | \beta_0, \sigma^2$$



σ^2 is set to 253.11. Prior for β_1 is $N(0, 10000)$.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Simulation-based MCMC algorithm

- Obtaining the joint distribution from individual conditional distributions.
 - $f(\beta_0, \beta_1, \sigma^2)$ from...

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Simulation-based MCMC algorithm

- Obtaining the joint distribution from individual conditional distributions.
 - $f(\beta_0, \beta_1, \sigma^2)$ from...
 - $f(\beta_0|\beta_1, \sigma^2)$, $f(\beta_1|\beta_0, \sigma^2)$ and $f(\sigma^2|\beta_0, \beta_1)$
 - 1 Choose a starting value $\beta_1^{(0)}$ and $\sigma^{2(0)}$
 - 2 Draw $\beta_0^{(1)}$ from $f(\beta_0|\beta_1^{(0)}, \sigma^{2(0)})$.
 - 3 Draw $\beta_1^{(1)}$ from $f(\beta_1|\beta_0^{(1)}, \sigma^{2(0)})$.
 - 4 Draw $\sigma^{2(1)}$ from $f(\sigma^2|\beta_0^{(1)}, \beta_1^{(1)})$.
 - 5 \vdots
 - 6 Draw $\beta_0^{(g)}$ from $f(\beta_0|\beta_1^{(g-1)}, \sigma^{2(g-1)})$.
 - 7 Draw $\beta_1^{(g)}$ from $f(\beta_1|\beta_0^{(g)}, \sigma^{2(g-1)})$.
 - 8 Draw $\sigma^{2(g)}$ from $f(\sigma^2|\beta_0^{(g)}, \beta_1^{(g)})$.

Gibbs sampling

Bivariate regression model

```
> chain
      beta0      beta1 sigma.sqr
starting.value      NA      0.000000      1.0000
Iteration 1      59.52691     -2.654881     847.7251
Iteration 2      62.76098     -4.367976     966.1888
Iteration 3      69.56556     -4.469316     877.0900
Iteration 4      63.97678     -4.759496     704.2265
Iteration 5      71.45450     -5.533939     576.5033
Iteration 6      75.19838     -7.264575     511.4502
Iteration 7      81.29282     -9.173630     772.7803
Iteration 8      92.61787    -14.100181     592.2868
Iteration 9     104.85203    -12.859007     395.7271
Iteration 10     99.47174    -13.724145     275.8063
Iteration 11     100.49510    -14.362180     552.9542
Iteration 12     108.28597    -17.344073     311.3477
Iteration 13     111.61257    -17.510766     281.4529
Iteration 14     111.68330    -15.617246     315.2950
Iteration 15     112.97822    -18.594780     358.8411
Iteration 16     115.27897    -19.137953     286.8063
Iteration 17     121.49736    -19.647173     205.9642
```

An example Markov Chain

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Gibbs sampling

Bivariate regression model

```
> chain
      beta0      beta1 sigma.sqr
starting.value      NA      0.000000      1.0000
Iteration 1      59.52691     -2.654881     847.7251
Iteration 2      62.76098     -4.367976     966.1888
Iteration 3      69.56556     -4.469316     877.0900
Iteration 4      63.97678     -4.759496     704.2265
Iteration 5      71.45450     -5.533939     576.5033
Iteration 6      75.19838     -7.264575     511.4502
Iteration 7      81.29282     -9.173630     772.7803
Iteration 8      92.61787    -14.100181     592.2868
Iteration 9     104.85203    -12.859007     395.7271
Iteration 10     99.47174    -13.724145     275.8063
Iteration 11     100.49510    -14.362180     552.9542
Iteration 12     108.28597    -17.344073     311.3477
Iteration 13     111.61257    -17.510766     281.4529
Iteration 14     111.68330    -15.617246     315.2950
Iteration 15     112.97822    -18.594780     358.8411
Iteration 16     115.27897    -19.137953     286.8063
Iteration 17     121.49736    -19.647173     205.9642
```

An example Markov Chain

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Gibbs sampling

Bivariate regression model



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Gibbs sampling

Bivariate regression model



After some iterations (burn-in), the chain converged to an invariant distribution.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Gibbs sampling

Bivariate regression model



After 100 iterations.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Metropolis-Hasting Algorithm?

- More general MCMC of the Gibbs sampling
- Differently from the Gibbs sampling, the full set of the conditional posterior for all parameters is not required.

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Basic idea

Random draws from a posterior $f(\beta|y)$. The MH algorithm can be described by the following iterative steps for $t = 1, \dots, T$:

- 1 Set $\beta = \beta^{(t-1)}$
- 2 Generate new candidate values β' from a proposal distribution $q(\beta'|\beta)$
- 3 Calculate
$$\alpha = \min \left(1, \frac{f(\beta'|y)q(\beta|\beta')}{f(\beta|y)q(\beta'|\beta)} \right) = \min \left(1, \frac{f(y|\beta')f(\beta')q(\beta|\beta')}{f(y|\beta)f(\beta)q(\beta'|\beta)} \right).$$
- 4 Update $\beta^{(t)} = \beta'$ with probability α (acceptance). Otherwise set $\beta^{(t)} = \beta$ (rejection of β').

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- You obtained as posterior (=product of prior and likelihood) the following density function:

$$f(\theta) = \frac{a^d}{\Gamma(d)} \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right)$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- You obtained as posterior (=product of prior and likelihood) the following density function:

$$f(\theta) = \frac{a^d}{\Gamma(d)} \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right)$$

- Which form has the distribution?

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- You obtained as posterior (=product of prior and likelihood) the following density function:

$$f(\theta) = \frac{a^d}{\Gamma(d)} \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right)$$

- Which form has the distribution?
- What is the expected value?

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- You obtained as posterior (=product of prior and likelihood) the following density function:

$$f(\theta) = \frac{a^d}{\Gamma(d)} \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right)$$

- Which form has the distribution?
- What is the expected value?
- What is the variance?
- ...

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- Parameter values: $a=2$, $d=3$
- Initial value: 30
- Proposal dist.: normal distribution with $sd=3$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up


Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

Markov Chain (after 200 iter)



Numerical and Analytical IG



- Parameter values: $a=2$, $d=3$

Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling


MCMC

Wrap up


Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

Markov Chain (after 200 iter)



Numerical and Analytical IG



- Parameter values: $a=2, d=3$
- Average of IG: $\frac{a}{d-1} = 1$

Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- MH also works even if we know only the following posterior

$$\begin{aligned} f(\theta) &= \frac{a^d}{\Gamma(d)} \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right) \\ &\propto \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right) \end{aligned}$$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Metropolis-Hasting Algorithm

Application: Inverse Gamma Distribution

- MH also works even if we know only the following posterior

$$\begin{aligned}f(\theta) &= \frac{a^d}{\Gamma(d)} \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right) \\&\propto \theta^{-d-1} \exp\left(-\frac{a}{\theta}\right)\end{aligned}$$

- $\frac{f(y|\beta')f(\beta')q(\beta|\beta')}{f(y|\beta)f(\beta)q(\beta'|\beta)}$ can cancel out the constant term!

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

A drawback of MH Algorithm

If a proposal distribution's variance of proposal distribution is

- too small: Markov chains converge very slowly.
- too large: acceptance rate is too small (inefficient).


Any algorithm which requires no proposal distribution?

Procedure

- Goal: Generation of random numbers from a pdf $f(x)$
- You have no program for $f(x)$.
 - 1 Choose a starting value x_0 for which $f(x_0) > 0$.
 - 2 Sample a y value uniformly between 0 and $f(x_0)$.
 - 3 Draw a horizontal line across the curve at this y position.
 - 4 Sample a point (x, y) from the line segments within the curve.
 - 5 Repeat from step 2 using the new x value.

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm


Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm



Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $IG(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm



Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $IG(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm



Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $IG(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm



Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $IG(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm



Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $\text{IG}(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm



Slice Sampling

MCMC

Wrap up

Slice Sampling

Example: $IG(a=2, d=3)$ with $x_0 = 1$



Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

- Tools for random number generation based on a certain distribution. . .

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

- Tools for random number generation based on a certain distribution. . .
- with many iterations.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

- Tools for random number generation based on a certain distribution. . .
- with many iterations.
- Random number generation in each iteration depends on the last iteration.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

- Tools for random number generation based on a certain distribution. . .
- with many iterations.
- Random number generation in each iteration depends on the last iteration.
- → Markov-Chain-Monte-Carlo

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

- Tools for **random number generation** based on a certain distribution. . .
- with many iterations.
- Random number generation in each iteration depends on the last iteration.
- → Markov-Chain-**Monte-Carlo**

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Gibbs, MH and Slice sampling

- Tools for random number generation based on a certain distribution. . .
- with many iterations.
- Random number generation in each iteration **depends on the last iteration**.
- → **Markov-Chain**-Monte-Carlo

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Markov chain

- A stochastic process satisfying the Markov property.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Markov chain

- A stochastic process satisfying the Markov property.
- Markov property: the probability distribution at $t + 1$ depends only on the state of the system at t .

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Markov chain

- A stochastic process satisfying the Markov property.
- Markov property: the probability distribution at $t + 1$ depends only on the state of the system at t .
- In a finite state space with all probabilities positive, there is a unique invariant distribution.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Markov chain

- A stochastic process satisfying the Markov property.
- Markov property: the probability distribution at $t + 1$ depends only on the state of the system at t .
- In a finite state space with all probabilities positive, there is a unique invariant distribution.
 - Condition: The chain has to be irreducible and aperiodic.
 - For large enough n , the initial state plays almost no role.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Markov-Chain-Monte-Carlo (MCMC)

Advantages: simple and straightforward interpretation!

- Quite easy since we generate a „dataset” of draws from MC.
- Simple calculation of any quantity of interest:
 - the expected value or the median of that posterior, just calculate those of the generated draws.
 - 95% credible intervals, just find the 2.5th and 97.5th percentiles of the draws.
- You need just descriptive statistics to describe MC.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Bayesian regression models

- In conjugacy analysis posterior can be derived analytically.
- The larger n /the smaller the dispersion of prior, the more similar results with the maximum likelihood.
- Alternative to conjugacy analysis:
 - Deriving conditional posterior
 - Run Gibbs sampling (one of MCMC)
 - Obtain joint and marginal posterior
- Gibbs sampling is one possible MCMC algorithm.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Introduction

Alternative
approach

Deriving the posterior

Gibbs sampling

Metropolis-Hasting
Algorithm

Slice Sampling

MCMC

Wrap up

Rejection Sampling or Acceptance Rejection Sampling

- Goal: Generation of random numbers from a pdf $f(x)$
- You have no program for $f(x)$, but for another distribution $g(x)$
 - ① You find a constant c so that $f(x) \leq cg(x)$ for all x .
 - ② You generate a random number x^* from $g(x)$.
 - ③ You generate a random number u from $U[0, 1]$.
 - ④ If $u \leq f(x^*)/cg(x^*)$ you accept x^* as random number from $f(x)$. Otherwise you reject it.
 - ⑤ Go to Step 2.

Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Rejection
Sampling

Markov Chain

Markov process

- A stochastic process X_t taking values...
 - in the finite set $S = \{1, 2, \dots, s\}$
 - t : time or iteration

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Markov process

- A stochastic process X_t taking values...
 - in the finite set $S = \{1, 2, \dots, s\}$
 - t : time or iteration
- p_{ij} is transition probabilities
 - $p_{ij} = \Pr(X_{t+1} = j | X_t = i), i, j \in S.$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Markov process

- A stochastic process X_t taking values...
 - in the finite set $S = \{1, 2, \dots, s\}$
 - t : time or iteration
- p_{ij} is transition probabilities
 - $p_{ij} = \Pr(X_{t+1} = j | X_t = i), i, j \in S.$
- Markov property: the probability distribution at $t + 1$ depends only on the state of the system at t .

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Markov process

- A stochastic process X_t taking values...
 - in the finite set $S = \{1, 2, \dots, s\}$
 - t : time or iteration
- p_{ij} is transition probabilities
 - $p_{ij} = \Pr(X_{t+1} = j | X_t = i), i, j \in S.$
- Markov property: the probability distribution at $t + 1$ depends only on the state of the system at t .
- Note that p_{ij} are constant over t .

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Transition matrix

- Transition probabilities: p_{ij}
 - $p_{ij} \geq 0$
 - $\sum_{j=1}^s p_{ij} = 1$
- $s \times s$ transition matrix $P = \{p_{ij}\}$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Transition matrix

- Transition probabilities: p_{ij}
 - $p_{ij} \geq 0$
 - $\sum_{j=1}^s p_{ij} = 1$
- $s \times s$ transition matrix $P = \{p_{ij}\}$
- Distribution at $t + 2$?

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Transition matrix

- Transition probabilities: p_{ij}
 - $p_{ij} \geq 0$
 - $\sum_{j=1}^s p_{ij} = 1$
- $s \times s$ transition matrix $P = \{p_{ij}\}$
- Distribution at $t + 2$?
 - $p_{ij}^{(2)} = \sum_k p_{ik} p_{kj}$

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Transition matrix

- Transition probabilities: p_{ij}
 - $p_{ij} \geq 0$
 - $\sum_{j=1}^s p_{ij} = 1$
- $s \times s$ transition matrix $P = \{p_{ij}\}$
- Distribution at $t + 2$?
 - $p_{ij}^{(2)} = \sum_k p_{ik} p_{kj}$
 - This is given by $PP \equiv P^2$

Invariant distribution

- The probability distribution $\pi = (\pi_1, \dots, \pi_s)'$ is invariant for P
- if $\pi' = \pi' P$,
- or $\pi_j = \sum_i \pi_i p_{ij}$ for $j = 1, \dots, s$.

Invariant distribution

- The probability distribution $\pi = (\pi_1, \dots, \pi_s)'$ is invariant for P
- if $\pi' = \pi' P$,
- or $\pi_j = \sum_i \pi_i p_{ij}$ for $j = 1, \dots, s$.
- Note that π' is an eigenvector of P .

Theorem

- Suppose S is finite and $p_{ij} > 0 \forall i, j$.
- Then there exists a unique probability distribution $\pi_j, j \in S$
- such that $\sum_i \pi_i p_{ij} = \pi_j \forall j \in S$.
- Further, $|p_{ij}^{(n)} - \pi_j| \leq r^n$,
- where $0 < r < 1$, for all i, j and $n \geq 1$.

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Theorem

- Suppose S is finite and $p_{ij} > 0 \forall i, j$.
- Then there exists a unique probability distribution $\pi_j, j \in S$
- such that $\sum_i \pi_i p_{ij} = \pi_j \forall j \in S$.
- Further, $|p_{ij}^{(n)} - \pi_j| \leq r^n$,
- where $0 < r < 1$, for all i, j and $n \geq 1$.

Substantive meaning of the theorem

- In a finite state space with all probabilities positive, there is a unique invariant distribution.
- For large enough n , the initial state plays almost no role.

More generalized Theorem

- Let P be irreducible and aperiodic over a finite state space.
- Then there exists a unique probability distribution $\pi_j, j \in S$
- such that $\sum_i \pi_i p_{ij} = \pi_j \forall j \in S$
- and $|p_{ij}^{(n)} - \pi_j| \leq r^{n/\nu},$
- where $0 < r < 1$, for all i, j and for some positive integer ν .

Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Rejection
Sampling

Markov Chain

More generalized Theorem

- Let P be irreducible and aperiodic over a finite state space.
- Then there exists a unique probability distribution $\pi_j, j \in S$
- such that $\sum_i \pi_i p_{ij} = \pi_j \forall j \in S$
- and $|p_{ij}^{(n)} - \pi_j| \leq r^{n/\nu}$,
- where $0 < r < 1$, for all i, j and for some positive integer ν .

- Irreducible: starting from state i , the process can reach any other state with positive probability.
- Aperiodic: if $p_{ii}^{(n)} > 0 \forall i$ and for sufficiently large n .

Bayesian
estimation of
bivariate
regression
models via MCMC
Susumu Shikano

Rejection
Sampling

Markov Chain

Markov Chains in Continuous Spaces

Bayesian
estimation of
bivariate
regression
models via MCMC

Susumu Shikano

Rejection
Sampling

Markov Chain

Markov-Process taking values in \mathbb{R}

- $f_{(X_1, \dots, X_n | X_0 = x_0)}(x_1, \dots, x_n) = p(x_0, x_1)p(x_1, x_2) \cdots p(x_{n-1}, x_n)$
- $p(x, y)$: Transitional kernel
- Invariant density: $\pi(y) = \int_{\mathbb{R}} \pi(x)p(x, y)dx$.