

Basics

Applied Bayesian Statistics
Winter Term 2018

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Susumu Shikano
GSDS

This session

Basics

Susumu Shikano

Goals

- Basic logic and concepts of Bayesian inference.
- Likelihood, prior and posterior
- Conjugate analysis

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

An example research question

Which ideological orientation does this student candidate have? Left or right?

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

An example research question

Which ideological orientation does this student candidate have? Left or right?



Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

An example research question

Which ideological orientation does this student candidate have? Left or right?



Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

An example research question

Which ideological orientation does this student candidate have? Left or right?



Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

An example research question

Which ideological orientation does this student candidate have? Left or right?



- Right (RCDS)



- Left (JUSO)

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

An example research question

Now assume...

- A person claims that he should be able to distinguish ideological orientation of others (left or right) only based on the facial characteristics in 7 out of 10 cases.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

An example research question

Now assume...

- A person claims that he should be able to distinguish ideological orientation of others (left or right) only based on the facial characteristics in 7 out of 10 cases.
- You are sceptical.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

An example research question

Now assume...

- A person claims that he should be able to distinguish ideological orientation of others (left or right) only based on the facial characteristics in 7 out of 10 cases.
- You are sceptical.
 - If you are frequentist: Let him judge the ideological orientation of many people.
 - Given: He could correctly judge the ideological orientation of 60 out of 100 persons.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

you take a model with the maximum likelihood.

- There are two models
 - H_1 : He can distinguish 7 out of 10.
 - H_0 : His answer is random.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

you take a model with the maximum likelihood.

- There are two models
 - H_1 : He can distinguish 7 out of 10.
 - H_0 : His answer is random.
- ... and you calculate how likely the data (y) is collected under different models.
- → Likelihood

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Computing Likelihood ($p(y|\theta)$)

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

Computing Likelihood ($p(y|\theta)$)

- Which probability distribution form has a percentage?

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Computing Likelihood ($p(y|\theta)$)

- Which probability distribution form has a percentage?
- Binomial distribution

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

Computing Likelihood ($p(y|\theta)$)

- Which probability distribution form has a percentage?
- Binomial distribution

Bernoulli distribution

Coin toss: If the coin is fair,

$$\Pr(Y_i = \text{"head"}) = 0.5$$

$$\Pr(Y_i = \text{"tail"}) = 0.5$$

$$\Pr(Y_i \neq \text{"head"}, \text{"tail"}) = 0$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

Bernoulli distribution

More generally

$$\Pr(Y_i = 1) = \pi$$

$$\Pr(Y_i = 0) = 1 - \pi$$

$$\Pr(Y_i \neq 0, 1) = 0$$

- parameter: π
- $\pi \in [0, 1]$

This is called Bernoulli distribution.

Formally

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1 - \pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

You repeat coin tossing for three times ($N=3$).

Important assumption: each trial is independent and identical.

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1 - \pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

You repeat coin tossing for three times ($N=3$).

Important assumption: each trial is independent and identical.

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1 - \pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

How likely do you have heads

- for three times?: $\Pr(Y_i = 3) = \pi^3$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

You repeat coin tossing for three times ($N=3$).

Important assumption: each trial is independent and identical.

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1 - \pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

How likely do you have heads

- for three times?: $\Pr(Y_i = 3) = \pi^3$
- never?: $\Pr(Y_i = 0) = (1 - \pi)^3$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

You repeat coin tossing for three times ($N=3$).

Important assumption: each trial is independent and identical.

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1 - \pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

How likely do you have heads

- for three times?: $\Pr(Y_i = 3) = \pi^3$
- never?: $\Pr(Y_i = 0) = (1 - \pi)^3$
- for twice?: $\Pr(Y_i = 2) = \pi^2(1 - \pi)?$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

You repeat coin tossing for three times ($N=3$).

Important assumption: each trial is independent and identical.

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1 - \pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

How likely do you have heads

- for three times?: $\Pr(Y_i = 3) = \pi^3$
- never?: $\Pr(Y_i = 0) = (1 - \pi)^3$
- for twice?: $\Pr(Y_i = 2) = \pi^2(1 - \pi)?$
- ... No! $\Pr(Y_i = 2) = 3\pi^2(1 - \pi)$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

You repeat coin tossing for three times ($N=3$).

Important assumption: each trial is independent and identical.

$$Y_i \sim f_{Bern}(y_i|\pi) = \begin{cases} \pi^{y_i}(1-\pi)^{1-y_i} & \text{for } y_i = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

How likely do you have heads

- for three times?: $\Pr(Y_i = 3) = \pi^3$
- never?: $\Pr(Y_i = 0) = (1 - \pi)^3$
- for twice?: $\Pr(Y_i = 2) = \pi^2(1 - \pi)?$
- ... No! $\Pr(Y_i = 2) = 3\pi^2(1 - \pi)$

$$3 = \frac{3 \times 2}{2 \times 1} = \binom{N}{y_i} = \frac{N!}{y_i!(N - y_i)!}$$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Binomial distribution

Generally...

$$\begin{aligned}f_b(y_i|\pi, N) &= \Pr(Y_i = y_i|\pi, N) \\&= \begin{cases} \binom{N}{y_i} \pi^{y_i} (1 - \pi)^{N-y_i} & \text{for } y_i = 0, 1, \dots, N \\ 0 & \text{otherwise} \end{cases}\end{aligned}$$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are frequentist...

Back to our example

- There are two models
 - H_1 : He can distinguish 7 out of 10. $\rightarrow \pi = 0.7$
 - H_0 : His answer is random. $\rightarrow \pi = 0.5$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

Back to our example

- There are two models
 - H_1 : He can distinguish 7 out of 10. $\rightarrow \pi = 0.7$
 - H_0 : His answer is random. $\rightarrow \pi = 0.5$
- ... and you calculate how likely the data (y) is collected under different models.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

Back to our example

- There are two models
 - H_1 : He can distinguish 7 out of 10. $\rightarrow \pi = 0.7$
 - H_0 : His answer is random. $\rightarrow \pi = 0.5$
- ... and you calculate how likely the data (y) is collected under different models.

$$\begin{aligned}\Pr(y|H_1) &= \binom{100}{60} \cdot 0.7^{60} \cdot 0.3^{40} \\ &\approx 0.0085\end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are frequentist...

Back to our example

- There are two models
 - H_1 : He can distinguish 7 out of 10. $\rightarrow \pi = 0.7$
 - H_0 : His answer is random. $\rightarrow \pi = 0.5$
- ... and you calculate how likely the data (y) is collected under different models.

$$\begin{aligned}\Pr(y|H_1) &= \binom{100}{60} \cdot 0.7^{60} \cdot 0.3^{40} \\ &\approx 0.0085\end{aligned}$$

$$\begin{aligned}\Pr(y|H_0) &= \binom{100}{60} \cdot 0.5^{60} \cdot 0.5^{40} \\ &\approx 0.0108 \rightarrow \text{larger likelihood!}\end{aligned}$$

* $y = \{60 \text{ times right; } 40 \text{ times wrong}\}$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

What is at stake?: Your belief about how likely the models are!!

- $\Pr(H)$ instead of $\Pr(y)$.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

What is at stake?: Your belief about how likely the models are!!

- $\Pr(H)$ instead of $\Pr(y)$.
- Bayes Theorem!

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H) p(H)}^{\text{likelihood}}}{p(y)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

What is at stake?: Your belief about how likely the models are!!

- $\Pr(H)$ instead of $\Pr(y)$.
- Bayes Theorem!

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H) p(H)}^{\text{likelihood}}}{p(y)}$$

- $\Pr(H)??$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

What is at stake?: Your belief about how likely the models are!!

- $\Pr(H)$ instead of $\Pr(y)$.
- Bayes Theorem!

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H) p(H)}^{\text{likelihood}}}{p(y)}$$

- $\Pr(H)??$
- The belief which you have about models **prior** to the data collection.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

What is at stake?: Your belief about how likely the models are!!

- $\Pr(H)$ instead of $\Pr(y)$.
- Bayes Theorem!

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H) p(H)}^{\text{likelihood}}}{p(y)}$$

- $\Pr(H)??$
- The belief which you have about models **prior** to the data collection.
- E.g.: Before testing, it is 50:50 whether he or you are right. There can be no further model ($\Pr(H_1) + \Pr(H_0) = 1$).
- $\rightarrow \Pr(H_1) = 0.5$ and $\Pr(H_0) = 0.5$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

$$\Pr(H_1) = 0.5 \text{ and } \Pr(H_0) = 0.5$$

- The first trial: successful judgement.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

$$\Pr(H_1) = 0.5 \text{ and } \Pr(H_0) = 0.5$$

- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} \end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

$$\Pr(H_1) = 0.5 \text{ and } \Pr(H_0) = 0.5$$

- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.5}{p(y)} = \frac{0.25}{p(y)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

$$\Pr(H_1) = 0.5 \text{ and } \Pr(H_0) = 0.5$$

- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \underbrace{0.5}_{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} = \frac{0.35}{0.35 + 0.25} \approx 0.583 \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.5}{p(y)} = \frac{0.25}{p(y)} = \frac{0.25}{0.35 + 0.25} \approx 0.417$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

$$\Pr(H_1) = 0.5 \text{ and } \Pr(H_0) = 0.5$$

- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} = \frac{0.35}{0.35 + 0.25} \approx 0.583 \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.5}{p(y)} = \frac{0.25}{p(y)} = \frac{0.25}{0.35 + 0.25} \approx 0.417$$

- Even after only one observation you can say something about the models.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

After first trial: $\Pr(H_1) = 0.583$ and $\Pr(H_0) = 0.417$

- The next trial: successful judgement again.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.7^1 \cdot 0.3^0 \times 0.583}{p(y)} \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.417}{p(y)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

After first trial: $\Pr(H_1) = 0.583$ and $\Pr(H_0) = 0.417$

- The next trial: successful judgement again.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.7^1 \cdot 0.3^0 \times 0.583}{p(y)} = \frac{0.408}{p(y)} = \frac{0.408}{0.408 + 0.209} \approx 0.661 \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.417}{p(y)} = \frac{0.209}{p(y)} = \frac{0.209}{0.408 + 0.209} \approx 0.339$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

After first trial: $\Pr(H_1) = 0.583$ and $\Pr(H_0) = 0.417$

- The next trial: successful judgement again.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.7^1 \cdot 0.3^0 \times 0.583}{p(y)} = \frac{0.408}{p(y)} = \frac{0.408}{0.408 + 0.209} \approx 0.661 \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.417}{p(y)} = \frac{0.209}{p(y)} = \frac{0.209}{0.408 + 0.209} \approx 0.339$$

- The posterior is different after the same observation (successful judgement).
- You are more sure that he is right (H_1 is correct).

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

After 100 trials with 60 correct judgements...

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.0085 \times 0.5}{p(y)} = \frac{0.0425}{p(y)} = \frac{0.0425}{0.0425 + 0.0504} \approx 0.439 \end{aligned}$$

$$p(H_0|y) = \frac{0.0108 \times 0.5}{p(y)} = \frac{0.0504}{p(y)} = \frac{0.0504}{0.0425 + 0.0504} \approx 0.561$$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are Bayesian...

After 100 trials with 60 correct judgements...

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.0085 \times 0.5}{p(y)} = \frac{0.0425}{p(y)} = \frac{0.0425}{0.0425 + 0.0504} \approx 0.439 \end{aligned}$$

$$\begin{aligned} p(H_0|y) &= \frac{0.0108 \times 0.5}{p(y)} = \frac{0.0504}{p(y)} = \frac{0.0504}{0.0425 + 0.0504} \approx 0.561 \end{aligned}$$

- You tend to believe more H_0 than H_1 .

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are Bayesian...

After 100 trials with 60 correct judgements...

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.0085 \times 0.5}{p(y)} = \frac{0.0425}{p(y)} = \frac{0.0425}{0.0425 + 0.0504} \approx 0.439 \end{aligned}$$

$$p(H_0|y) = \frac{0.0108 \times 0.5}{p(y)} = \frac{0.0504}{p(y)} = \frac{0.0504}{0.0425 + 0.0504} \approx 0.561$$

- You tend to believe more H_0 than H_1 .
- What if we have further models besides H_0 and H_1 ?
 - He can distinguish 6 out of 10, 8 out of 10,...

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are Bayesian...

After 100 trials with 60 correct judgements...

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{0.0085 \times 0.5}{p(y)} = \frac{0.0425}{p(y)} = \frac{0.0425}{0.0425 + 0.0504} \approx 0.439 \end{aligned}$$

$$p(H_0|y) = \frac{0.0108 \times 0.5}{p(y)} = \frac{0.0504}{p(y)} = \frac{0.0504}{0.0425 + 0.0504} \approx 0.561$$

- You tend to believe more H_0 than H_1 .
- What if we have further models besides H_0 and H_1 ?
 - He can distinguish 6 out of 10, 8 out of 10,...
- You can extend the calculation so many times as you like.

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem
(revisited)

Summary

If you are Bayesian...

Not only two models, but more models



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

Not only two models, but more models



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

Not only two models, but more models



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

Not only two models, but more models



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

Not only two models, but more models



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

Not only two models, but more models



- However, it is often the case that we have uncountable number of models (or better: parameter values)...

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

If you are Bayesian...

Not only two models, but more models



- However, it is often the case that we have uncountable number of models (or better: parameter values)...
- We can solve the problem analytically (conjugate analysis).

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis


Bayes Theorem (revisited)

Summary

Conjugate analysis

One can use beta distribution as continuous prior...

Beta distribution: $f_{\beta}(\pi|a, b)$



Why?

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis


Bayes Theorem
(revisited)

Summary

Conjugate analysis

One can use beta distribution as continuous prior...

Beta distribution: $f_{\beta}(\pi|a, b)$



Why?

- The relevant range (between 0 and 1)
- Flexible shapes

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

Beta distribution: $f_{\beta}(\pi|a, b)$

$$f_{\beta}(\pi|a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1}$$

$$0 \leq \pi \leq 1, a, b > 0$$

$$E(\pi) = \frac{a}{a+b}$$

$$Var(\pi) = \frac{ab}{(a+b)^2(a+b+1)}$$

* If n is a positive integer:

$$\Gamma(n) = (n-1)!$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis


Bayes Theorem (revisited)

Summary

Conjugate analysis

If one has a continuous prior based on beta distribution...

- Using beta priors
- and data (60 out of 100 are correct judgements),
- posterior can be analytically calculated.



Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis


Bayes Theorem
(revisited)

Summary

Conjugate analysis

If one has a continuous prior based on beta distribution...

- Using beta priors
- and data (60 out of 100 are correct judgements),
- posterior can be analytically calculated.



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned}f_{\beta}(\pi|a, b) &= \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \\f_b(y_i|\pi, N) &= \Pr(Y_i = y_i|\pi, N) \\&= \begin{cases} \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i} & \text{for } y_i = 0, 1, \dots, N \\ 0 & \text{otherwise} \end{cases}\end{aligned}$$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned}f_{\beta}(\pi|a, b) &= \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \\f_b(y_i|\pi, N) &= \Pr(Y_i = y_i|\pi, N) \\&= \begin{cases} \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i} & \text{for } y_i = 0, 1, \dots, N \\ 0 & \text{otherwise} \end{cases}\end{aligned}$$

- Both functions have a similar component.

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned}f_{\beta}(\pi|a, b) &= \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \\f_b(y_i|\pi, N) &= \Pr(Y_i = y_i|\pi, N) \\&= \begin{cases} \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i} & \text{for } y_i = 0, 1, \dots, N \\ 0 & \text{otherwise} \end{cases}\end{aligned}$$

- Both functions have a similar component.
- Beta distribution has normalizing constant.

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$f(\pi|y_i) = \frac{f_b(y_i|\pi) \cdot f_\beta(\pi)}{f(y_i)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned} f(\pi|y_i) &= \frac{f_b(y_i|\pi) \cdot f_\beta(\pi)}{f(y_i)} \\ &= \frac{\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \cdot \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i}}{f(y_i)} \end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned}f(\pi|y_i) &= \frac{f_b(y_i|\pi) \cdot f_\beta(\pi)}{f(y_i)} \\&= \frac{\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \cdot \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i}}{f(y_i)} \\&= \mathcal{C} \cdot \pi^{a+y_i-1} (1-\pi)^{b+N-y_i-1}\end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned}f(\pi|y_i) &= \frac{f_b(y_i|\pi) \cdot f_\beta(\pi)}{f(y_i)} \\&= \frac{\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \cdot \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i}}{f(y_i)} \\&= \mathcal{C} \cdot \pi^{a+y_i-1} (1-\pi)^{b+N-y_i-1} \\&= \frac{\Gamma(a+b+N)}{\Gamma(a+y_i)\Gamma(b+N-y_i)} \pi^{a+y_i-1} (1-\pi)^{b+N-y_i-1}\end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

Calculating posterior by multiplying beta distribution and binomial distribution?

$$\begin{aligned}f(\pi|y_i) &= \frac{f_b(y_i|\pi) \cdot f_\beta(\pi)}{f(y_i)} \\&= \frac{\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \pi^{a-1} (1-\pi)^{b-1} \cdot \binom{N}{y_i} \pi^{y_i} (1-\pi)^{N-y_i}}{f(y_i)} \\&= \mathcal{C} \cdot \pi^{a+y_i-1} (1-\pi)^{b+N-y_i-1} \\&= \frac{\Gamma(a+b+N)}{\Gamma(a+y_i)\Gamma(b+N-y_i)} \pi^{a+y_i-1} (1-\pi)^{b+N-y_i-1} \\&= f_\beta(\pi|a+y_i, b+N-y_i)\end{aligned}$$

- Prior and posterior have a same distribution form → Conjugacy

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

If one has a continuous prior based on beta distribution...



Or...

- prior: $p(\pi) = f_{\beta}(a, b)$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

If one has a continuous prior based on beta distribution...



Or...

- prior: $p(\pi) = f_{\beta}(a, b)$
- likelihood: $f_b(60|\pi, 100)$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

If one has a continuous prior based on beta distribution...



Or...

- prior: $p(\pi) = f_{\beta}(a, b)$
- likelihood: $f_b(60|\pi, 100)$
- posterior: $p(\pi|y) = f_{\beta}(a + 60, b + 100 - 60)$

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Conjugate analysis

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugacy

- Prior and posterior have a same distribution form.
- Easy to compute.

Conjugate analysis

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)


Summary

Conjugacy

- Prior and posterior have a same distribution form.
- Easy to compute.
- Use of posterior as prior in subsequent analysis

Conjugate analysis

What we can observe here:



- The more certain $p(\theta)$ is,

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What we can observe here:



- The more certain $p(\theta)$ is,

... the more the prior distribution influences the posterior.

→ A more different result from ML.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...


Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What we can observe here:



- The more certain $p(\theta)$ is,
- The smaller n is,*

... the more the prior distribution influences the posterior.

→ A more different result from ML.

* Here, 6 out of 10 are correct judgement.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis


Bayes Theorem (revisited)

Summary


Conjugate analysis

What kind of statement?

Posterior for Prior: $f(\pi|1,1)$



Posterior for Prior: $f(\pi|3,1.5)$



Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...



Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What kind of statement?



- You can report the posterior percentage of your model(s).

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...



Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What kind of statement?



- You can report the posterior percentage of your model(s).
- But how can we calculate the percentage?

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...



Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What kind of statement?



- You can report the posterior percentage of your model(s).
- But how can we calculate the percentage?
- You have to integrate the posterior.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...



Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What kind of statement?



- You can report the posterior percentage of your model(s).
- But how can we calculate the percentage?
- You have to integrate the posterior.
- Sometimes difficult or unfeasible. . .

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist. . .

If you are Bayesian. . .



Conjugate analysis

Bayes Theorem (revisited)

Summary

Conjugate analysis

What kind of statement?



- You can report the posterior percentage of your model(s).
- But how can we calculate the percentage?
- You have to integrate the posterior.
- Sometimes difficult or unfeasible. . .
- Solution: Markov-Chain Monte-Carlo

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist. . .

If you are Bayesian. . .

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H)}^{\text{likelihood}} \overbrace{p(H)}^{\text{prior}}}{p(y)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H)}^{\text{likelihood}} \overbrace{p(H)}^{\text{prior}}}{p(y)}$$
$$p(H|y) \propto p(y|H)p(H)$$

- $\Pr(y)$ can be ignored as constant for H .

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H)}^{\text{likelihood}} \overbrace{p(H)}^{\text{prior}}}{p(y)}$$
$$p(H|y) \propto p(y|H)p(H)$$

- $\Pr(y)$ can be ignored as constant for H .
- But: What is $\Pr(y)$? Probability of data?

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H)}^{\text{likelihood}} \overbrace{p(H)}^{\text{prior}}}{p(y)}$$
$$p(H|y) \propto p(y|H)p(H)$$

- $\Pr(y)$ can be ignored as constant for H .
- But: What is $\Pr(y)$? Probability of data?

If there are only two models: H_1 and H_0 ,

$$p(y|H_1)p(H_1) + p(y|H_0)p(H_0) = p(y)$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\underbrace{p(H|y)}_{\text{posterior}} = \frac{\overbrace{p(y|H)}^{\text{likelihood}} \overbrace{p(H)}^{\text{prior}}}{p(y)}$$
$$p(H|y) \propto p(y|H)p(H)$$

- $\Pr(y)$ can be ignored as constant for H .
- But: What is $\Pr(y)$? Probability of data?

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...
If you are Bayesian...
Conjugate analysis

Bayes Theorem (revisited)

Summary

If there are only two models: H_1 and H_0 ,

$$p(y|H_1)p(H_1) + p(y|H_0)p(H_0) = p(y)$$

In general:

$$\sum_H p(y|H)p(H) = p(y)$$

Bayes Theorem (revisited)

The first trial

- Prior: $\Pr(H_1) = 0.5$ and $\Pr(H_0) = 0.5$
- The first trial: successful judgement.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

The first trial

- Prior: $\Pr(H_1) = 0.5$ and $\Pr(H_0) = 0.5$
- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} \end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

The first trial

- Prior: $\Pr(H_1) = 0.5$ and $\Pr(H_0) = 0.5$
- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.5}{p(y)} = \frac{0.25}{p(y)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

The first trial

- Prior: $\Pr(H_1) = 0.5$ and $\Pr(H_0) = 0.5$
- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} = \frac{0.35}{0.35 + 0.25} \approx 0.583 \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.5}{p(y)} = \frac{0.25}{p(y)} = \frac{0.25}{0.35 + 0.25} \approx 0.417$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

The first trial

- Prior: $\Pr(H_1) = 0.5$ and $\Pr(H_0) = 0.5$
- The first trial: successful judgement.

$$\begin{aligned} p(H_1|y) &= \frac{p(y|H_1)p(H_1)}{p(y)} \\ &= \frac{\overbrace{0.7^1 \cdot 0.3^0}^{\text{Likelihood}} \times \overbrace{0.5}^{\text{prior}}}{p(y)} = \frac{0.35}{p(y)} = \frac{0.35}{0.35 + 0.25} \approx 0.583 \end{aligned}$$

$$p(H_0|y) = \frac{0.5^1 \cdot 0.5^0 \times 0.5}{p(y)} = \frac{0.25}{p(y)} = \frac{0.25}{0.35 + 0.25} \approx 0.417$$

- $p(y)$: normalizing constant ensuring $p(H_1|y) + p(H_0|y) = 1$.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$p(H|y) = \frac{p(y|H)p(H)}{p(y)}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\begin{aligned} p(H|y) &= \frac{p(y|H)p(H)}{p(y)} \\ &= \frac{p(y|H)p(H)}{\sum_H p(y|H)p(H)} \end{aligned}$$

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Bayes Theorem (revisited)

Bayes Theorem

$$\begin{aligned} p(H|y) &= \frac{p(y|H)p(H)}{p(y)} \\ &= \frac{p(y|H)p(H)}{\sum_H p(y|H)p(H)} \end{aligned}$$

In terms of densities:

$$\begin{aligned} f(\theta|y) &= \frac{f(y|\theta)f(\theta)}{f(y)} \\ &= \frac{f(y|\theta)f(\theta)}{\int f(y|\theta)f(\theta)d\theta} \end{aligned}$$

Here again: integration is required!

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Short summary

- In Bayesian inference the (posterior) probability of individual models is at stake.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Short summary

- In Bayesian inference the (posterior) probability of individual models is at stake.
- You can obtain the posterior even after observing one single case.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Short summary

- In Bayesian inference the (posterior) probability of individual models is at stake.
- You can obtain the posterior even after observing one single case.
- The posterior also depends on your prior belief.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Short summary

- In Bayesian inference the (posterior) probability of individual models is at stake.
- You can obtain the posterior even after observing one single case.
- The posterior also depends on your prior belief.
- For a larger number of observations frequentists and Bayesian have similar results.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Short summary

- In Bayesian inference the (posterior) probability of individual models is at stake.
- You can obtain the posterior even after observing one single case.
- The posterior also depends on your prior belief.
- For a larger number of observations frequentists and Bayesian have similar results.
- Conjugacy of the beta distribution with the likelihood based on the binomial distribution.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Short summary

- In Bayesian inference the (posterior) probability of individual models is at stake.
- You can obtain the posterior even after observing one single case.
- The posterior also depends on your prior belief.
- For a larger number of observations frequentists and Bayesian have similar results.
- Conjugacy of the beta distribution with the likelihood based on the binomial distribution.
- For more details: Susumu Shikano (2014), Bayesian estimation of regression models. Henning Best and Christof Wolf, eds. *The SAGE Handbook of Regression Analysis and Causal Inference*, Sage. p.31-54.

Basics

Susumu Shikano

Bayesian Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem (revisited)

Summary

Back to the first question

Are people capable to estimate the ideological orientation based on facial traits?

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Back to the first question

Are people capable to estimate the ideological orientation based on facial traits?

- 164 students of the University of Konstanz judged 35 pictures of student candidates unknown to them.

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Back to the first question

Are people capable to estimate the ideological orientation based on facial traits?

- 164 students of the University of Konstanz judged 35 pictures of student candidates unknown to them.
- Average correct judgement: 60%.

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Back to the first question

Are people capable to estimate the ideological orientation based on facial traits?

- 164 students of the University of Konstanz judged 35 pictures of student candidates unknown to them.
- Average correct judgement: 60%.
- But: A significant part of the successful judgements is due to the attractive halo effect.

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary

Back to the first question

Are people capable to estimate the ideological orientation based on facial traits?

- 164 students of the University of Konstanz judged 35 pictures of student candidates unknown to them.
- Average correct judgement: 60%.
- But: A significant part of the successful judgements is due to the attractive halo effect.
- For more details: Michael Herrmann & Susumu Shikano (2016) Attractiveness and facial competence bias face-based inferences of candidate ideology. *Political Psychology*. 37(3): pp.401-17.

<http://onlinelibrary.wiley.com/doi/10.1111/pops.12256/abstract>

Basics

Susumu Shikano

Bayesian
Inference?

If you are frequentist...

If you are Bayesian...

Conjugate analysis

Bayes Theorem
(revisited)

Summary