

Introduction to Multiple Linear Regression

Author: Nicholas G Reich

This material is part of the **statsTeachR** project

Derivative of OpenIntro slides, released under a CC BY-NC-SA license.

Outline

Introduction to multiple regression

Many variables in a model

Adjusted R^2

Multiple regression

- ▶ Simple linear regression: Bivariate - two variables: y and x
- ▶ Multiple linear regression: Multiple variables: y and x_1, x_2, \dots

Weights of books


	weight (g)	volume (cm ³)	cover
1	800	885	hc
2	950	1016	hc
3	1050	1125	hc
4	350	239	hc
5	750	701	hc
6	600	641	hc
7	1075	1228	hc
8	250	412	pb
9	700	953	pb
10	650	929	pb
11	975	1492	pb
12	350	419	pb
13	950	1010	pb
14	425	595	pb
15	725	1034	pb



From: Maindonald, J.H. and Braun, W.J. (2nd ed., 2007) "Data Analysis and Graphics Using R"

Weights of books (cont.)


The scatterplot shows the relationship between weights and volumes of books as well as the regression output. Which of the below is correct?



- (a) Weights of 80% of the books can be predicted accurately using this model.
- (b) We would expect a book that is 10 cm³ bigger than another expected to weigh 7 g more.
- (c) The correlation between weight and volume is $R = 0.80^2 = 0.64$.
- (d) The model underestimates the weight of the book with the highest volume.

Weights of books (cont.)

The scatterplot shows the relationship between weights and volumes of books as well as the regression output. Which of the below is correct?



- (a) Weights of 80% of the books can be predicted accurately using this model.
- (b) *We would expect a book that is 10 cm³ bigger than another expected to weigh 7 g more.*
- (c) The correlation between weight and volume is $R = 0.80^2 = 0.64$.
- (d) The model underestimates the weight of the book with the highest volume.

Modeling weights of books using volume

somewhat abbreviated output...

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	107.67931	88.37758	1.218	0.245
volume	0.70864	0.09746	7.271	6.26e-06


Residual standard error: 123.9 on 13 degrees of freedom

Multiple R-squared: 0.8026, Adjusted R-squared: 0.7875

F-statistic: 52.87 on 1 and 13 DF, p-value: 6.262e-06

Weights of hardcover and paperback books


Can you identify a trend in the relationship between volume and weight of hardcover and paperback books?



Weights of hardcover and paperback books

Can you identify a trend in the relationship between volume and weight of hardcover and paperback books?

Paperbacks generally weigh less than hardcover books after controlling for the book's volume.



Modeling weights of books using volume and cover type

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	197.96284	59.19274	3.344	0.005841	**
volume	0.71795	0.06153	11.669	6.6e-08	***
cover:pb	-184.04727	40.49420	-4.545	0.000672	***

Residual standard error: 78.2 on 12 degrees of freedom

Multiple R-squared: 0.9275, Adjusted R-squared: 0.9154

F-statistic: 76.73 on 2 and 12 DF, p-value: 1.455e-07

Visualising the linear model



Determining the reference level

Based on the regression output below, which level of cover is the reference level? Note that pb: paperback.

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.9628	59.1927	3.34	0.0058
volume	0.7180	0.0615	11.67	0.0000
cover:pb	-184.0473	40.4942	-4.55	0.0007

(a) paperback

(b) hardcover

Determining the reference level

Based on the regression output below, which level of cover is the reference level? Note that pb: paperback.

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.9628	59.1927	3.34	0.0058
volume	0.7180	0.0615	11.67	0.0000
cover:pb	-184.0473	40.4942	-4.55	0.0007

(a) paperback

(b) *hardcover*

Determining the reference level

Which of the below correctly describes the roles of variables in this regression model?

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.9628	59.1927	3.34	0.0058
volume	0.7180	0.0615	11.67	0.0000
cover:pb	-184.0473	40.4942	-4.55	0.0007

- (a) **response**: weight; **explanatory**: volume, paperback cover
- (b) **response**: weight; **explanatory**: volume, hardcover cover
- (c) **response**: volume; **explanatory**: weight, cover type
- (d) **response**: weight; **explanatory**: volume, cover type

Determining the reference level

Which of the below correctly describes the roles of variables in this regression model?

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.9628	59.1927	3.34	0.0058
volume	0.7180	0.0615	11.67	0.0000
cover:pb	-184.0473	40.4942	-4.55	0.0007

- (a) **response:** weight; **explanatory:** volume, paperback cover
- (b) **response:** weight; **explanatory:** volume, hardcover cover
- (c) **response:** volume; **explanatory:** weight, cover type
- (d) **response:** *weight*; **explanatory:** *volume, cover type*

Linear model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

Linear model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover : pb}$$

Linear model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover : pb}$$

1. For *hardcover* books: plug in 0 for cover

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \times 0$$

Linear model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover : pb}$$

1. For *hardcover* books: plug in 0 for cover

$$\begin{aligned}\widehat{weight} &= 197.96 + 0.72 \text{ volume} - 184.05 \times 0 \\ &= 197.96 + 0.72 \text{ volume}\end{aligned}$$

Linear model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover : pb}$$

1. For *hardcover* books: plug in *0* for cover

$$\begin{aligned}\widehat{weight} &= 197.96 + 0.72 \text{ volume} - 184.05 \times 0 \\ &= 197.96 + 0.72 \text{ volume}\end{aligned}$$

2. For *paperback* books: plug in *1* for cover

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \times 1$$

Linear model

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover : pb}$$


1. For *hardcover* books: plug in *0* for cover

$$\begin{aligned}\widehat{weight} &= 197.96 + 0.72 \text{ volume} - 184.05 \times 0 \\ &= 197.96 + 0.72 \text{ volume}\end{aligned}$$

2. For *paperback* books: plug in *1* for cover

$$\begin{aligned}\widehat{weight} &= 197.96 + 0.72 \text{ volume} - 184.05 \times 1 \\ &= 13.91 + 0.72 \text{ volume}\end{aligned}$$

Visualising the linear model



Interpretation of the regression coefficients

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

Interpretation of the regression coefficients

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

- *Slope of volume:* All else held constant, books that are 1 more cubic centimeter in volume tend to weigh about 0.72 grams more.

Interpretation of the regression coefficients

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

- ▶ *Slope of volume:* All else held constant, books that are 1 more cubic centimeter in volume tend to weigh about 0.72 grams more.
- ▶ *Slope of cover:* All else held constant, the model predicts that paperback books weigh 184 grams lower than hardcover books.

Interpretation of the regression coefficients

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

- ▶ *Slope of volume:* All else held constant, books that are 1 more cubic centimeter in volume tend to weigh about 0.72 grams more.
- ▶ *Slope of cover:* All else held constant, the model predicts that paperback books weigh 184 grams lower than hardcover books.
- ▶ *Intercept:* Hardcover books with no volume are expected on average to weigh 198 grams.

Interpretation of the regression coefficients

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

- ▶ *Slope of volume:* All else held constant, books that are 1 more cubic centimeter in volume tend to weigh about 0.72 grams more.
- ▶ *Slope of cover:* All else held constant, the model predicts that paperback books weigh 184 grams lower than hardcover books.
- ▶ *Intercept:* Hardcover books with no volume are expected on average to weigh 198 grams.
 - ▶ Obviously, the intercept does not make sense in context. It only serves to adjust the height of the line.

Prediction

Which of the following is the correct calculation for the predicted weight of a paperback book that is 600 cm³?

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

- (a) $197.96 + 0.72 * 600 - 184.05 * 1$
- (b) $184.05 + 0.72 * 600 - 197.96 * 1$
- (c) $197.96 + 0.72 * 600 - 184.05 * 0$
- (d) $197.96 + 0.72 * 1 - 184.05 * 600$

Prediction

Which of the following is the correct calculation for the predicted weight of a paperback book that is 600 cm³?

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

- (a) $197.96 + 0.72 * 600 - 184.05 * 1 = 445.91 \text{ grams}$
- (b) $184.05 + 0.72 * 600 - 197.96 * 1$
- (c) $197.96 + 0.72 * 600 - 184.05 * 0$
- (d) $197.96 + 0.72 * 1 - 184.05 * 600$

Another example: Modeling kid's test scores


Predicting cognitive test scores of 434 three- and four-year-old children using characteristics of their mothers. Data are from a survey of adult American women and their children - a subsample from the National Longitudinal Survey of Youth.

```
library(rstanarm)
data("kidiq")
head(kidiq)
```

##	kid_score	mom_hs	mom_iq	mom_age
## 1	65	1	121.11753	27
## 2	98	1	89.36188	25
## 3	85	1	115.44316	27
## 4	83	1	99.44964	25
## 5	115	1	92.74571	27
## 6	98	0	107.90184	18

Exploratory analysis

```
library(GGally)
kidiq$mom_hs <- factor(kidiq$mom_hs, levels=c(0,1), labels=c("no", "yes"))
ggpairs(kidiq)
```



What is a reasonable model

In generic model syntax

In regression formula syntax

Interpreting the slope

What is the correct interpretation of the slope for mom's IQ?

```
fm <- lm(kid_score ~ mom_hs + mom_iq + mom_age, data=kidiq)
round(summary(fm)$coef, 3)
```

##	Estimate	Std. Error	t value	Pr(> t)
## (Intercept)	20.985	9.130	2.298	0.022
## mom_hsys	5.647	2.258	2.501	0.013
## mom_iq	0.563	0.061	9.276	0.000
## mom_age	0.225	0.331	0.680	0.497

, kids with mothers whose IQs are one point higher tend to score on average 0.56 points higher.

Interpreting the slope

What is the correct interpretation of the intercept?

```
round(summary(fm)$coef, 3)
```

##	Estimate	Std. Error	t value	Pr(> t)
## (Intercept)	20.985	9.130	2.298	0.022
## mom_hsyas	5.647	2.258	2.501	0.013
## mom_iq	0.563	0.061	9.276	0.000
## mom_age	0.225	0.331	0.680	0.497

Interpreting the slope

What is the correct interpretation of the intercept?

```
round(summary(fm)$coef, 3)
```

##	Estimate	Std. Error	t value	Pr(> t)
## (Intercept)	20.985	9.130	2.298	0.022
## mom_hsyas	5.647	2.258	2.501	0.013
## mom_iq	0.563	0.061	9.276	0.000
## mom_age	0.225	0.331	0.680	0.497

Kids whose moms haven't gone to HS, whose moms have an IQ of 0, and who are 0 yrs old are expected on average to score 20.98. Obviously, the intercept does not make any sense in context.

Interpreting the slope

What is the correct interpretation of the slope for mom_hs?

```
round(summary(fm)$coef, 3)
```

##	Estimate	Std. Error	t value	Pr(> t)
## (Intercept)	20.985	9.130	2.298	0.022
## mom_hsyas	5.647	2.258	2.501	0.013
## mom_iq	0.563	0.061	9.276	0.000
## mom_age	0.225	0.331	0.680	0.497

All else being equal, kids whose moms graduated from high school are estimated to score than those whose moms did not work.


Modeling poverty

Description: Data for 3083 counties in the United States, including variables for demographic, financial, education, and other characteristics.

Source: Census website.


- ▶ FIPS: FIPS code.
- ▶ poverty: Percent below poverty level (2006-2010).
- ▶ pop2010: 2010 county population.
- ▶ female_house: Percent of population that lives in a female-owned house (2010).
- ▶ metro_res: Percent of population living in metropolitan area.
- ▶ hs_grad: Percent of population that is a high school graduate (2006-2010).
- ▶ ...

Modeling poverty



Predicting poverty using % female householder

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.31	1.90	1.74	0.09
female_house	0.69	0.16	4.32	0.00



$$R = 0.53$$

$$R^2 = 0.53^2 = 0.28$$

Another look at R^2

R^2 can be calculated in three ways:

Another look at R^2

R^2 can be calculated in three ways:

1. square the correlation coefficient of x and y (how we have been calculating it)

Another look at R^2

R^2 can be calculated in three ways:

1. square the correlation coefficient of x and y (how we have been calculating it)
2. square the correlation coefficient of y and \hat{y}

Another look at R^2

R^2 can be calculated in three ways:

1. square the correlation coefficient of x and y (how we have been calculating it)
2. square the correlation coefficient of y and \hat{y}
3. based on definition:

$$R^2 = \frac{\text{explained variability in } y}{\text{total variability in } y}$$

Another look at R^2

R^2 can be calculated in three ways:

1. square the correlation coefficient of x and y (how we have been calculating it)
2. square the correlation coefficient of y and \hat{y}
3. based on definition:

$$R^2 = \frac{\text{explained variability in } y}{\text{total variability in } y}$$

Using **ANOVA** we can calculate the explained variability and total variability in y .

Sum of squares

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.68	0.00
Residuals	49	347.68	7.10		
Total	50	480.25			

Sum of squares

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.68	0.00
Residuals	49	347.68	7.10		
Total	50	480.25			

Sum of squares of y : $SS_{Total} = \sum (y - \bar{y})^2 = 480.25 \rightarrow \text{total variability}$

Sum of squares

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.68	0.00
Residuals	49	347.68	7.10		
Total	50	480.25			

Sum of squares of y : $SS_{Total} = \sum (y - \bar{y})^2 = 480.25 \rightarrow \text{total variability}$

Sum of squares of residuals: $SS_{Error} = \sum e_i^2 = 347.68 \rightarrow \text{unexplained variability}$

Sum of squares

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.68	0.00
Residuals	49	347.68	7.10		
Total	50	480.25			

Sum of squares of y : $SS_{Total} = \sum (y - \bar{y})^2 = 480.25 \rightarrow \text{total variability}$

Sum of squares of residuals: $SS_{Error} = \sum e_i^2 = 347.68 \rightarrow \text{unexplained variability}$

Sum of squares of x : $SS_{Model} = SS_{Total} - SS_{Error} \rightarrow \text{explained variability}$
 $= 480.25 - 347.68 = 132.57$

Sum of squares

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.68	0.00
Residuals	49	347.68	7.10		
Total	50	480.25			

Sum of squares of y : $SS_{Total} = \sum (y - \bar{y})^2 = 480.25 \rightarrow \text{total variability}$

Sum of squares of residuals: $SS_{Error} = \sum e_i^2 = 347.68 \rightarrow \text{unexplained variability}$

Sum of squares of x : $SS_{Model} = SS_{Total} - SS_{Error} \rightarrow \text{explained variability}$
 $= 480.25 - 347.68 = 132.57$

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{132.57}{480.25} = 0.28 \checkmark$$

Why bother?

Why bother with another approach for calculating R^2 when we had a perfectly good way to calculate it as the correlation coefficient squared?

Why bother?

Why bother with another approach for calculating R^2 when we had a perfectly good way to calculate it as the correlation coefficient squared?

- ▶ *For single-predictor linear regression, having three ways to calculate the same value may seem like overkill.*
- ▶ *However, in multiple linear regression, we can't calculate R^2 as the square of the correlation between x and y because we have multiple x s.*
- ▶ *And next we'll learn another measure of explained variability, **adjusted R^2** , that requires the use of the third approach, ratio of explained and unexplained variability.*

Predicting poverty using % female hh + % white

<i>Linear model:</i>	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-2.58	5.78	-0.45	0.66
female_house	0.89	0.24	3.67	0.00
white	0.04	0.04	1.08	0.29

<i>ANOVA:</i>	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.00
white	1	8.21	8.21	1.16	0.29
Residuals	48	339.47	7.07		
Total	50	480.25			


Predicting poverty using % female hh + % white

<i>Linear model:</i>	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-2.58	5.78	-0.45	0.66
female_house	0.89	0.24	3.67	0.00
white	0.04	0.04	1.08	0.29

<i>ANOVA:</i>	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.00
white	1	8.21	8.21	1.16	0.29
Residuals	48	339.47	7.07		
Total	50	480.25			

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{132.57 + 8.21}{480.25} = 0.29$$

Does adding the variable `white` to the model add valuable information that wasn't provided by `female_house`?



R^2 vs. adjusted R^2

	R^2	Adjusted R^2
Model 1 (Single-predictor)	0.28	0.26
Model 2 (Multiple)	0.29	0.26

R^2 vs. adjusted R^2

	R^2	Adjusted R^2
Model 1 (Single-predictor)	0.28	0.26
Model 2 (Multiple)	0.29	0.26

- ▶ When any variable is added to the model R^2 increases.

R^2 vs. adjusted R^2

	R^2	Adjusted R^2
Model 1 (Single-predictor)	0.28	0.26
Model 2 (Multiple)	0.29	0.26

- ▶ When any variable is added to the model R^2 increases.
- ▶ But if the added variable doesn't really provide any new information, or is completely unrelated, adjusted R^2 does not increase.

Adjusted R^2

Adjusted R^2

$$R_{adj}^2 = 1 - \left(\frac{SS_{Error}}{SS_{Total}} \times \frac{n - 1}{n - p - 1} \right)$$

where n is the number of cases and p is the number of predictors (explanatory variables) in the model.

- ▶ Because p is never negative, R_{adj}^2 will always be smaller than R^2 .
- ▶ R_{adj}^2 applies a penalty for the number of predictors included in the model.
- ▶ Therefore, we choose models with higher R_{adj}^2 over others.

Calculate adjusted R^2

ANOVA:	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.0001
white	1	8.21	8.21	1.16	0.2868
Residuals	48	339.47	7.07		
Total	50	480.25			

$$R_{adj}^2 = 1 - \left(\frac{SS_{Error}}{SS_{Total}} \times \frac{n - 1}{n - p - 1} \right)$$

Calculate adjusted R^2

ANOVA:	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.0001
white	1	8.21	8.21	1.16	0.2868
Residuals	48	339.47	7.07		
Total	50	480.25			

$$\begin{aligned} R_{adj}^2 &= 1 - \left(\frac{SS_{Error}}{SS_{Total}} \times \frac{n - 1}{n - p - 1} \right) \\ &= 1 - \left(\frac{339.47}{480.25} \times \frac{51 - 1}{51 - 2 - 1} \right) \end{aligned}$$

Calculate adjusted R^2

ANOVA:	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.0001
white	1	8.21	8.21	1.16	0.2868
Residuals	48	339.47	7.07		
Total	50	480.25			

$$\begin{aligned}R_{adj}^2 &= 1 - \left(\frac{SS_{Error}}{SS_{Total}} \times \frac{n - 1}{n - p - 1} \right) \\&= 1 - \left(\frac{339.47}{480.25} \times \frac{51 - 1}{51 - 2 - 1} \right) \\&= 1 - \left(\frac{339.47}{480.25} \times \frac{50}{48} \right)\end{aligned}$$

Calculate adjusted R^2

ANOVA:	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.0001
white	1	8.21	8.21	1.16	0.2868
Residuals	48	339.47	7.07		
Total	50	480.25			

$$\begin{aligned}R_{adj}^2 &= 1 - \left(\frac{SS_{Error}}{SS_{Total}} \times \frac{n - 1}{n - p - 1} \right) \\&= 1 - \left(\frac{339.47}{480.25} \times \frac{51 - 1}{51 - 2 - 1} \right) \\&= 1 - \left(\frac{339.47}{480.25} \times \frac{50}{48} \right) \\&= 1 - 0.74\end{aligned}$$

Calculate adjusted R^2

ANOVA:	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.0001
white	1	8.21	8.21	1.16	0.2868
Residuals	48	339.47	7.07		
Total	50	480.25			

$$\begin{aligned}R_{adj}^2 &= 1 - \left(\frac{SS_{Error}}{SS_{Total}} \times \frac{n - 1}{n - p - 1} \right) \\&= 1 - \left(\frac{339.47}{480.25} \times \frac{51 - 1}{51 - 2 - 1} \right) \\&= 1 - \left(\frac{339.47}{480.25} \times \frac{50}{48} \right) \\&= 1 - 0.74 \\&= 0.26\end{aligned}$$

On your own

Play around with some regression models and ANOVAS.

```
load(url("http://www.openintro.org/stat/data/cc.RData"))
dim(countyComplete)
colnames(countyComplete)
fm <- lm(poverty ~ white_not_hispanic + female, data=countyComplete)
anova(fm)
```

Note: the actual dataset has slightly different variable names than those in the slides. More details on the dataset can be found here <https://www.openintro.org/stat/data/?data=cc>.