# Introduction to simple linear regression

#### Nicholas Reich, UMass-Amherst Biostatistics

Derivative of OpenIntro slides, released under a CC BY-NC-SA license

### Outline

Line fitting, residuals, and correlation


Fitting a line by least squares regression

Types of outliers in linear regression

# Modeling numerical variables


In this unit we will learn to quantify the relationship between two numerical variables, as well as modeling numerical response variables using a numerical or categorical explanatory variable.

The *scatterplot* below shows the relationship between HS graduate rate in all 50 US states and DC and the % of residents who live below the poverty line (income below \$23,050 for a family of 4 in 2012).




Response variable?

The *scatterplot* below shows the relationship between HS graduate rate in all 50 US states and DC and the % of residents who live below the poverty line (income below \$23,050 for a family of 4 in 2012).




Response variable?
% in poverty

The *scatterplot* below shows the relationship between HS graduate rate in all 50 US states and DC and the % of residents who live below the poverty line (income below \$23,050 for a family of 4 in 2012).




Response variable?
% in poverty
Explanatory variable?

The *scatterplot* below shows the relationship between HS graduate rate in all 50 US states and DC and the % of residents who live below the poverty line (income below \$23,050 for a family of 4 in 2012).




Response variable?
% in poverty
Explanatory variable?
% HS grad

The *scatterplot* below shows the relationship between HS graduate rate in all 50 US states and DC and the % of residents who live below the poverty line (income below \$23,050 for a family of 4 in 2012).



Response variable?
% in poverty
Explanatory variable?
% HS grad
Relationship?

The *scatterplot* below shows the relationship between HS graduate rate in all 50 US states and DC and the % of residents who live below the poverty line (income below \$23,050 for a family of 4 in 2012).



Response variable?

% in poverty

Explanatory variable?

% HS grad

Relationship?

linear, negative, moderately strong

# Quantifying the relationship

Correlation describes the strength of the linear association between two variables.

## Quantifying the relationship


- Correlation describes the strength of the linear association between two variables.
- It takes values between -1 (perfect negative) and +1 (perfect positive).

# Quantifying the relationship

- Correlation describes the strength of the linear association between two variables.
- It takes values between -1 (perfect negative) and +1 (perfect positive).
- A value of 0 indicates no linear association.




Which of the following is the best guess for the correlation between % in poverty and % HS grad?




Which of the following is the best guess for the correlation between % in poverty and % HS grad?

- (a) 0.6
- (b) -0.75
- (c) -0.1
- (d) 0.02
- (e) -1.5




Which of the following is the best guess for the correlation between % in poverty and <del>% HS grad?</del>

- (a) 0.1
- (b) -0.6
- (c) -0.4
- (d) 0.9
  - (e) 0.5




% female householder, no husband present

Which of the following is the best guess for the correlation between % in poverty and % HS grad?




## Assessing the correlation

Which of the following is has the strongest correlation, i.e. correlation coefficient closest to +1 or -1?



# Assessing the correlation

Which of the following is has the strongest correlation, i.e. correlation coefficient closest to +1 or -1?



(b) →
correlation
means <u>linear</u>
association

### Outline

Line fitting, residuals, and correlation

#### Fitting a line by least squares regression

Eyeballing the line

Residuals

Best line

The least squares line

Recap: Interpreting the slope and the intercept

Prediction & extrapolation

Conditions for the least squares line


 $R^2$ 

Categorical explanatory variables

Types of outliers in linear regression

## Eyeballing the line


Which of the following appears to be the line that best fits the linear relationship between % in poverty and % HS grad? Choose one.



## Eyeballing the line


Which of the following appears to be the line that best fits the linear relationship between % in poverty and % HS grad? Choose one.





### Residuals

Residuals are the leftovers from the model fit: Data = Fit + Residual




# Residuals (cont.)

#### Residual

Residual is the difference between the observed  $(y_i)$  and predicted  $\hat{y}_i$ .

$$e_i = y_i - \hat{y}_i$$




# Residuals (cont.)

#### Residual

Residual is the difference between the observed  $(y_i)$  and predicted  $\hat{y}_i$ .

$$e_i = y_i - \hat{y}_i$$




% living in poverty in DC is 5.44% more than predicted.

# Residuals (cont.)

#### Residual

Residual is the difference between the observed  $(y_i)$  and predicted  $\hat{y}_i$ .

$$e_i = y_i - \hat{y}_i$$



- % living in poverty in DC is 5.44% more than predicted.
- % living in poverty in RI is 4.16% less than predicted.

▶ We want a line that has small residuals:

- We want a line that has small residuals:
  - 1. Option 1: Minimize the sum of magnitudes (absolute values) of residuals

$$|e_1| + |e_2| + \cdots + |e_n|$$

- We want a line that has small residuals:
  - Option 1: Minimize the sum of magnitudes (absolute values) of residuals

$$|e_1| + |e_2| + \cdots + |e_n|$$

2. Option 2: Minimize the sum of squared residuals – *least squares* 

$$e_1^2 + e_2^2 + \dots + e_n^2$$

- We want a line that has small residuals:
  - Option 1: Minimize the sum of magnitudes (absolute values) of residuals

$$|e_1| + |e_2| + \cdots + |e_n|$$


2. Option 2: Minimize the sum of squared residuals – *least squares* 

$$e_1^2 + e_2^2 + \dots + e_n^2$$


- Why least squares?
  - Most commonly used
  - 2. Easier to compute by hand and using software
  - In many applications, a residual twice as large as another is usually more than twice as bad
  - 4. Squared error estimation has good statistical properties.

## Geometric interpretation

Minimizing the  $RSS(\beta)$  is equivalent to minimizing the Euclidean distance between y and all possible linear combinations of X.



## The least squares line



#### **Notation:**

Intercept:

Parameter:  $\beta_0$ 


Point estimate: b<sub>0</sub>

Slope:

Parameter: β<sub>1</sub>

Point estimate: b<sub>1</sub>

# Given...



	% HS grad	% in poverty
	(x)	(y)
mean	$\bar{x} = 86.01$	$\bar{y} = 11.35$
sd	$s_x = 3.73$	$s_y = 3.1$
	correlation	R = -0.75

# Slope

Slope

The slope of the regression can be calculated as

$$b_1 = \frac{s_y}{s_x} R$$

# Slope

### Slope

The slope of the regression can be calculated as

$$b_{1} = \frac{s_{y}}{s_{x}}R = \underbrace{\frac{c_{0} \vee (X_{3}Y)}{S_{x}^{2}}}$$

$$b_{1} = \underbrace{\frac{3.1}{3.73}} \times -0.75 = -0.62$$

In context...

# Slope

### Slope

The slope of the regression can be calculated as

$$b_1 = \frac{s_y}{s_x} R$$

In context...

$$b_1 = \frac{3.1}{3.73} \times -0.75 = -0.62$$

### Interpretation

For each additional % point in HS graduate rate, we would expect the % living in poverty to be lower on average by 0.62% points.

### Intercept

### Intercept

The intercept is where the regression line intersects the *y*-axis. The calculation of the intercept uses the fact the a regression line always passes through  $(\bar{x}, \bar{y})$ .


$$b_0 = \bar{y} - b_1 \bar{x}$$

## Intercept

### Intercept

The intercept is where the regression line intersects the *y*-axis. The calculation of the intercept uses the fact the a regression line always passes through  $(\bar{x}, \bar{y})$ .

$$b_0 = \bar{y} - b_1 \bar{x}$$




### Intercept

#### Intercept

The intercept is where the regression line intersects the y-axis. The calculation of the intercept uses the fact the a regression line always passes through  $(\bar{x}, \bar{y})$ .

$$b_0 = \bar{y} - b_1 \bar{x}$$



$$b_0 = 11.35 - (-0.62) \times 86.01$$
  
= 64.68

### Which of the following is the correct interpretation of the intercept?


- (a) For each % point increase in HS graduate rate, % living in poverty is expected to increase on average by 64.68%.
- (b) For each % point decrease in HS graduate rate, % living in poverty is expected to increase on average by 64.68%.
- Having no HS graduates leads to 64.68% of residents living below the poverty line.
- (d) States with no HS graduates are expected on average to have 64.68% of residents living below the poverty line.
- (e) In states with no HS graduates % living in poverty is expected to increase on average by 64.68%.

### Which of the following is the correct interpretation of the intercept?

- (a) For each % point increase in HS graduate rate, % living in poverty is expected to increase on average by 64.68%.
- (b) For each % point decrease in HS graduate rate, % living in poverty is expected to increase on average by 64.68%.
- (c) Having no HS graduates leads to 64.68% of residents living below the poverty line.
- (d) States with no HS graduates are expected on average to have 64.68% of residents living below the poverty line.
- (e) In states with no HS graduates % living in poverty is expected to increase on average by 64.68%.


## More on the intercept

Since there are no states in the dataset with no HS graduates, the intercept is of no interest, not very useful, and also not reliable since the predicted value of the intercept is so far from the bulk of the data.




## Regression line

% 
$$\widehat{in\ poverty} = 64.68 - 0.62$$
 % HS grad



## Interpretation of slope and intercept


- Intercept: When x = 0, y is expected to equal the intercept.
- Slope: For each unit in x, y is expected to increase / decrease on average by the slope.



Note: These statements are not causal, unless the study is a randomized controlled experiment.


#### Prediction

- ▶ Using the linear model to predict the value of the response variable for a given value of the explanatory variable is called prediction, simply by plugging in the value of x in the linear model equation.
- There will be some uncertainty associated with the predicted value.




## Extrapolation

- Applying a model estimate to values outside of the realm of the original data is called extrapolation.
- Sometimes the intercept might be an extrapolation.



# Examples of extrapolation



## Examples of extrapolation



Watch One-Minute World News



Last Updated: Thursday, 30 September, 2004, 04:04 GMT 05:04 UK

■ E-mail this to a friend 

→ Printable version

#### Women 'may outsprint men by 2156'

Women sprinters may be outrunning men in the 2156 Olympics if they continue to close the gap at the rate they are doing, according to scientists.

UK An Oxford University study
England found that women are running faster than they have ever done over 100m.



sprinters

At their current rate of improvement, they should overtake men within 150 years, said Dr Andrew Tatem.

The study, comparing winning times for the Olympic 100m since 1900, is published in the journal Nature.

However, former British Olympic sprinter Derek Redmond told the BBC: "I find it difficult to believe.

"I can see the gap closing between men and women but I can't necessarily see it being overtaken because mens' times are also going to improve."

## Examples of extrapolation

# Momentous sprint at the 2156 Olympics?

Women sprinters are closing the gap on men and may one day overtake them.




Figure 1 The winning Olympic 100-metre sprint times for men (blue points) and women (red points), with superimposed best-fit linear regression lines (solid black lines) and coefficients of determination. The regression lines are extrapolated (broken blue and red lines for men and women, respectively) and 95% confidence intervals (dotted black lines) based on the available points are superimposed. The projections intervals to the transfer the 2156 Olympics, when the winning women's 100-metre sprint time of 8,079 s will be faster than the men's at 0.098 s.

# Conditions for the least squares line

1. Linearity

## Conditions for the least squares line

- 1. Linearity
- 2. Nearly normal residuals

# Conditions for the least squares line

- Linearity
- 2. Nearly normal residuals
- 3. Constant variability

4 Indepence of observations

Lytime series

Ly clustered observations

cause harder to see u/plots

## Conditions: (1) Linearity


► The relationship between the explanatory and the response variable should be linear.

## Conditions: (1) Linearity


- ► The relationship between the explanatory and the response variable should be linear.
- Methods for fitting a model to non-linear relationships exist, but are beyond the scope of this class. If this topic is of interest, an Online Extra is available on openintro.org covering new techniques.

## Conditions: (1) Linearity

- ► The relationship between the explanatory and the response variable should be linear.
- Methods for fitting a model to non-linear relationships exist, but are beyond the scope of this class. If this topic is of interest, an Online Extra is available on openintro.org covering new techniques.
- Check using a scatterplot of the data, or a residuals plot.




## Anatomy of a residuals plot



#### ▲ RI:

% HS grad = 81 % in poverty = 10.3  
% in poverty = 
$$64.68 - 0.62 * 81 = 14.46$$
  
 $e = \%$  in poverty - % in poverty  
=  $10.3 - 14.46 = -4.16$ 

## Anatomy of a residuals plot



#### **▲** RI:

% HS grad = 81 % in poverty = 10.3  
% in poverty = 
$$64.68 - 0.62 * 81 = 14.46$$
  
 $e = \%$  in poverty -  $\%$  in poverty  
=  $10.3 - 14.46 = -4.16$ 

DC:

% HS grad = 86 % in poverty = 16.8  
% in poverty = 
$$64.68 - 0.62 * 86 = 11.36$$
  
 $e = \%$  in poverty -  $\%$  in poverty  
=  $16.8 - 11.36 = 5.44$ 

## Conditions: (2) Nearly normal residuals


The residuals should be nearly normal.


## Conditions: (2) Nearly normal residuals

- The residuals should be nearly normal.
- This condition may not be satisfied when there are unusual observations that don't follow the trend of the rest of the data.


## Conditions: (2) Nearly normal residuals

- The residuals should be nearly normal.
- This condition may not be satisfied when there are unusual observations that don't follow the trend of the rest of the data.
- Check using a histogram or normal probability plot of residuals.







The variability of points around the least squares line should be roughly constant.



- The variability of points around the least squares line should be roughly constant.
- This implies that the variability of residuals around the 0 line should be roughly constant as well.





- The variability of points around the least squares line should be roughly constant.
- This implies that the variability of residuals around the 0 line should be roughly constant as well.
- Also called homoscedasticity.




- The variability of points around the least squares line should be roughly constant.
- This implies that the variability of residuals around the 0 line should be roughly constant as well.
- Also called homoscedasticity.
- Check using a histogram or normal probability plot of residuals.


- (a) Constant variability
- (b) Linear relationship
- (c) Normal residuals
- (d) No extreme outliers







- (a) Constant variability
- (b) Linear relationship
- (c) Normal residuals
- (d) No extreme outliers







- (a) Constant variability
- (b) Linear relationship
- (c) Normal residuals
- (d) No extreme outliers





- (a) Constant variability
- (b) Linear relationship
- (c) Normal residuals
- (d) No extreme outliers





#### $R^2$

► The strength of the fit of a linear model is most commonly evaluated using  $\mathbb{R}^2$ .

#### $R^2$

- ► The strength of the fit of a linear model is most commonly evaluated using  $\mathbb{R}^2$ .
- $R^2$  is calculated as the square of the correlation coefficient.

- ► The strength of the fit of a linear model is most commonly evaluated using  $R^2$ .
- $ightharpoonup R^2$  is calculated as the square of the correlation coefficient.
- It tells us what percent of variability in the response variable is explained by the model.

- The strength of the fit of a linear model is most commonly evaluated using R<sup>2</sup>.
- $ightharpoonup R^2$  is calculated as the square of the correlation coefficient.
- ▶ It tells us what percent of variability in the response variable is explained by the model.
- The remainder of the variability is explained by variables not included in the model or by inherent randomness in the data.


good Riveres by Freld.

- ► The strength of the fit of a linear model is most commonly evaluated using  $R^2$ .
- $ightharpoonup R^2$  is calculated as the square of the correlation coefficient.
- ▶ It tells us what percent of variability in the response variable is explained by the model.
- The remainder of the variability is explained by variables not included in the model or by inherent randomness in the data.
- For the model we've been working with,  $R^2 = -0.62^2 = 0.38$ .

# Interpretation of $R^2$

# Which of the below is the correct interpretation of R = -0.62, $R^2 = 0.38$ ?


- (a) 38% of the variability in the % of HG graduates among the 51 states is explained by the model.
- (b) 38% of the variability in the % of residents living in poverty among the 51 states is explained by the model.
- (c) 38% of the time % HS graduates predict % living in poverty correctly.
- (d) 62% of the variability in the % of residents living in poverty among the 51 states is explained by the model.



# Interpretation of $R^2$

#### Which of the below is the correct interpretation of R = -0.62, $R^2 = 0.38$ ?

- (a) 38% of the variability in the % of HG graduates among the 51 states is explained by the model.
- (b) 38% of the variability in the % of residents living in poverty among the 51 states is explained by the model.
- (c) 38% of the time % HS graduates predict % living in poverty correctly.
- (d) 62% of the variability in the % of residents living in poverty among the 51 states is explained by the model.



$$\widehat{poverty} = 11.17 + 0.38 \times west$$

- Explanatory variable: region, reference level: east
- Intercept: The estimated average poverty percentage in eastern states is 11.17%

$$\widehat{poverty} = 11.17 + 0.38 \times west$$

- Explanatory variable: region, reference level: east
- ► Intercept: The estimated average poverty percentage in eastern states is 11.17%
  - ► This is the value we get if we plug in 0 for the explanatory variable

$$\widehat{poverty} = 11.17 + 0.38 \times west$$

- Explanatory variable: region, reference level: east
- Intercept: The estimated average poverty percentage in eastern states is 11.17%
  - ► This is the value we get if we plug in 0 for the explanatory variable
- ► *Slope:* The estimated average poverty percentage in western states is 0.38% higher than eastern states.

$$\widehat{poverty} = 11.17 + 0.38 \times west$$

- Explanatory variable: region, reference level: east
- Intercept: The estimated average poverty percentage in eastern states is 11.17%
  - ► This is the value we get if we plug in 0 for the explanatory variable
- ► *Slope:* The estimated average poverty percentage in western states is 0.38% higher than eastern states.
  - Then, the estimated average poverty percentage in western states is 11.17 + 0.38 = 11.55%.

$$\widehat{poverty} = 11.17 + 0.38 \times west$$

- Explanatory variable: region, reference level: east
- Intercept: The estimated average poverty percentage in eastern states is 11.17%
  - ► This is the value we get if we plug in 0 for the explanatory variable
- ► *Slope:* The estimated average poverty percentage in western states is 0.38% higher than eastern states.
  - Then, the estimated average poverty percentage in western states is 11.17 + 0.38 = 11.55%.
  - ► This is the value we get if we plug in 1 for the explanatory variable

Which region (northeast, midwest, west, or south) is the reference level?

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.50	0.87	10.94	0.00
region4midwest	0.03	1.15	0.02	0.98
region4west	1.79	1.13	1.59	0.12
region4south	4.16	1.07	3.87	0.00

- (a) northeast
- (b) midwest
- (c) west
- (d) south
- (e) cannot tell

Which region (northeast, midwest, west, or south) is the reference level?

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.50	0.87	10.94	0.00
region4midwest	0.03	1.15	0.02	0.98
region4west	1.79	1.13	1.59	0.12
region4south	4.16	1.07	3.87	0.00

- (a) northeast
- (b) midwest
- (c) west
- (d) south
- (e) cannot tell

Which region (northeast, midwest, west, or south) has the lowest poverty percentage?

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.50	0.87	10.94	0.00
region4midwest	0.03	1.15	0.02	0.98
region4west	1.79	1.13	1.59	0.12
region4south	4.16	1.07	3.87	0.00

- (a) northeast
- (b) midwest
- (c) west
- (d) south
- (e) cannot tell

Which region (northeast, midwest, west, or south) has the lowest poverty percentage?

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.50	0.87	10.94	0.00
region4midwest	0.03	1.15	0.02	0.98
region4west	1.79	1.13	1.59	0.12
region4south	4.16	1.07	3.87	0.00

- (a) northeast
- (b) midwest
- (c) west
- (d) south
- (e) cannot tell

#### Outline


Line fitting, residuals, and correlation

Fitting a line by least squares regression


Types of outliers in linear regression

How do outliers influence the least squares line in this plot?

To answer this question think of where the regression line would be with and without the outlier(s). Without the outliers the regression line would be steeper, and lie closer to the larger group of observations. With the outliers the line is pulled up and away from some of the observations in the larger group.




How do outliers influence the least squares line in this plot?



How do outliers influence the least squares line in this plot?

Without the outlier there is no evident relationship between *x* and *y*.



Outliers are points that lie away from the cloud of points.


- Outliers are points that lie away from the cloud of points.
- Outliers that lie horizontally away from the center of the cloud are called *high leverage* points.

- Outliers are points that lie away from the cloud of points.
- Outliers that lie horizontally away from the center of the cloud are called *high leverage* points.
- High leverage points that actually influence the <u>slope</u> of the regression line are called *influential* points.

- Outliers are points that lie away from the cloud of points.
- Outliers that lie horizontally away from the center of the cloud are called *high leverage* points.
- ► High leverage points that actually influence the <u>slope</u> of the regression line are called *influential* points.
- In order to determine if a point is influential, visualize the regression line with and without the point. Does the slope of the line change considerably? If so, then the point is influential. If not, then itÕs not an influential point.

# Influential points


Data are available on the log of the surface temperature and the log of the light intensity of 47 stars in the star cluster CYG OB1.






# Which of the below best describes the outlier?

- (a) influential
- (b) high leverage
- (c) none of the above
- (d) there are no outliers




Which of the below best describes the outlier?

- (a) influential
- (b) high leverage
- (c) none of the above
- (d) there are no outliers




Does this outlier influence the slope of the regression line?



Does this outlier influence the slope of the regression line?

Not much...



#### Recap

#### Which of following is true?

- (a) Influential points always change the intercept of the regression line.
- (b) Influential points always reduce  $R^2$ .
- (c) It is much more likely for a low leverage point to be influential, than a high leverage point.
- (d) When the data set includes an influential point, the relationship between the explanatory variable and the response variable is always nonlinear.
- (e) None of the above.

#### Recap

#### Which of following is true?

- (a) Influential points always change the intercept of the regression line.
- (b) Influential points always reduce  $R^2$ .
- (c) It is much more likely for a low leverage point to be influential, than a high leverage point.
- (d) When the data set includes an influential point, the relationship between the explanatory variable and the response variable is always nonlinear.
- (e) None of the above.

# Recap (cont.)

