

# Warm-Up

Econ 350: The University of Chicago

This Draft: December 11, 2013

## Abstract

The objective of this exercise is for you to become familiar with the basic requirements to proceed with the rest of the empirical exercise. This lets you make sure that your software is installed and that the maximization package is working. Also, it teaches you how to do numerical integration, which is useful in the rest of the empirical project. Be sure to get this exercise correct so that you can be confident to proceed with the rest of the project.

## 1 How to Calculate a Numerical Integral

This exercise is based on [Langtangen \(2011\)](#). This is a useful source for the rest of the course. Try to solve the exercises first and then look at the book.

### 1.1 Interpolation

The information that a real-valued continuous function contains is actually infinite (in the not-countable sense). The computer, however, works in  $\aleph_0$ . This is not trivial, the computer can go ahead and count forever. How does a computer plots functions?

**Exercise 1.1** (*Plot a basic continuous function through a grid*) Create a grid of 1,000 points to plot one cycle in the unitary circle of the sine function. Hint: use Google.

Roughly, with 1,000 points, the computer: (i) evaluates the sine function, (ii) puts together coordinates, (iii) scatters them, (iv) joins them through lines. This method has a name: *linear interpolation*. This is how linear interpolation happens. A given point  $x$  lies in the interval  $[x_k, x_{k+1}]$ , for an arbitrary  $k$ . In that interval, the computer defines a function that passes through the points  $(x_k, s_k)$  and  $(x_{k+1}, s_{k+1})$ :

$$S_k(x) = s_k + \frac{s_{k+1} - s_k}{x_{k+1} - x_k}(x - x_k). \quad (1)$$

$S_k(x)$  is identical to  $\sin(x)$  at  $x_k$  and  $x_{k+1}$ ; between these points,  $S_k(x)$  is linear.

### 1.2 Trapezoidal Integration

Some functions have *primitive* form integrals and some do not. How to integrate the sine function on the interval  $(0, \pi)$  without a primitive form? For example,

$$\int_0^{\pi} \sin(x) dx. \quad (2)$$

From basic calculus it is possible to partition this integral in an arbitrary sum of integrals:

$$\int_0^{\pi} \sin(x) dx = \sum_{k=0}^{n-1} \int_{x_k}^{x_{k+1}} \sin(x) dx \quad (3)$$

where  $n$  is the number of partitions and  $x_n \equiv \pi$ . This is useful because it is more precise to approximate it “by pieces”. Precisely, the task is to compute the following:

$$\int_{x_k}^{x_{k+1}} \sin(x) dx \quad (4)$$

... but  $\sin(x) \approx S_k(x)$ . Then,

$$\int_{x_k}^{x_{k+1}} \sin(x) dx = \int_{x_k}^{x_{k+1}} S_k(x) dx. \quad (5)$$

Now, note that: (i) a definite integral is the surface under the curve in certain interval; (ii) the interpolation function is linear. Thus, the approximation of the integral in the interval of interest implies the calculation of the area of a trapezoid. In particular:

$$\int_{x_k}^{x_{k+1}} \sin(x) dx \approx \frac{1}{2} (s_{k+1} + s_k) (x_{k+1} - x_k). \quad (6)$$

It is almost always the case that the spacing between the points in the grid in which functions are worked out is uniform. Thus,  $(x_{k+1} - x_k) = h$  for some elected  $h$  and for  $k = 0, \dots, n-1$ . The integral becomes a sum:

$$\begin{aligned} \int_0^{\pi} \sin(x) dx &\approx \sum_{k=0}^{n-1} (s_{k+1} + s_k) \\ &= \frac{h}{2} \left[ s_0 + 2 \sum_{k=1}^{n-1} s_k + s_n \right] \end{aligned} \quad (7)$$

where you can show that the second equality holds by the principle of mathematical induction.

**Clue 1.2** Do you remember the name of the functions for which the sum on (8) converges to a finite value?

### 1.2.1 Generalization

It is possible to generalize this reasoning as follows. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function in the interval  $[a, b]$ . The “grid version” of  $f$  is given by the sets of lines that connect the sequence of coordinates  $(x_i, y_i)_{i=0}^n$  where  $x_i = a + ih$  and  $y_i = f(x_i)$  for  $i = 0, \dots, n$ ,  $n \geq 1$ ,  $h = (b - a)/n$ . Then,

**Definition 1.3** (Trapezoidal Integral)

$$\int_a^b f(x) dx \approx \frac{h}{2} \left[ y_0 + 2 \sum_{k=1}^{n-1} y_k + y_n \right]. \quad (8)$$

**Exercise 1.4** (*Trapezoidal Integration*) Write down a code that has a general function for trapezoidal integration.

**Exercise 1.5** (*Trapezoidal Integration of the Sine Function*) Integrate (2) with  $n = 10, 50, 100, 500$  using your code of Exercise 1.4. How do these values differ from the values that you can get in your calculator. Hint: switch to radians.

### 1.3 Monte-Carlo Integration

Take the same framework as in Section 1.2.1. The first mean value theorem for integrals establishes that the integral of a function over the interval  $[a, b]$  is the mean value of  $f$  over the interval  $[a, b]$  multiplied by the length of the interval,  $b - a$ . Then, a simulated integral is calculated as follows: (i) take  $n$  draws from a uniform distribution over  $[a, b]$ ; (ii) evaluate  $f$  at each of them; (iii) take an average; (iv) multiply the average by the length of the interval. Put differently, if  $x_i$  is a typical random draw,

$$\int_a^b f(x)dx \approx (b - a) \frac{1}{n} \sum_{i=1}^n f(x_i). \quad (9)$$

This approach to numerical integrations is called Monte Carlo integration.

**Exercise 1.6** (*Monte Carlo Integration*) Write down a code that has a general function for Monte Carlo Integration.

**Exercise 1.7** (*Monte Carlo Integration of the Sine Function*) Integrate (2) with  $n = 10, 50, 100, 500$  using your code of Exercise 1.6. How do these values differ from the values that you can get in your calculator and in Exercise 1.4. Why?

## 2 How to Maximize a Likelihood

Consider a sample of  $N$  i.i.d. observations:

$$x_i \sim \mathcal{N}(0, \theta) \quad (10)$$

for  $i = 1, \dots, N$ .  $\theta$  is the variance and you want to estimate it.

**Exercise 2.1** (*Basics*) (i) Write down the individual likelihood, the sample likelihood, and the sample log-likelihood; (ii) write down the sample score; (iii) calculate the Maximum Likelihood Estimator (MLE) of  $\theta$  as a function the data.

**Exercise 2.2** (*Estimation I*) In Exercise 2.1 you find that the MLE estimator for  $\theta$  has a closed and easy functional form. Actually, you do not need numeric maximization to obtain it. Use the data set “warmup.csv” to estimate  $\theta$  by MLE. The file “readmewarmup.txt” describes this data. Do not use a maximization routine. Use sums and multiplications.

**Exercise 2.3** (*Estimation II*) Use the data set “warmup.csv” to estimate  $\theta$  by MLE. The file “readmewarmup.txt” describes this data. Use a the maximization routine discussed in the introductory session of Computational Econometrics.

**Exercise 2.4** Compare the estimate in Exercise 2.2 to the estimate in Exercise 2.3. Do they differ? Why? Which one is better?

## References

Langtangen, H. P. (2011). *A primer on scientific programming with Python*, Volume 6. Springer.