

Metric Spaces

Motivation

- ▶ In political science, our actors rank actions or outcomes so we need **ordered sets**.

Motivation

- ▶ In political science, our actors rank actions or outcomes so we need **ordered sets**.
- ▶ We also often think about distance in our theories.

Motivation

- ▶ In political science, our actors rank actions or outcomes so we need **ordered sets**.
- ▶ We also often think about distance in our theories.
- ▶ To think spatially in a rigorous manner, we need **metric spaces**

Distance

- ▶ We want a mathematical concept of distance that satisfies some intuitive properties.

Distance

- ▶ We want a mathematical concept of distance that satisfies some intuitive properties.
- ▶ Distance should be positive

Distance

- ▶ We want a mathematical concept of distance that satisfies some intuitive properties.
- ▶ Distance should be positive
- ▶ Distance from Princeton to New York should equal distance from New York to Princeton.

Distance

- ▶ We want a mathematical concept of distance that satisfies some intuitive properties.
- ▶ Distance should be positive
- ▶ Distance from Princeton to New York should equal distance from New York to Princeton.
- ▶ Shortest distance between two points should be the direct route.

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$
 2. $d(x, y) = 0$ if and only if $x = y$

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$
 2. $d(x, y) = 0$ if and only if $x = y$
 3. $d(x, y) = d(y, x)$ (symmetry)

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$
 2. $d(x, y) = 0$ if and only if $x = y$
 3. $d(x, y) = d(y, x)$ (symmetry)
 4. $d(x, y) \leq d(x, z) + d(z, y)$ (triangle inequality)

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$
 2. $d(x, y) = 0$ if and only if $x = y$
 3. $d(x, y) = d(y, x)$ (symmetry)
 4. $d(x, y) \leq d(x, z) + d(z, y)$ (triangle inequality)
- ▶ A **metric space** is a pair (X, d) where X is a set and d is a metric.

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$
 2. $d(x, y) = 0$ if and only if $x = y$
 3. $d(x, y) = d(y, x)$ (symmetry)
 4. $d(x, y) \leq d(x, z) + d(z, y)$ (triangle inequality)
- ▶ A **metric space** is a pair (X, d) where X is a set and d is a metric.
- ▶ Elements of a metric space are usually called **points**.

Metric Space: Definition

- ▶ A **metric** on a set X associates all $x, y \in X$ with a real number, $d(x, y)$ satisfying the following properties:
 1. $d(x, y) \geq 0$
 2. $d(x, y) = 0$ if and only if $x = y$
 3. $d(x, y) = d(y, x)$ (symmetry)
 4. $d(x, y) \leq d(x, z) + d(z, y)$ (triangle inequality)
- ▶ A **metric space** is a pair (X, d) where X is a set and d is a metric.
- ▶ Elements of a metric space are usually called **points**.
- ▶ The most familiar metric space is (\mathbb{R}, d) where $d(x, y) = |x - y|$.

Metric Space: Example

- ▶ Consider how we might define the distance between policy bundles represented as points in \mathbb{R}^2

Metric Space: Example

- ▶ Consider how we might define the distance between policy bundles represented as points in \mathbb{R}^2
- ▶ Manhattan metric:

$$d_1(x, y) = |x_1 - y_1| + |x_2 - y_2|$$

Metric Space: Example

- ▶ Consider how we might define the distance between policy bundles represented as points in \mathbb{R}^2
- ▶ Manhattan metric:

$$d_1(x, y) = |x_1 - y_1| + |x_2 - y_2|$$

- ▶ Euclidean metric:

$$d_2(x, y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

Metric Space: Example

- ▶ Consider how we might define the distance between policy bundles represented as points in \mathbb{R}^2
- ▶ Manhattan metric:

$$d_1(x, y) = |x_1 - y_1| + |x_2 - y_2|$$

- ▶ Euclidean metric:

$$d_2(x, y) = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$$

- ▶ Sup metric:

$$d_\infty(x, y) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

Metric Subspaces

- ▶ We will occasionally use objects called **metric subspaces**

Metric Subspaces

- ▶ We will occasionally use objects called **metric subspaces**
- ▶ Let (X, d) be a metric space and let Y be a nonempty subset of X .

Metric Subspaces

- ▶ We will occasionally use objects called **metric subspaces**
- ▶ Let (X, d) be a metric space and let Y be a nonempty subset of X .
- ▶ Y can be thought of as a metric space itself using the distance function induced by d on Y , $d|_{Y \times Y}$.

Metric Subspaces

- ▶ We will occasionally use objects called **metric subspaces**
- ▶ Let (X, d) be a metric space and let Y be a nonempty subset of X .
- ▶ Y can be thought of as a metric space itself using the distance function induced by d on Y , $d|_{Y \times Y}$.
- ▶ We then say that $(Y, d|_{Y \times Y})$ is a metric subspace of (X, d) .

Metric Subspaces

- ▶ We will occasionally use objects called **metric subspaces**
- ▶ Let (X, d) be a metric space and let Y be a nonempty subset of X .
- ▶ Y can be thought of as a metric space itself using the distance function induced by d on Y , $d|_{Y \times Y}$.
- ▶ We then say that $(Y, d|_{Y \times Y})$ is a metric subspace of (X, d) .
- ▶ $[0, 1]$ can be thought of as a metric subspace of \mathbb{R}^2 where the distance between $x, y \in [0, 1]$ is calculated by viewing x and y as points in \mathbb{R} : $d_1(x, y) = |x - y|$

Open Balls

- ▶ We want to have a concept of proximity when we use metric spaces.

Open Balls

- ▶ We want to have a concept of proximity when we use metric spaces.
- ▶ The set of points in close proximity to a given point x_0 is called a “ball” about x_0 .

Open Balls

- ▶ We want to have a concept of proximity when we use metric spaces.
- ▶ The set of points in close proximity to a given point x_0 is called a “ball” about x_0 .
- ▶ Formally, an **open ball** about x_0 with radius r is the set of points

$$B_r(x_0) = \{x \in X : d(x, x_0) < r\}$$


Open Balls

- ▶ We want to have a concept of proximity when we use metric spaces.
- ▶ The set of points in close proximity to a given point x_0 is called a “ball” about x_0 .
- ▶ Formally, an **open ball** about x_0 with radius r is the set of points


$$B_r(x_0) = \{x \in X : d(x, x_0) < r\}$$

- ▶ Open balls are not necessarily spherical: their shape depends on the metric.


Open Balls



$$d(x, y) < L$$



$$d_{\infty}(x, y) < L$$



$$d_1(x, y) < L$$

Neighborhood

- ▶ A set $N(x_0) \subseteq S \subseteq X$ is called a **neighborhood** of x_0 if $B_r(x_0) \subseteq N(x_0)$ for some $r > 0$.

Neighborhood

- ▶ A set $N(x_0) \subseteq S \subseteq X$ is called a **neighborhood** of x_0 if $B_r(x_0) \subseteq N(x_0)$ for some $r > 0$.
- ▶ A point $x_0 \in S$ is called an **interior point** of S if there exists a neighborhood $N(x_0) \subseteq S$.

Neighborhood

- ▶ A set $N(x_0) \subseteq S \subseteq X$ is called a **neighborhood** of x_0 if $B_r(x_0) \subseteq N(x_0)$ for some $r > 0$.
- ▶ A point $x_0 \in S$ is called an **interior point** of S if there exists a neighborhood $N(x_0) \subseteq S$.
- ▶ The set of interior points of a set S is called the **interior** of S , denoted $\text{int } S$.

Neighborhood

- ▶ A set $N(x_0) \subseteq S \subseteq X$ is called a **neighborhood** of x_0 if $B_r(x_0) \subseteq N(x_0)$ for some $r > 0$.
- ▶ A point $x_0 \in S$ is called an **interior point** of S if there exists a neighborhood $N(x_0) \subseteq S$.
- ▶ The set of interior points of a set S is called the **interior** of S , denoted $\text{int } S$.
- ▶ A point $x_0 \in X$ is called a **boundary point** of $S \subseteq X$ if every neighborhood of x_0 contains points of S and S^c .

Neighborhood

- ▶ A set $N(x_0) \subseteq S \subseteq X$ is called a **neighborhood** of x_0 if $B_r(x_0) \subseteq N(x_0)$ for some $r > 0$.
- ▶ A point $x_0 \in S$ is called an **interior point** of S if there exists a neighborhood $N(x_0) \subseteq S$.
- ▶ The set of interior points of a set S is called the **interior** of S , denoted $\text{int } S$.
- ▶ A point $x_0 \in X$ is called a **boundary point** of $S \subseteq X$ if every neighborhood of x_0 contains points of S and S^c .
- ▶ The **boundary** is the set of all boundary points of S , $b(S)$.

Neighborhood

- ▶ A set $N(x_0) \subseteq S \subseteq X$ is called a **neighborhood** of x_0 if $B_r(x_0) \subseteq N(x_0)$ for some $r > 0$.
- ▶ A point $x_0 \in S$ is called an **interior point** of S if there exists a neighborhood $N(x_0) \subseteq S$.
- ▶ The set of interior points of a set S is called the **interior** of S , denoted $\text{int } S$.
- ▶ A point $x_0 \in X$ is called a **boundary point** of $S \subseteq X$ if every neighborhood of x_0 contains points of S and S^c .
- ▶ The **boundary** is the set of all boundary points of S , $b(S)$.
- ▶ The **closure** \overline{S} of S is the union of S with its boundary, i.e. $\overline{S} = S \cup b(S)$.

Open and Closed Sets

- ▶ A set S is **open** if $S = \text{int } S$.

Open and Closed Sets

- ▶ A set S is **open** if $S = \text{int } S$.
- ▶ A set S is **closed** if $S = \overline{S}$

Sequences

- ▶ A **sequence** in a metric space X is a list of particular elements x^1, x^2, \dots of X denoted (x^n) .

Sequences

- ▶ A **sequence** in a metric space X is a list of particular elements x^1, x^2, \dots of X denoted (x^n) .
- ▶ We often encounter sequences in political science.

Sequences

- ▶ A **sequence** in a metric space X is a list of particular elements x^1, x^2, \dots of X denoted (x^n) .
- ▶ We often encounter sequences in political science.
- ▶ Time series data

Sequences

- ▶ A **sequence** in a metric space X is a list of particular elements x^1, x^2, \dots of X denoted (x^n) .
- ▶ We often encounter sequences in political science.
- ▶ Time series data
- ▶ Sequence of moves in a game

Sequences

- ▶ A **sequence** in a metric space X is a list of particular elements x^1, x^2, \dots of X denoted (x^n) .
- ▶ We often encounter sequences in political science.
- ▶ Time series data
- ▶ Sequence of moves in a game
- ▶ Asymptotic properties of an estimator identified by analyzing sequence of estimates of population parameter

Convergence

- ▶ (x^n) converges to x if for all $\epsilon > 0$, there exists some N such that for all $n \geq N$, $d(x^n, x) < \epsilon$

Convergence

- ▶ (x^n) converges to x if for all $\epsilon > 0$, there exists some N such that for all $n \geq N$, $d(x^n, x) < \epsilon$
- ▶ We denote convergence by $x^n \rightarrow x$ or $\lim x^n = x$

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)
- ▶ This sequence converges to $3/7$

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)
- ▶ This sequence converges to $3/7$
- ▶ Let $\epsilon > 0$ and let $N = \frac{19}{49\epsilon} + \frac{4}{7}$

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)
- ▶ This sequence converges to $3/7$
- ▶ Let $\epsilon > 0$ and let $N = \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ $n > N$ implies $n > \frac{19}{49\epsilon} + \frac{4}{7}$

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)
- ▶ This sequence converges to $3/7$
- ▶ Let $\epsilon > 0$ and let $N = \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ $n > N$ implies $n > \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ Rearranging yields $7n > \frac{19}{7\epsilon} + 4 \implies \epsilon > \frac{19}{7(7n-4)}$

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)
- ▶ This sequence converges to $3/7$
- ▶ Let $\epsilon > 0$ and let $N = \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ $n > N$ implies $n > \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ Rearranging yields $7n > \frac{19}{7\epsilon} + 4 \implies \epsilon > \frac{19}{7(7n-4)}$
- ▶ Hence $\frac{19}{7(7n-4)} = \frac{3n+1}{7n-4} - \frac{3}{7} < \epsilon$

Example

- ▶ Let $(x^n) = \frac{3n+1}{7n-4}$ be a sequence in (\mathbb{R}, d_1)
- ▶ This sequence converges to $3/7$
- ▶ Let $\epsilon > 0$ and let $N = \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ $n > N$ implies $n > \frac{19}{49\epsilon} + \frac{4}{7}$
- ▶ Rearranging yields $7n > \frac{19}{7\epsilon} + 4 \implies \epsilon > \frac{19}{7(7n-4)}$
- ▶ Hence $\frac{19}{7(7n-4)} = \frac{3n+1}{7n-4} - \frac{3}{7} < \epsilon$
- ▶ Therefore for $n > N$, $|\frac{3n+1}{7n-4} - \frac{3}{7}| < \epsilon$

Convergence

- ▶ If a sequence converges, it has a unique limit.

Convergence

- ▶ If a sequence converges, it has a unique limit.
- ▶ To see this, assume $x^n \rightarrow x$ and $x^n \rightarrow y$.

Convergence

- ▶ If a sequence converges, it has a unique limit.
- ▶ To see this, assume $x^n \rightarrow x$ and $x^n \rightarrow y$.
- ▶ By the triangle equality, $d(x, y) \leq d(x, x^n) + d(x^n, y)$ for all n

Convergence

- ▶ If a sequence converges, it has a unique limit.
- ▶ To see this, assume $x^n \rightarrow x$ and $x^n \rightarrow y$.
- ▶ By the triangle equality, $d(x, y) \leq d(x, x^n) + d(x^n, y)$ for all n
- ▶ Because the sequence converges,

$$[d(x, x^n) \rightarrow 0] \wedge [d(x^n, y) \rightarrow 0]$$

Convergence

- ▶ If a sequence converges, it has a unique limit.
- ▶ To see this, assume $x^n \rightarrow x$ and $x^n \rightarrow y$.
- ▶ By the triangle equality, $d(x, y) \leq d(x, x^n) + d(x^n, y)$ for all n
- ▶ Because the sequence converges,

$$[d(x, x^n) \rightarrow 0] \wedge [d(x^n, y) \rightarrow 0]$$

▶

$$\implies d(x, y) \rightarrow 0$$

Convergence

- ▶ If a sequence converges, it has a unique limit.
- ▶ To see this, assume $x^n \rightarrow x$ and $x^n \rightarrow y$.
- ▶ By the triangle equality, $d(x, y) \leq d(x, x^n) + d(x^n, y)$ for all n
- ▶ Because the sequence converges,

$$[d(x, x^n) \rightarrow 0] \wedge [d(x^n, y) \rightarrow 0]$$

▶

$$\implies d(x, y) \rightarrow 0$$

- ▶ Since x and y are constant, we must have $d(x, y) = 0$

Convergence

- ▶ If a sequence converges, it has a unique limit.
- ▶ To see this, assume $x^n \rightarrow x$ and $x^n \rightarrow y$.
- ▶ By the triangle equality, $d(x, y) \leq d(x, x^n) + d(x^n, y)$ for all n
- ▶ Because the sequence converges,

$$[d(x, x^n) \rightarrow 0] \wedge [d(x^n, y) \rightarrow 0]$$

▶

$$\implies d(x, y) \rightarrow 0$$

- ▶ Since x and y are constant, we must have $d(x, y) = 0$
- ▶ This implies $x = y$.

Bounded Sequence

- ▶ A sequence is bounded if there exists some $K \in \mathbb{R}$ such that $x^n \leq K$ for all n .

Bounded Sequence

- ▶ A sequence is bounded if there exists some $K \in \mathbb{R}$ such that $x^n \leq K$ for all n .
- ▶ To show that a sequence is bounded, it is sufficient to show that $d(x^n, y) \leq L$ for all n where $L \in \mathbb{R}$ and y is a point in X .

Bounded Sequence

- ▶ A sequence is bounded if there exists some $K \in \mathbb{R}$ such that $x^n \leq K$ for all n .
- ▶ To show that a sequence is bounded, it is sufficient to show that $d(x^n, y) \leq L$ for all n where $L \in \mathbb{R}$ and y is a point in X .
- ▶ Every convergent sequence is bounded.

Subsequences

- ▶ We will also occasionally use **subsequences** to prove properties of sequences.

Subsequences

- ▶ We will also occasionally use **subsequences** to prove properties of sequences.
- ▶ A subsequence of (x^n) is a sequence (y^k) where $y^k = x^{n_k}$ where $n_1 < n_2 < \dots$ is an increasing sequence of indices.

Subsequences

- ▶ We will also occasionally use **subsequences** to prove properties of sequences.
- ▶ A subsequence of (x^n) is a sequence (y^k) where $y^k = x^{n_k}$ where $n_1 < n_2 < \dots$ is an increasing sequence of indices.
- ▶ We often denote a subsequence of (x^n) with (x^{n_k})

Subsequences

- ▶ We will also occasionally use **subsequences** to prove properties of sequences.
- ▶ A subsequence of (x^n) is a sequence (y^k) where $y^k = x^{n_k}$ where $n_1 < n_2 < \dots$ is an increasing sequence of indices.
- ▶ We often denote a subsequence of (x^n) with (x^{n_k})
- ▶ *Example:* Let $(x^n) = 1, 1/2, 1/3, 1/4, 1/5, \dots$

Subsequences

- ▶ We will also occasionally use **subsequences** to prove properties of sequences.
- ▶ A subsequence of (x^n) is a sequence (y^k) where $y^k = x^{n_k}$ where $n_1 < n_2 < \dots$ is an increasing sequence of indices.
- ▶ We often denote a subsequence of (x^n) with (x^{n_k})
- ▶ *Example:* Let $(x^n) = 1, 1/2, 1/3, 1/4, 1/5, \dots$
- ▶ The sequence $(y^k) = 1, 1/3, 1/5, \dots$ is a subsequence of (x^n)

Sequences and Closed Sets

- ▶ Sequences allow an alternative characterization of closed sets from the definition given earlier.

Sequences and Closed Sets

- ▶ Sequences allow an alternative characterization of closed sets from the definition given earlier.
- ▶ A set S in a metric space X is closed if and only if every sequence in S that converges in X converges to a point in S

Proof (\implies)

- First we prove that closed implies that the limit of a convergent sequence is in S by contradiction.

Proof (\implies)

- ▶ First we prove that closed implies that the limit of a convergent sequence is in S by contradiction.
- ▶ Let S be a closed subset of X and take $(x^n) \in S$ with $x^n \rightarrow x$ for some $x \in X$

Proof (\implies)

- ▶ First we prove that closed implies that the limit of a convergent sequence is in S by contradiction.
- ▶ Let S be a closed subset of X and take $(x^n) \in S$ with $x^n \rightarrow x$ for some $x \in X$
- ▶ If $x \in X \setminus S$ then we can find an $r > 0$ such that $B_r(x) \subseteq X \setminus S$ because $X \setminus S$ is open.

Proof (\implies)

- ▶ First we prove that closed implies that the limit of a convergent sequence is in S by contradiction.
- ▶ Let S be a closed subset of X and take $(x^n) \in S$ with $x^n \rightarrow x$ for some $x \in X$
- ▶ If $x \in X \setminus S$ then we can find an $r > 0$ such that $B_r(x) \subseteq X \setminus S$ because $X \setminus S$ is open.
- ▶ But since $d(x^n, x) \rightarrow 0$, there must exist an N such that $x^N \in B_r(x)$

Proof (\implies)

- ▶ First we prove that closed implies that the limit of a convergent sequence is in S by contradiction.
- ▶ Let S be a closed subset of X and take $(x^n) \in S$ with $x^n \rightarrow x$ for some $x \in X$
- ▶ If $x \in X \setminus S$ then we can find an $r > 0$ such that $B_r(x) \subseteq X \setminus S$ because $X \setminus S$ is open.
- ▶ But since $d(x^n, x) \rightarrow 0$, there must exist an N such that $x^N \in B_r(x)$
- ▶ This contradicts the assertion that all terms of the sequence (x^n) are in S

Proof (\Leftarrow)

- Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”
- ▶ Suppose S is not closed

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”
- ▶ Suppose S is not closed
- ▶ Then $X \setminus S$ is not open

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”
- ▶ Suppose S is not closed
- ▶ Then $X \setminus S$ is not open
- ▶ Therefore we can find an $x \in X \setminus S$ such that every open ball around x intersects S

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”
- ▶ Suppose S is not closed
- ▶ Then $X \setminus S$ is not open
- ▶ Therefore we can find an $x \in X \setminus S$ such that every open ball around x intersects S
- ▶ Thus for any $n = 1, 2, \dots$, there is an $x^n \in B_{\frac{1}{n}}(x) \cap S$

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”
- ▶ Suppose S is not closed
- ▶ Then $X \setminus S$ is not open
- ▶ Therefore we can find an $x \in X \setminus S$ such that every open ball around x intersects S
- ▶ Thus for any $n = 1, 2, \dots$, there is an $x^n \in B_{\frac{1}{n}}(x) \cap S$
- ▶ But then $(x^n) \in S$ and $\lim x^n = x$ yet $x \notin S$

Proof (\Leftarrow)

- ▶ Now we prove that all convergent sequences in S converging to a point in S implies that S is closed by contraposition.
- ▶ That is, we show that “not closed” implies “there exists convergent sequence in S that does not converge to point in S ”
- ▶ Suppose S is not closed
- ▶ Then $X \setminus S$ is not open
- ▶ Therefore we can find an $x \in X \setminus S$ such that every open ball around x intersects S
- ▶ Thus for any $n = 1, 2, \dots$, there is an $x^n \in B_{\frac{1}{n}}(x) \cap S$
- ▶ But then $(x^n) \in S$ and $\lim_{n \rightarrow \infty} x^n = x$ yet $x \notin S$
- ▶ Thus if S is not closed, there would exist at least one sequence in S that converges to a point outside of S

Compactness

- ▶ Compactness is one of the most fundamental concepts in real analysis and plays an important role in optimization theory.

Compactness

- ▶ Compactness is one of the most fundamental concepts in real analysis and plays an important role in optimization theory.
- ▶ You've probably encountered a definition of compact that is loosely something like "a set that is closed and bounded."

Compactness

- ▶ Compactness is one of the most fundamental concepts in real analysis and plays an important role in optimization theory.
- ▶ You've probably encountered a definition of compact that is loosely something like "a set that is closed and bounded."
- ▶ This is true in some metric spaces but is not a proper definition of compact.

Compactness

- ▶ Let X be a metric space and $S \subseteq X$

Compactness

- ▶ Let X be a metric space and $S \subseteq X$
- ▶ A class \mathcal{O} of subsets of X is said to **cover** S if $S \subseteq \bigcup \mathcal{O}$

Compactness

- ▶ Let X be a metric space and $S \subseteq X$
- ▶ A class \mathcal{O} of subsets of X is said to **cover** S if $S \subseteq \bigcup \mathcal{O}$
- ▶ If all members of such a class \mathcal{O} are open in X , then we say that \mathcal{O} is an **open cover**.

Compactness

- ▶ Let X be a metric space and $S \subseteq X$
- ▶ A class \mathcal{O} of subsets of X is said to **cover** S if $S \subseteq \bigcup \mathcal{O}$
- ▶ If all members of such a class \mathcal{O} are open in X , then we say that \mathcal{O} is an **open cover**.
- ▶ A subset of \mathcal{O} that also covers S is called a **subcover**.

Compactness

- ▶ Let X be a metric space and $S \subseteq X$
- ▶ A class \mathcal{O} of subsets of X is said to **cover** S if $S \subseteq \bigcup \mathcal{O}$
- ▶ If all members of such a class \mathcal{O} are open in X , then we say that \mathcal{O} is an **open cover**.
- ▶ A subset of \mathcal{O} that also covers S is called a **subcover**.
- ▶ A metric space X is **compact** if every open cover of X has a finite subcover.

Example

- ▶ Consider the interval $(0, 1)$ and the collection $\mathcal{O} = \{(\frac{1}{i}, 1) : i = 1, 2, \dots\}$

Example

- ▶ Consider the interval $(0, 1)$ and the collection $\mathcal{O} = \{(\frac{1}{i}, 1) : i = 1, 2, \dots\}$
- ▶ Note that $(0, 1) = (\frac{1}{2}, 1) \cup (\frac{1}{3}, 1) \cup \dots$

Example

- ▶ Consider the interval $(0, 1)$ and the collection $\mathcal{O} = \{(\frac{1}{i}, 1) : i = 1, 2, \dots\}$
- ▶ Note that $(0, 1) = (\frac{1}{2}, 1) \cup (\frac{1}{3}, 1) \cup \dots$
- ▶ Therefore \mathcal{O} is an open cover of $(0, 1)$

Example

- ▶ Consider the interval $(0, 1)$ and the collection $\mathcal{O} = \{(\frac{1}{i}, 1) : i = 1, 2, \dots\}$
- ▶ Note that $(0, 1) = (\frac{1}{2}, 1) \cup (\frac{1}{3}, 1) \cup \dots$
- ▶ Therefore \mathcal{O} is an open cover of $(0, 1)$
- ▶ However, \mathcal{O} does not have a finite subset that covers \mathcal{O}

Example

- ▶ Consider the interval $(0, 1)$ and the collection $\mathcal{O} = \{(\frac{1}{i}, 1) : i = 1, 2, \dots\}$
- ▶ Note that $(0, 1) = (\frac{1}{2}, 1) \cup (\frac{1}{3}, 1) \cup \dots$
- ▶ Therefore \mathcal{O} is an open cover of $(0, 1)$
- ▶ However, \mathcal{O} does not have a finite subset that covers \mathcal{O}
- ▶ Therefore $(0, 1)$ is not compact.

Example

- ▶ $[0, 1]$ is a compact subset of \mathbb{R}

Example

- ▶ $[0, 1]$ is a compact subset of \mathbb{R}
- ▶ Suppose there exists an open cover \mathcal{O} of $[0, 1]$ such that no finite subset of \mathcal{O} covers $[0, 1]$.

Example

- ▶ $[0, 1]$ is a compact subset of \mathbb{R}
- ▶ Suppose there exists an open cover \mathcal{O} of $[0, 1]$ such that no finite subset of \mathcal{O} covers $[0, 1]$.
- ▶ This implies that either $[0, \frac{1}{2}]$ or $[\frac{1}{2}, 1]$ is not covered by any finite subset of \mathcal{O} .

Example

- ▶ $[0, 1]$ is a compact subset of \mathbb{R}
- ▶ Suppose there exists an open cover \mathcal{O} of $[0, 1]$ such that no finite subset of \mathcal{O} covers $[0, 1]$.
- ▶ This implies that either $[0, \frac{1}{2}]$ or $[\frac{1}{2}, 1]$ is not covered by any finite subset of \mathcal{O} .
- ▶ Pick any one of the intervals with this property and call it $[a_1, b_1]$.

Example

- ▶ $[0, 1]$ is a compact subset of \mathbb{R}
- ▶ Suppose there exists an open cover \mathcal{O} of $[0, 1]$ such that no finite subset of \mathcal{O} covers $[0, 1]$.
- ▶ This implies that either $[0, \frac{1}{2}]$ or $[\frac{1}{2}, 1]$ is not covered by any finite subset of \mathcal{O} .
- ▶ Pick any one of the intervals with this property and call it $[a_1, b_1]$.
- ▶ Then either $[a_1, \frac{1}{2}(a_1 + b_1)]$ or $[\frac{1}{2}(b_1 + a_1), b_1]$ is not covered by any finite subset of \mathcal{O} .

Example

- ▶ $[0, 1]$ is a compact subset of \mathbb{R}
- ▶ Suppose there exists an open cover \mathcal{O} of $[0, 1]$ such that no finite subset of \mathcal{O} covers $[0, 1]$.
- ▶ This implies that either $[0, \frac{1}{2}]$ or $[\frac{1}{2}, 1]$ is not covered by any finite subset of \mathcal{O} .
- ▶ Pick any one of the intervals with this property and call it $[a_1, b_1]$.
- ▶ Then either $[a_1, \frac{1}{2}(a_1 + b_1)]$ or $[\frac{1}{2}(b_1 + a_1), b_1]$ is not covered by any finite subset of \mathcal{O} .
- ▶ Pick any one of these intervals with this property and call it $[a_2, b_2]$.

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$
 3. $[a_m, b_m]$ is not covered by any finite subset of \mathcal{O}

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$
 3. $[a_m, b_m]$ is not covered by any finite subset of \mathcal{O}
- ▶ Properties (1) and (2) allow us to find a real number c with $\{c\} = \bigcap^{\infty} [a_i, b_i]$

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$
 3. $[a_m, b_m]$ is not covered by any finite subset of \mathcal{O}
- ▶ Properties (1) and (2) allow us to find a real number c with $\{c\} = \bigcap^{\infty} [a_i, b_i]$
- ▶ Now take any $O \in \mathcal{O}$ which contains c

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$
 3. $[a_m, b_m]$ is not covered by any finite subset of \mathcal{O}
- ▶ Properties (1) and (2) allow us to find a real number c with $\{c\} = \bigcap_{i=1}^{\infty} [a_i, b_i]$
- ▶ Now take any $O \in \mathcal{O}$ which contains c
- ▶ Since O is open and $\lim a_m = \lim b_m = c$, we must have $[a_m, b_m] \subset O$ for a large enough m

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$
 3. $[a_m, b_m]$ is not covered by any finite subset of \mathcal{O}
- ▶ Properties (1) and (2) allow us to find a real number c with $\{c\} = \bigcap^{\infty} [a_i, b_i]$
- ▶ Now take any $O \in \mathcal{O}$ which contains c
- ▶ Since O is open and $\lim a_m = \lim b_m = c$, we must have $[a_m, b_m] \subset O$ for a large enough m
- ▶ But this contradicts condition (3)!

Example (cont.)

- ▶ Continue this inductively to obtain two sequences, (a_m) and (b_m) in $[0, 1]$ such that for each $m = 1, 2, \dots$
 1. $a_m \leq a_{m+1} < b_{m+1} \leq b_m$
 2. $b_m - a_m = \frac{1}{2^m}$
 3. $[a_m, b_m]$ is not covered by any finite subset of \mathcal{O}
- ▶ Properties (1) and (2) allow us to find a real number c with $\{c\} = \bigcap_{i=1}^{\infty} [a_i, b_i]$
- ▶ Now take any $O \in \mathcal{O}$ which contains c
- ▶ Since O is open and $\lim a_m = \lim b_m = c$, we must have $[a_m, b_m] \subset O$ for a large enough m
- ▶ But this contradicts condition (3)!
- ▶ We conclude that $[0, 1]$ is compact

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.
- ▶ This is primarily due to the fact that we are used to Euclidean space.

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.
- ▶ This is primarily due to the fact that we are used to Euclidean space.
- ▶ The Heine-Borel Theorem is to thank for this result

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.
- ▶ This is primarily due to the fact that we are used to Euclidean space.
- ▶ The Heine-Borel Theorem is to thank for this result
- ▶ The theorem establishes that the following two statements are equivalent for a subset S of (\mathbb{R}^n, d_2) :

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.
- ▶ This is primarily due to the fact that we are used to Euclidean space.
- ▶ The Heine-Borel Theorem is to thank for this result
- ▶ The theorem establishes that the following two statements are equivalent for a subset S of (\mathbb{R}^n, d_2) :
 1. S is closed and bounded

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.
- ▶ This is primarily due to the fact that we are used to Euclidean space.
- ▶ The Heine-Borel Theorem is to thank for this result
- ▶ The theorem establishes that the following two statements are equivalent for a subset S of (\mathbb{R}^n, d_2) :
 1. S is closed and bounded
 2. S is compact, i.e. every open cover of S has a finite subcover

Heine-Borel Theorem

- ▶ Of course, we are familiar with a simpler version of compactness.
- ▶ This is primarily due to the fact that we are used to Euclidean space.
- ▶ The Heine-Borel Theorem is to thank for this result
- ▶ The theorem establishes that the following two statements are equivalent for a subset S of (\mathbb{R}^n, d_2) :
 1. S is closed and bounded
 2. S is compact, i.e. every open cover of S has a finite subcover
- ▶ In general, compactness implies closed and bounded but the reverse is guaranteed to hold only in Euclidean space.

Compactness

- ▶ An additional pair of facts will help us find other compact sets.

Compactness

- ▶ An additional pair of facts will help us find other compact sets.
- ▶ First, any closed subset of a compact metric space is compact.

Compactness

- ▶ An additional pair of facts will help us find other compact sets.
- ▶ First, any closed subset of a compact metric space is compact.
- ▶ Second, the product of two compact metric spaces is compact (Tychonoff's theorem).

Compactness

- ▶ An additional pair of facts will help us find other compact sets.
- ▶ First, any closed subset of a compact metric space is compact.
- ▶ Second, the product of two compact metric spaces is compact (Tychonoff's theorem).
- ▶ This latter fact will be very useful in game theory when we want to determine whether a strategy space $S = S_1 \times S_2 \times \dots \times S_n$ is compact.

Compactness

- ▶ An additional pair of facts will help us find other compact sets.
- ▶ First, any closed subset of a compact metric space is compact.
- ▶ Second, the product of two compact metric spaces is compact (Tychonoff's theorem).
- ▶ This latter fact will be very useful in game theory when we want to determine whether a strategy space $S = S_1 \times S_2 \times \dots \times S_n$ is compact.
- ▶ If each player $i = 1, 2, \dots, n$ has a compact strategy space, the strategy space is compact.

Compactness

- ▶ An additional pair of facts will help us find other compact sets.
- ▶ First, any closed subset of a compact metric space is compact.
- ▶ Second, the product of two compact metric spaces is compact (Tychonoff's theorem).
- ▶ This latter fact will be very useful in game theory when we want to determine whether a strategy space $S = S_1 \times S_2 \times \dots \times S_n$ is compact.
- ▶ If each player $i = 1, 2, \dots, n$ has a compact strategy space, the strategy space is compact.
- ▶ A compact strategy space will be a key component of the fixed point theorems we use to prove the existence of a Nash equilibrium.

Cauchy Sequences

- ▶ A sequence (x^n) in a metric space X is called a **Cauchy sequence** if for any $\epsilon > 0$, there exists an $N \in \mathbb{R}$ such that $d(x^k, x^l) < \epsilon$ for all $k, l \geq N$.

Cauchy Sequences

- ▶ A sequence (x^n) in a metric space X is called a **Cauchy sequence** if for any $\epsilon > 0$, there exists an $N \in \mathbb{R}$ such that $d(x^k, x^l) < \epsilon$ for all $k, l \geq N$.
- ▶ *Example:* $(x^n) = 1/2^n$ is a Cauchy sequence in (\mathbb{R}, d_1) .

Cauchy Sequences

- ▶ A sequence (x^n) in a metric space X is called a **Cauchy sequence** if for any $\epsilon > 0$, there exists an $N \in \mathbb{R}$ such that $d(x^k, x^l) < \epsilon$ for all $k, l \geq N$.
- ▶ *Example:* $(x^n) = 1/2^n$ is a Cauchy sequence in (\mathbb{R}, d_1) .
- ▶ Take $\epsilon > 0$ and choose N so large that $2^{-N} < \epsilon$.

Cauchy Sequences

- ▶ A sequence (x^n) in a metric space X is called a **Cauchy sequence** if for any $\epsilon > 0$, there exists an $N \in \mathbb{R}$ such that $d(x^k, x^l) < \epsilon$ for all $k, l \geq N$.
- ▶ *Example:* $(x^n) = 1/2^n$ is a Cauchy sequence in (\mathbb{R}, d_1) .
- ▶ Take $\epsilon > 0$ and choose N so large that $2^{-N} < \epsilon$.
- ▶ Then if $n, m > N$, we have

$$|x^n - x^m| = |1/2^n - 1/2^m| \leq 1/2^n + 1/2^m \leq 1/2^N + 1/2^N < \epsilon \quad \blacksquare$$

Properties of Cauchy Sequences

1. If (x^n) is convergent, then it is Cauchy.

Properties of Cauchy Sequences

1. If (x^n) is convergent, then it is Cauchy.
2. If (x^n) is Cauchy, then (x^n) is bounded but need not converge in X .

Properties of Cauchy Sequences

1. If (x^n) is convergent, then it is Cauchy.
2. If (x^n) is Cauchy, then (x^n) is bounded but need not converge in X .
3. If (x^n) is Cauchy and has a subsequence that converges in X , then (x^n) converges in X as well.

Properties of Cauchy Sequences

- ▶ Property 1 and the first part of Property 2 are exercises.

Properties of Cauchy Sequences

- ▶ Property 1 and the first part of Property 2 are exercises.
- ▶ To see the second part of Property 2, consider the sequence $(x^n) = (1, 1/2, 1/3, \dots)$ in $(0, 1]$

Properties of Cauchy Sequences

- ▶ Property 1 and the first part of Property 2 are exercises.
- ▶ To see the second part of Property 2, consider the sequence $(x^n) = (1, 1/2, 1/3, \dots)$ in $(0, 1]$
- ▶ This sequence is Cauchy but does not converge in this space.

Properties of Cauchy Sequences

- ▶ Property 1 and the first part of Property 2 are exercises.
- ▶ To see the second part of Property 2, consider the sequence $(x^n) = (1, 1/2, 1/3, \dots)$ in $(0, 1]$
- ▶ This sequence is Cauchy but does not converge in this space.
- ▶ For Property 3, say (x^n) is Cauchy and has a convergent subsequence $(x^{n_k}) \rightarrow x$

Properties of Cauchy Sequences

- ▶ Property 1 and the first part of Property 2 are exercises.
- ▶ To see the second part of Property 2, consider the sequence $(x^n) = (1, 1/2, 1/3, \dots)$ in $(0, 1]$
- ▶ This sequence is Cauchy but does not converge in this space.
- ▶ For Property 3, say (x^n) is Cauchy and has a convergent subsequence $(x^{n_k}) \rightarrow x$
- ▶

$$d(x^n, x) \leq d(x^n, x^{n_k}) + d(x^{n_k}, x)$$

Properties of Cauchy Sequences

- ▶ Property 1 and the first part of Property 2 are exercises.
- ▶ To see the second part of Property 2, consider the sequence $(x^n) = (1, 1/2, 1/3, \dots)$ in $(0, 1]$
- ▶ This sequence is Cauchy but does not converge in this space.
- ▶ For Property 3, say (x^n) is Cauchy and has a convergent subsequence $(x^{n_k}) \rightarrow x$



$$d(x^n, x) \leq d(x^n, x^{n_k}) + d(x^{n_k}, x)$$

- ▶ $d(x^n, x^{n_k}) + d(x^{n_k}, x) \rightarrow 0$

Completeness

- ▶ Compactness generalizes the concept of finiteness and is one of two fundamental properties of metric spaces.

Completeness

- ▶ Compactness generalizes the concept of finiteness and is one of two fundamental properties of metric spaces.
- ▶ The other property, **completeness**, generalizes the idea of richness.

Completeness

- ▶ Compactness generalizes the concept of finiteness and is one of two fundamental properties of metric spaces.
- ▶ The other property, **completeness**, generalizes the idea of richness.
- ▶ An incomplete space lacks certain necessary elements.

Completeness

- ▶ Compactness generalizes the concept of finiteness and is one of two fundamental properties of metric spaces.
- ▶ The other property, **completeness**, generalizes the idea of richness.
- ▶ An incomplete space lacks certain necessary elements.
- ▶ A **complete metric space** is defined as a metric space (X, d) in which every Cauchy sequence in X converges to a point in X .

Completeness

- ▶ Compactness generalizes the concept of finiteness and is one of two fundamental properties of metric spaces.
- ▶ The other property, **completeness**, generalizes the idea of richness.
- ▶ An incomplete space lacks certain necessary elements.
- ▶ A **complete metric space** is defined as a metric space (X, d) in which every Cauchy sequence in X converges to a point in X .
- ▶ We saw earlier that $(0, 1]$ with the d_1 metric is not complete.

Completeness

- ▶ It is natural to think about a relationship between completeness and closedness.

Completeness

- ▶ It is natural to think about a relationship between completeness and closedness.
- ▶ Let X be a metric space and Y a metric subspace of X .

Completeness

- ▶ It is natural to think about a relationship between completeness and closedness.
- ▶ Let X be a metric space and Y a metric subspace of X .
- ▶ If Y is complete, then it is closed in X .

Completeness

- ▶ It is natural to think about a relationship between completeness and closedness.
- ▶ Let X be a metric space and Y a metric subspace of X .
- ▶ If Y is complete, then it is closed in X .
- ▶ Conversely, if Y is closed in X and X is complete, then Y is complete.

Proof: Complete Subspace \implies Subspace Closed

- ▶ Let Y be complete and take any $(x^n) \in Y$ that converges in X .

Proof: Complete Subspace \implies Subspace Closed

- ▶ Let Y be complete and take any $(x^n) \in Y$ that converges in X .
- ▶ Because (x^n) converges, it is Cauchy and thus $\lim x^n \in Y$.

Proof: Complete Subspace \implies Subspace Closed

- ▶ Let Y be complete and take any $(x^n) \in Y$ that converges in X .
- ▶ Because (x^n) converges, it is Cauchy and thus $\lim x^n \in Y$.
- ▶ We saw earlier that if a sequence in Y that converges in X converges to a point in Y , then Y is closed.

Proof: Closed Subspace of Complete Space \implies Subspace Complete

- ▶ Assume X is complete and Y is closed in X

Proof: Closed Subspace of Complete Space \implies Subspace Complete

- ▶ Assume X is complete and Y is closed in X
- ▶ If (x^n) is a Cauchy sequence in Y , then by the completeness of X , it must converge in X

Proof: Closed Subspace of Complete Space \implies Subspace Complete

- ▶ Assume X is complete and Y is closed in X
- ▶ If (x^n) is a Cauchy sequence in Y , then by the completeness of X , it must converge in X
- ▶ Since Y is closed, $\lim x^n$ must be an element of Y .

Proof: Closed Subspace of Complete Space \implies Subspace Complete

- ▶ Assume X is complete and Y is closed in X
- ▶ If (x^n) is a Cauchy sequence in Y , then by the completeness of X , it must converge in X
- ▶ Since Y is closed, $\lim x^n$ must be an element of Y .
- ▶ Therefore Y is complete.

Completeness

- ▶ Two more facts about completeness will be useful

Completeness

- ▶ Two more facts about completeness will be useful
- ▶ First, a metric subspace of a complete metric space is complete if and only if it is closed in X .

Completeness

- ▶ Two more facts about completeness will be useful
- ▶ First, a metric subspace of a complete metric space is complete if and only if it is closed in X .
- ▶ This is a corollary of the proposition we just proved.

Completeness

- ▶ Two more facts about completeness will be useful
- ▶ First, a metric subspace of a complete metric space is complete if and only if it is closed in X .
- ▶ This is a corollary of the proposition we just proved.
- ▶ Second, every compact metric space is complete.

Intro to Fixed Point Theory

- ▶ Why do we care about completeness?

Intro to Fixed Point Theory

- ▶ Why do we care about completeness?
- ▶ It turns out that certain types of mappings on a complete metric space possess a very desirable “fixed point” property.

Intro to Fixed Point Theory

- ▶ Why do we care about completeness?
- ▶ It turns out that certain types of mappings on a complete metric space possess a very desirable “fixed point” property.
- ▶ A **self-map** is a function, $f(\cdot)$, whose domain and codomain are identical.

Intro to Fixed Point Theory

- ▶ Why do we care about completeness?
- ▶ It turns out that certain types of mappings on a complete metric space possess a very desirable “fixed point” property.
- ▶ A **self-map** is a function, $f(\cdot)$, whose domain and codomain are identical.
- ▶ A **fixed point** of a self-map is a point that satisfies the property $f(x) = x$.

Intro to Fixed Point Theory

- ▶ Why do we care about completeness?
- ▶ It turns out that certain types of mappings on a complete metric space possess a very desirable “fixed point” property.
- ▶ A **self-map** is a function, $f(\cdot)$, whose domain and codomain are identical.
- ▶ A **fixed point** of a self-map is a point that satisfies the property $f(x) = x$.
- ▶ Fixed points will be extremely useful in political science theory for ensuring that a model has an equilibrium.

Contractions

- ▶ One type of self-map is called a **contraction**.

Contractions

- ▶ One type of self-map is called a **contraction**.
- ▶ Let (X, d) be a metric space.

Contractions

- ▶ One type of self-map is called a **contraction**.
- ▶ Let (X, d) be a metric space.
- ▶ A self-map f on X is a contraction if there exists a real number $K \in (0, 1)$ such that

$$d(f(x), f(y)) \leq Kd(x, y)$$

for all $x, y \in X$.

Contractions

- ▶ One type of self-map is called a **contraction**.
- ▶ Let (X, d) be a metric space.
- ▶ A self-map f on X is a contraction if there exists a real number $K \in (0, 1)$ such that

$$d(f(x), f(y)) \leq Kd(x, y)$$

for all $x, y \in X$.

- ▶ *Example:* Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = \frac{x}{2}$.

Contractions

- ▶ One type of self-map is called a **contraction**.
- ▶ Let (X, d) be a metric space.
- ▶ A self-map f on X is a contraction if there exists a real number $K \in (0, 1)$ such that

$$d(f(x), f(y)) \leq Kd(x, y)$$

for all $x, y \in X$.

- ▶ *Example:* Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = \frac{x}{2}$.
- ▶ To verify that this is a contraction, consider two arbitrary points, $x, y \in \mathbb{R}$

Contractions

- ▶ One type of self-map is called a **contraction**.
- ▶ Let (X, d) be a metric space.
- ▶ A self-map f on X is a contraction if there exists a real number $K \in (0, 1)$ such that

$$d(f(x), f(y)) \leq Kd(x, y)$$

for all $x, y \in X$.

- ▶ *Example:* Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = \frac{x}{2}$.
- ▶ To verify that this is a contraction, consider two arbitrary points, $x, y \in \mathbb{R}$
- ▶ It is easy to check that $d(f(x), f(y)) \leq Kd(x, y)$ for some $K \in (0, 1)$:

$$\begin{aligned} d(f(x), f(y)) &= |f(x) - f(y)| = \left| \frac{x}{2} - \frac{y}{2} \right| \\ &= \left| \frac{1}{2}(x - y) \right| \leq \frac{1}{2}|x - y| = \frac{1}{2}d(x, y) \end{aligned}$$

Intro to Fixed Point Theory

- ▶ When a contraction is defined on a complete metric space, a contraction must map a point to itself.

Intro to Fixed Point Theory

- ▶ When a contraction is defined on a complete metric space, a contraction must map a point to itself.
- ▶ In other words, it *must* have a fixed point.

Intro to Fixed Point Theory

- ▶ When a contraction is defined on a complete metric space, a contraction must map a point to itself.
- ▶ In other words, it *must* have a fixed point.
- ▶ In fact, we can show that it must have a *unique* fixed point!

Intro to Fixed Point Theory

- ▶ When a contraction is defined on a complete metric space, a contraction must map a point to itself.
- ▶ In other words, it *must* have a fixed point.
- ▶ In fact, we can show that it must have a *unique* fixed point!
- ▶ Formally, the **Banach Fixed Point Theorem** or **Contraction Mapping Theorem** states that if (X, d) is a complete metric space and f is a contraction defined on X , then there exists a unique $x^* \in X$ such that $f(x^*) = x^*$.

Proof

- ▶ Let (X, d) be a complete metric space and let $f : X \rightarrow X$ be a contraction.

Proof

- ▶ Let (X, d) be a complete metric space and let $f : X \rightarrow X$ be a contraction.
- ▶ To show the existence of a fixed point for f , pick any $x \in X$ and define $(x^n) \in X$ such that $x^{n+1} = f(x^n)$ for $n = 0, 1, \dots$

Proof

- ▶ Let (X, d) be a complete metric space and let $f : X \rightarrow X$ be a contraction.
- ▶ To show the existence of a fixed point for f , pick any $x \in X$ and define $(x^n) \in X$ such that $x^{n+1} = f(x^n)$ for $n = 0, 1, \dots$
- ▶ We can show that the sequence is Cauchy

Proof

- ▶ Let (X, d) be a complete metric space and let $f : X \rightarrow X$ be a contraction.
- ▶ To show the existence of a fixed point for f , pick any $x \in X$ and define $(x^n) \in X$ such that $x^{n+1} = f(x^n)$ for $n = 0, 1, \dots$
- ▶ We can show that the sequence is Cauchy
- ▶ By construction,

$$\begin{aligned}d(x^{n+1}, x^{n+2}) &\leq Kd(x^n, x^{n+1}) \leq K^2d(x^{n-1}, x^n) \leq \\ &\dots \leq K^nd(x^0, x^1)\end{aligned}$$

Proof

- ▶ Let (X, d) be a complete metric space and let $f : X \rightarrow X$ be a contraction.
- ▶ To show the existence of a fixed point for f , pick any $x \in X$ and define $(x^n) \in X$ such that $x^{n+1} = f(x^n)$ for $n = 0, 1, \dots$
- ▶ We can show that the sequence is Cauchy
- ▶ By construction,

$$\begin{aligned}d(x^{n+1}, x^{n+2}) &\leq Kd(x^n, x^{n+1}) \leq K^2d(x^{n-1}, x^n) \leq \\ &\dots \leq K^nd(x^0, x^1)\end{aligned}$$

- ▶ That is, $d(x^{n+1}, x^n) \leq K^nd(x^1, x^0)$ for all $n = 1, 2, \dots$

Proof (cont.)

- ▶ Thus for any $k > l + 1$,

$$\begin{aligned}d(x^k, x^l) &\leq d(x^k, x^{k-1}) + \dots + d(x^{l+1}, x^l) \\&\leq (K^{k-1} + \dots + K^l)d(x^1, x^0) \\&= \frac{K^l(1 - K^{k-l})}{1 - K}d(x^1, x^0)\end{aligned}$$

Proof (cont.)

- ▶ Thus for any $k > l + 1$,

$$\begin{aligned}d(x^k, x^l) &\leq d(x^k, x^{k-1}) + \dots + d(x^{l+1}, x^l) \\&\leq (K^{k-1} + \dots + K^l)d(x^1, x^0) \\&= \frac{K^l(1 - K^{k-l})}{1 - K}d(x^1, x^0)\end{aligned}$$

- ▶ Therefore $d(x^k, x^l) < \frac{K^l}{1-K}d(x^1, x^0)$

Proof (cont.)

- ▶ Thus for any $k > l + 1$,

$$\begin{aligned}d(x^k, x^l) &\leq d(x^k, x^{k-1}) + \dots + d(x^{l+1}, x^l) \\&\leq (K^{k-1} + \dots + K^l)d(x^1, x^0) \\&= \frac{K^l(1 - K^{k-l})}{1 - K}d(x^1, x^0)\end{aligned}$$

- ▶ Therefore $d(x^k, x^l) < \frac{K^l}{1-K}d(x^1, x^0)$
- ▶ This implies that (x^n) is a Cauchy sequence.

Proof (cont.)

- ▶ Since X is complete, our Cauchy sequence (x^n) must converge to some $x^* \in X$

Proof (cont.)

- ▶ Since X is complete, our Cauchy sequence (x^n) must converge to some $x^* \in X$
- ▶ Then for any $\epsilon > 0$, there must exist some N such that $d(x^*, x^n) < \frac{\epsilon}{2}$ for all $n = N, N + 1, \dots$

Proof (cont.)

- ▶ Since X is complete, our Cauchy sequence (x^n) must converge to some $x^* \in X$
- ▶ Then for any $\epsilon > 0$, there must exist some N such that $d(x^*, x^n) < \frac{\epsilon}{2}$ for all $n = N, N + 1, \dots$
- ▶ Therefore

$$\begin{aligned}d(f(x^*), x^*) &\leq d(f(x^*), x^{n+1}) + d(x^{n+1}, x^*) \\&= d(f(x^*), f(x^n)) + d(x^{n+1}, x^*) \\&\leq Kd(x^*, x^n) + d(x^{n+1}, x^*) \\&< \frac{\epsilon}{2} + \frac{\epsilon}{2}\end{aligned}$$

Proof (cont.)

- ▶ Since X is complete, our Cauchy sequence (x^n) must converge to some $x^* \in X$
- ▶ Then for any $\epsilon > 0$, there must exist some N such that $d(x^*, x^n) < \frac{\epsilon}{2}$ for all $n = N, N + 1, \dots$
- ▶ Therefore

$$\begin{aligned}d(f(x^*), x^*) &\leq d(f(x^*), x^{n+1}) + d(x^{n+1}, x^*) \\&= d(f(x^*), f(x^n)) + d(x^{n+1}, x^*) \\&\leq Kd(x^*, x^n) + d(x^{n+1}, x^*) \\&< \frac{\epsilon}{2} + \frac{\epsilon}{2}\end{aligned}$$

- ▶ Since $\epsilon > 0$ is arbitrary, we must have $d(f(x^*), x^*) = 0$ which is possible only if $f(x^*) = x^*$.

Proof (cont.)

- To prove uniqueness, note that if $x \in X$ was another fixed point, we would have

$$d(x, x^*) = d(f(x), f(x^*)) \leq Kd(x, x^*)$$

which is possible only if $x = x^*$.