

# Monotone, Linear, and Convex Functions

# Motivation

- ▶ We saw in the last lecture that continuous functions preserve the *geometric structure* of the sets they associate.

# Motivation

- ▶ We saw in the last lecture that continuous functions preserve the *geometric structure* of the sets they associate.
- ▶ Monotone functions preserve the *order* structure of their domains.

# Motivation

- ▶ We saw in the last lecture that continuous functions preserve the *geometric structure* of the sets they associate.
- ▶ Monotone functions preserve the *order* structure of their domains.
- ▶ Linear functions preserve the *algebraic structure* of the spaces they link.

# Motivation

- ▶ We saw in the last lecture that continuous functions preserve the *geometric structure* of the sets they associate.
- ▶ Monotone functions preserve the *order* structure of their domains.
- ▶ Linear functions preserve the *algebraic structure* of the spaces they link.
- ▶ These different types of functions have different properties we can exploit when building theories and analyzing data.

# Monotone Functions

- ▶ A function between ordered sets  $X$  and  $Y$  is called **monotone** if it respects the order of  $X$  and  $Y$ .

# Monotone Functions

- ▶ A function between ordered sets  $X$  and  $Y$  is called **monotone** if it respects the order of  $X$  and  $Y$ .
- ▶  $f$  is **increasing** if it preserves the ordering so that

$$x_2 \preceq_X x_1 \implies f(x_2) \preceq_Y f(x_1)$$

where  $\preceq_i$  is the order on set  $i$ .

# Monotone Functions

- ▶ A function between ordered sets  $X$  and  $Y$  is called **monotone** if it respects the order of  $X$  and  $Y$ .
- ▶  $f$  is **increasing** if it preserves the ordering so that

$$x_2 \succsim_X x_1 \implies f(x_2) \succsim_Y f(x_1)$$

where  $\succsim_i$  is the order on set  $i$ .

- ▶ A function is **strictly increasing** if the order relation is strict.



# Monotone Functions

- ▶ A function between ordered sets  $X$  and  $Y$  is called **monotone** if it respects the order of  $X$  and  $Y$ .
- ▶  $f$  is **increasing** if it preserves the ordering so that

$$x_2 \preceq_X x_1 \implies f(x_2) \preceq_Y f(x_1)$$

where  $\preceq_i$  is the order on set  $i$ .

- ▶ A function is **strictly increasing** if the order relation is strict.
- ▶ Decreasing and strictly decreasing functions are defined analogously.

# Monotone Functions

- ▶ A function between ordered sets  $X$  and  $Y$  is called **monotone** if it respects the order of  $X$  and  $Y$ .
- ▶  $f$  is **increasing** if it preserves the ordering so that

$$x_2 \preceq_X x_1 \implies f(x_2) \preceq_Y f(x_1)$$

where  $\preceq_i$  is the order on set  $i$ .

- ▶ A function is **strictly increasing** if the order relation is strict.
- ▶ Decreasing and strictly decreasing functions are defined analogously.
- ▶ A function  $f$  is monotone if it is either increasing or decreasing.

## Examples

- ▶ In practice we will often be using functions that map from  $\mathbb{R}$  into  $\mathbb{R}$  where each set is ordered with  $\geq$ .

# Examples

- ▶ In practice we will often be using functions that map from  $\mathbb{R}$  into  $\mathbb{R}$  where each set is ordered with  $\geq$ .
- ▶ The function  $\ln(x)$  is a monotone function: for any  $x_2 \geq x_1$  in  $\mathbb{R}_+$ ,  $\ln(x_2) \geq \ln(x_1)$  in  $\mathbb{R}$

# Examples

- ▶ In practice we will often be using functions that map from  $\mathbb{R}$  into  $\mathbb{R}$  where each set is ordered with  $\geq$ .
- ▶ The function  $\ln(x)$  is a monotone function: for any  $x_2 \geq x_1$  in  $\mathbb{R}_+$ ,  $\ln(x_2) \geq \ln(x_1)$  in  $\mathbb{R}$
- ▶ We also use utility functions that map from  $\mathbb{R}^n$  into  $\mathbb{R}$ .

# Examples

- ▶ In practice we will often be using functions that map from  $\mathbb{R}$  into  $\mathbb{R}$  where each set is ordered with  $\geq$ .
- ▶ The function  $\ln(x)$  is a monotone function: for any  $x_2 \geq x_1$  in  $\mathbb{R}_+$ ,  $\ln(x_2) \geq \ln(x_1)$  in  $\mathbb{R}$
- ▶ We also use utility functions that map from  $\mathbb{R}^n$  into  $\mathbb{R}$ .
- ▶ A function that maps into  $\mathbb{R}$  is called a **functional**

# Examples

- ▶ In practice we will often be using functions that map from  $\mathbb{R}$  into  $\mathbb{R}$  where each set is ordered with  $\geq$ .
- ▶ The function  $\ln(x)$  is a monotone function: for any  $x_2 \geq x_1$  in  $\mathbb{R}_+$ ,  $\ln(x_2) \geq \ln(x_1)$  in  $\mathbb{R}$
- ▶ We also use utility functions that map from  $\mathbb{R}^n$  into  $\mathbb{R}$ .
- ▶ A function that maps into  $\mathbb{R}$  is called a **functional**
- ▶ A Cobb-Douglas utility functional,  $f(x, y) = x^a y^b$  for  $a, b > 0$ , is monotone on the domain  $(\mathbb{R}_+^2, \geq)$

# Examples

- ▶ In practice we will often be using functions that map from  $\mathbb{R}$  into  $\mathbb{R}$  where each set is ordered with  $\geq$ .
- ▶ The function  $\ln(x)$  is a monotone function: for any  $x_2 \geq x_1$  in  $\mathbb{R}_+$ ,  $\ln(x_2) \geq \ln(x_1)$  in  $\mathbb{R}$
- ▶ We also use utility functions that map from  $\mathbb{R}^n$  into  $\mathbb{R}$ .
- ▶ A function that maps into  $\mathbb{R}$  is called a **functional**
- ▶ A Cobb-Douglas utility functional,  $f(x, y) = x^a y^b$  for  $a, b > 0$ , is monotone on the domain  $(\mathbb{R}_+^2, \geq)$
- ▶  $(x_2, y_2) \geq (x_1, y_1)$  implies  $x_2 \geq x_1$  and  $y_2 \geq y_1$



# Monotonic Transformations

- ▶ Given any functional  $f$  on  $X$  and a strictly increasing functional  $g : \mathbb{R} \rightarrow \mathbb{R}$ , their composition  $g \circ f : X \rightarrow \mathbb{R}$  is called a **monotonic transformation**.

# Monotonic Transformations

- ▶ Given any functional  $f$  on  $X$  and a strictly increasing functional  $g : \mathbb{R} \rightarrow \mathbb{R}$ , their composition  $g \circ f : X \rightarrow \mathbb{R}$  is called a **monotonic transformation**.
- ▶ A monotonic transformation preserves the ordering implied by  $f$ .

# Monotonic Transformations

- ▶ Given any functional  $f$  on  $X$  and a strictly increasing functional  $g : \mathbb{R} \rightarrow \mathbb{R}$ , their composition  $g \circ f : X \rightarrow \mathbb{R}$  is called a **monotonic transformation**.
- ▶ A monotonic transformation preserves the ordering implied by  $f$ .
- ▶ *Example:* Frequently it is easier to work with a log transformation of a function.

# Monotonic Transformations

- ▶ Given any functional  $f$  on  $X$  and a strictly increasing functional  $g : \mathbb{R} \rightarrow \mathbb{R}$ , their composition  $g \circ f : X \rightarrow \mathbb{R}$  is called a **monotonic transformation**.
- ▶ A monotonic transformation preserves the ordering implied by  $f$ .
- ▶ *Example:* Frequently it is easier to work with a log transformation of a function.
- ▶ The log of a Cobb-Douglass function is
$$\ln f(x) = a \ln(x) + b \ln(y)$$

# Linear Functions

- ▶ A function  $f : X \rightarrow Y$  between two linear spaces is **linear** if it preserves the linearity of the sets  $X$  and  $Y$ .

# Linear Functions

- ▶ A function  $f : X \rightarrow Y$  between two linear spaces is **linear** if it preserves the linearity of the sets  $X$  and  $Y$ .
- ▶ For all  $x_1, x_2 \in X$  and  $\alpha \in \mathbb{R}$ , we have

# Linear Functions

- ▶ A function  $f : X \rightarrow Y$  between two linear spaces is **linear** if it preserves the linearity of the sets  $X$  and  $Y$ .
- ▶ For all  $x_1, x_2 \in X$  and  $\alpha \in \mathbb{R}$ , we have
- ▶ **additivity**:  $f(x_1 + x_2) = f(x_1) + f(x_2)$

# Linear Functions

- ▶ A function  $f : X \rightarrow Y$  between two linear spaces is **linear** if it preserves the linearity of the sets  $X$  and  $Y$ .
- ▶ For all  $x_1, x_2 \in X$  and  $\alpha \in \mathbb{R}$ , we have
- ▶ **additivity**:  $f(x_1 + x_2) = f(x_1) + f(x_2)$
- ▶ **homogeneity**:  $f(\alpha x_1) = \alpha f(x_1)$



## Examples

- ▶  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by  $f(x_1, x_2, x_3) = (x_1, x_2, 0)$  is a linear function

## Examples

- ▶  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by  $f(x_1, x_2, x_3) = (x_1, x_2, 0)$  is a linear function
- ▶ Additivity:

$$\begin{aligned} f(x_1 + y_1, x_2 + y_2, x_3 + y_3) &= (x_1 + y_1, x_2 + y_2, 0) \\ &= (x_1, x_2, 0) + (y_1, y_2, 0) \end{aligned}$$

## Examples

- ▶  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by  $f(x_1, x_2, x_3) = (x_1, x_2, 0)$  is a linear function
- ▶ Additivity:

$$\begin{aligned} f(x_1 + y_1, x_2 + y_2, x_3 + y_3) &= (x_1 + y_1, x_2 + y_2, 0) \\ &= (x_1, x_2, 0) + (y_1, y_2, 0) \end{aligned}$$

- ▶ Homogeneity:

$$f(\alpha x_1, \alpha x_2, \alpha x_3) = (\alpha x_1, \alpha x_2, 0) = \alpha(x_1, x_2, 0)$$

## Examples

- ▶  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by  $f(x_1, x_2, x_3) = (x_1, x_2, 0)$  is a linear function
- ▶ Additivity:

$$\begin{aligned} f(x_1 + y_1, x_2 + y_2, x_3 + y_3) &= (x_1 + y_1, x_2 + y_2, 0) \\ &= (x_1, x_2, 0) + (y_1, y_2, 0) \end{aligned}$$

- ▶ Homogeneity:

$$f(\alpha x_1, \alpha x_2, \alpha x_3) = (\alpha x_1, \alpha x_2, 0) = \alpha(x_1, x_2, 0)$$

- ▶  $\ln(x)$  is not a linear mapping:  $\ln(1) = 0$ ,  $\ln(2) \approx .7$ ,  $\ln(3) \approx 1.1$

## Example

- ▶ Can you show that  $f(x) = 2x + 3$  is a linear function?

## Example

- ▶ Can you show that  $f(x) = 2x + 3$  is a linear function?
- ▶ No.

## Example

- ▶ Can you show that  $f(x) = 2x + 3$  is a linear function?
- ▶ No.
- ▶  $f(x+y) = 2(x+y)+3 = 2x+2y+3 \neq 2x+2y+6 = f(x)+f(y)$

## Example

- ▶ Can you show that  $f(x) = 2x + 3$  is a linear function?
- ▶ No.
- ▶  $f(x+y) = 2(x+y)+3 = 2x+2y+3 \neq 2x+2y+6 = f(x)+f(y)$
- ▶  $f(\alpha x) = 2\alpha x + 3 \neq 2\alpha x + 3\alpha = \alpha f(x)$



## Example

- ▶ Can you show that  $f(x) = 2x + 3$  is a linear function?
- ▶ No.
- ▶  $f(x+y) = 2(x+y)+3 = 2x+2y+3 \neq 2x+2y+6 = f(x)+f(y)$
- ▶  $f(\alpha x) = 2\alpha x + 3 \neq 2\alpha x + 3\alpha = \alpha f(x)$
- ▶  $f$  is an example of an **affine function** which relate to linear functions in the same way as subspaces relate to affine sets

## Example

- ▶ Can you show that  $f(x) = 2x + 3$  is a linear function?
- ▶ No.
- ▶  $f(x+y) = 2(x+y)+3 = 2x+2y+3 \neq 2x+2y+6 = f(x)+f(y)$
- ▶  $f(\alpha x) = 2\alpha x + 3 \neq 2\alpha x + 3\alpha = \alpha f(x)$
- ▶  $f$  is an example of an **affine function** which relate to linear functions in the same way as subspaces relate to affine sets
- ▶ Affine functions preserve affine sets (lines and planes).

# Matrices as Linear Functions

- ▶ Any  $m \times n$  matrix  $A = (a_{ij})$  defines a linear mapping from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  defined by

$$f(x) = \begin{bmatrix} \sum_{j=1}^n a_{1j}x_j \\ \sum_{j=1}^n a_{2j}x_j \\ \dots \\ \sum_{j=1}^n a_{mj}x_j \end{bmatrix}$$

# Properties of Linear Functions

- ▶ Every linear function  $f : X \rightarrow Y$  maps the zero vector in  $X$  into the zero vector in  $Y$ .

# Properties of Linear Functions

- ▶ Every linear function  $f : X \rightarrow Y$  maps the zero vector in  $X$  into the zero vector in  $Y$ .
- ▶ Every composition of linear functions is linear.

# Properties of Linear Functions

- ▶ Every linear function  $f : X \rightarrow Y$  maps the zero vector in  $X$  into the zero vector in  $Y$ .
- ▶ Every composition of linear functions is linear.
- ▶ A linear function (such as a matrix) that has an inverse  $f^{-1} : Y \rightarrow X$  is said to be **nonsingular**. The inverse of a nonsingular linear function is also linear.

# Bilinear Functions

- ▶ A function  $f : X \times Y \rightarrow Z$  between linear spaces  $X$ ,  $Y$ , and  $Z$  is **bilinear** if it is linear in each factor separately:

$$f(x_1 + x_2, y) = f(x_1, y) + f(x_2, y)$$

$$f(x, y_1 + y_2) = f(x, y_1) + f(x, y_2)$$

$$f(\alpha x, y) = \alpha f(x, y) = f(x, \alpha y)$$

for all  $\alpha \in \mathbb{R}$

## Example

- ▶ The function  $f(x, y) = xy$  is bilinear:



## Example

- ▶ The function  $f(x, y) = xy$  is bilinear:
- ▶  $f(x_1 + x_2, y) = (x_1 + x_2)y = x_1y + x_2y = f(x_1, y) + f(x_2, y)$

## Example

- ▶ The function  $f(x, y) = xy$  is bilinear:
- ▶  $f(x_1 + x_2, y) = (x_1 + x_2)y = x_1y + x_2y = f(x_1, y) + f(x_2, y)$
- ▶  $f(\alpha x, y) = (\alpha x)y = \alpha xy = \alpha f(x, y)$

## Example

- ▶ Any  $m \times n$  matrix  $A = (a_{ij})$  defines a bilinear functional on  $\mathbb{R}^m \times \mathbb{R}^n$  by

$$f(x, y) = \sum_{i=1}^m \sum_{j=1}^n a_{ij} x_i y_j$$

## Example

- ▶ Any  $m \times n$  matrix  $A = (a_{ij})$  defines a bilinear functional on  $\mathbb{R}^m \times \mathbb{R}^n$  by

$$f(x, y) = \sum_{i=1}^m \sum_{j=1}^n a_{ij} x_i y_j$$

- ▶ In the quantitative sequence will often have an  $n \times n$  weighting matrix  $W$  and use the operation  $x^T W x$

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$
- ▶ **positive definite** if  $f(x, x) > 0$  for every  $x \in X$



# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$
- ▶ **positive definite** if  $f(x, x) > 0$  for every  $x \in X$
- ▶ A symmetric, positive definite bilinear functional on a linear space  $X$  is called an **inner product**, denoted  $x^T y$ .

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$
- ▶ **positive definite** if  $f(x, x) > 0$  for every  $x \in X$
- ▶ A symmetric, positive definite bilinear functional on a linear space  $X$  is called an **inner product**, denoted  $x^T y$ .
- ▶ A linear space equipped with an inner product is called an **inner product space**.

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$
- ▶ **positive definite** if  $f(x, x) > 0$  for every  $x \in X$
- ▶ A symmetric, positive definite bilinear functional on a linear space  $X$  is called an **inner product**, denoted  $x^T y$ .
- ▶ A linear space equipped with an inner product is called an **inner product space**.
- ▶ Every inner product defines a norm given by  $\|x\| = \sqrt{x^T x}$ .

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$
- ▶ **positive definite** if  $f(x, x) > 0$  for every  $x \in X$
- ▶ A symmetric, positive definite bilinear functional on a linear space  $X$  is called an **inner product**, denoted  $x^T y$ .
- ▶ A linear space equipped with an inner product is called an **inner product space**.
- ▶ Every inner product defines a norm given by  $\|x\| = \sqrt{x^T x}$ .
- ▶ Therefore every inner product space is a normed linear space

# Bilinear Functionals and Inner Product Spaces

- ▶ A bilinear functional  $f$  on  $X \times X$  is called
- ▶ **symmetric** if  $f(x, y) = f(y, x)$  for all  $x, y \in X$
- ▶ **positive semidefinite** if  $f(x, x) \geq 0$  for every  $x \in X$
- ▶ **positive definite** if  $f(x, x) > 0$  for every  $x \in X$
- ▶ A symmetric, positive definite bilinear functional on a linear space  $X$  is called an **inner product**, denoted  $x^T y$ .
- ▶ A linear space equipped with an inner product is called an **inner product space**.
- ▶ Every inner product defines a norm given by  $\|x\| = \sqrt{x^T x}$ .
- ▶ Therefore every inner product space is a normed linear space
- ▶ We spend most of our time in Euclidean space  $\mathbb{R}^n$  with the inner product  $x^T y = \sum_{i=1}^n x_i y_i$ .

# Inner Product Space Properties

- ▶ In any inner product space, the **Cauchy-Schwarz inequality** holds:

$$|x^T y| \leq \|x\| \|y\|$$

# Inner Product Space Properties

- ▶ In any inner product space, the **Cauchy-Schwarz inequality** holds:

$$|x^T y| \leq \|x\| \|y\|$$

- ▶ A nonempty compact convex set in an inner product space has at least one extreme point.

# Linear Operators

- ▶ A **linear operator** is a linear function  $f : X \rightarrow X$ .



# Linear Operators

- ▶ A **linear operator** is a linear function  $f : X \rightarrow X$ .
- ▶ Every linear operator on a finite-dimensional space can be represented by a square matrix.

# Linear Operators

- ▶ A **linear operator** is a linear function  $f : X \rightarrow X$ .
- ▶ Every linear operator on a finite-dimensional space can be represented by a square matrix.
- ▶ Every linear operator  $f : X \rightarrow X$  has at least one fixed point.

# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .

# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .
- ▶ If  $X$  is finite-dimensional, there is a unique functional  $\det$  called the **determinant** on  $L(X, X)$  with the following properties:

# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .
- ▶ If  $X$  is finite-dimensional, there is a unique functional  $\det$  called the **determinant** on  $L(X, X)$  with the following properties:
- ▶  $\det(f \circ g) = \det(f) \det(g)$  for all  $g, f \in L(X, X)$

# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .
- ▶ If  $X$  is finite-dimensional, there is a unique functional  $\det$  called the **determinant** on  $L(X, X)$  with the following properties:
  - ▶  $\det(f \circ g) = \det(f) \det(g)$  for all  $g, f \in L(X, X)$
  - ▶  $\det(I) = 1$  where  $I$  is the identity matrix

# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .
- ▶ If  $X$  is finite-dimensional, there is a unique functional  $\det$  called the **determinant** on  $L(X, X)$  with the following properties:
  - ▶  $\det(f \circ g) = \det(f) \det(g)$  for all  $g, f \in L(X, X)$
  - ▶  $\det(I) = 1$  where  $I$  is the identity matrix
  - ▶  $\det(f) = 0$  if and only if  $f$  is nonsingular

# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .
- ▶ If  $X$  is finite-dimensional, there is a unique functional  $\det$  called the **determinant** on  $L(X, X)$  with the following properties:
  - ▶  $\det(f \circ g) = \det(f) \det(g)$  for all  $g, f \in L(X, X)$
  - ▶  $\det(I) = 1$  where  $I$  is the identity matrix
  - ▶  $\det(f) = 0$  if and only if  $f$  is nonsingular
- ▶ The last property will be particularly useful when we want to check whether an operator is nonsingular.



# Linear Operators and the Determinant

- ▶ The set of all linear operators on a given space  $X$  is denoted  $L(X, X)$ .
- ▶ If  $X$  is finite-dimensional, there is a unique functional  $\det$  called the **determinant** on  $L(X, X)$  with the following properties:
  - ▶  $\det(f \circ g) = \det(f) \det(g)$  for all  $g, f \in L(X, X)$
  - ▶  $\det(I) = 1$  where  $I$  is the identity matrix
  - ▶  $\det(f) = 0$  if and only if  $f$  is nonsingular
  - ▶ The last property will be particularly useful when we want to check whether an operator is nonsingular.
- ▶  $\det$  itself is not a linear functional

# Quadratic Forms

- ▶ Let  $X$  be a Euclidean space.

# Quadratic Forms

- ▶ Let  $X$  be a Euclidean space.
- ▶ A functional  $Q : X \rightarrow \mathbb{R}$  is called a **quadratic form** if there exists a symmetric linear operator  $f : X \rightarrow X$  such that  $Q(x) = x^T f(x)$  for every  $x \in X$ .

# Quadratic Forms

- ▶ Let  $X$  be a Euclidean space.
- ▶ A functional  $Q : X \rightarrow \mathbb{R}$  is called a **quadratic form** if there exists a symmetric linear operator  $f : X \rightarrow X$  such that  $Q(x) = x^T f(x)$  for every  $x \in X$ .
- ▶ Quadratic forms are among the simplest nonlinear functionals we encounter.

## Example

- ▶ The nonlinear function  $Q(x, y) = x^2 + 4xy + y^2$  is a quadratic form on  $\mathbb{R}^2$ .

## Example

- ▶ The nonlinear function  $Q(x, y) = x^2 + 4xy + y^2$  is a quadratic form on  $\mathbb{R}^2$ .
- ▶ The matrix

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

defines a symmetric linear operator on  $\mathbb{R}^2$ :

$$f(x, y) = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ 2x + y \end{bmatrix}$$

$$[x, y] \begin{bmatrix} x + 2y \\ 2x + y \end{bmatrix} = x^2 + 4xy + y^2 = Q(x, y)$$

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .
- ▶ A symmetric matrix  $A$  is called



# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .
- ▶ A symmetric matrix  $A$  is called
- ▶ **positive definite** if  $x^T A x > 0$  for all  $x \neq 0 \in X$

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .
- ▶ A symmetric matrix  $A$  is called
- ▶ **positive definite** if  $x^T A x > 0$  for all  $x \neq 0 \in X$
- ▶ **positive semidefinite** if  $x^T A x \geq 0$  for all  $x \neq 0 \in X$

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .
- ▶ A symmetric matrix  $A$  is called
- ▶ **positive definite** if  $x^T A x > 0$  for all  $x \neq 0 \in X$
- ▶ **positive semidefinite** if  $x^T A x \geq 0$  for all  $x \neq 0 \in X$
- ▶ **negative definite** if  $x^T A x < 0$  for all  $x \neq 0 \in X$

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .
- ▶ A symmetric matrix  $A$  is called
- ▶ **positive definite** if  $x^T A x > 0$  for all  $x \neq 0 \in X$
- ▶ **positive semidefinite** if  $x^T A x \geq 0$  for all  $x \neq 0 \in X$
- ▶ **negative definite** if  $x^T A x < 0$  for all  $x \neq 0 \in X$
- ▶ **negative semidefinite** if  $x^T A x \leq 0$  for all  $x \neq 0 \in X$

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T A x$ .
- ▶ A symmetric matrix  $A$  is called
- ▶ **positive definite** if  $x^T A x > 0$  for all  $x \neq 0 \in X$
- ▶ **positive semidefinite** if  $x^T A x \geq 0$  for all  $x \neq 0 \in X$
- ▶ **negative definite** if  $x^T A x < 0$  for all  $x \neq 0 \in X$
- ▶ **negative semidefinite** if  $x^T A x \leq 0$  for all  $x \neq 0 \in X$
- ▶ All definite matrices are nonsingular.

# Quadratic Forms

- ▶ Any  $n \times n$  symmetric matrix,  $A$ , defines a quadratic form  $Q(x) = x^T Ax$ .
- ▶ A symmetric matrix  $A$  is called
- ▶ **positive definite** if  $x^T Ax > 0$  for all  $x \neq 0 \in X$
- ▶ **positive semidefinite** if  $x^T Ax \geq 0$  for all  $x \neq 0 \in X$
- ▶ **negative definite** if  $x^T Ax < 0$  for all  $x \neq 0 \in X$
- ▶ **negative semidefinite** if  $x^T Ax \leq 0$  for all  $x \neq 0 \in X$
- ▶ All definite matrices are nonsingular.
- ▶ A positive (negative) definite matrix has a positive (negative) diagonal.

# Convex Functions

- ▶ For many purposes in political science, linearity is too restrictive of an assumption.

# Convex Functions

- ▶ For many purposes in political science, linearity is too restrictive of an assumption.
- ▶ Convex functions generalize some of the properties of linear functions while providing more suitable functional forms.



# Convex Functions

- ▶ For many purposes in political science, linearity is too restrictive of an assumption.
- ▶ Convex functions generalize some of the properties of linear functions while providing more suitable functional forms.
- ▶ A real-valued function  $f$  defined on a convex set of a linear space  $X$  is **convex** if for every  $x_1, x_2 \in S$ ,

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \alpha f(x_1) + (1 - \alpha)f(x_2)$$


# Convex Functions

- ▶ For many purposes in political science, linearity is too restrictive of an assumption.
- ▶ Convex functions generalize some of the properties of linear functions while providing more suitable functional forms.
- ▶ A real-valued function  $f$  defined on a convex set of a linear space  $X$  is **convex** if for every  $x_1, x_2 \in S$ ,

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \alpha f(x_1) + (1 - \alpha)f(x_2)$$

- ▶ A function is **strictly convex** if the inequality is strict and **concave** if the inequality is reversed.

## Example



# Properties of Convex Functions

- ▶ If  $f$  is an invertible function, then  $f$  is concave if and only if  $f^{-1}$  is convex

# Properties of Convex Functions

- ▶ If  $f$  is an invertible function, then  $f$  is concave if and only if  $f^{-1}$  is convex
- ▶ If  $f$  and  $g$  are convex,  $f + g$  is convex

# Properties of Convex Functions

- ▶ If  $f$  is an invertible function, then  $f$  is concave if and only if  $f^{-1}$  is convex
- ▶ If  $f$  and  $g$  are convex,  $f + g$  is convex
- ▶ If  $f$  is convex, then  $\alpha f$  is convex for every  $\alpha \geq 0$

# Properties of Convex Functions

- ▶ If  $f$  is an invertible function, then  $f$  is concave if and only if  $f^{-1}$  is convex
- ▶ If  $f$  and  $g$  are convex,  $f + g$  is convex
- ▶ If  $f$  is convex, then  $\alpha f$  is convex for every  $\alpha \geq 0$
- ▶ Logarithmic transformations preserve concavity because  $\ln$  is concave and increasing

# Properties of Convex Functions

- ▶ If  $f$  is an invertible function, then  $f$  is concave if and only if  $f^{-1}$  is convex
- ▶ If  $f$  and  $g$  are convex,  $f + g$  is convex
- ▶ If  $f$  is convex, then  $\alpha f$  is convex for every  $\alpha \geq 0$
- ▶ Logarithmic transformations preserve concavity because  $\ln$  is concave and increasing
- ▶ For a twice differentiable function on an open interval  $S \subseteq \mathbb{R}$ ,  $f : S \rightarrow \mathbb{R}$  is convex if and only if  $f''(x) \geq 0$  for every  $x \in S$  and concave if and only if  $f''(x) \leq 0$  for every  $x \in S$ .



## Second Derivative and Convexity in $\mathbb{R}^n$

- ▶ In the next lecture, we will see how to differentiate functions defined on higher dimensional domains.

## Second Derivative and Convexity in $\mathbb{R}^n$

- ▶ In the next lecture, we will see how to differentiate functions defined on higher dimensional domains.
- ▶ The generalization of the second derivative on a one dimensional domain is given by a symmetric square matrix that is referred to as a **Hessian**,  $H_f$ .

## Second Derivative and Convexity in $\mathbb{R}^n$

- ▶ In the next lecture, we will see how to differentiate functions defined on higher dimensional domains.
- ▶ The generalization of the second derivative on a one dimensional domain is given by a symmetric square matrix that is referred to as a **Hessian**,  $H_f$ .
- ▶ In higher dimensions, we would like to exploit the sign of the derivative to identify whether a function is convex or concave at some point.

## Second Derivative and Convexity in $\mathbb{R}^n$

- ▶ In the next lecture, we will see how to differentiate functions defined on higher dimensional domains.
- ▶ The generalization of the second derivative on a one dimensional domain is given by a symmetric square matrix that is referred to as a **Hessian**,  $H_f$ .
- ▶ In higher dimensions, we would like to exploit the sign of the derivative to identify whether a function is convex or concave at some point.
- ▶ The concept of definiteness that we saw earlier will generalize the idea of positive and negative into higher dimensions.

## Second Derivative and Convexity in $\mathbb{R}^n$

- ▶ In the next lecture, we will see how to differentiate functions defined on higher dimensional domains.
- ▶ The generalization of the second derivative on a one dimensional domain is given by a symmetric square matrix that is referred to as a **Hessian**,  $H_f$ .
- ▶ In higher dimensions, we would like to exploit the sign of the derivative to identify whether a function is convex or concave at some point.
- ▶ The concept of definiteness that we saw earlier will generalize the idea of positive and negative into higher dimensions.
- ▶  $H_f$  defines a quadratic form on  $\mathbb{R}^n$ :  $x^T H_f x \equiv H_f(x)$

## Second Derivative and Convexity in $\mathbb{R}^n$

- ▶ In the next lecture, we will see how to differentiate functions defined on higher dimensional domains.
- ▶ The generalization of the second derivative on a one dimensional domain is given by a symmetric square matrix that is referred to as a **Hessian**,  $H_f$ .
- ▶ In higher dimensions, we would like to exploit the sign of the derivative to identify whether a function is convex or concave at some point.
- ▶ The concept of definiteness that we saw earlier will generalize the idea of positive and negative into higher dimensions.
- ▶  $H_f$  defines a quadratic form on  $\mathbb{R}^n$ :  $x^T H_f x \equiv H_f(x)$
- ▶  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is convex at  $x$  iff and only if  $H_f(x)$  is positive semidefinite and concave iff  $H_f(x)$  is negative semidefinite.

## Quasiconcavity

- ▶ A functional  $f$  on a convex set  $S$  of a linear space is **quasiconvex** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \max \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

# Quasiconcavity

- ▶ A functional  $f$  on a convex set  $S$  of a linear space is **quasiconvex** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \max \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ A function is **strictly quasiconvex** if the inequality is strict



# Quasiconcavity

- ▶ A functional  $f$  on a convex set  $S$  of a linear space is **quasiconvex** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \max \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ A function is **strictly quasiconvex** if the inequality is strict
- ▶ A function is **quasiconcave** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \geq \min \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

# Quasiconcavity

- ▶ A functional  $f$  on a convex set  $S$  of a linear space is **quasiconvex** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \max \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ A function is **strictly quasiconvex** if the inequality is strict
- ▶ A function is **quasiconcave** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \geq \min \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ We often encounter quasiconcave functions rather than quasiconvex functions.

# Quasiconcavity

- ▶ A functional  $f$  on a convex set  $S$  of a linear space is **quasiconvex** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \max \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ A function is **strictly quasiconvex** if the inequality is strict
- ▶ A function is **quasiconcave** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \geq \min \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ We often encounter quasiconcave functions rather than quasiconvex functions.
- ▶ Geometrically a function is quasiconcave if the function along a line joining any two points in the domain lies above at least one of the endpoints.

# Quasiconcavity

- ▶ A functional  $f$  on a convex set  $S$  of a linear space is **quasiconvex** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \leq \max \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ A function is **strictly quasiconvex** if the inequality is strict
- ▶ A function is **quasiconcave** if

$$f(\alpha x_1 + (1 - \alpha)x_2) \geq \min \{f(x_1), f(x_2)\}$$

for every  $x_1, x_2 \in S$  and  $\alpha \in [0, 1]$

- ▶ We often encounter quasiconcave functions rather than quasiconvex functions.
- ▶ Geometrically a function is quasiconcave if the function along a line joining any two points in the domain lies above at least one of the endpoints.
- ▶ Any monotone function is both quasiconcave and quasiconvex.

## Example

