

# Comparative Statics

# Motivation

- ▶ We are often particularly interested in how the behavior of a set of actors changes as some feature of the political environment changes.

# Motivation

- ▶ We are often particularly interested in how the behavior of a set of actors changes as some feature of the political environment changes.
- ▶ “H1: A rise in  $X$  leads to a fall in  $Y$ ”

# Motivation

- ▶ We are often particularly interested in how the behavior of a set of actors changes as some feature of the political environment changes.
- ▶ “H1: A rise in  $X$  leads to a fall in  $Y$ ”
- ▶ Before asserting “H1” and going to the data, it is necessary first to demonstrate that a hypothesized relationship between two variables follows logically from a set of explicit assumptions.

# Motivation

- ▶ We are often particularly interested in how the behavior of a set of actors changes as some feature of the political environment changes.
- ▶ “H1: A rise in  $X$  leads to a fall in  $Y$ ”
- ▶ Before asserting “H1” and going to the data, it is necessary first to demonstrate that a hypothesized relationship between two variables follows logically from a set of explicit assumptions.
- ▶ This exercise is referred to as “comparative statics.”

# Parameterized Optimization

- ▶ Let's suppose we want to build a theory of how much time a member of Congress spends fundraising.

# Parameterized Optimization

- ▶ Let's suppose we want to build a theory of how much time a member of Congress spends fundraising.
- ▶ What does the member want?

# Parameterized Optimization

- ▶ Let's suppose we want to build a theory of how much time a member of Congress spends fundraising.
- ▶ What does the member want?
- ▶ How does his choice of fundraising affect his achievement of that?



# Parameterized Optimization

- ▶ Let's suppose we want to build a theory of how much time a member of Congress spends fundraising.
- ▶ What does the member want?
- ▶ How does his choice of fundraising affect his achievement of that?
- ▶ This will presumably depend on some feature of the political environment.

# Parameterized Optimization

- ▶ Let's suppose we want to build a theory of how much time a member of Congress spends fundraising.
- ▶ What does the member want?
- ▶ How does his choice of fundraising affect his achievement of that?
- ▶ This will presumably depend on some feature of the political environment.
- ▶ These assumptions yield a **parameterized objective function**,  $f(x; \theta)$

# Parameterized Optimization

- ▶ Let's suppose we want to build a theory of how much time a member of Congress spends fundraising.
- ▶ What does the member want?
- ▶ How does his choice of fundraising affect his achievement of that?
- ▶ This will presumably depend on some feature of the political environment.
- ▶ These assumptions yield a **parameterized objective function**,  $f(x; \theta)$
- ▶ To find out how much time a legislator spends fundraising given  $\theta$ , we solve

$$\max_x f(x; \theta)$$

## Parameterized Optimization

- ▶ We are ultimately interested in how something in  $\theta$  such as electoral danger affects time spent fundraising.

# Parameterized Optimization

- ▶ We are ultimately interested in how something in  $\theta$  such as electoral danger affects time spent fundraising.
- ▶ This will allow us to derive a hypothesis that we then test in the empirical portion of the paper.

# Parameterized Optimization

- ▶ We are ultimately interested in how something in  $\theta$  such as electoral danger affects time spent fundraising.
- ▶ This will allow us to derive a hypothesis that we then test in the empirical portion of the paper.
- ▶ Formally, we are interested how

$$x^*(\theta) \equiv \arg \max_x f(x; \theta)$$

changes with  $\theta$

# Parameterized Optimization

- ▶ We are ultimately interested in how something in  $\theta$  such as electoral danger affects time spent fundraising.
- ▶ This will allow us to derive a hypothesis that we then test in the empirical portion of the paper.
- ▶ Formally, we are interested how

$$x^*(\theta) \equiv \arg \max_x f(x; \theta)$$

changes with  $\theta$

- ▶ That is, we want to know the sign of  $\frac{\partial}{\partial \theta} x^*(\theta)$ .

# Parameterized Optimization

- ▶ We are ultimately interested in how something in  $\theta$  such as electoral danger affects time spent fundraising.
- ▶ This will allow us to derive a hypothesis that we then test in the empirical portion of the paper.
- ▶ Formally, we are interested how

$$x^*(\theta) \equiv \arg \max_x f(x; \theta)$$

changes with  $\theta$

- ▶ That is, we want to know the sign of  $\frac{\partial}{\partial \theta} x^*(\theta)$ .
- ▶ The set of maximizers,  $x^*(\theta)$ , is referred to as the **solution correspondence**.



# Parameterized Optimization

- ▶ If we have a closed-form solution for  $x^*(\theta)$ , finding how  $x^*(\theta)$  changes in  $\theta$  is very straightforward.

# Parameterized Optimization

- ▶ If we have a closed-form solution for  $x^*(\theta)$ , finding how  $x^*(\theta)$  changes in  $\theta$  is very straightforward.
- ▶ For example, if

$$f(x; \theta) = \theta \ln(x) - x^2$$

for  $\theta \geq 0$ , then by first order conditions,

$$\frac{\theta}{x} - 2x = 0$$

which yields

$$x^*(\theta) = \sqrt{\frac{\theta}{2}}$$

# Parameterized Optimization

- ▶ If we have a closed-form solution for  $x^*(\theta)$ , finding how  $x^*(\theta)$  changes in  $\theta$  is very straightforward.
- ▶ For example, if

$$f(x; \theta) = \theta \ln(x) - x^2$$

for  $\theta \geq 0$ , then by first order conditions,

$$\frac{\theta}{x} - 2x = 0$$

which yields

$$x^*(\theta) = \sqrt{\frac{\theta}{2}}$$

- ▶ We see clearly then that  $x^*(\theta)$  is increasing in  $\theta$ .

# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.

## Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?

# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?

## Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?
- ▶ Legislator pays cost that increases in time spent fundraising (at an increasing rate).

# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?
- ▶ Legislator pays cost that increases in time spent fundraising (at an increasing rate).
- ▶ Each hour spent fundraising more valuable in close races.



# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?
- ▶ Legislator pays cost that increases in time spent fundraising (at an increasing rate).
- ▶ Each hour spent fundraising more valuable in close races.
- ▶ Utility from funds raised increasing with diminishing returns.

# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?
- ▶ Legislator pays cost that increases in time spent fundraising (at an increasing rate).
- ▶ Each hour spent fundraising more valuable in close races.
- ▶ Utility from funds raised increasing with diminishing returns.
- ▶ This then gives us a general utility function

$$f(x; \theta) = v(x; \theta) - c(x)$$

# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?
- ▶ Legislator pays cost that increases in time spent fundraising (at an increasing rate).
- ▶ Each hour spent fundraising more valuable in close races.
- ▶ Utility from funds raised increasing with diminishing returns.
- ▶ This then gives us a general utility function

$$f(x; \theta) = v(x; \theta) - c(x)$$

- ▶ Now we cannot find a closed-form solution for  $x^*(\theta)$  to help us.

# Parameterized Optimization

- ▶ Typically we want to make weak and general assumptions.
- ▶ It is reasonable to believe that  $f(x; \theta) = \theta \ln(x) - x^2$ ?
- ▶ What is?
- ▶ Legislator pays cost that increases in time spent fundraising (at an increasing rate).
- ▶ Each hour spent fundraising more valuable in close races.
- ▶ Utility from funds raised increasing with diminishing returns.
- ▶ This then gives us a general utility function

$$f(x; \theta) = v(x; \theta) - c(x)$$

- ▶ Now we cannot find a closed-form solution for  $x^*(\theta)$  to help us.
- ▶ Fortunately, we have an extremely powerful tool to help us with this problem.

# Implicit Function Theorem

- ▶ The **implicit function theorem** states that as long as a certain condition is satisfied, we can use differential calculus to characterize how  $x^*(\theta)$  varies in response to small changes in  $\theta$ .

# Implicit Function Theorem (Light)

## Theorem

*Let  $x^* \in \mathbb{R}$  solve  $g(x, y) = 0$  at  $y^* \in \mathbb{R}$ . If  $g(\cdot, \cdot)$  is continuously differentiable and  $\frac{\partial}{\partial x}g(x^*, y^*) \neq 0$  then for some open set  $A$  containing  $x^*$  and an open set  $B$  containing  $y^*$ , there exists a continuously differentiable function  $h : B \rightarrow A$  with  $g(h(y), y) = 0$ . The derivative of this function at  $y^*$  is given by*

$$\frac{\partial h(y^*)}{\partial y} = -\left(\frac{\partial g(x^*, y^*)}{\partial y}\right)\left(\frac{\partial g(x^*, y^*)}{\partial x}\right)^{-1}$$

# Implicit Function Theorem

- ▶ To see how this applies to comparative statics, note that the first-order conditions of a parameterized optimization problem give us the condition  $g(x, y) = 0$

# Implicit Function Theorem

- ▶ To see how this applies to comparative statics, note that the first-order conditions of a parameterized optimization problem give us the condition  $g(x, y) = 0$
- ▶ The problem must have a solution to begin with.



# Implicit Function Theorem

- ▶ To see how this applies to comparative statics, note that the first-order conditions of a parameterized optimization problem give us the condition  $g(x, y) = 0$
- ▶ The problem must have a solution to begin with.
- ▶ In the next lecture we will explore how verify that a general program has a solution.

# Implicit Function Theorem

- ▶ To see how this applies to comparative statics, note that the first-order conditions of a parameterized optimization problem give us the condition  $g(x, y) = 0$
- ▶ The problem must have a solution to begin with.
- ▶ In the next lecture we will explore how verify that a general program has a solution.
- ▶ Here we assume that all problems have solutions, i.e. that  $x^*(\theta)$  is always nonempty.

## Example

- ▶ Returning to the example, we want to solve

$$\max_x v(x; \theta) - c(x)$$

## Example

- ▶ Returning to the example, we want to solve

$$\max_x v(x; \theta) - c(x)$$

- ▶ From FOCs,  $\frac{\partial f(x^*; \theta)}{\partial x} = 0$

## Example

- ▶ Returning to the example, we want to solve

$$\max_x v(x; \theta) - c(x)$$

- ▶ From FOCs,  $\frac{\partial f(x^*; \theta)}{\partial x} = 0$
- ▶ Now as long as  $\frac{\partial^2 f(x^*; \theta)}{\partial x^2} \neq 0$ , then

$$\begin{aligned} \frac{\partial x^*(\theta)}{\partial \theta} = \\ -\frac{\partial^2 f(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 f(x; \theta)}{\partial x^2} \right)^{-1} = -\frac{\partial^2 v(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 v(x; \theta)}{\partial x^2} - \frac{\partial^2 c(x)}{\partial x^2} \right)^{-1} \end{aligned}$$

## Example

- ▶ Returning to the example, we want to solve

$$\max_x v(x; \theta) - c(x)$$

- ▶ From FOCs,  $\frac{\partial f(x^*; \theta)}{\partial x} = 0$
- ▶ Now as long as  $\frac{\partial^2 f(x^*; \theta)}{\partial x^2} \neq 0$ , then

$$\begin{aligned} \frac{\partial x^*(\theta)}{\partial \theta} = \\ -\frac{\partial^2 f(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 f(x; \theta)}{\partial x^2} \right)^{-1} = -\frac{\partial^2 v(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 v(x; \theta)}{\partial x^2} - \frac{\partial^2 c(x)}{\partial x^2} \right)^{-1} \end{aligned}$$

- ▶ We assumed that  $v(x; \theta)$  is increasing and concave in  $x$  so  $\frac{\partial^2 v(x; \theta)}{\partial x^2} < 0$ .

## Example

- ▶ Returning to the example, we want to solve

$$\max_x v(x; \theta) - c(x)$$

- ▶ From FOCs,  $\frac{\partial f(x^*; \theta)}{\partial x} = 0$
- ▶ Now as long as  $\frac{\partial^2 f(x^*; \theta)}{\partial x^2} \neq 0$ , then

$$\begin{aligned} \frac{\partial x^*(\theta)}{\partial \theta} = \\ -\frac{\partial^2 f(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 f(x; \theta)}{\partial x^2} \right)^{-1} = -\frac{\partial^2 v(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 v(x; \theta)}{\partial x^2} - \frac{\partial^2 c(x)}{\partial x^2} \right)^{-1} \end{aligned}$$

- ▶ We assumed that  $v(x; \theta)$  is increasing and concave in  $x$  so  $\frac{\partial^2 v(x; \theta)}{\partial x^2} < 0$ .
- ▶ If we assume that marginal cost is non-decreasing in  $x$ , then  $\frac{\partial^2 c(x)}{\partial x^2} \geq 0$

## Example

- ▶ Returning to the example, we want to solve

$$\max_x v(x; \theta) - c(x)$$

- ▶ From FOCs,  $\frac{\partial f(x^*; \theta)}{\partial x} = 0$
- ▶ Now as long as  $\frac{\partial^2 f(x^*; \theta)}{\partial x^2} \neq 0$ , then

$$\begin{aligned} \frac{\partial x^*(\theta)}{\partial \theta} = \\ -\frac{\partial^2 f(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 f(x; \theta)}{\partial x^2} \right)^{-1} = -\frac{\partial^2 v(x; \theta)}{\partial x \partial \theta} \left( \frac{\partial^2 v(x; \theta)}{\partial x^2} - \frac{\partial^2 c(x)}{\partial x^2} \right)^{-1} \end{aligned}$$

- ▶ We assumed that  $v(x; \theta)$  is increasing and concave in  $x$  so  $\frac{\partial^2 v(x; \theta)}{\partial x^2} < 0$ .
- ▶ If we assume that marginal cost is non-decreasing in  $x$ , then  $\frac{\partial^2 c(x)}{\partial x^2} \geq 0$
- ▶ The sign of  $\frac{\partial^2 v(x; \theta)}{\partial x \partial \theta}$  will therefore tell us how  $x^*(\theta)$  changes with  $\theta$ .



# Implicit Function Theorem

The implicit function theorem generalizes to higher dimensions with  $n$  choice variables and  $m$  parameters.

## Theorem

*Let  $x^* \in \mathbb{R}^n$  solve  $f(x, y) = 0$  at  $y \in \mathbb{R}^m$ . If  $f_1(\cdot)$  through  $f_n(\cdot)$  are continuously differentiable in each coordinate of  $x$  and  $y$  and the Jacobian matrix with respect to the endogenous variables is nonsingular, then for some open set  $A$  containing  $x^*$  and an open set  $B$  containing  $y^*$ , there exists a continuously differentiable function  $\phi : B \rightarrow A$  with  $f(\phi(y), y) = 0$ . The derivative of this function at  $y^*$  is given by the  $n \times m$  matrix*

$$D_y \phi(y^*) = -[D_x f(x^*, y^*)]^{-1} D_y f(x^*, y^*)$$

# Implicit Function Theorem

- ▶ Returning to our application, let  $x = (x_1, x_2, \dots, x_n)$  and  $\theta = (\theta_1, \theta_2, \dots, \theta_m)$ .

# Implicit Function Theorem

- ▶ Returning to our application, let  $x = (x_1, x_2, \dots, x_n)$  and  $\theta = (\theta_1, \theta_2, \dots, \theta_m)$ .
- ▶ We are interested in finding the  $m \times n$  Jacobian matrix of  $x^*(\theta)$ ,  $D_\theta x^*(\theta)$

# Implicit Function Theorem

- ▶ Returning to our application, let  $x = (x_1, x_2, \dots, x_n)$  and  $\theta = (\theta_1, \theta_2, \dots, \theta_m)$ .
- ▶ We are interested in finding the  $m \times n$  Jacobian matrix of  $x^*(\theta)$ ,  $D_\theta x^*(\theta)$
- ▶ The implicit function theorem tells us that

$$D_\theta x^*(\theta) = -[H_x f(x, \theta)]^{-1} D_\theta f(x, \theta)$$

where  $H_x f(x^*, \theta)$  is the  $n \times n$  Hessian matrix of  $f(x^*, \theta)$  with respect to the choice variables and  $D_\theta f(x^*, \theta)$  is the  $n \times m$  Jacobian of  $f(x^*, \theta)$  with respect to the parameters.

# Implicit Function Theorem

- ▶ Returning to our application, let  $x = (x_1, x_2, \dots, x_n)$  and  $\theta = (\theta_1, \theta_2, \dots, \theta_m)$ .
- ▶ We are interested in finding the  $m \times n$  Jacobian matrix of  $x^*(\theta)$ ,  $D_\theta x^*(\theta)$
- ▶ The implicit function theorem tells us that

$$D_\theta x^*(\theta) = -[H_x f(x, \theta)]^{-1} D_\theta f(x, \theta)$$

where  $H_x f(x^*, \theta)$  is the  $n \times n$  Hessian matrix of  $f(x^*, \theta)$  with respect to the choice variables and  $D_\theta f(x^*, \theta)$  is the  $n \times m$  Jacobian of  $f(x^*, \theta)$  with respect to the parameters.

- ▶ Note that the implicit function theorem requires that  $H$  be nonsingular.

## Example

- Let

$$f(x, y; \theta) = v(x) + \theta w(y) - c(x, y)$$

where  $v(x)$  and  $w(y)$  are increasing and concave and  $c(x, y)$  is increasing and convex in both arguments.

## Example

- Let

$$f(x, y; \theta) = v(x) + \theta w(y) - c(x, y)$$

where  $v(x)$  and  $w(y)$  are increasing and concave and  $c(x, y)$  is increasing and convex in both arguments.

- We want to find  $\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix}$

## Example

- ▶ Let

$$f(x, y; \theta) = v(x) + \theta w(y) - c(x, y)$$

where  $v(x)$  and  $w(y)$  are increasing and concave and  $c(x, y)$  is increasing and convex in both arguments.

- ▶ We want to find  $\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix}$
- ▶ By the implicit function theorem,

$$\begin{aligned} \begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} &= - \begin{bmatrix} \frac{\partial^2 f(x, y; \theta)}{\partial x^2} & \frac{\partial^2 f(x, y; \theta)}{\partial x \partial y} \\ \frac{\partial^2 f(x, y; \theta)}{\partial x \partial y} & \frac{\partial^2 f(x, y; \theta)}{\partial y^2} \end{bmatrix}^{-1} \begin{bmatrix} \frac{\partial^2 f(x, y; \theta)}{\partial x \partial \theta} \\ \frac{\partial^2 f(x, y; \theta)}{\partial y \partial \theta} \end{bmatrix} \\ &= - \begin{bmatrix} \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x, y)}{\partial x^2} & -\frac{\partial^2 c(x, y)}{\partial x \partial y} \\ -\frac{\partial^2 c(x, y)}{\partial x \partial y} & \frac{\partial^2 \theta w(y)}{\partial y^2} - \frac{\partial^2 c(x, y)}{\partial y^2} \end{bmatrix}^{-1} \begin{bmatrix} 0 \\ \frac{\partial w(y)}{\partial y} \end{bmatrix} \end{aligned}$$



## Example

$$\begin{aligned}
 & - \begin{bmatrix} \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} & -\frac{\partial^2 c(x,y)}{\partial x \partial y} \\ -\frac{\partial^2 c(x,y)}{\partial x \partial y} & \frac{\partial^2 \theta w(y)}{\partial y^2} - \frac{\partial^2 c(x,y)}{\partial y^2} \end{bmatrix}^{-1} \begin{bmatrix} 0 \\ \frac{\partial w(y)}{\partial y} \end{bmatrix} \\
 &= -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 \theta w(y)}{\partial y^2} - \frac{\partial^2 c(x,y)}{\partial y^2} & \frac{\partial^2 c(x,y)}{\partial x \partial y} \\ \frac{\partial^2 c(x,y)}{\partial x \partial y} & \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \end{bmatrix} \begin{bmatrix} 0 \\ \frac{\partial w(y)}{\partial y} \end{bmatrix} \\
 &= -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}
 \end{aligned}$$

## Example

$$\blacktriangleright \begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$

## Example

- ▶ 
$$\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$
- ▶ By second order conditions,  $\det(H) > 0$ .

## Example

- ▶ 
$$\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$
- ▶ By second order conditions,  $\det(H) > 0$ .
- ▶ By the concavity of  $v(\cdot)$  and convexity of  $c(\cdot, \cdot)$ ,  $\left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) < 0$ .

## Example

- ▶ 
$$\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$
- ▶ By second order conditions,  $\det(H) > 0$ .
- ▶ By the concavity of  $v(\cdot)$  and convexity of  $c(\cdot, \cdot)$ ,  $\left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) < 0$ .
- ▶ By assumption  $\frac{\partial w(y)}{\partial y} > 0$ .

## Example

- ▶ 
$$\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$
- ▶ By second order conditions,  $\det(H) > 0$ .
- ▶ By the concavity of  $v(\cdot)$  and convexity of  $c(\cdot, \cdot)$ ,  $\left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) < 0$ .
- ▶ By assumption  $\frac{\partial w(y)}{\partial y} > 0$ .
- ▶ Therefore  $y^*(\theta) > 0$ .

## Example

- ▶ 
$$\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$
- ▶ By second order conditions,  $\det(H) > 0$ .
- ▶ By the concavity of  $v(\cdot)$  and convexity of  $c(\cdot, \cdot)$ ,  
$$\left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) < 0.$$
- ▶ By assumption  $\frac{\partial w(y)}{\partial y} > 0$ .
- ▶ Therefore  $y^*(\theta) > 0$ .
- ▶ We have not yet made any assumptions about  $\frac{\partial^2 c(x,y)}{\partial x \partial y}$ .

## Example

- ▶ 
$$\begin{bmatrix} \frac{\partial x^*(\theta)}{\partial \theta} \\ \frac{\partial y^*(\theta)}{\partial \theta} \end{bmatrix} = -\frac{1}{\det(H)} \begin{bmatrix} \frac{\partial^2 c(x,y)}{\partial x \partial y} \frac{\partial w(y)}{\partial y} \\ \left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) \frac{\partial w(y)}{\partial y} \end{bmatrix}$$
- ▶ By second order conditions,  $\det(H) > 0$ .
- ▶ By the concavity of  $v(\cdot)$  and convexity of  $c(\cdot, \cdot)$ ,  
$$\left( \frac{\partial^2 v(x)}{\partial x^2} - \frac{\partial^2 c(x,y)}{\partial x^2} \right) < 0.$$
- ▶ By assumption  $\frac{\partial w(y)}{\partial y} > 0$ .
- ▶ Therefore  $y^*(\theta) > 0$ .
- ▶ We have not yet made any assumptions about  $\frac{\partial^2 c(x,y)}{\partial x \partial y}$ .
- ▶ Since  $-\frac{\partial w(y)}{\partial y} < 0$ , if  $\frac{\partial^2 c(x,y)}{\partial x \partial y} \geq 0$ , then  $x^*(\theta) \leq 0$ . Otherwise,  $x^*(\theta) > 0$ .



## Caveats

- ▶ In all of the examples,  $x^*(\theta)$  is nonempty and singleton.

## Caveats

- ▶ In all of the examples,  $x^*(\theta)$  is nonempty and singleton.
- ▶ In general,  $x^*(\theta)$  may be empty or multi-valued. You need to separately verify that a solution exists.

# Caveats

- ▶ In all of the examples,  $x^*(\theta)$  is nonempty and singleton.
- ▶ In general,  $x^*(\theta)$  may be empty or multi-valued. You need to separately verify that a solution exists.
- ▶ When  $x^*(\theta)$  is multi-valued, the implicit function works locally for every element of  $x^*(\theta)$ .

## Caveats

- ▶ In all of the examples,  $x^*(\theta)$  is nonempty and singleton.
- ▶ In general,  $x^*(\theta)$  may be empty or multi-valued. You need to separately verify that a solution exists.
- ▶ When  $x^*(\theta)$  is multi-valued, the implicit function works locally for every element of  $x^*(\theta)$ .
- ▶ But the direction of  $\frac{\partial x^*(\theta)}{\partial \theta}$  be different at different  $x^* \in x^*(\theta)$ .

## Caveats

- ▶ In all of the examples,  $x^*(\theta)$  is nonempty and singleton.
- ▶ In general,  $x^*(\theta)$  may be empty or multi-valued. You need to separately verify that a solution exists.
- ▶ When  $x^*(\theta)$  is multi-valued, the implicit function works locally for every element of  $x^*(\theta)$ .
- ▶ But the direction of  $\frac{\partial x^*(\theta)}{\partial \theta}$  be different at different  $x^* \in x^*(\theta)$ .
- ▶ The IFT requires  $f$  to be differentiable.

## Caveats

- ▶ In all of the examples,  $x^*(\theta)$  is nonempty and singleton.
- ▶ In general,  $x^*(\theta)$  may be empty or multi-valued. You need to separately verify that a solution exists.
- ▶ When  $x^*(\theta)$  is multi-valued, the implicit function works locally for every element of  $x^*(\theta)$ .
- ▶ But the direction of  $\frac{\partial x^*(\theta)}{\partial \theta}$  be different at different  $x^* \in x^*(\theta)$ .
- ▶ The IFT requires  $f$  to be differentiable.
- ▶ For discontinuous or continuous but nondifferentiable objective functions, we need a different tool.

# Value Functions

- ▶ So far we have been analyzing the behavior of the maximizers of a parameterized optimization problem.

# Value Functions

- ▶ So far we have been analyzing the behavior of the maximizers of a parameterized optimization problem.
- ▶ We often will also want to analyze how the value of a parameterized optimization problem changes with  $\theta$ .



# Value Functions

- ▶ So far we have been analyzing the behavior of the maximizers of a parameterized optimization problem.
- ▶ We often will also want to analyze how the value of a parameterized optimization problem changes with  $\theta$ .
- ▶ The function

$$V(\theta) \equiv f(x^*(\theta); \theta)$$

is referred to as the **value function** or **indirect utility function** for the problem

$$\max_x f(x; \theta)$$

# Value Functions

- ▶ So far we have been analyzing the behavior of the maximizers of a parameterized optimization problem.
- ▶ We often will also want to analyze how the value of a parameterized optimization problem changes with  $\theta$ .
- ▶ The function

$$V(\theta) \equiv f(x^*(\theta); \theta)$$

is referred to as the **value function** or **indirect utility function** for the problem

$$\max_x f(x; \theta)$$

- ▶ The sign of  $\frac{\partial V(\theta)}{\partial \theta}$  tells us how an actor's utility changes for a small rise in  $\theta$  given that she is maximizing.

# Value Functions

- ▶ So far we have been analyzing the behavior of the maximizers of a parameterized optimization problem.
- ▶ We often will also want to analyze how the value of a parameterized optimization problem changes with  $\theta$ .
- ▶ The function

$$V(\theta) \equiv f(x^*(\theta); \theta)$$

is referred to as the **value function** or **indirect utility function** for the problem

$$\max_x f(x; \theta)$$

- ▶ The sign of  $\frac{\partial V(\theta)}{\partial \theta}$  tells us how an actor's utility changes for a small rise in  $\theta$  given that she is maximizing.
- ▶ It is not obvious how to do this as  $\theta$  has two effects on  $V(\theta)$ , a direct effect and an indirect effect that operates through  $x^*(\theta)$ .

# The Envelope Theorem

## Theorem

*For some  $\theta$ , suppose the unique maximizer  $x^*(\theta)$  is locally characterized by first order conditions*

$$\frac{\partial f(x^*(\theta), \theta)}{\partial x} = 0$$

*Suppose that the conditions of the implicit function theorem hold locally so that  $x^*(\cdot)$  is a differentiable function of  $\theta$ . Then*

$$\frac{\partial V(\theta)}{\partial \theta} = \frac{\partial f(x^*(\theta), \theta)}{\partial \theta}$$

# The Envelope Theorem

- ▶ The proof of the envelope theorem is remarkably simple.

# The Envelope Theorem

- ▶ The proof of the envelope theorem is remarkably simple.
- ▶ By first order conditions,  $\frac{\partial f(x;\theta)}{\partial x} = 0$  at the optimum.

# The Envelope Theorem

- ▶ The proof of the envelope theorem is remarkably simple.
- ▶ By first order conditions,  $\frac{\partial f(x;\theta)}{\partial x} = 0$  at the optimum.
- ▶ By the chain rule,

$$\frac{\partial V(\theta)}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial x} \frac{\partial x^*(\theta)}{\partial \theta} + \frac{\partial f(x^*(\theta); \theta)}{\partial \theta} \frac{\partial \theta}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial \theta}$$

# The Envelope Theorem

- ▶ The proof of the envelope theorem is remarkably simple.
- ▶ By first order conditions,  $\frac{\partial f(x;\theta)}{\partial x} = 0$  at the optimum.
- ▶ By the chain rule,

$$\frac{\partial V(\theta)}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial x} \frac{\partial x^*(\theta)}{\partial \theta} + \frac{\partial f(x^*(\theta); \theta)}{\partial \theta} \frac{\partial \theta}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial \theta}$$

- ▶ This is a powerful result. As  $\theta$  changes, so does  $x^*(\theta)$ . Therefore we need to take into account both the direct effect of  $\theta$  on utility but also the change that comes indirectly from the change in  $x^*(\theta)$ .



# The Envelope Theorem

- ▶ The proof of the envelope theorem is remarkably simple.
- ▶ By first order conditions,  $\frac{\partial f(x;\theta)}{\partial x} = 0$  at the optimum.
- ▶ By the chain rule,

$$\frac{\partial V(\theta)}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial x} \frac{\partial x^*(\theta)}{\partial \theta} + \frac{\partial f(x^*(\theta); \theta)}{\partial \theta} \frac{\partial \theta}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial \theta}$$

- ▶ This is a powerful result. As  $\theta$  changes, so does  $x^*(\theta)$ . Therefore we need to take into account both the direct effect of  $\theta$  on utility but also the change that comes indirectly from the change in  $x^*(\theta)$ .
- ▶ What the envelope theorem says is that the indirect effect does not matter!

# The Envelope Theorem

- ▶ The proof of the envelope theorem is remarkably simple.
- ▶ By first order conditions,  $\frac{\partial f(x;\theta)}{\partial x} = 0$  at the optimum.
- ▶ By the chain rule,

$$\frac{\partial V(\theta)}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial x} \frac{\partial x^*(\theta)}{\partial \theta} + \frac{\partial f(x^*(\theta); \theta)}{\partial \theta} \frac{\partial \theta}{\partial \theta} = \frac{\partial f(x^*(\theta); \theta)}{\partial \theta}$$

- ▶ This is a powerful result. As  $\theta$  changes, so does  $x^*(\theta)$ . Therefore we need to take into account both the direct effect of  $\theta$  on utility but also the change that comes indirectly from the change in  $x^*(\theta)$ .
- ▶ What the envelope theorem says is that the indirect effect does not matter!
- ▶ Only the direct effect of the exogenous parameter needs to be considered even though the exogenous parameter may enter the value function indirectly.

## Example

- ▶ Let's revisit the earlier fundraising example where  $f(x; \theta) = \theta \ln(x) - x^2$  and  $x^*(\theta) = \sqrt{\frac{\theta}{2}}$ .

## Example

- ▶ Let's revisit the earlier fundraising example where  $f(x; \theta) = \theta \ln(x) - x^2$  and  $x^*(\theta) = \sqrt{\frac{\theta}{2}}$ .
- ▶ This gives us  $V(\theta) = \theta \ln(\sqrt{\frac{\theta}{2}}) - \frac{\theta}{2}$ .

$$\frac{\partial V(\theta)}{\partial \theta} = \ln(\sqrt{\frac{\theta}{2}}) - \frac{1}{2} + \theta \frac{\partial}{\partial \theta} \ln(\sqrt{\frac{\theta}{2}})$$

$$\frac{\partial}{\partial \theta} \theta \ln(\sqrt{\frac{\theta}{2}}) = \theta \sqrt{\frac{2}{\theta}} \frac{\partial}{\partial \theta} \sqrt{\frac{\theta}{2}} = \theta \sqrt{\frac{2}{\theta}} \frac{1}{\sqrt{2}} \frac{\partial}{\partial \theta} \sqrt{\theta} = \frac{1}{2} \frac{\sqrt{\theta}}{\sqrt{\theta}} = \frac{1}{2}$$

## Example

- ▶ Let's revisit the earlier fundraising example where  $f(x; \theta) = \theta \ln(x) - x^2$  and  $x^*(\theta) = \sqrt{\frac{\theta}{2}}$ .
- ▶ This gives us  $V(\theta) = \theta \ln(\sqrt{\frac{\theta}{2}}) - \frac{\theta}{2}$ .

$$\frac{\partial V(\theta)}{\partial \theta} = \ln(\sqrt{\frac{\theta}{2}}) - \frac{1}{2} + \theta \frac{\partial}{\partial \theta} \ln(\sqrt{\frac{\theta}{2}})$$

$$\frac{\partial}{\partial \theta} \theta \ln(\sqrt{\frac{\theta}{2}}) = \theta \sqrt{\frac{2}{\theta}} \frac{\partial}{\partial \theta} \sqrt{\frac{\theta}{2}} = \theta \sqrt{\frac{2}{\theta}} \frac{1}{\sqrt{2}} \frac{\partial}{\partial \theta} \sqrt{\theta} = \frac{1}{2} \frac{\sqrt{\theta}}{\sqrt{\theta}} = \frac{1}{2}$$

- ▶ Therefore

$$\frac{\partial V(\theta)}{\partial \theta} = \ln(\sqrt{\frac{\theta}{2}}) = \frac{\partial f(x, \theta)}{\partial \theta} \Big|_{x^*(\theta)}$$

## Example

- ▶ Let  $f(x, y; \theta) = v(x) + \theta w(y) - c(x, y)$


## Example


- ▶ Let  $f(x, y; \theta) = v(x) + \theta w(y) - c(x, y)$
- ▶  $V(\theta) = v(x^*(\theta)) + \theta w(y^*(\theta)) - c(x^*(\theta), y^*(\theta))$


## Example


- ▶ Let  $f(x, y; \theta) = v(x) + \theta w(y) - c(x, y)$
- ▶  $V(\theta) = v(x^*(\theta)) + \theta w(y^*(\theta)) - c(x^*(\theta), y^*(\theta))$
- ▶ By the envelope theorem,  $\frac{\partial V(\theta)}{\partial \theta} = w(y^*(\theta)) \geq 0$ .














# Correspondences

- ▶ We have so far been assuming that  $x^*(\theta)$  is a singleton for all  $\theta$ .

# Correspondences

- ▶ We have so far been assuming that  $x^*(\theta)$  is a singleton for all  $\theta$ .
- ▶ We have similarly been assuming that  $x \in \mathbb{R}$  and ignoring the possibility of constraints.

# Correspondences

- ▶ We have so far been assuming that  $x^*(\theta)$  is a singleton for all  $\theta$ .
- ▶ We have similarly been assuming that  $x \in \mathbb{R}$  and ignoring the possibility of constraints.
- ▶ We typically assume that  $x \in G(\theta)$  where  $G$  is called a **feasible set**.



# Correspondences

- ▶ We have so far been assuming that  $x^*(\theta)$  is a singleton for all  $\theta$ .
- ▶ We have similarly been assuming that  $x \in \mathbb{R}$  and ignoring the possibility of constraints.
- ▶ We typically assume that  $x \in G(\theta)$  where  $G$  is called a **feasible set**.
- ▶ If  $G(\theta)$  depends on  $\theta$ , then unless  $G(\theta)$  is a singleton, then we need a more general concept than function to think about the properties of  $G(\theta)$  and  $x^*(\theta)$  (why?)

# Correspondences

- ▶ Given two sets  $X$  and  $Y$ , **correspondence**,  $G : X \rightrightarrows Y$  is a rule that assigns to every element  $x \in X$  a subset  $G(x)$  of  $Y$ .

# Correspondences

- ▶ Given two sets  $X$  and  $Y$ , **correspondence**,  $G : X \rightrightarrows Y$  is a rule that assigns to every element  $x \in X$  a subset  $G(x)$  of  $Y$ .
- ▶ Correspondences are therefore sometimes referred to as “set valued functions.”

# Correspondences

- ▶ Given two sets  $X$  and  $Y$ , **correspondence**,  $G : X \rightrightarrows Y$  is a rule that assigns to every element  $x \in X$  a subset  $G(x)$  of  $Y$ .
- ▶ Correspondences are therefore sometimes referred to as “set valued functions.”
- ▶ *Example:* Let  $X = \mathbb{R}_+$  and  $Y = \mathbb{R}_+$ .  $G(x) = [0, x]$  is a correspondence.

# Correspondences

- ▶ Given two sets  $X$  and  $Y$ , **correspondence**,  $G : X \rightrightarrows Y$  is a rule that assigns to every element  $x \in X$  a subset  $G(x)$  of  $Y$ .
- ▶ Correspondences are therefore sometimes referred to as “set valued functions.”
- ▶ *Example:* Let  $X = \mathbb{R}_+$  and  $Y = \mathbb{R}_+$ .  $G(x) = [0, x]$  is a correspondence.
- ▶ *Example:* Let  $X = \{1, 2, 3, 4, 5, 6\}$  and let  $G(x) = \{z \in X : z \geq x\}$ .

# Correspondences

- ▶ Given two sets  $X$  and  $Y$ , **correspondence**,  $G : X \rightrightarrows Y$  is a rule that assigns to every element  $x \in X$  a subset  $G(x)$  of  $Y$ .
- ▶ Correspondences are therefore sometimes referred to as “set valued functions.”
- ▶ *Example:* Let  $X = \mathbb{R}_+$  and  $Y = \mathbb{R}_+$ .  $G(x) = [0, x]$  is a correspondence.
- ▶ *Example:* Let  $X = \{1, 2, 3, 4, 5, 6\}$  and let  $G(x) = \{z \in X : z \geq x\}$ .
- ▶ This correspondence returns all the elements of  $x$  that are greater than or equal to the input e.g.  $G(4) = \{4, 5, 6\}$ .

# Correspondences

- ▶ Given two sets  $X$  and  $Y$ , **correspondence**,  $G : X \rightrightarrows Y$  is a rule that assigns to every element  $x \in X$  a subset  $G(x)$  of  $Y$ .
- ▶ Correspondences are therefore sometimes referred to as “set valued functions.”
- ▶ *Example:* Let  $X = \mathbb{R}_+$  and  $Y = \mathbb{R}_+$ .  $G(x) = [0, x]$  is a correspondence.
- ▶ *Example:* Let  $X = \{1, 2, 3, 4, 5, 6\}$  and let  $G(x) = \{z \in X : z \geq x\}$ .
- ▶ This correspondence returns all the elements of  $x$  that are greater than or equal to the input e.g.  $G(4) = \{4, 5, 6\}$ .
- ▶ We refer to  $x^*(\theta)$  as a solution correspondence because there may be more than one solution to a problem given  $\theta$ .

# Correspondences

- ▶ A correspondence is **compact valued** if  $G(x)$  is a compact set.



# Correspondences

- ▶ A correspondence is **compact valued** if  $G(x)$  is a compact set.
- ▶ It is called **convex valued** if  $G(x)$  is convex.

# Correspondences

- ▶ A correspondence is **compact valued** if  $G(x)$  is a compact set.
- ▶ It is called **convex valued** if  $G(x)$  is convex.
- ▶ It is called **closed valued** if  $G(x)$  is closed.

# Correspondences

- ▶ A correspondence is **compact valued** if  $G(x)$  is a compact set.
- ▶ It is called **convex valued** if  $G(x)$  is convex.
- ▶ It is called **closed valued** if  $G(x)$  is closed.
- ▶ A **fixed point** of a correspondence is a point  $x \in X$  such that  $x \in G(x)$ .

# Correspondences

- ▶ A correspondence is **compact valued** if  $G(x)$  is a compact set.
- ▶ It is called **convex valued** if  $G(x)$  is convex.
- ▶ It is called **closed valued** if  $G(x)$  is closed.
- ▶ A **fixed point** of a correspondence is a point  $x \in X$  such that  $x \in G(x)$ .
- ▶ A correspondence  $G : X \rightrightarrows Y$  is **upper hemicontinuous at**  $x \in X$  if for every open subset  $V$  of  $Y$  with  $G(x) \subseteq V$ , there exists a  $\delta > 0$  such that  $G(N_\delta(x)) \subseteq V$ .

# Correspondences

- ▶ A correspondence is **compact valued** if  $G(x)$  is a compact set.
- ▶ It is called **convex valued** if  $G(x)$  is convex.
- ▶ It is called **closed valued** if  $G(x)$  is closed.
- ▶ A **fixed point** of a correspondence is a point  $x \in X$  such that  $x \in G(x)$ .
- ▶ A correspondence  $G : X \rightrightarrows Y$  is **upper hemicontinuous at**  $x \in X$  if for every open subset  $V$  of  $Y$  with  $G(x) \subseteq V$ , there exists a  $\delta > 0$  such that  $G(N_\delta(x)) \subseteq V$ .
- ▶  $G$  is called **upper hemicontinuous** if it is upper hemicontinuous on all of  $X$ .

# Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.

# Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.
- ▶ Often, however, we can appeal to properties of the objective function and constraint set to infer properties of the constraint correspondence.

## Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.
- ▶ Often, however, we can appeal to properties of the objective function and constraint set to infer properties of the constraint correspondence.
- ▶ So far we've been exploiting the differentiability of an objective function to learn about the properties of locally smooth solution correspondences and value functions.



# Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.
- ▶ Often, however, we can appeal to properties of the objective function and constraint set to infer properties of the constraint correspondence.
- ▶ So far we've been exploiting the differentiability of an objective function to learn about the properties of locally smooth solution correspondences and value functions.
- ▶ Our problems, however, will not always be so well behaved.

## Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.
- ▶ Often, however, we can appeal to properties of the objective function and constraint set to infer properties of the constraint correspondence.
- ▶ So far we've been exploiting the differentiability of an objective function to learn about the properties of locally smooth solution correspondences and value functions.
- ▶ Our problems, however, will not always be so well behaved.
- ▶ A common utility function in spatial models is the **absolute loss function**,  $u(x; z) = -|x - z|$ .

# Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.
- ▶ Often, however, we can appeal to properties of the objective function and constraint set to infer properties of the constraint correspondence.
- ▶ So far we've been exploiting the differentiability of an objective function to learn about the properties of locally smooth solution correspondences and value functions.
- ▶ Our problems, however, will not always be so well behaved.
- ▶ A common utility function in spatial models is the **absolute loss function**,  $u(x; z) = -|x - z|$ .
- ▶ We cannot use the implicit function for comparative statics here.

## Non-Smooth Functions

- ▶ As we have seen, it is rare that we will be able to find an explicit solution correspondence.
- ▶ Often, however, we can appeal to properties of the objective function and constraint set to infer properties of the constraint correspondence.
- ▶ So far we've been exploiting the differentiability of an objective function to learn about the properties of locally smooth solution correspondences and value functions.
- ▶ Our problems, however, will not always be so well behaved.
- ▶ A common utility function in spatial models is the **absolute loss function**,  $u(x; z) = -|x - z|$ .
- ▶ We cannot use the implicit function for comparative statics here.
- ▶ Intuitively, however, we should be able to exploit the continuity of the objective function to learn about the solution correspondence.

# The Theorem of the Maximum (Light)

## Theorem

*For a parameterized optimization problem, if  $f : X \times \Theta \rightarrow \mathbb{R}$  is continuous and the feasible set  $G$  is compact, then  $V(\theta)$  is continuous and  $x^*(\theta)$  is nonempty, compact-valued, and upper hemicontinuous.*

- ▶ The result of the theorem is that if the elements of an optimization problem are sufficiently continuous, then some, but not all, of that continuity is preserved in the solutions.

# Monotone Comparative Statics

- ▶ Occasionally, even continuity of an objective function may be too strong an assumption.

# Monotone Comparative Statics

- ▶ Occasionally, even continuity of an objective function may be too strong an assumption.
- ▶ A powerful and underutilized tool in political research, **monotone comparative statics**, allows us to study the properties of  $x^*(\theta)$  even when  $f$  is discontinuous.

# Monotone Comparative Statics

- ▶ Occasionally, even continuity of an objective function may be too strong an assumption.
- ▶ A powerful and underutilized tool in political research, **monotone comparative statics**, allows us to study the properties of  $x^*(\theta)$  even when  $f$  is discontinuous.
- ▶ We say that  $f(x, \theta)$  satisfies the **single-crossing property** if for all  $x > x'$  and  $\theta > \theta'$ ,

$$f(x, \theta') - f(x', \theta') \geq 0 \implies f(x, \theta) - f(x', \theta) \geq 0$$

and

$$f(x, \theta') - f(x', \theta') > 0 \implies f(x, \theta) - f(x', \theta) > 0$$



# Monotone Comparative Statics

- ▶ Occasionally, even continuity of an objective function may be too strong an assumption.
- ▶ A powerful and underutilized tool in political research, **monotone comparative statics**, allows us to study the properties of  $x^*(\theta)$  even when  $f$  is discontinuous.
- ▶ We say that  $f(x, \theta)$  satisfies the **single-crossing property** if for all  $x > x'$  and  $\theta > \theta'$ ,


$$f(x, \theta') - f(x', \theta') \geq 0 \implies f(x, \theta) - f(x', \theta) \geq 0$$

and

$$f(x, \theta') - f(x', \theta') > 0 \implies f(x, \theta) - f(x', \theta) > 0$$

- ▶ Intuitively if the two objective functions for different  $\theta$  cross only once, the function satisfies the single crossing property.

# Single-Crossing



# Increasing Differences

- ▶ A sufficient condition for single crossing is **increasing differences**

# Increasing Differences

- ▶ A sufficient condition for single crossing is **increasing differences**
- ▶ A function displays increasing differences if

$$f(x, \cdot) - f(x', \cdot)$$

is weakly increasing in  $\theta$ .

# Increasing Differences

- ▶ A sufficient condition for single crossing is **increasing differences**
- ▶ A function displays increasing differences if

$$f(x, \cdot) - f(x', \cdot)$$


is weakly increasing in  $\theta$ .

- ▶ If  $f(x, \theta') - f(x', \theta') \geq 0$ , then

$$f(x, \theta) - f(x', \theta) \geq f(x, \theta') - f(x', \theta') \geq 0$$

and  $f$  is single crossing

# Increasing Differences



# Monotone Comparative Statics

- ▶ The central result of monotone comparative statics is that if the single crossing condition holds, then  $x^*(\theta)$  is weakly increasing in  $\theta$  (Milgrom and Shannon (1994)).

# Monotone Comparative Statics

- ▶ The central result of monotone comparative statics is that if the single crossing condition holds, then  $x^*(\theta)$  is weakly increasing in  $\theta$  (Milgrom and Shannon (1994)).
- ▶ This result generalizes to higher dimensions.



# Supermodularity

- ▶ If  $X \subseteq \mathbb{R}^n$  and  $\Theta \subseteq \mathbb{R}^m$ , the set  $X \times \Theta$  is a **product set** if it can be represented by the Cartesian product of subsets of  $\mathbb{R}$ .

# Supermodularity

- ▶ If  $X \subseteq \mathbb{R}^n$  and  $\Theta \subseteq \mathbb{R}^m$ , the set  $X \times \Theta$  is a **product set** if it can be represented by the Cartesian product of subsets of  $\mathbb{R}$ .
- ▶  $[a, b] \times [a, b]$  is a product set but not  $\{a, b : x \geq 0, y \geq 0, x + y \leq 1\}$

# Supermodularity

- ▶ If  $X \subseteq \mathbb{R}^n$  and  $\Theta \subseteq \mathbb{R}^m$ , the set  $X \times \Theta$  is a **product set** if it can be represented by the Cartesian product of subsets of  $\mathbb{R}$ .
- ▶  $[a, b] \times [a, b]$  is a product set but not  $\{a, b : x \geq 0, y \geq 0, x + y \leq 1\}$
- ▶ A product set is a special type of chain.

# Supermodularity

- ▶ If  $X \subseteq \mathbb{R}^n$  and  $\Theta \subseteq \mathbb{R}^m$ , the set  $X \times \Theta$  is a **product set** if it can be represented by the Cartesian product of subsets of  $\mathbb{R}$ .
- ▶  $[a, b] \times [a, b]$  is a product set but not  $\{a, b : x \geq 0, y \geq 0, x + y \leq 1\}$
- ▶ A product set is a special type of chain.
- ▶ A function  $f : X \times \Theta \rightarrow \mathbb{R}$  is **supermodular** if  $X \times \Theta$  is a product set and  $f$  has increasing differences for all pairs of arguments of the function.

# Supermodularity

- ▶ If  $X \subseteq \mathbb{R}^n$  and  $\Theta \subseteq \mathbb{R}^m$ , the set  $X \times \Theta$  is a **product set** if it can be represented by the Cartesian product of subsets of  $\mathbb{R}$ .
- ▶  $[a, b] \times [a, b]$  is a product set but not  $\{a, b : x \geq 0, y \geq 0, x + y \leq 1\}$
- ▶ A product set is a special type of chain.
- ▶ A function  $f : X \times \Theta \rightarrow \mathbb{R}$  is **supermodular** if  $X \times \Theta$  is a product set and  $f$  has increasing differences for all pairs of arguments of the function.
- ▶ Supermodularity generalizes the notion of “complementarity.”

# Topkis' Theorem

## Theorem

*Consider the problem*

$$\max_{x \in X} f(x, \theta)$$

*where  $\theta \in \Theta$  is a parameter. If  $f$  is supermodular, then every component of  $x^*(\theta)$  is weakly increasing in  $\theta$*

# Topkis' Theorem

## Theorem

*Consider the problem*

$$\max_{x \in X} f(x, \theta)$$

*where  $\theta \in \Theta$  is a parameter. If  $f$  is supermodular, then every component of  $x^*(\theta)$  is weakly increasing in  $\theta$*

- ▶ A rise in  $\theta$  leads the decision maker to directly raise each  $x_i$ .

# Topkis' Theorem

## Theorem

*Consider the problem*

$$\max_{x \in X} f(x, \theta)$$

*where  $\theta \in \Theta$  is a parameter. If  $f$  is supermodular, then every component of  $x^*(\theta)$  is weakly increasing in  $\theta$*

- ▶ A rise in  $\theta$  leads the decision maker to directly raise each  $x_i$ .
- ▶ Because the  $x_i$  are all complementary to one another, all indirect effects are also positive.



# Topkis' Theorem

## Theorem

*Consider the problem*

$$\max_{x \in X} f(x, \theta)$$

*where  $\theta \in \Theta$  is a parameter. If  $f$  is supermodular, then every component of  $x^*(\theta)$  is weakly increasing in  $\theta$*

- ▶ A rise in  $\theta$  leads the decision maker to directly raise each  $x_i$ .
- ▶ Because the  $x_i$  are all complementary to one another, all indirect effects are also positive.
- ▶ Without supermodularity, indirect effects may cancel out direct effects.

# Topkis' Theorem

## Theorem

*Consider the problem*

$$\max_{x \in X} f(x, \theta)$$

*where  $\theta \in \Theta$  is a parameter. If  $f$  is supermodular, then every component of  $x^*(\theta)$  is weakly increasing in  $\theta$*

- ▶ A rise in  $\theta$  leads the decision maker to directly raise each  $x_i$ .
- ▶ Because the  $x_i$  are all complementary to one another, all indirect effects are also positive.
- ▶ Without supermodularity, indirect effects may cancel out direct effects.
- ▶ Hence the “monotone” in monotone comparative statics.