

Continuous Functions

Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.

Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.
- ▶ Now we reintroduce functions as relationships between elements in two or more spaces.

Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.
- ▶ Now we reintroduce functions as relationships between elements in two or more spaces.
- ▶ “Sets and spaces provide the basic characters of mathematical analysis, functions provide the plot” (Carter 2001).

Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.
- ▶ Now we reintroduce functions as relationships between elements in two or more spaces.
- ▶ “Sets and spaces provide the basic characters of mathematical analysis, functions provide the plot” (Carter 2001).
- ▶ In this lecture we study functions that preserve the geometric structure of the sets they associate, continuous functions.

Functions

- ▶ A **function** is a rule $f : X \rightarrow Y$ that assigns every element x of a set X to a single element of a set Y .

Functions

- ▶ A **function** is a rule $f : X \rightarrow Y$ that assigns every element x of a set X to a single element of a set Y .
- ▶ X is called the **domain**

Functions

- ▶ A **function** is a rule $f : X \rightarrow Y$ that assigns every element x of a set X to a single element of a set Y .
- ▶ X is called the **domain**
- ▶ Y is called the **codomain**

Functions

- ▶ A **function** is a rule $f : X \rightarrow Y$ that assigns every element x of a set X to a single element of a set Y .
- ▶ X is called the **domain**
- ▶ Y is called the **codomain**
- ▶ A mapping is denoted $y = f(x)$ and y is referred to as the **image** of x under f .

Functions

- ▶ A **function** is a rule $f : X \rightarrow Y$ that assigns every element x of a set X to a single element of a set Y .
- ▶ X is called the **domain**
- ▶ Y is called the **codomain**
- ▶ A mapping is denoted $y = f(x)$ and y is referred to as the **image** of x under f .
- ▶ We typically refer to x as an **independent variable** and $y = f(x)$ as a **dependent variable**.

Functions

- ▶ A **function** is a rule $f : X \rightarrow Y$ that assigns every element x of a set X to a single element of a set Y .
- ▶ X is called the **domain**
- ▶ Y is called the **codomain**
- ▶ A mapping is denoted $y = f(x)$ and y is referred to as the **image** of x under f .
- ▶ We typically refer to x as an **independent variable** and $y = f(x)$ as a **dependent variable**.
- ▶ The **range** or “image” of $f : X \rightarrow Y$ is the set of all elements in Y that are images of elements in X , denoted

$$f(X) = \{y \in Y : y = f(x) \text{ for some } x \in X\}$$

Functions

- ▶ If $f(X) = Y$, then f maps X **onto** Y .

Functions

- ▶ If $f(X) = Y$, then f maps X **onto** Y .
- ▶ If $f(x) = f(x')$ implies $x = x'$, then f is **one-to-one**.

Functions

- ▶ If $f(X) = Y$, then f maps X **onto** Y .
- ▶ If $f(x) = f(x')$ implies $x = x'$, then f is **one-to-one**.
- ▶ A function f has an **inverse** $f^{-1} : Y \rightarrow X$ if and only if f is one-to-one and onto.

Functions

- ▶ If $f(X) = Y$, then f maps X **onto** Y .
- ▶ If $f(x) = f(x')$ implies $x = x'$, then f is **one-to-one**.
- ▶ A function f has an **inverse** $f^{-1} : Y \rightarrow X$ if and only if f is one-to-one and onto.
- ▶ A **graph** of a function

$$\text{graph}(f) = \{(x, y) \in X \times Y : y = f(x), x \in X\}$$

Example

- ▶ Let $f(x) = x^2$ for $X = Y = \mathbb{R}$.

Example

- ▶ Let $f(x) = x^2$ for $X = Y = \mathbb{R}$.
- ▶ The range of f is $f(X) = \mathbb{R}_+$.

Example

- ▶ Let $f(x) = x^2$ for $X = Y = \mathbb{R}$.
- ▶ The range of f is $f(X) = \mathbb{R}_+$.
- ▶ The function is not onto.

Example

- ▶ Let $f(x) = x^2$ for $X = Y = \mathbb{R}$.
- ▶ The range of f is $f(X) = \mathbb{R}_+$.
- ▶ The function is not onto.
- ▶ It is also not one-to-one: $-x^2 = x^2$.

Example

- ▶ Let $f(x) = x^3$ for $X = Y = \mathbb{R}$.

Example

- ▶ Let $f(x) = x^3$ for $X = Y = \mathbb{R}$.
- ▶ The range of f is $f(X) = \mathbb{R}$.

Example

- ▶ Let $f(x) = x^3$ for $X = Y = \mathbb{R}$.
- ▶ The range of f is $f(X) = \mathbb{R}$.
- ▶ Therefore the function is onto.

Example

- ▶ Let $f(x) = x^3$ for $X = Y = \mathbb{R}$.
- ▶ The range of f is $f(X) = \mathbb{R}$.
- ▶ Therefore the function is onto.
- ▶ It is also one-to-one.

Example

- ▶ Let $X = \{\text{members of the department}\}$ and $Y = \{\text{days of the year}\}$.

Example

- ▶ Let $X = \{\text{members of the department}\}$ and $Y = \{\text{days of the year}\}$.
- ▶ Let $f : X \rightarrow Y$ be a rule that assigns each member of the department his or her birthday.

Example

- ▶ Let $X = \{\text{members of the department}\}$ and $Y = \{\text{days of the year}\}$.
- ▶ Let $f : X \rightarrow Y$ be a rule that assigns each member of the department his or her birthday.
- ▶ Note that this is a function: each member of the department has a unique birthday.

Example

- ▶ Let $X = \{\text{members of the department}\}$ and $Y = \{\text{days of the year}\}$.
- ▶ Let $f : X \rightarrow Y$ be a rule that assigns each member of the department his or her birthday.
- ▶ Note that this is a function: each member of the department has a unique birthday.
- ▶ Under what conditions is f onto?

Example

- ▶ Let $X = \{\text{members of the department}\}$ and $Y = \{\text{days of the year}\}$.
- ▶ Let $f : X \rightarrow Y$ be a rule that assigns each member of the department his or her birthday.
- ▶ Note that this is a function: each member of the department has a unique birthday.
- ▶ Under what conditions is f onto?
- ▶ One-to-one?

Functionals

- ▶ We will often work with a special type of function in political science that maps into \mathbb{R} .

Functionals

- ▶ We will often work with a special type of function in political science that maps into \mathbb{R} .
- ▶ A **functional** is a real-valued function $f : X \rightarrow \mathbb{R}$.

Functionals

- ▶ We will often work with a special type of function in political science that maps into \mathbb{R} .
- ▶ A **functional** is a real-valued function $f : X \rightarrow \mathbb{R}$.
- ▶ Utility functions such as $-(x - z)^2$ are functionals.

Composition of Functions

- ▶ We also will often use compositions of functions in our research.

Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions** $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is denoted $g \circ f$ or $g(f(x))$ for $x \in X$ and

$$g \circ f : X \rightarrow Z$$

Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions** $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is denoted $g \circ f$ or $g(f(x))$ for $x \in X$ and

$$g \circ f : X \rightarrow Z$$

- ▶ *Example* Let $f(x, y) = x^\alpha y^\beta$ and let $g(\cdot) = \ln(\cdot)$.

Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions** $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is denoted $g \circ f$ or $g(f(x))$ for $x \in X$ and

$$g \circ f : X \rightarrow Z$$

- ▶ *Example* Let $f(x, y) = x^\alpha y^\beta$ and let $g(\cdot) = \ln(\cdot)$.
- ▶ Note

$$f : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$$

and

$$g : \mathbb{R}_{++} \rightarrow \mathbb{R}$$

Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions** $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is denoted $g \circ f$ or $g(f(x))$ for $x \in X$ and

$$g \circ f : X \rightarrow Z$$

- ▶ *Example* Let $f(x, y) = x^\alpha y^\beta$ and let $g(\cdot) = \ln(\cdot)$.
- ▶ Note

$$f : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$$

and

$$g : \mathbb{R}_{++} \rightarrow \mathbb{R}$$

- ▶ The composition of these two functions is

$$g(f(x)) = \alpha \ln(x) + \beta \ln(y)$$

with $g(f(x)) : \mathbb{R}_+^2 \rightarrow \mathbb{R}$

Inverse Image

- ▶ An additional definition will prove useful in defining continuity.

Inverse Image

- ▶ An additional definition will prove useful in defining continuity.
- ▶ The **inverse image** of a function of a set $T \subseteq Y$ is the set of $x \in X$ that are mapped to a elements $y \in T$

$$f^{-1}(T) = \{x \in X | f(x) \in T\}$$

Inverse Image

- ▶ An additional definition will prove useful in defining continuity.
- ▶ The **inverse image** of a function of a set $T \subseteq Y$ is the set of $x \in X$ that are mapped to a elements $y \in T$

$$f^{-1}(T) = \{x \in X | f(x) \in T\}$$

- ▶ *Example:* Let $f(x) = x^2$ and let $T = [0, 4] \subset \mathbb{R} = Y$.

Inverse Image

- ▶ An additional definition will prove useful in defining continuity.
- ▶ The **inverse image** of a function of a set $T \subseteq Y$ is the set of $x \in X$ that are mapped to a elements $y \in T$

$$f^{-1}(T) = \{x \in X | f(x) \in T\}$$

- ▶ *Example:* Let $f(x) = x^2$ and let $T = [0, 4] \subset \mathbb{R} = Y$.
- ▶ The inverse image $f^{-1}(T)$ is $[0, 2] \subset \mathbb{R} = X$.

Continuity

- ▶ Informally, a function is continuous if small changes in x produce only small changes in y .

Continuity

- ▶ Informally, a function is continuous if small changes in x produce only small changes in y .
- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

Continuity

- ▶ Informally, a function is continuous if small changes in x produce only small changes in y .

- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

- ▶ In a two-candidate race, $f(x) = 0$ for $x < 1/2$ and $f(x) = 1$ for $x > 1/2$.

Continuity

- ▶ Informally, a function is continuous if small changes in x produce only small changes in y .

- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

- ▶ In a two-candidate race, $f(x) = 0$ for $x < 1/2$ and $f(x) = 1$ for $x > 1/2$.
- ▶ A small change in x in the neighborhood of $1/2$ results in a large change in y .

Continuity

- ▶ Informally, a function is continuous if small changes in x produce only small changes in y .

- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

- ▶ In a two-candidate race, $f(x) = 0$ for $x < 1/2$ and $f(x) = 1$ for $x > 1/2$.
- ▶ A small change in x in the neighborhood of $1/2$ results in a large change in y .
- ▶ This is a discontinuous function.

Continuity

- ▶ Now consider the function

$$f(x, y, z) = \beta_0 + \beta_1 x + \beta_2 y + \beta_3 z + \epsilon$$

for $\beta_i, \epsilon, x, y, z \in \mathbb{R}$

Continuity

- ▶ Now consider the function

$$f(x, y, z) = \beta_0 + \beta_1 x + \beta_2 y + \beta_3 z + \epsilon$$

for $\beta_i, \epsilon, x, y, z \in \mathbb{R}$

- ▶ Note that a small change in any variable results in only a small change in $f(x)$.

Continuity

- ▶ Now consider the function

$$f(x, y, z) = \beta_0 + \beta_1 x + \beta_2 y + \beta_3 z + \epsilon$$

for $\beta_i, \epsilon, x, y, z \in \mathbb{R}$

- ▶ Note that a small change in any variable results in only a small change in $f(x)$.
- ▶ This is a continuous function.

Continuity

- ▶ Recall that our concept of distance (and hence a formal notion of “small difference”) is defined for metric spaces.

Continuity

- ▶ Recall that our concept of distance (and hence a formal notion of “small difference”) is defined for metric spaces.
- ▶ A continuous function preserves the geometric structure of the sets it associates i.e. the metric spaces it associates.

Continuity

- ▶ Recall that our concept of distance (and hence a formal notion of “small difference”) is defined for metric spaces.
- ▶ A continuous function preserves the geometric structure of the sets it associates i.e. the metric spaces it associates.
- ▶ For the rest of this lecture we will assume that X and Y are metric spaces.

Continuity: Epsilon-Delta Characterization

- ▶ A function $f : X \rightarrow Y$ is continuous at x_0 if for all $\epsilon > 0$, there exists a $\delta > 0$ such that for every $x \in X$,

$$d(x, x_0) < \delta \implies d(f(x), f(x_0)) < \epsilon$$

Continuity: Epsilon-Delta Characterization

- ▶ A function $f : X \rightarrow Y$ is continuous at x_0 if for all $\epsilon > 0$, there exists a $\delta > 0$ such that for every $x \in X$,

$$d(x, x_0) < \delta \implies d(f(x), f(x_0)) < \epsilon$$

- ▶ If a function is continuous for all x_0 in its domain, we call the function continuous.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?
- ▶ Let $\epsilon > 0$.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?
- ▶ Let $\epsilon > 0$.
- ▶ Notice that $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?
- ▶ Let $\epsilon > 0$.
- ▶ Notice that $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$.
- ▶ Note that $f(x) \leq x^2$ for all x because the range of $\sin(\cdot)$ is $[-1, 1]$.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?
- ▶ Let $\epsilon > 0$.
- ▶ Notice that $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$.
- ▶ Note that $f(x) \leq x^2$ for all x because the range of $\sin(\cdot)$ is $[-1, 1]$.
- ▶ Let $\delta = \sqrt{\epsilon}$.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?
- ▶ Let $\epsilon > 0$.
- ▶ Notice that $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$.
- ▶ Note that $f(x) \leq x^2$ for all x because the range of $\sin(\cdot)$ is $[-1, 1]$.
- ▶ Let $\delta = \sqrt{\epsilon}$.
- ▶ Now note that $|x - 0| < \delta$ implies $x^2 < \delta^2 = \epsilon$.

Example

- ▶ Let $f(x) = x^2 \sin(\frac{1}{x})$ for $x \neq 0$ and $f(0) = 0$.
- ▶ Is $f(x)$ continuous at $x = 0$?
- ▶ Let $\epsilon > 0$.
- ▶ Notice that $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$.
- ▶ Note that $f(x) \leq x^2$ for all x because the range of $\sin(\cdot)$ is $[-1, 1]$.
- ▶ Let $\delta = \sqrt{\epsilon}$.
- ▶ Now note that $|x - 0| < \delta$ implies $x^2 < \delta^2 = \epsilon$.
- ▶ Therefore $|x - 0| < \delta$ implies $|f(x) - f(0)| < \epsilon$ so f is continuous at $x = 0$.

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$
- ▶ We need to show that $d(f(x_0), f(x)) < \epsilon$ for $d(x_0, x)$ sufficiently small where x is an element of the domain.

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$
- ▶ We need to show that $d(f(x_0), f(x)) < \epsilon$ for $d(x_0, x)$ sufficiently small where x is an element of the domain.
- ▶ Note that $d(x, x_0) = |x - x_0|$ and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$
- ▶ We need to show that $d(f(x_0), f(x)) < \epsilon$ for $d(x_0, x)$ sufficiently small where x is an element of the domain.
- ▶ Note that $d(x, x_0) = |x - x_0|$ and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if $|x - x_0| < k$, then $|x| - |x_0| < k$ or $|x| < |x_0| + k$.

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$
- ▶ We need to show that $d(f(x_0), f(x)) < \epsilon$ for $d(x_0, x)$ sufficiently small where x is an element of the domain.
- ▶ Note that $d(x, x_0) = |x - x_0|$ and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if $|x - x_0| < k$, then $|x| - |x_0| < k$ or $|x| < |x_0| + k$.
- ▶ By the triangle inequality, $|x + x_0| \leq |x| + |x_0| < 2|x_0| + k$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$
- ▶ We need to show that $d(f(x_0), f(x)) < \epsilon$ for $d(x_0, x)$ sufficiently small where x is an element of the domain.
- ▶ Note that $d(x, x_0) = |x - x_0|$ and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if $|x - x_0| < k$, then $|x| - |x_0| < k$ or $|x| < |x_0| + k$.
- ▶ By the triangle inequality, $|x + x_0| \leq |x| + |x_0| < 2|x_0| + k$
- ▶ Therefore

$$|f(x) - f(x_0)| \leq 2|x - x_0|(2|x_0| + k)$$

if $|x - x_0| < k$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Take an arbitrary x_0 and $\epsilon > 0$
- ▶ We need to show that $d(f(x_0), f(x)) < \epsilon$ for $d(x_0, x)$ sufficiently small where x is an element of the domain.
- ▶ Note that $d(x, x_0) = |x - x_0|$ and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if $|x - x_0| < k$, then $|x| - |x_0| < k$ or $|x| < |x_0| + k$.
- ▶ By the triangle inequality, $|x + x_0| \leq |x| + |x_0| < 2|x_0| + k$
- ▶ Therefore

$$|f(x) - f(x_0)| \leq 2|x - x_0|(2|x_0| + k)$$

if $|x - x_0| < k$

- ▶ For $2|x - x_0|(2|x_0| + k) < \epsilon$ to hold, it is sufficient that $|x - x_0| < \frac{\epsilon}{2(2|x_0| + k)}$ and $|x - x_0| < k$

Example (cont.)

- So let

$$\delta = \min\left\{k, \frac{\epsilon}{2(2|x_0| + k)}\right\}$$

Example (cont.)

- So let

$$\delta = \min\left\{k, \frac{\epsilon}{2(2|x_0| + k)}\right\}$$

- Therefore $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let $\epsilon = 1/2$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let $\epsilon = 1/2$.
- ▶ If f is continuous, then I can find a δ such that $|x| < \delta$ implies $|f(x)| < \epsilon$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let $\epsilon = 1/2$.
- ▶ If f is continuous, then I can find a δ such that $|x| < \delta$ implies $|f(x)| < \epsilon$.
- ▶ This is equivalent to finding an open interval $D = (-\delta, \delta)$ such that $x \in D$ implies $|f(x)| < \epsilon$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let $\epsilon = 1/2$.
- ▶ If f is continuous, then I can find a δ such that $|x| < \delta$ implies $|f(x)| < \epsilon$.
- ▶ This is equivalent to finding an open interval $D = (-\delta, \delta)$ such that $x \in D$ implies $|f(x)| < \epsilon$.
- ▶ But note that however small we make this interval, if $x \in D$, then $-x \in D$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let $\epsilon = 1/2$.
- ▶ If f is continuous, then I can find a δ such that $|x| < \delta$ implies $|f(x)| < \epsilon$.
- ▶ This is equivalent to finding an open interval $D = (-\delta, \delta)$ such that $x \in D$ implies $|f(x)| < \epsilon$.
- ▶ But note that however small we make this interval, if $x \in D$, then $-x \in D$.
- ▶ If x is positive, $f(x) = 0$ and $f(-x) = 1 > 1/2 = \epsilon$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$
- ▶ For $x = 0$ note that $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let $\epsilon = 1/2$.
- ▶ If f is continuous, then I can find a δ such that $|x| < \delta$ implies $|f(x)| < \epsilon$.
- ▶ This is equivalent to finding an open interval $D = (-\delta, \delta)$ such that $x \in D$ implies $|f(x)| < \epsilon$.
- ▶ But note that however small we make this interval, if $x \in D$, then $-x \in D$.
- ▶ If x is positive, $f(x) = 0$ and $f(-x) = 1 > 1/2 = \epsilon$.
- ▶ Therefore the function is not continuous.

Continuity: Sequence Convergence Characterization

- ▶ The $\epsilon - \delta$ definition of continuity is difficult to work with. Often an alternative and equivalent characterization of continuity is easier:

Continuity: Sequence Convergence Characterization

- ▶ The $\epsilon - \delta$ definition of continuity is difficult to work with. Often an alternative and equivalent characterization of continuity is easier:
- ▶ A function $f : X \rightarrow Y$ is continuous if and only if $f(x) = \lim_{n \rightarrow \infty} f(x^n)$ for every sequence $x^n \rightarrow x$.

Continuity: Sequence Convergence Characterization

- ▶ The $\epsilon - \delta$ definition of continuity is difficult to work with. Often an alternative and equivalent characterization of continuity is easier:
- ▶ A function $f : X \rightarrow Y$ is continuous if and only if $f(x) = \lim_{n \rightarrow \infty} f(x^n)$ for every sequence $x^n \rightarrow x$.
- ▶ This alternative definition implies that values of $f(x)$ are close to $f(x_0)$ when values of x are close to x_0 .

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Suppose we take an arbitrary sequence in X such that $\lim x^n = x_0$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Suppose we take an arbitrary sequence in X such that $\lim x^n = x_0$
- ▶ We have

$$\lim(2x^2 + 1) = 2(\lim x)^2 + 1 = 2x_0^2 + 1 = f(x_0)$$

Example

- ▶ Let $f(x) = 2x^2 + 1$ for $X = \mathbb{R}$
- ▶ Suppose we take an arbitrary sequence in X such that $\lim x^n = x_0$
- ▶ We have

$$\lim(2x^2 + 1) = 2(\lim x)^2 + 1 = 2x_0^2 + 1 = f(x_0)$$

- ▶ Therefore $f(x)$ is continuous.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$.
- ▶ To show that f is discontinuous, it is sufficient to find single sequence $x^n \rightarrow x_0$ in X such that $\lim f(x^n) \neq f(x_0)$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$.
- ▶ To show that f is discontinuous, it is sufficient to find single sequence $x^n \rightarrow x_0$ in X such that $\lim f(x^n) \neq f(x_0)$.
- ▶ Consider the sequence $x^n = -\frac{1}{n}$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$.
- ▶ To show that f is discontinuous, it is sufficient to find single sequence $x^n \rightarrow x_0$ in X such that $\lim f(x^n) \neq f(x_0)$.
- ▶ Consider the sequence $x^n = -\frac{1}{n}$.
- ▶ Note that $\lim x^n = 0$ and $x^n < 0$ for all x^n .

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$.
- ▶ To show that f is discontinuous, it is sufficient to find single sequence $x^n \rightarrow x_0$ in X such that $\lim f(x^n) \neq f(x_0)$.
- ▶ Consider the sequence $x^n = -\frac{1}{n}$.
- ▶ Note that $\lim x^n = 0$ and $x^n < 0$ for all x^n .
- ▶ We therefore have $\lim f(x^n) = \lim 1 = 1 \neq f(x^0) = 0$.

Example

- ▶ Let $f(x) = 1$ for $x < 0$ and $f(x) = 0$ for $x \geq 0$.
- ▶ To show that f is discontinuous, it is sufficient to find single sequence $x^n \rightarrow x_0$ in X such that $\lim f(x^n) \neq f(x_0)$.
- ▶ Consider the sequence $x^n = -\frac{1}{n}$.
- ▶ Note that $\lim x^n = 0$ and $x^n < 0$ for all x^n .
- ▶ We therefore have $\lim f(x^n) = \lim 1 = 1 \neq f(x^0) = 0$.
- ▶ Therefore f is discontinuous at $x = 0$.

Continuity: Inverse Image Characterization

- ▶ A third characterization of continuity is often useful in proofs that require or seek to establish continuity.

Continuity: Inverse Image Characterization

- ▶ A third characterization of continuity is often useful in proofs that require or seek to establish continuity.
- ▶ A function $f : X \rightarrow Y$ is continuous if and only if the inverse image of any open (closed) subset of Y is an open (closed) subset of X .

Example

- ▶ Let $f(x) = x^3$

Example

- ▶ Let $f(x) = x^3$
- ▶ Consider the subset of its codomain $(0, 2)$

Example

- ▶ Let $f(x) = x^3$
- ▶ Consider the subset of its codomain $(0, 2)$
- ▶ The inverse image $f^{-1}((0, 8))$ is the set of all points x in the domain of x^3 such that $x^3 \in (0, 8)$.

Example

- ▶ Let $f(x) = x^3$
- ▶ Consider the subset of its codomain $(0, 2)$
- ▶ The inverse image $f^{-1}((0, 8))$ is the set of all points x in the domain of x^3 such that $x^3 \in (0, 8)$.
- ▶ Therefore $f^{-1}((0, 2)) = (0, 2)$.

Example

- ▶ Let $f(x) = x^3$
- ▶ Consider the subset of its codomain $(0, 2)$
- ▶ The inverse image $f^{-1}((0, 8))$ is the set of all points x in the domain of x^3 such that $x^3 \in (0, 8)$.
- ▶ Therefore $f^{-1}((0, 2)) = (0, 2)$.
- ▶ Note that this is an open subset of $X = \mathbb{R}$.

Example

- ▶ Let $f(x) = 1$ for all x .

Example

- ▶ Let $f(x) = 1$ for all x .
- ▶ Consider a subset of the codomain, $(-1, 1)$.

Example

- ▶ Let $f(x) = 1$ for all x .
- ▶ Consider a subset of the codomain, $(-1, 1)$.
- ▶ The inverse image under $(-1, 1)$ is \emptyset which is open.

Example

- ▶ Let $f(x) = 1$ for all x .
- ▶ Consider a subset of the codomain, $(-1, 1)$.
- ▶ The inverse image under $(-1, 1)$ is \emptyset which is open.
- ▶ Now consider $(0, 2)$.

Example

- ▶ Let $f(x) = 1$ for all x .
- ▶ Consider a subset of the codomain, $(-1, 1)$.
- ▶ The inverse image under $(-1, 1)$ is \emptyset which is open.
- ▶ Now consider $(0, 2)$.
- ▶ The inverse image under $(0, 2)$ is \mathbb{R} .

Continuity: Inverse Image Characterization

- ▶ Using this result to show that a function is continuous can be difficult because we have to evaluate every subset of the codomain.

Continuity: Inverse Image Characterization

- ▶ Using this result to show that a function is continuous can be difficult because we have to evaluate every subset of the codomain.
- ▶ It is easier to use the definition to show discontinuity because we only need to find a single example of a set that violates the result.

Example

- ▶ Let $f(x) = x$ for $x \leq 0$ and let $f(x) = x + 1$ for $x > 0$.

Example

- ▶ Let $f(x) = x$ for $x \leq 0$ and let $f(x) = x + 1$ for $x > 0$.
- ▶ Consider the open subset of the codomain $(-1, 1) \subset Y$.

Example

- ▶ Let $f(x) = x$ for $x \leq 0$ and let $f(x) = x + 1$ for $x > 0$.
- ▶ Consider the open subset of the codomain $(-1, 1) \subset Y$.
- ▶ The inverse image under $(-1, 1)$ is $(-1, 1]$ which is not open.

Example

- ▶ Let $f(x) = x$ for $x \leq 0$ and let $f(x) = x + 1$ for $x > 0$.
- ▶ Consider the open subset of the codomain $(-1, 1) \subset Y$.
- ▶ The inverse image under $(-1, 1)$ is $(-1, 1]$ which is not open.
- ▶ Therefore $f(x)$ is not continuous.

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Note that $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$.

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Note that $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$.
- ▶ Therefore kf is continuous at x_0 .

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Note that $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$.
- ▶ Therefore kf is continuous at x_0 .
- ▶ Now consider $|f|$ at x_0 .

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Note that $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$.
- ▶ Therefore kf is continuous at x_0 .
- ▶ Now consider $|f|$ at x_0 .
- ▶ By the reverse triangle inequality,

$$||f(x_n)| - |f(x_0)|| \leq |f(x_n) - f(x_0)|$$

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Note that $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$.
- ▶ Therefore kf is continuous at x_0 .
- ▶ Now consider $|f|$ at x_0 .
- ▶ By the reverse triangle inequality,

$$||f(x_n)| - |f(x_0)|| \leq |f(x_n) - f(x_0)|$$

- ▶ Now consider $\epsilon > 0$

Properties of Continuous Functions

- ▶ Let f be a real-valued function with domain $X \subseteq \mathbb{R}$. If f is continuous at x_0 in X , then $|f|$ and kf , $k \in \mathbb{R}$ are continuous at x_0 .
- ▶ **Proof:** Consider the sequence $x_n \rightarrow x_0$ in the domain of f .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Note that $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$.
- ▶ Therefore kf is continuous at x_0 .
- ▶ Now consider $|f|$ at x_0 .
- ▶ By the reverse triangle inequality,

$$||f(x_n)| - |f(x_0)|| \leq |f(x_n) - f(x_0)|$$

- ▶ Now consider $\epsilon > 0$
- ▶ Because $\lim f(x_n) = f(x_0)$, there exists an N such that $n > N$ implies $||f(x_n)| - |f(x_0)|| < \epsilon$ and thus $\lim |f(x_n)| = |f(x_0)|$.

Properties of Continuous Functions

- ▶ If f and g are real-valued functions and continuous at x_0 , then

Properties of Continuous Functions

- ▶ If f and g are real-valued functions and continuous at x_0 , then
- ▶ i) $f + g$ is continuous at x_0

Properties of Continuous Functions

- ▶ If f and g are real-valued functions and continuous at x_0 , then
- ▶ i) $f + g$ is continuous at x_0
- ▶ ii) fg is continuous at x_0

Properties of Continuous Functions

- ▶ If f and g are real-valued functions and continuous at x_0 , then
- ▶ i) $f + g$ is continuous at x_0
- ▶ ii) fg is continuous at x_0
- ▶ iii) $\frac{f}{g}$ is continuous at x_0 if $g(x_0) \neq 0$.

Proof of $f + g$ continuity at x_0

- ▶ x_0 is in the domain of f and g .

Proof of $f + g$ continuity at x_0

- ▶ x_0 is in the domain of f and g .
- ▶ Let x_n be a sequence in the intersection of their domains that converges to x_0 .

Proof of $f + g$ continuity at x_0

- ▶ x_0 is in the domain of f and g .
- ▶ Let x_n be a sequence in the intersection of their domains that converges to x_0 .
- ▶ Because each is continuous, we have $\lim f(x_n) = f(x_0)$ and $\lim g(x_n) = g(x_0)$.

Proof of $f + g$ continuity at x_0

- ▶ x_0 is in the domain of f and g .
- ▶ Let x_n be a sequence in the intersection of their domains that converges to x_0 .
- ▶ Because each is continuous, we have $\lim f(x_n) = f(x_0)$ and $\lim g(x_n) = g(x_0)$.
- ▶ Therefore

$$\begin{aligned}\lim(f + g)(x_n) &= \lim[f(x_n) + g(x_n)] = \lim f(x_n) + \lim g(x_n) \\ &= f(x_0) + g(x_0) = (f + g)(x_0)\end{aligned}$$

Proof of $f + g$ continuity at x_0

- ▶ x_0 is in the domain of f and g .
- ▶ Let x_n be a sequence in the intersection of their domains that converges to x_0 .
- ▶ Because each is continuous, we have $\lim f(x_n) = f(x_0)$ and $\lim g(x_n) = g(x_0)$.
- ▶ Therefore

$$\begin{aligned}\lim(f + g)(x_n) &= \lim[f(x_n) + g(x_n)] = \lim f(x_n) + \lim g(x_n) \\ &= f(x_0) + g(x_0) = (f + g)(x_0)\end{aligned}$$

- ▶ Therefore $f + g$ is continuous at x_0 .

Proof of $f + g$ continuity at x_0

- ▶ x_0 is in the domain of f and g .
- ▶ Let x_n be a sequence in the intersection of their domains that converges to x_0 .
- ▶ Because each is continuous, we have $\lim f(x_n) = f(x_0)$ and $\lim g(x_n) = g(x_0)$.
- ▶ Therefore

$$\begin{aligned}\lim(f + g)(x_n) &= \lim[f(x_n) + g(x_n)] = \lim f(x_n) + \lim g(x_n) \\ &= f(x_0) + g(x_0) = (f + g)(x_0)\end{aligned}$$

- ▶ Therefore $f + g$ is continuous at x_0 .
- ▶ A similar proof establishes that fg and f/g are continuous at x_0 .

Properties of Continuous Functions

- ▶ If f is continuous at x_0 and g is continuous at $f(x_0)$, then $g(f(x))$ is continuous at x_0 .

Properties of Continuous Functions

- ▶ If f is continuous at x_0 and g is continuous at $f(x_0)$, then $g(f(x))$ is continuous at x_0 .
- ▶ **Proof:** x_0 is in the domain of f and $f(x_0)$ is in the domain of g

Properties of Continuous Functions

- ▶ If f is continuous at x_0 and g is continuous at $f(x_0)$, then $g(f(x))$ is continuous at x_0 .
- ▶ **Proof:** x_0 is in the domain of f and $f(x_0)$ is in the domain of g
- ▶ Let x_n be a sequence such that x_n is in the domain of f and $f(x_n)$ is in the domain of g converging to x_0 .

Properties of Continuous Functions

- ▶ If f is continuous at x_0 and g is continuous at $f(x_0)$, then $g(f(x))$ is continuous at x_0 .
- ▶ **Proof:** x_0 is in the domain of f and $f(x_0)$ is in the domain of g
- ▶ Let x_n be a sequence such that x_n is in the domain of f and $f(x_n)$ is in the domain of g converging to x_0 .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.

Properties of Continuous Functions

- ▶ If f is continuous at x_0 and g is continuous at $f(x_0)$, then $g(f(x))$ is continuous at x_0 .
- ▶ **Proof:** x_0 is in the domain of f and $f(x_0)$ is in the domain of g
- ▶ Let x_n be a sequence such that x_n is in the domain of f and $f(x_n)$ is in the domain of g converging to x_0 .
- ▶ Because f is continuous at x_0 , $\lim f(x_n) = f(x_0)$.
- ▶ Because $f(x_n)$ converges to $f(x_0)$ and g is continuous at $f(x_0)$, we have

$$\lim g(f(x_n)) = g(f(x_0))$$

Properties of Continuous Functions

- ▶ A continuous functional $f : X \rightarrow Y$ on a compact set achieves a maximum and a minimum (Weierstrass theorem).

Properties of Continuous Functions

- ▶ A continuous functional $f : X \rightarrow Y$ on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because X is bounded, $f(x)$ has a supremum.

Properties of Continuous Functions

- ▶ A continuous functional $f : X \rightarrow Y$ on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because X is bounded, $f(x)$ has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.

Properties of Continuous Functions

- ▶ A continuous functional $f : X \rightarrow Y$ on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because X is bounded, $f(x)$ has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.
- ▶ Let $M = \sup_{x \in X} f(x)$.

Properties of Continuous Functions

- ▶ A continuous functional $f : X \rightarrow Y$ on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because X is bounded, $f(x)$ has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.
- ▶ Let $M = \sup_{x \in X} f(x)$.
- ▶ Take a sequence $f(x_n) \in Y$ with $f(x_n) \rightarrow M$.

Properties of Continuous Functions

- ▶ A continuous functional $f : X \rightarrow Y$ on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because X is bounded, $f(x)$ has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.
- ▶ Let $M = \sup_{x \in X} f(x)$.
- ▶ Take a sequence $f(x_n) \in Y$ with $f(x_n) \rightarrow M$.
- ▶ Then $x_n \in X$.

Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set $S \subseteq \mathbb{R}^n$ has a convergent subsequence which converges to a point in S .

Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set $S \subseteq \mathbb{R}^n$ has a convergent subsequence which converges to a point in S .
- ▶ Therefore because X is compact, we can find a convergent subsequence $x_{n_k} \rightarrow x_0$ where $f(x_{n_k}) \rightarrow M$.

Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set $S \subseteq \mathbb{R}^n$ has a convergent subsequence which converges to a point in S .
- ▶ Therefore because X is compact, we can find a convergent subsequence $x_{n_k} \rightarrow x_0$ where $f(x_{n_k}) \rightarrow M$.
- ▶ Applying the limit definition of continuity, we have $\lim f(x_{n_k}) = f(x_0) = M$.

Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set $S \subseteq \mathbb{R}^n$ has a convergent subsequence which converges to a point in S .
- ▶ Therefore because X is compact, we can find a convergent subsequence $x_{n_k} \rightarrow x_0$ where $f(x_{n_k}) \rightarrow M$.
- ▶ Applying the limit definition of continuity, we have $\lim f(x_{n_k}) = f(x_0) = M$.
- ▶ Therefore $f(x_0) = M$

Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set $S \subseteq \mathbb{R}^n$ has a convergent subsequence which converges to a point in S .
- ▶ Therefore because X is compact, we can find a convergent subsequence $x_{n_k} \rightarrow x_0$ where $f(x_{n_k}) \rightarrow M$.
- ▶ Applying the limit definition of continuity, we have $\lim f(x_{n_k}) = f(x_0) = M$.
- ▶ Therefore $f(x_0) = M$
- ▶ The same proof can be applied to the infimum of the set.