

Logic and Proofs

Statements and Their Truth Values

- ▶ Let A be a **statement**

Statements and Their Truth Values

- ▶ Let A be a **statement**
- ▶ A statement is sentence that can be true or false.

Statements and Their Truth Values

- ▶ Let A be a **statement**
- ▶ A statement is sentence that can be true or false.
- ▶ Let X be a set and let A be the statement " $x \in X$ "

Statements and Their Truth Values

- ▶ Let A be a **statement**
- ▶ A statement is sentence that can be true or false.
- ▶ Let X be a set and let A be the statement " $x \in X$ "
- ▶ A can be true or false.

Statements and Their Truth Values

- ▶ Let A be a **statement**
- ▶ A statement is sentence that can be true or false.
- ▶ Let X be a set and let A be the statement " $x \in X$ "
- ▶ A can be true or false.
- ▶ If x is an irrational number and X is \mathbb{Z} , then A is false.

Statements and Their Truth Values

- ▶ Let A be a **statement**
- ▶ A statement is sentence that can be true or false.
- ▶ Let X be a set and let A be the statement " $x \in X$ "
- ▶ A can be true or false.
- ▶ If x is an irrational number and X is \mathbb{Z} , then A is false.
- ▶ If x is a banana and X is the set of all types of fruit, then A is true.

Conditional Statements

- ▶ Sufficiency: $A \implies B$

Conditional Statements

- ▶ Sufficiency: $A \implies B$
- ▶ Necessity and Sufficiency: $A \iff B$

Conditional Statements

- ▶ Sufficiency: $A \implies B$
- ▶ Necessity and Sufficiency: $A \iff B$
- ▶ Necessity: $\neg A \implies \neg B$

Connectives

- ▶ New statements can be constructed from other statements by using the connectives “and” or “or.”

Connectives

- ▶ New statements can be constructed from other statements by using the connectives “and” or “or.”
- ▶ $A \vee B$ is read “ A or B ” and corresponds to the concept of union in set theory.

Connectives

- ▶ New statements can be constructed from other statements by using the connectives “and” or “or.”
- ▶ $A \vee B$ is read “ A or B ” and corresponds to the concept of union in set theory.
- ▶ $A \wedge B$ is read “ A and B ” and corresponds to the concept of intersection in set theory.

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$
- ▶ $B \implies C$

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$
- ▶ $B \implies C$
- ▶ $\neg A \implies \neg C$

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$
- ▶ $B \implies C$
- ▶ $\neg A \implies \neg C$
- ▶ $\neg B \implies \neg C$

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$
- ▶ $B \implies C$
- ▶ $\neg A \implies \neg C$
- ▶ $\neg B \implies \neg C$
- ▶ $A \wedge B \implies C$

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$
- ▶ $B \implies C$
- ▶ $\neg A \implies \neg C$
- ▶ $\neg B \implies \neg C$
- ▶ $A \wedge B \implies C$
- ▶ $A \vee B \implies C$

Example

- ▶ Let A be the statement “the child is a boy,” B be the statement “the child is a twin,” and C be the statement “the child is a brother.”
- ▶ Are the statements below true or false?
- ▶ $A \implies C$
- ▶ $B \implies C$
- ▶ $\neg A \implies \neg C$
- ▶ $\neg B \implies \neg C$
- ▶ $A \wedge B \implies C$
- ▶ $A \vee B \implies C$
- ▶ $C \implies A$

Logical Principles

- ▶ **Contrapositive:** $C \implies A$ iff $\neg A \implies \neg C$ is true.

Logical Principles

- ▶ **Contrapositive:** $C \implies A$ iff $\neg A \implies \neg C$ is true.
- ▶ **Noncontradiction:** $(A \wedge \neg A)$ is always false

Proofs

- ▶ Mathematics has well-defined procedures for verifying that a given statement is true or false.

Proof by Deduction

- ▶ Assume too that we know that there exists some C such that $C \implies B$.

Proof by Deduction

- ▶ Assume too that we know that there exists some C such that $C \implies B$.
- ▶ If we can show that $A \implies C$, then we have proven that $A \implies B$.

Example

- ▶ Claim: $x^2 - 4x + 9$ is always positive.

Example

- ▶ Claim: $x^2 - 4x + 9$ is always positive.
- ▶ Proof: We know that for any $x \in \mathbb{R}$, $x^2 \geq 0$.

Example

- ▶ Claim: $x^2 - 4x + 9$ is always positive.
- ▶ Proof: We know that for any $x \in \mathbb{R}$, $x^2 \geq 0$.
- ▶ Therefore $x^2 + y$ for any $y > 0$ is always positive.

Example

- ▶ Claim: $x^2 - 4x + 9$ is always positive.
- ▶ Proof: We know that for any $x \in \mathbb{R}$, $x^2 \geq 0$.
- ▶ Therefore $x^2 + y$ for any $y > 0$ is always positive.
- ▶ Now note that $x^2 - 4x + 9$ can be rewritten as $(x - 2)^2 + 5$ by completing the square.

Example

- ▶ Claim: $x^2 - 4x + 9$ is always positive.
- ▶ Proof: We know that for any $x \in \mathbb{R}$, $x^2 \geq 0$.
- ▶ Therefore $x^2 + y$ for any $y > 0$ is always positive.
- ▶ Now note that $x^2 - 4x + 9$ can be rewritten as $(x - 2)^2 + 5$ by completing the square.
- ▶ Therefore $x^2 - 4x + 9$ is always positive. ■

Example

- ▶ Claim: if $f(x)$ is even, then it is not one-to-one.

Example

- ▶ Claim: if $f(x)$ is even, then it is not one-to-one.
- ▶ Proof: Recall that $f(x)$ is even if $f(-x) = f(x)$ for all x and is one-to-one if for all x , $f(x)$ is unique.

Example

- ▶ Claim: if $f(x)$ is even, then it is not one-to-one.
- ▶ Proof: Recall that $f(x)$ is even if $f(-x) = f(x)$ for all x and is one-to-one if for all x , $f(x)$ is unique.
- ▶ If $f(x)$ is even, then $f(x) = f(-x)$.

Example

- ▶ Claim: if $f(x)$ is even, then it is not one-to-one.
- ▶ Proof: Recall that $f(x)$ is even if $f(-x) = f(x)$ for all x and is one-to-one if for all x , $f(x)$ is unique.
- ▶ If $f(x)$ is even, then $f(x) = f(-x)$.
- ▶ Thus there exists an a and a b such that $f(a) = f(b)$.

Example

- ▶ Claim: if $f(x)$ is even, then it is not one-to-one.
- ▶ Proof: Recall that $f(x)$ is even if $f(-x) = f(x)$ for all x and is one-to-one if for all x , $f(x)$ is unique.
- ▶ If $f(x)$ is even, then $f(x) = f(-x)$.
- ▶ Thus there exists an a and a b such that $f(a) = f(b)$.
- ▶ Therefore $f(x)$ is not one-to-one. ■

Proof by Contradiction

- ▶ A second proof strategy exploits the fact that $(\neg B \wedge B)$ is logically invalid.

Proof by Contradiction

- ▶ A second proof strategy exploits the fact that $(\neg B \wedge B)$ is logically invalid.
- ▶ Prove A by showing that $\neg A \implies (\neg B \wedge B)$.

Proof by Contradiction

- ▶ A second proof strategy exploits the fact that $(\neg B \wedge B)$ is logically invalid.
- ▶ Prove A by showing that $\neg A \implies (\neg B \wedge B)$.
- ▶ B can be any statement, not necessarily one that we are trying to prove or disprove.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.
- ▶ Therefore b must be odd.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.
- ▶ Therefore b must be odd.
- ▶ Since a is even, there must be some integer c such that $a = 2c$

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.
- ▶ Therefore b must be odd.
- ▶ Since a is even, there must be some integer c such that $a = 2c$
- ▶ This yields $(2c)^2 = 2b^2$ so $4c^2 = 2b^2$ and hence $b^2 = 2c^2$

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.
- ▶ Therefore b must be odd.
- ▶ Since a is even, there must be some integer c such that $a = 2c$
- ▶ This yields $(2c)^2 = 2b^2$ so $4c^2 = 2b^2$ and hence $b^2 = 2c^2$
- ▶ This implies that b^2 is even which implies that b is also even.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.
- ▶ Therefore b must be odd.
- ▶ Since a is even, there must be some integer c such that $a = 2c$
- ▶ This yields $(2c)^2 = 2b^2$ so $4c^2 = 2b^2$ and hence $b^2 = 2c^2$
- ▶ This implies that b^2 is even which implies that b is also even.
- ▶ But we just deduced that b is odd. Therefore we have a contradiction: b is both even and odd.

Example

- ▶ Claim: $\sqrt{2}$ is irrational.
- ▶ Proof: Suppose that $\sqrt{2}$ is rational.
- ▶ Then there exist two integers, a and b such that $\frac{a}{b} = \sqrt{2}$.
- ▶ Let the fraction be fully reduced. This implies that a and b are not both even (why?).
- ▶ Our assumption implies that $a^2 = 2b^2$.
- ▶ We thus know that a^2 must be even which implies that a is even.
- ▶ Therefore b must be odd.
- ▶ Since a is even, there must be some integer c such that $a = 2c$
- ▶ This yields $(2c)^2 = 2b^2$ so $4c^2 = 2b^2$ and hence $b^2 = 2c^2$
- ▶ This implies that b^2 is even which implies that b is also even.
- ▶ But we just deduced that b is odd. Therefore we have a contradiction: b is both even and odd.
- ▶ Therefore our presumption that $\sqrt{2}$ is rational must be false.



Proof by Contradiction

- ▶ Contradiction is often a good strategy for proving statements of the form “for all x , A is true of x .”

Proof by Contradiction

- ▶ Contradiction is often a good strategy for proving statements of the form “for all x , A is true of x .”
- ▶ The setup for contradiction involves assuming that “there exists an x such that A is not true of x .”

Proof by Contradiction

- ▶ Contradiction is often a good strategy for proving statements of the form “for all x , A is true of x .”
- ▶ The setup for contradiction involves assuming that “there exists an x such that A is not true of x .”
- ▶ This gives us a specific x for which A is false which is often enough to produce a contradiction.

Example

- ▶ Claim: There exists an integer $n > 0$ such that $n^2 + n + 17$ is not a prime number.

Example

- ▶ Claim: There exists an integer $n > 0$ such that $n^2 + n + 17$ is not a prime number.
- ▶ Proof: Assume that for all integers $n > 0$, $n^2 + n + 17$ is a prime number.

Example

- ▶ Claim: There exists an integer $n > 0$ such that $n^2 + n + 17$ is not a prime number.
- ▶ Proof: Assume that for all integers $n > 0$, $n^2 + n + 17$ is a prime number.
- ▶ This implies that $n + 1 + 17/n$ is not an integer for all n (the sum is greater than 1 for all integers).

Example

- ▶ Claim: There exists an integer $n > 0$ such that $n^2 + n + 17$ is not a prime number.
- ▶ Proof: Assume that for all integers $n > 0$, $n^2 + n + 17$ is a prime number.
- ▶ This implies that $n + 1 + 17/n$ is not an integer for all n (the sum is greater than 1 for all integers).
- ▶ This therefore implies that $17/n$ is not an integer for all n which implies that 1 is not an integer which is false. ■

Proof by Contradiction

- ▶ We typically are interested in proving conditional statements in political science.

Proof by Contradiction

- ▶ We typically are interested in proving conditional statements in political science.
- ▶ To prove $A \implies B$, we assume $\neg(A \implies B)$.

Proof by Contradiction

- ▶ We typically are interested in proving conditional statements in political science.
- ▶ To prove $A \implies B$, we assume $\neg(A \implies B)$.
- ▶ That is, we assume A is true while B is false (why?).

Proof by Contradiction

- ▶ We typically are interested in proving conditional statements in political science.
- ▶ To prove $A \implies B$, we assume $\neg(A \implies B)$.
- ▶ That is, we assume A is true while B is false (why?).
- ▶ Assume $(A \wedge \neg B)$ and show that $(C \wedge \neg C)$ for some statement C .

Example

- ▶ Claim: Assume $a \in \mathbb{Z}$. If a^2 is even, then a is even.

Example

- ▶ Claim: Assume $a \in \mathbb{Z}$. If a^2 is even, then a is even.
- ▶ Proof: Assume a is odd and a^2 is even.

Example

- ▶ Claim: Assume $a \in \mathbb{Z}$. If a^2 is even, then a is even.
- ▶ Proof: Assume a is odd and a^2 is even.
- ▶ Since a is odd, there exists an integer c for which $a = 2c + 1$.

Example

- ▶ Claim: Assume $a \in \mathbb{Z}$. If a^2 is even, then a is even.
- ▶ Proof: Assume a is odd and a^2 is even.
- ▶ Since a is odd, there exists an integer c for which $a = 2c + 1$.
- ▶ Then $a^2 = 2(2c^2 + 2c) + 1$ which implies that a^2 is odd, a contradiction.

Example

- ▶ Claim: Assume $a \in \mathbb{Z}$. If a^2 is even, then a is even.
- ▶ Proof: Assume a is odd and a^2 is even.
- ▶ Since a is odd, there exists an integer c for which $a = 2c + 1$.
- ▶ Then $a^2 = 2(2c^2 + 2c) + 1$ which implies that a^2 is odd, a contradiction.
- ▶ Therefore a must be even. ■

Proof by Induction

- ▶ Some statements describe a property of an index number n and may be written as $A(n)$.

Proof by Induction

- ▶ Some statements describe a property of an index number n and may be written as $A(n)$.
- ▶ One way to prove that $A(n)$ is true for all natural numbers n is to demonstrate that $A(1)$ is true and that if $A(n)$ is true then $A(n + 1)$ must be true.

Example

► Claim: $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$

Example

- ▶ Claim: $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
- ▶ Proof: First show that the equality holds for $n = 1$.
 $1 = 1(1 + 1)/2 = 1.$

Example

- ▶ Claim: $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
- ▶ Proof: First show that the equality holds for $n = 1$.
 $1 = 1(1 + 1)/2 = 1$.
- ▶ Now assume $n = k$ is true (inductive hypothesis)

Example

- ▶ Claim: $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
- ▶ Proof: First show that the equality holds for $n = 1$.
 $1 = 1(1 + 1)/2 = 1$.
- ▶ Now assume $n = k$ is true (inductive hypothesis)
- ▶ That is, we assume $1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Example

- ▶ Claim: $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
- ▶ Proof: First show that the equality holds for $n = 1$.
 $1 = 1(1 + 1)/2 = 1$.
- ▶ Now assume $n = k$ is true (inductive hypothesis)
- ▶ That is, we assume $1 + 2 + \dots + k = \frac{k(k+1)}{2}$
- ▶ Now we just need show that $n = k + 1$ holds:

$$1 + 2 + \dots + k + (k + 1) = \frac{(k + 1)((k + 1) + 1)}{2}$$

Example (cont.)

- ▶ We need to show $1 + 2 + \dots + k + (k + 1) = \frac{(k+1)((k+1)+1)}{2}$

Example (cont.)

- ▶ We need to show $1 + 2 + \dots + k + (k + 1) = \frac{(k+1)((k+1)+1)}{2}$
- ▶ Start with the left side of the equation. By the inductive hypothesis,

$$1 + 2 + \dots + k + (k + 1) = \frac{k(k + 1)}{2} + (k + 1)$$

$$= \frac{k(k + 1) + 2(k + 1)}{2}$$

$$= \frac{(k + 2)(k + 1)}{2}$$

$$= \frac{(k + 1)(k + 2)}{2}$$

$$= \frac{(k + 1)((k + 1) + 1)}{2} \quad \blacksquare$$

Proof by Induction

- ▶ Algorithm for proof by induction is simple.

Proof by Induction

- ▶ Algorithm for proof by induction is simple.
- ▶ First prove the statement for a base case.

Proof by Induction

- ▶ Algorithm for proof by induction is simple.
- ▶ First prove the statement for a base case.
- ▶ Then assume the statement is true for some n .

Proof by Induction

- ▶ Algorithm for proof by induction is simple.
- ▶ First prove the statement for a base case.
- ▶ Then assume the statement is true for some n .
- ▶ Then show that given the inductive hypothesis (step 2), the statement holds for $n + 1$.

Proof by Induction

- ▶ Algorithm for proof by induction is simple.
- ▶ First prove the statement for a base case.
- ▶ Then assume the statement is true for some n .
- ▶ Then show that given the inductive hypothesis (step 2), the statement holds for $n + 1$.
- ▶ While the algorithm is simple, intuition for why inductive proofs are valid may take a while to understand.

Proof by Contraposition

- ▶ Recall $A \implies B$ is equivalent to $\neg B \implies \neg A$

Proof by Contraposition

- ▶ Recall $A \implies B$ is equivalent to $\neg B \implies \neg A$
- ▶ Proof by contraposition exploits this fact to prove $A \implies B$

Proof by Contraposition

- ▶ Recall $A \implies B$ is equivalent to $\neg B \implies \neg A$
- ▶ Proof by contraposition exploits this fact to prove $A \implies B$
- ▶ Often $A \implies B$ is too hard to prove by deduction, contradiction, or induction while $\neg B \implies \neg A$ is relatively simple to prove by one of these techniques.

Example

- ▶ Claim: if $7m$ is an odd integer, then m is an odd integer for $m > 1$.

Example

- ▶ Claim: if $7m$ is an odd integer, then m is an odd integer for $m > 1$.
- ▶ Proof: We will prove that m is even implies $7m$ is even.

Example

- ▶ Claim: if $7m$ is an odd integer, then m is an odd integer for $m > 1$.
- ▶ Proof: We will prove that m is even implies $7m$ is even.
- ▶ If m is even, then $m = 2k$ for some integer $k \implies 7m = 7(2k) \implies 7m = 2(7k) \implies 7m = 2n$ for some integer $n \implies 7m$ is even. ■

If and Only If

- ▶ So far we have been focusing on proving simple statements and conditional statements of the form $A \implies B$.

If and Only If

- ▶ So far we have been focusing on proving simple statements and conditional statements of the form $A \implies B$.
- ▶ To prove a conditional statement of the form $A \iff B$, we have to prove $A \implies B$ and $B \implies A$.

If and Only If

- ▶ So far we have been focusing on proving simple statements and conditional statements of the form $A \implies B$.
- ▶ To prove a conditional statement of the form $A \iff B$, we have to prove $A \implies B$ and $B \implies A$.
- ▶ We can use different proof techniques to prove both sides of the statement.