

# Continuous Functions

# Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.

# Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.
- ▶ Now we reintroduce functions as relationships between elements in two or more spaces.

# Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.
- ▶ Now we reintroduce functions as relationships between elements in two or more spaces.
- ▶ “Sets and spaces provide the basic characters of mathematical analysis, functions provide the plot” (Carter 2001).

# Motivation

- ▶ The last several lectures introduced the spaces in which we carry out research in political science.
- ▶ Now we reintroduce functions as relationships between elements in two or more spaces.
- ▶ “Sets and spaces provide the basic characters of mathematical analysis, functions provide the plot” (Carter 2001).
- ▶ In this lecture we study functions that preserve the geometric structure of the sets they associate, continuous functions.

# Functions

- ▶ A **function** is a rule  $f : X \rightarrow Y$  that assigns every element  $x$  of a set  $X$  to a single element of a set  $Y$ .

# Functions

- ▶ A **function** is a rule  $f : X \rightarrow Y$  that assigns every element  $x$  of a set  $X$  to a single element of a set  $Y$ .
- ▶  $X$  is called the **domain**

# Functions

- ▶ A **function** is a rule  $f : X \rightarrow Y$  that assigns every element  $x$  of a set  $X$  to a single element of a set  $Y$ .
- ▶  $X$  is called the **domain**
- ▶  $Y$  is called the **codomain**



# Functions

- ▶ A **function** is a rule  $f : X \rightarrow Y$  that assigns every element  $x$  of a set  $X$  to a single element of a set  $Y$ .
- ▶  $X$  is called the **domain**
- ▶  $Y$  is called the **codomain**
- ▶ A mapping is denoted  $y = f(x)$  and  $y$  is referred to as the **image** of  $x$  under  $f$ .

# Functions

- ▶ A **function** is a rule  $f : X \rightarrow Y$  that assigns every element  $x$  of a set  $X$  to a single element of a set  $Y$ .
- ▶  $X$  is called the **domain**
- ▶  $Y$  is called the **codomain**
- ▶ A mapping is denoted  $y = f(x)$  and  $y$  is referred to as the **image** of  $x$  under  $f$ .
- ▶ We typically refer to  $x$  as an **independent variable** and  $y = f(x)$  as a **dependent variable**.

# Functions

- ▶ A **function** is a rule  $f : X \rightarrow Y$  that assigns every element  $x$  of a set  $X$  to a single element of a set  $Y$ .
- ▶  $X$  is called the **domain**
- ▶  $Y$  is called the **codomain**
- ▶ A mapping is denoted  $y = f(x)$  and  $y$  is referred to as the **image** of  $x$  under  $f$ .
- ▶ We typically refer to  $x$  as an **independent variable** and  $y = f(x)$  as a **dependent variable**.
- ▶ The **range** or “image” of  $f : X \rightarrow Y$  is the set of all elements in  $Y$  that are images of elements in  $X$ , denoted

$$f(X) = \{y \in Y : y = f(x) \text{ for some } x \in X\}$$

# Functions

- ▶ If  $f(X) = Y$ , then  $f$  maps  $X$  **onto**  $Y$ .

# Functions

- ▶ If  $f(X) = Y$ , then  $f$  maps  $X$  **onto**  $Y$ .
- ▶ If  $f(x) = f(x')$  implies  $x = x'$ , then  $f$  is **one-to-one**.

# Functions

- ▶ If  $f(X) = Y$ , then  $f$  maps  $X$  **onto**  $Y$ .
- ▶ If  $f(x) = f(x')$  implies  $x = x'$ , then  $f$  is **one-to-one**.
- ▶ A function  $f$  has an **inverse**  $f^{-1} : Y \rightarrow X$  if and only if  $f$  is one-to-one and onto.

# Functions

- ▶ If  $f(X) = Y$ , then  $f$  maps  $X$  **onto**  $Y$ .
- ▶ If  $f(x) = f(x')$  implies  $x = x'$ , then  $f$  is **one-to-one**.
- ▶ A function  $f$  has an **inverse**  $f^{-1} : Y \rightarrow X$  if and only if  $f$  is one-to-one and onto.
- ▶ A **graph** of a function

$$\text{graph}(f) = \{(x, y) \in X \times Y : y = f(x), x \in X\}$$

## Example

- ▶ Let  $f(x) = x^2$  for  $X = Y = \mathbb{R}$ .



## Example

- ▶ Let  $f(x) = x^2$  for  $X = Y = \mathbb{R}$ .
- ▶ The range of  $f$  is  $f(X) = \mathbb{R}_+$ .

## Example

- ▶ Let  $f(x) = x^2$  for  $X = Y = \mathbb{R}$ .
- ▶ The range of  $f$  is  $f(X) = \mathbb{R}_+$ .
- ▶ The function is not onto.

## Example

- ▶ Let  $f(x) = x^2$  for  $X = Y = \mathbb{R}$ .
- ▶ The range of  $f$  is  $f(X) = \mathbb{R}_+$ .
- ▶ The function is not onto.
- ▶ It is also not one-to-one:  $-x^2 = x^2$ .

## Example

- ▶ Let  $f(x) = x^3$  for  $X = Y = \mathbb{R}$ .

## Example

- ▶ Let  $f(x) = x^3$  for  $X = Y = \mathbb{R}$ .
- ▶ The range of  $f$  is  $f(X) = \mathbb{R}$ .

## Example

- ▶ Let  $f(x) = x^3$  for  $X = Y = \mathbb{R}$ .
- ▶ The range of  $f$  is  $f(X) = \mathbb{R}$ .
- ▶ Therefore the function is onto.

## Example

- ▶ Let  $f(x) = x^3$  for  $X = Y = \mathbb{R}$ .
- ▶ The range of  $f$  is  $f(X) = \mathbb{R}$ .
- ▶ Therefore the function is onto.
- ▶ It is also one-to-one.

## Example

- ▶ Let  $X = \{\text{members of the department}\}$  and  $Y = \{\text{days of the year}\}$ .



## Example

- ▶ Let  $X = \{\text{members of the department}\}$  and  $Y = \{\text{days of the year}\}$ .
- ▶ Let  $f : X \rightarrow Y$  be a rule that assigns each member of the department his or her birthday.

## Example

- ▶ Let  $X = \{\text{members of the department}\}$  and  $Y = \{\text{days of the year}\}$ .
- ▶ Let  $f : X \rightarrow Y$  be a rule that assigns each member of the department his or her birthday.
- ▶ Note that this is a function: each member of the department has a unique birthday.

## Example

- ▶ Let  $X = \{\text{members of the department}\}$  and  $Y = \{\text{days of the year}\}$ .
- ▶ Let  $f : X \rightarrow Y$  be a rule that assigns each member of the department his or her birthday.
- ▶ Note that this is a function: each member of the department has a unique birthday.
- ▶ Under what conditions is  $f$  onto?

## Example

- ▶ Let  $X = \{\text{members of the department}\}$  and  $Y = \{\text{days of the year}\}$ .
- ▶ Let  $f : X \rightarrow Y$  be a rule that assigns each member of the department his or her birthday.
- ▶ Note that this is a function: each member of the department has a unique birthday.
- ▶ Under what conditions is  $f$  onto?
- ▶ One-to-one?

# Functionals

- ▶ We will often work with a special type of function in political science that maps into  $\mathbb{R}$ .

# Functionals

- ▶ We will often work with a special type of function in political science that maps into  $\mathbb{R}$ .
- ▶ A **functional** is a real-valued function  $f : X \rightarrow \mathbb{R}$ .

# Functionals

- ▶ We will often work with a special type of function in political science that maps into  $\mathbb{R}$ .
- ▶ A **functional** is a real-valued function  $f : X \rightarrow \mathbb{R}$ .
- ▶ Utility functions such as  $-(x - z)^2$  are functionals.

# Composition of Functions

- ▶ We also will often use compositions of functions in our research.



# Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions**  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  is denoted  $g \circ f$  or  $g(f(x))$  for  $x \in X$  and

$$g \circ f : X \rightarrow Z$$

# Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions**  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  is denoted  $g \circ f$  or  $g(f(x))$  for  $x \in X$  and

$$g \circ f : X \rightarrow Z$$

- ▶ *Example* Let  $f(x, y) = x^\alpha y^\beta$  and let  $g(\cdot) = \ln(\cdot)$ .

# Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions**  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  is denoted  $g \circ f$  or  $g(f(x))$  for  $x \in X$  and

$$g \circ f : X \rightarrow Z$$

- ▶ *Example* Let  $f(x, y) = x^\alpha y^\beta$  and let  $g(\cdot) = \ln(\cdot)$ .
- ▶ Note

$$f : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$$

and

$$g : \mathbb{R}_{++} \rightarrow \mathbb{R}$$

# Composition of Functions

- ▶ We also will often use compositions of functions in our research.
- ▶ A **composition of functions**  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  is denoted  $g \circ f$  or  $g(f(x))$  for  $x \in X$  and

$$g \circ f : X \rightarrow Z$$

- ▶ *Example* Let  $f(x, y) = x^\alpha y^\beta$  and let  $g(\cdot) = \ln(\cdot)$ .
- ▶ Note

$$f : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$$

and

$$g : \mathbb{R}_{++} \rightarrow \mathbb{R}$$

- ▶ The composition of these two functions is

$$g(f(x)) = \alpha \ln(x) + \beta \ln(y)$$

with  $g(f(x)) : \mathbb{R}_+^2 \rightarrow \mathbb{R}$

## Inverse Image

- ▶ An additional definition will prove useful in defining continuity.

# Inverse Image

- ▶ An additional definition will prove useful in defining continuity.
- ▶ The **inverse image** of a function of a set  $T \subseteq Y$  is the set of  $x \in X$  that are mapped to a elements  $y \in T$

$$f^{-1}(T) = \{x \in X | f(x) \in T\}$$

# Inverse Image

- ▶ An additional definition will prove useful in defining continuity.
- ▶ The **inverse image** of a function of a set  $T \subseteq Y$  is the set of  $x \in X$  that are mapped to a elements  $y \in T$

$$f^{-1}(T) = \{x \in X | f(x) \in T\}$$

- ▶ *Example:* Let  $f(x) = x^2$  and let  $T = [0, 4] \subset \mathbb{R} = Y$ .

# Inverse Image

- ▶ An additional definition will prove useful in defining continuity.
- ▶ The **inverse image** of a function of a set  $T \subseteq Y$  is the set of  $x \in X$  that are mapped to a elements  $y \in T$

$$f^{-1}(T) = \{x \in X | f(x) \in T\}$$

- ▶ *Example:* Let  $f(x) = x^2$  and let  $T = [0, 4] \subset \mathbb{R} = Y$ .
- ▶ The inverse image  $f^{-1}(T)$  is  $[0, 2] \subset \mathbb{R} = X$ .



# Continuity

- ▶ Informally, a function is continuous if small changes in  $x$  produce only small changes in  $y$ .

# Continuity

- ▶ Informally, a function is continuous if small changes in  $x$  produce only small changes in  $y$ .
- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

# Continuity

- ▶ Informally, a function is continuous if small changes in  $x$  produce only small changes in  $y$ .

- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

- ▶ In a two-candidate race,  $f(x) = 0$  for  $x < 1/2$  and  $f(x) = 1$  for  $x > 1/2$ .

# Continuity

- ▶ Informally, a function is continuous if small changes in  $x$  produce only small changes in  $y$ .

- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

- ▶ In a two-candidate race,  $f(x) = 0$  for  $x < 1/2$  and  $f(x) = 1$  for  $x > 1/2$ .
- ▶ A small change in  $x$  in the neighborhood of  $1/2$  results in a large change in  $y$ .

# Continuity

- ▶ Informally, a function is continuous if small changes in  $x$  produce only small changes in  $y$ .

- ▶ Let

$$f : [0, 1] \rightarrow \{0, 1\}$$

map a candidate's vote share in an election to a dummy variable that identifies whether or not he or she won (1) or lost (0) the election.

- ▶ In a two-candidate race,  $f(x) = 0$  for  $x < 1/2$  and  $f(x) = 1$  for  $x > 1/2$ .
- ▶ A small change in  $x$  in the neighborhood of  $1/2$  results in a large change in  $y$ .
- ▶ This is a discontinuous function.

# Continuity

- Now consider the function

$$f(x, y, z) = \beta_0 + \beta_1 x + \beta_2 y + \beta_3 z + \epsilon$$

for  $\beta_i, \epsilon, x, y, z \in \mathbb{R}$

# Continuity

- ▶ Now consider the function

$$f(x, y, z) = \beta_0 + \beta_1 x + \beta_2 y + \beta_3 z + \epsilon$$

for  $\beta_i, \epsilon, x, y, z \in \mathbb{R}$

- ▶ Note that a small change in any variable results in only a small change in  $f(x)$ .

# Continuity

- ▶ Now consider the function

$$f(x, y, z) = \beta_0 + \beta_1 x + \beta_2 y + \beta_3 z + \epsilon$$

for  $\beta_i, \epsilon, x, y, z \in \mathbb{R}$

- ▶ Note that a small change in any variable results in only a small change in  $f(x)$ .
- ▶ This is a continuous function.



# Continuity

- ▶ Recall that our concept of distance (and hence a formal notion of “small difference”) is defined for metric spaces.

# Continuity

- ▶ Recall that our concept of distance (and hence a formal notion of “small difference”) is defined for metric spaces.
- ▶ A continuous function preserves the geometric structure of the sets it associates i.e. the metric spaces it associates.

# Continuity

- ▶ Recall that our concept of distance (and hence a formal notion of “small difference”) is defined for metric spaces.
- ▶ A continuous function preserves the geometric structure of the sets it associates i.e. the metric spaces it associates.
- ▶ For the rest of this lecture we will assume that  $X$  and  $Y$  are metric spaces.

## Continuity: Epsilon-Delta Characterization

- ▶ A function  $f : X \rightarrow Y$  is continuous at  $x_0$  if for all  $\epsilon > 0$ , there exists a  $\delta > 0$  such that for every  $x \in X$ ,

$$d(x, x_0) < \delta \implies d(f(x), f(x_0)) < \epsilon$$

## Continuity: Epsilon-Delta Characterization

- ▶ A function  $f : X \rightarrow Y$  is continuous at  $x_0$  if for all  $\epsilon > 0$ , there exists a  $\delta > 0$  such that for every  $x \in X$ ,

$$d(x, x_0) < \delta \implies d(f(x), f(x_0)) < \epsilon$$

- ▶ If a function is continuous for all  $x_0$  in its domain, we call the function continuous.

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?
- ▶ Let  $\epsilon > 0$ .



## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?
- ▶ Let  $\epsilon > 0$ .
- ▶ Notice that  $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$ .

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?
- ▶ Let  $\epsilon > 0$ .
- ▶ Notice that  $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$ .
- ▶ Note that  $f(x) \leq x^2$  for all  $x$  because the range of  $\sin(\cdot)$  is  $[-1, 1]$ .

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?
- ▶ Let  $\epsilon > 0$ .
- ▶ Notice that  $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$ .
- ▶ Note that  $f(x) \leq x^2$  for all  $x$  because the range of  $\sin(\cdot)$  is  $[-1, 1]$ .
- ▶ Let  $\delta = \sqrt{\epsilon}$ .

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?
- ▶ Let  $\epsilon > 0$ .
- ▶ Notice that  $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$ .
- ▶ Note that  $f(x) \leq x^2$  for all  $x$  because the range of  $\sin(\cdot)$  is  $[-1, 1]$ .
- ▶ Let  $\delta = \sqrt{\epsilon}$ .
- ▶ Now note that  $|x - 0| < \delta$  implies  $x^2 < \delta^2 = \epsilon$ .

## Example

- ▶ Let  $f(x) = x^2 \sin(\frac{1}{x})$  for  $x \neq 0$  and  $f(0) = 0$ .
- ▶ Is  $f(x)$  continuous at  $x = 0$ ?
- ▶ Let  $\epsilon > 0$ .
- ▶ Notice that  $d(f(x), f(0)) = |f(x) - f(0)| = |f(x)|$ .
- ▶ Note that  $f(x) \leq x^2$  for all  $x$  because the range of  $\sin(\cdot)$  is  $[-1, 1]$ .
- ▶ Let  $\delta = \sqrt{\epsilon}$ .
- ▶ Now note that  $|x - 0| < \delta$  implies  $x^2 < \delta^2 = \epsilon$ .
- ▶ Therefore  $|x - 0| < \delta$  implies  $|f(x) - f(0)| < \epsilon$  so  $f$  is continuous at  $x = 0$ .

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$
- ▶ We need to show that  $d(f(x_0), f(x)) < \epsilon$  for  $d(x_0, x)$  sufficiently small where  $x$  is an element of the domain.



## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$
- ▶ We need to show that  $d(f(x_0), f(x)) < \epsilon$  for  $d(x_0, x)$  sufficiently small where  $x$  is an element of the domain.
- ▶ Note that  $d(x, x_0) = |x - x_0|$  and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$
- ▶ We need to show that  $d(f(x_0), f(x)) < \epsilon$  for  $d(x_0, x)$  sufficiently small where  $x$  is an element of the domain.
- ▶ Note that  $d(x, x_0) = |x - x_0|$  and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if  $|x - x_0| < k$ , then  $|x| - |x_0| < k$  or  $|x| < |x_0| + k$ .

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$
- ▶ We need to show that  $d(f(x_0), f(x)) < \epsilon$  for  $d(x_0, x)$  sufficiently small where  $x$  is an element of the domain.
- ▶ Note that  $d(x, x_0) = |x - x_0|$  and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if  $|x - x_0| < k$ , then  $|x| - |x_0| < k$  or  $|x| < |x_0| + k$ .
- ▶ By the triangle inequality,  $|x + x_0| \leq |x| + |x_0| < 2|x_0| + k$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$
- ▶ We need to show that  $d(f(x_0), f(x)) < \epsilon$  for  $d(x_0, x)$  sufficiently small where  $x$  is an element of the domain.
- ▶ Note that  $d(x, x_0) = |x - x_0|$  and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if  $|x - x_0| < k$ , then  $|x| - |x_0| < k$  or  $|x| < |x_0| + k$ .
- ▶ By the triangle inequality,  $|x + x_0| \leq |x| + |x_0| < 2|x_0| + k$
- ▶ Therefore

$$|f(x) - f(x_0)| \leq 2|x - x_0|(2|x_0| + k)$$

$$\text{if } |x - x_0| < k$$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Take an arbitrary  $x_0$  and  $\epsilon > 0$
- ▶ We need to show that  $d(f(x_0), f(x)) < \epsilon$  for  $d(x_0, x)$  sufficiently small where  $x$  is an element of the domain.
- ▶ Note that  $d(x, x_0) = |x - x_0|$  and observe that

$$|f(x) - f(x_0)| = |2x^2 + 1 - (2x_0^2 + 1)| = |2x^2 - 2x_0^2| = 2|x - x_0| \cdot |x + x_0|$$

- ▶ Note that if  $|x - x_0| < k$ , then  $|x| - |x_0| < k$  or  $|x| < |x_0| + k$ .
- ▶ By the triangle inequality,  $|x + x_0| \leq |x| + |x_0| < 2|x_0| + k$
- ▶ Therefore

$$|f(x) - f(x_0)| \leq 2|x - x_0|(2|x_0| + k)$$

if  $|x - x_0| < k$

- ▶ For  $2|x - x_0|(2|x_0| + k) < \epsilon$  to hold, it is sufficient that  $|x - x_0| < \frac{\epsilon}{2(2|x_0| + k)}$  and  $|x - x_0| < k$

## Example (cont.)

- So let

$$\delta = \min\left\{k, \frac{\epsilon}{2(2|x_0| + k)}\right\}$$

## Example (cont.)

- So let

$$\delta = \min\left\{k, \frac{\epsilon}{2(2|x_0| + k)}\right\}$$

- Therefore  $|x - x_0| < \delta$  implies  $|f(x) - f(x_0)| < \epsilon$

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$



## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let  $\epsilon = 1/2$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let  $\epsilon = 1/2$ .
- ▶ If  $f$  is continuous, then I can find a  $\delta$  such that  $|x| < \delta$  implies  $|f(x)| < \epsilon$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let  $\epsilon = 1/2$ .
- ▶ If  $f$  is continuous, then I can find a  $\delta$  such that  $|x| < \delta$  implies  $|f(x)| < \epsilon$ .
- ▶ This is equivalent to finding an open interval  $D = (-\delta, \delta)$  such that  $x \in D$  implies  $|f(x)| < \epsilon$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let  $\epsilon = 1/2$ .
- ▶ If  $f$  is continuous, then I can find a  $\delta$  such that  $|x| < \delta$  implies  $|f(x)| < \epsilon$ .
- ▶ This is equivalent to finding an open interval  $D = (-\delta, \delta)$  such that  $x \in D$  implies  $|f(x)| < \epsilon$ .
- ▶ But note that however small we make this interval, if  $x \in D$ , then  $-x \in D$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let  $\epsilon = 1/2$ .
- ▶ If  $f$  is continuous, then I can find a  $\delta$  such that  $|x| < \delta$  implies  $|f(x)| < \epsilon$ .
- ▶ This is equivalent to finding an open interval  $D = (-\delta, \delta)$  such that  $x \in D$  implies  $|f(x)| < \epsilon$ .
- ▶ But note that however small we make this interval, if  $x \in D$ , then  $-x \in D$ .
- ▶ If  $x$  is positive,  $f(x) = 0$  and  $f(-x) = 1 > 1/2 = \epsilon$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$
- ▶ For  $x = 0$  note that  $d(f(0), f(x)) = d(0, f(x)) = |f(x)|$
- ▶ Let  $\epsilon = 1/2$ .
- ▶ If  $f$  is continuous, then I can find a  $\delta$  such that  $|x| < \delta$  implies  $|f(x)| < \epsilon$ .
- ▶ This is equivalent to finding an open interval  $D = (-\delta, \delta)$  such that  $x \in D$  implies  $|f(x)| < \epsilon$ .
- ▶ But note that however small we make this interval, if  $x \in D$ , then  $-x \in D$ .
- ▶ If  $x$  is positive,  $f(x) = 0$  and  $f(-x) = 1 > 1/2 = \epsilon$ .
- ▶ Therefore the function is not continuous.

# Continuity: Sequence Convergence Characterization

- ▶ The  $\epsilon - \delta$  definition of continuity is difficult to work with. Often an alternative and equivalent characterization of continuity is easier:



# Continuity: Sequence Convergence Characterization

- ▶ The  $\epsilon - \delta$  definition of continuity is difficult to work with. Often an alternative and equivalent characterization of continuity is easier:
- ▶ A function  $f : X \rightarrow Y$  is continuous if and only if  $f(x) = \lim_{n \rightarrow \infty} f(x^n)$  for every sequence  $x^n \rightarrow x$ .

# Continuity: Sequence Convergence Characterization

- ▶ The  $\epsilon - \delta$  definition of continuity is difficult to work with. Often an alternative and equivalent characterization of continuity is easier:
- ▶ A function  $f : X \rightarrow Y$  is continuous if and only if  $f(x) = \lim_{n \rightarrow \infty} f(x^n)$  for every sequence  $x^n \rightarrow x$ .
- ▶ This alternative definition implies that values of  $f(x)$  are close to  $f(x_0)$  when values of  $x$  are close to  $x_0$ .

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Suppose we take an arbitrary sequence in  $X$  such that  $\lim x^n = x_0$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Suppose we take an arbitrary sequence in  $X$  such that  $\lim x^n = x_0$
- ▶ We have

$$\lim(2x^2 + 1) = 2(\lim x)^2 + 1 = 2x_0^2 + 1 = f(x_0)$$

## Example

- ▶ Let  $f(x) = 2x^2 + 1$  for  $X = \mathbb{R}$
- ▶ Suppose we take an arbitrary sequence in  $X$  such that  $\lim x^n = x_0$
- ▶ We have

$$\lim(2x^2 + 1) = 2(\lim x)^2 + 1 = 2x_0^2 + 1 = f(x_0)$$

- ▶ Therefore  $f(x)$  is continuous.

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$ .
- ▶ To show that  $f$  is discontinuous, it is sufficient to find single sequence  $x^n \rightarrow x_0$  in  $X$  such that  $\lim f(x^n) \neq f(x_0)$ .



## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$ .
- ▶ To show that  $f$  is discontinuous, it is sufficient to find single sequence  $x^n \rightarrow x_0$  in  $X$  such that  $\lim f(x^n) \neq f(x_0)$ .
- ▶ Consider the sequence  $x^n = -\frac{1}{n}$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$ .
- ▶ To show that  $f$  is discontinuous, it is sufficient to find single sequence  $x^n \rightarrow x_0$  in  $X$  such that  $\lim f(x^n) \neq f(x_0)$ .
- ▶ Consider the sequence  $x^n = -\frac{1}{n}$ .
- ▶ Note that  $\lim x^n = 0$  and  $x^n < 0$  for all  $x^n$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$ .
- ▶ To show that  $f$  is discontinuous, it is sufficient to find single sequence  $x^n \rightarrow x_0$  in  $X$  such that  $\lim f(x^n) \neq f(x_0)$ .
- ▶ Consider the sequence  $x^n = -\frac{1}{n}$ .
- ▶ Note that  $\lim x^n = 0$  and  $x^n < 0$  for all  $x^n$ .
- ▶ We therefore have  $\lim f(x^n) = \lim 1 = 1 \neq f(x^0) = 0$ .

## Example

- ▶ Let  $f(x) = 1$  for  $x < 0$  and  $f(x) = 0$  for  $x \geq 0$ .
- ▶ To show that  $f$  is discontinuous, it is sufficient to find single sequence  $x^n \rightarrow x_0$  in  $X$  such that  $\lim f(x^n) \neq f(x_0)$ .
- ▶ Consider the sequence  $x^n = -\frac{1}{n}$ .
- ▶ Note that  $\lim x^n = 0$  and  $x^n < 0$  for all  $x^n$ .
- ▶ We therefore have  $\lim f(x^n) = \lim 1 = 1 \neq f(x^0) = 0$ .
- ▶ Therefore  $f$  is discontinuous at  $x = 0$ .

## Continuity: Inverse Image Characterization

- ▶ A third characterization of continuity is often useful in proofs that require or seek to establish continuity.

## Continuity: Inverse Image Characterization

- ▶ A third characterization of continuity is often useful in proofs that require or seek to establish continuity.
- ▶ A function  $f : X \rightarrow Y$  is continuous if and only if the inverse image of any open (closed) subset of  $Y$  is an open (closed) subset of  $X$ .

## Example

- ▶ Let  $f(x) = x^3$

## Example

- ▶ Let  $f(x) = x^3$
- ▶ Consider the subset of its codomain  $(0, 2)$



## Example

- ▶ Let  $f(x) = x^3$
- ▶ Consider the subset of its codomain  $(0, 2)$
- ▶ The inverse image  $f^{-1}((0, 8))$  is the set of all points  $x$  in the domain of  $x^3$  such that  $x^3 \in (0, 8)$ .

## Example

- ▶ Let  $f(x) = x^3$
- ▶ Consider the subset of its codomain  $(0, 2)$
- ▶ The inverse image  $f^{-1}((0, 8))$  is the set of all points  $x$  in the domain of  $x^3$  such that  $x^3 \in (0, 8)$ .
- ▶ Therefore  $f^{-1}((0, 2)) = (0, 2)$ .

## Example

- ▶ Let  $f(x) = x^3$
- ▶ Consider the subset of its codomain  $(0, 2)$
- ▶ The inverse image  $f^{-1}((0, 8))$  is the set of all points  $x$  in the domain of  $x^3$  such that  $x^3 \in (0, 8)$ .
- ▶ Therefore  $f^{-1}((0, 2)) = (0, 2)$ .
- ▶ Note that this is an open subset of  $X = \mathbb{R}$ .

## Example

- ▶ Let  $f(x) = 1$  for all  $x$ .

## Example

- ▶ Let  $f(x) = 1$  for all  $x$ .
- ▶ Consider a subset of the codomain,  $(-1, 1)$ .

## Example

- ▶ Let  $f(x) = 1$  for all  $x$ .
- ▶ Consider a subset of the codomain,  $(-1, 1)$ .
- ▶ The inverse image under  $(-1, 1)$  is  $\emptyset$  which is open.

## Example

- ▶ Let  $f(x) = 1$  for all  $x$ .
- ▶ Consider a subset of the codomain,  $(-1, 1)$ .
- ▶ The inverse image under  $(-1, 1)$  is  $\emptyset$  which is open.
- ▶ Now consider  $(0, 2)$ .

## Example

- ▶ Let  $f(x) = 1$  for all  $x$ .
- ▶ Consider a subset of the codomain,  $(-1, 1)$ .
- ▶ The inverse image under  $(-1, 1)$  is  $\emptyset$  which is open.
- ▶ Now consider  $(0, 2)$ .
- ▶ The inverse image under  $(0, 2)$  is  $\mathbb{R}$ .



## Continuity: Inverse Image Characterization

- ▶ Using this result to show that a function is continuous can be difficult because we have to evaluate every subset of the codomain.

## Continuity: Inverse Image Characterization

- ▶ Using this result to show that a function is continuous can be difficult because we have to evaluate every subset of the codomain.
- ▶ It is easier to use the definition to show discontinuity because we only need to find a single example of a set that violates the result.

## Example

- ▶ Let  $f(x) = x$  for  $x \leq 0$  and let  $f(x) = x + 1$  for  $x > 0$ .

## Example

- ▶ Let  $f(x) = x$  for  $x \leq 0$  and let  $f(x) = x + 1$  for  $x > 0$ .
- ▶ Consider the open subset of the codomain  $(-1, 1) \subset Y$ .

## Example

- ▶ Let  $f(x) = x$  for  $x \leq 0$  and let  $f(x) = x + 1$  for  $x > 0$ .
- ▶ Consider the open subset of the codomain  $(-1, 1) \subset Y$ .
- ▶ The inverse image under  $(-1, 1)$  is  $(-1, 1]$  which is not open.

## Example

- ▶ Let  $f(x) = x$  for  $x \leq 0$  and let  $f(x) = x + 1$  for  $x > 0$ .
- ▶ Consider the open subset of the codomain  $(-1, 1) \subset Y$ .
- ▶ The inverse image under  $(-1, 1)$  is  $(-1, 1]$  which is not open.
- ▶ Therefore  $f(x)$  is not continuous.

## Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .



# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Note that  $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$ .

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Note that  $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$ .
- ▶ Therefore  $kf$  is continuous at  $x_0$ .

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Note that  $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$ .
- ▶ Therefore  $kf$  is continuous at  $x_0$ .
- ▶ Now consider  $|f|$  at  $x_0$ .

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Note that  $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$ .
- ▶ Therefore  $kf$  is continuous at  $x_0$ .
- ▶ Now consider  $|f|$  at  $x_0$ .
- ▶ By the reverse triangle inequality,

$$||f(x_n)| - |f(x_0)|| \leq |f(x_n) - f(x_0)|$$

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Note that  $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$ .
- ▶ Therefore  $kf$  is continuous at  $x_0$ .
- ▶ Now consider  $|f|$  at  $x_0$ .
- ▶ By the reverse triangle inequality,

$$||f(x_n)| - |f(x_0)|| \leq |f(x_n) - f(x_0)|$$

- ▶ Now consider  $\epsilon > 0$

# Properties of Continuous Functions

- ▶ Let  $f$  be a real-valued function with domain  $X \subseteq \mathbb{R}$ . If  $f$  is continuous at  $x_0$  in  $X$ , then  $|f|$  and  $kf$ ,  $k \in \mathbb{R}$  are continuous at  $x_0$ .
- ▶ **Proof:** Consider the sequence  $x_n \rightarrow x_0$  in the domain of  $f$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Note that  $\lim kf(x_n) = k \lim f(x_n) = kf(x_0)$ .
- ▶ Therefore  $kf$  is continuous at  $x_0$ .
- ▶ Now consider  $|f|$  at  $x_0$ .
- ▶ By the reverse triangle inequality,

$$||f(x_n)| - |f(x_0)|| \leq |f(x_n) - f(x_0)|$$

- ▶ Now consider  $\epsilon > 0$
- ▶ Because  $\lim f(x_n) = f(x_0)$ , there exists an  $N$  such that  $n > N$  implies  $||f(x_n)| - |f(x_0)|| < \epsilon$  and thus  $\lim |f(x_n)| = |f(x_0)|$ .

# Properties of Continuous Functions

- ▶ If  $f$  and  $g$  are real-valued functions and continuous at  $x_0$ , then



# Properties of Continuous Functions

- ▶ If  $f$  and  $g$  are real-valued functions and continuous at  $x_0$ , then
- ▶ i)  $f + g$  is continuous at  $x_0$

# Properties of Continuous Functions

- ▶ If  $f$  and  $g$  are real-valued functions and continuous at  $x_0$ , then
- ▶ i)  $f + g$  is continuous at  $x_0$
- ▶ ii)  $fg$  is continuous at  $x_0$

# Properties of Continuous Functions

- ▶ If  $f$  and  $g$  are real-valued functions and continuous at  $x_0$ , then
- ▶ i)  $f + g$  is continuous at  $x_0$
- ▶ ii)  $fg$  is continuous at  $x_0$
- ▶ iii)  $\frac{f}{g}$  is continuous at  $x_0$  if  $g(x_0) \neq 0$ .

## Proof of $f + g$ continuity at $x_0$

- ▶  $x_0$  is in the domain of  $f$  and  $g$ .

## Proof of $f + g$ continuity at $x_0$

- ▶  $x_0$  is in the domain of  $f$  and  $g$ .
- ▶ Let  $x_n$  be a sequence in the intersection of their domains that converges to  $x_0$ .

## Proof of $f + g$ continuity at $x_0$

- ▶  $x_0$  is in the domain of  $f$  and  $g$ .
- ▶ Let  $x_n$  be a sequence in the intersection of their domains that converges to  $x_0$ .
- ▶ Because each is continuous, we have  $\lim f(x_n) = f(x_0)$  and  $\lim g(x_n) = g(x_0)$ .

## Proof of $f + g$ continuity at $x_0$

- ▶  $x_0$  is in the domain of  $f$  and  $g$ .
- ▶ Let  $x_n$  be a sequence in the intersection of their domains that converges to  $x_0$ .
- ▶ Because each is continuous, we have  $\lim f(x_n) = f(x_0)$  and  $\lim g(x_n) = g(x_0)$ .
- ▶ Therefore

$$\begin{aligned}\lim(f + g)(x_n) &= \lim[f(x_n) + g(x_n)] = \lim f(x_n) + \lim g(x_n) \\ &= f(x_0) + g(x_0) = (f + g)(x_0)\end{aligned}$$

## Proof of $f + g$ continuity at $x_0$

- ▶  $x_0$  is in the domain of  $f$  and  $g$ .
- ▶ Let  $x_n$  be a sequence in the intersection of their domains that converges to  $x_0$ .
- ▶ Because each is continuous, we have  $\lim f(x_n) = f(x_0)$  and  $\lim g(x_n) = g(x_0)$ .
- ▶ Therefore

$$\begin{aligned}\lim(f + g)(x_n) &= \lim[f(x_n) + g(x_n)] = \lim f(x_n) + \lim g(x_n) \\ &= f(x_0) + g(x_0) = (f + g)(x_0)\end{aligned}$$

- ▶ Therefore  $f + g$  is continuous at  $x_0$ .



## Proof of $f + g$ continuity at $x_0$

- ▶  $x_0$  is in the domain of  $f$  and  $g$ .
- ▶ Let  $x_n$  be a sequence in the intersection of their domains that converges to  $x_0$ .
- ▶ Because each is continuous, we have  $\lim f(x_n) = f(x_0)$  and  $\lim g(x_n) = g(x_0)$ .
- ▶ Therefore

$$\begin{aligned}\lim(f + g)(x_n) &= \lim[f(x_n) + g(x_n)] = \lim f(x_n) + \lim g(x_n) \\ &= f(x_0) + g(x_0) = (f + g)(x_0)\end{aligned}$$

- ▶ Therefore  $f + g$  is continuous at  $x_0$ .
- ▶ A similar proof establishes that  $fg$  and  $f/g$  are continuous at  $x_0$ .

# Properties of Continuous Functions

- ▶ If  $f$  is continuous at  $x_0$  and  $g$  is continuous at  $f(x_0)$ , then  $g(f(x))$  is continuous at  $x_0$ .

# Properties of Continuous Functions

- ▶ If  $f$  is continuous at  $x_0$  and  $g$  is continuous at  $f(x_0)$ , then  $g(f(x))$  is continuous at  $x_0$ .
- ▶ **Proof:**  $x_0$  is in the domain of  $f$  and  $f(x_0)$  is in the domain of  $g$

# Properties of Continuous Functions

- ▶ If  $f$  is continuous at  $x_0$  and  $g$  is continuous at  $f(x_0)$ , then  $g(f(x))$  is continuous at  $x_0$ .
- ▶ **Proof:**  $x_0$  is in the domain of  $f$  and  $f(x_0)$  is in the domain of  $g$
- ▶ Let  $x_n$  be a sequence such that  $x_n$  is in the domain of  $f$  and  $f(x_n)$  is in the domain of  $g$  converging to  $x_0$ .

# Properties of Continuous Functions

- ▶ If  $f$  is continuous at  $x_0$  and  $g$  is continuous at  $f(x_0)$ , then  $g(f(x))$  is continuous at  $x_0$ .
- ▶ **Proof:**  $x_0$  is in the domain of  $f$  and  $f(x_0)$  is in the domain of  $g$
- ▶ Let  $x_n$  be a sequence such that  $x_n$  is in the domain of  $f$  and  $f(x_n)$  is in the domain of  $g$  converging to  $x_0$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .

# Properties of Continuous Functions

- ▶ If  $f$  is continuous at  $x_0$  and  $g$  is continuous at  $f(x_0)$ , then  $g(f(x))$  is continuous at  $x_0$ .
- ▶ **Proof:**  $x_0$  is in the domain of  $f$  and  $f(x_0)$  is in the domain of  $g$
- ▶ Let  $x_n$  be a sequence such that  $x_n$  is in the domain of  $f$  and  $f(x_n)$  is in the domain of  $g$  converging to  $x_0$ .
- ▶ Because  $f$  is continuous at  $x_0$ ,  $\lim f(x_n) = f(x_0)$ .
- ▶ Because  $f(x_n)$  converges to  $f(x_0)$  and  $g$  is continuous at  $f(x_0)$ , we have

$$\lim g(f(x_n)) = g(f(x_0))$$

# Properties of Continuous Functions

- ▶ A continuous functional  $f : X \rightarrow Y$  on a compact set achieves a maximum and a minimum (Weierstrass theorem).

# Properties of Continuous Functions

- ▶ A continuous functional  $f : X \rightarrow Y$  on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because  $X$  is bounded,  $f(x)$  has a supremum.



# Properties of Continuous Functions

- ▶ A continuous functional  $f : X \rightarrow Y$  on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because  $X$  is bounded,  $f(x)$  has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.

# Properties of Continuous Functions

- ▶ A continuous functional  $f : X \rightarrow Y$  on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because  $X$  is bounded,  $f(x)$  has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.
- ▶ Let  $M = \sup_{x \in X} f(x)$ .

# Properties of Continuous Functions

- ▶ A continuous functional  $f : X \rightarrow Y$  on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because  $X$  is bounded,  $f(x)$  has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.
- ▶ Let  $M = \sup_{x \in X} f(x)$ .
- ▶ Take a sequence  $f(x_n) \in Y$  with  $f(x_n) \rightarrow M$ .

# Properties of Continuous Functions

- ▶ A continuous functional  $f : X \rightarrow Y$  on a compact set achieves a maximum and a minimum (Weierstrass theorem).
- ▶ **Proof:** Note that because  $X$  is bounded,  $f(x)$  has a supremum.
- ▶ We need to show that the supremum of the function's image is contained within the image.
- ▶ Let  $M = \sup_{x \in X} f(x)$ .
- ▶ Take a sequence  $f(x_n) \in Y$  with  $f(x_n) \rightarrow M$ .
- ▶ Then  $x_n \in X$ .

# Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set  $S \subseteq \mathbb{R}^n$  has a convergent subsequence which converges to a point in  $S$ .

# Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set  $S \subseteq \mathbb{R}^n$  has a convergent subsequence which converges to a point in  $S$ .
- ▶ Therefore because  $X$  is compact, we can find a convergent subsequence  $x_{n_k} \rightarrow x_0$  where  $f(x_{n_k}) \rightarrow M$ .

# Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set  $S \subseteq \mathbb{R}^n$  has a convergent subsequence which converges to a point in  $S$ .
- ▶ Therefore because  $X$  is compact, we can find a convergent subsequence  $x_{n_k} \rightarrow x_0$  where  $f(x_{n_k}) \rightarrow M$ .
- ▶ Applying the limit definition of continuity, we have  $\lim f(x_{n_k}) = f(x_0) = M$ .

# Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set  $S \subseteq \mathbb{R}^n$  has a convergent subsequence which converges to a point in  $S$ .
- ▶ Therefore because  $X$  is compact, we can find a convergent subsequence  $x_{n_k} \rightarrow x_0$  where  $f(x_{n_k}) \rightarrow M$ .
- ▶ Applying the limit definition of continuity, we have  $\lim f(x_{n_k}) = f(x_0) = M$ .
- ▶ Therefore  $f(x_0) = M$



# Properties of Continuous Functions

- ▶ Recall the Bolzano-Weierstrass Theorem: every sequence in a compact set  $S \subseteq \mathbb{R}^n$  has a convergent subsequence which converges to a point in  $S$ .
- ▶ Therefore because  $X$  is compact, we can find a convergent subsequence  $x_{n_k} \rightarrow x_0$  where  $f(x_{n_k}) \rightarrow M$ .
- ▶ Applying the limit definition of continuity, we have  $\lim f(x_{n_k}) = f(x_0) = M$ .
- ▶ Therefore  $f(x_0) = M$
- ▶ The same proof can be applied to the infimum of the set.