

Normed Linear Spaces

Motivation

- ▶ Often when carrying out theoretical and empirical political analysis, we want to work in sets that are simultaneously linear spaces and metric spaces.

Motivation

- ▶ Often when carrying out theoretical and empirical political analysis, we want to work in sets that are simultaneously linear spaces and metric spaces.
- ▶ In particular, we want to work in spaces where the algebraic structure of the space works is consistent with its geometric structure.

Motivation

- ▶ Often when carrying out theoretical and empirical political analysis, we want to work in sets that are simultaneously linear spaces and metric spaces.
- ▶ In particular, we want to work in spaces where the algebraic structure of the space works is consistent with its geometric structure.
- ▶ A normed linear space accomplishes this.

Motivation

- ▶ Often when carrying out theoretical and empirical political analysis, we want to work in sets that are simultaneously linear spaces and metric spaces.
- ▶ In particular, we want to work in spaces where the algebraic structure of the space works is consistent with its geometric structure.
- ▶ A normed linear space accomplishes this.
- ▶ In the quantitative sequence, you will be working almost exclusively in normed linear spaces.

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:
 - ▶ $d(x, y) \geq 0$
 - ▶ $d(x, y) = 0 \iff x = y$
 - ▶ $d(x, y) = d(y, x)$
 - ▶ $d(x, z) \leq d(x, y) + d(y, z)$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:
 - ▶ $d(x, y) \geq 0$
 - ▶ $d(x, y) = 0 \iff x = y$
 - ▶ $d(x, y) = d(y, x)$
 - ▶ $d(x, z) \leq d(x, y) + d(y, z)$
- ▶ Proof

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:
 - ▶ $d(x, y) \geq 0$
 - ▶ $d(x, y) = 0 \iff x = y$
 - ▶ $d(x, y) = d(y, x)$
 - ▶ $d(x, z) \leq d(x, y) + d(y, z)$
- ▶ Proof
 - ▶ $d(x, y) = \|x - y\| \geq 0$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:
 - ▶ $d(x, y) \geq 0$
 - ▶ $d(x, y) = 0 \iff x = y$
 - ▶ $d(x, y) = d(y, x)$
 - ▶ $d(x, z) \leq d(x, y) + d(y, z)$
- ▶ Proof
 - ▶ $d(x, y) = \|x - y\| \geq 0$
 - ▶ $d(x, y) = 0 \iff \|x - y\| = 0 \iff x - y = 0 \iff x = y$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:
 - ▶ $d(x, y) \geq 0$
 - ▶ $d(x, y) = 0 \iff x = y$
 - ▶ $d(x, y) = d(y, x)$
 - ▶ $d(x, z) \leq d(x, y) + d(y, z)$
- ▶ Proof
 - ▶ $d(x, y) = \|x - y\| \geq 0$
 - ▶ $d(x, y) = 0 \iff \|x - y\| = 0 \iff x - y = 0 \iff x = y$
 - ▶ $d(x, y) = \|x - y\| = \|(-1)(y - x)\| = (-1)\|y - x\| = d(x, y)$

Norms

- ▶ For any linear space X , a **norm**, denoted $\|x\|$, is a measure of the size of the elements satisfying the following properties:
 - ▶ 1. $\|x\| \geq 0$
 - ▶ 2. $\|x\| = 0$ if and only if $x = 0$
 - ▶ 3. $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{R}$
 - ▶ 4. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- ▶ A norm on X induces a metric on X , $d(x, y) = \|x - y\|$
- ▶ Recall properties of metric:
 - ▶ $d(x, y) \geq 0$
 - ▶ $d(x, y) = 0 \iff x = y$
 - ▶ $d(x, y) = d(y, x)$
 - ▶ $d(x, z) \leq d(x, y) + d(y, z)$
- ▶ Proof
 - ▶ $d(x, y) = \|x - y\| \geq 0$
 - ▶ $d(x, y) = 0 \iff \|x - y\| = 0 \iff x - y = 0 \iff x = y$
 - ▶ $d(x, y) = \|x - y\| = \|(-1)(y - x)\| = (-1)\|y - x\| = d(x, y)$
 - ▶ $d(x, z) = \|x - z\| = \|x - y + y - z\| \leq \|x - y\| + \|y - z\| = d(x, y) + d(y, z)$

Normed Linear Spaces

- ▶ A linear space with a norm, $(X, \|\cdot\|)$, is called a **normed linear space**.

Normed Linear Spaces

- ▶ A linear space with a norm, $(X, \|\cdot\|)$, is called a **normed linear space**.
- ▶ A normed linear space is a subclass of **metric linear spaces** which is a linear space equipped with a metric.

Normed Linear Spaces

- ▶ A linear space with a norm, $(X, \|\cdot\|)$, is called a **normed linear space**.
- ▶ A normed linear space is a subclass of **metric linear spaces** which is a linear space equipped with a metric.
- ▶ It is a special type of metric linear space with rich interaction of the algebraic and linear structures.

Example

- ▶ Say a government agency is devising a proposal to build a new highway.

Example

- ▶ Say a government agency is devising a proposal to build a new highway.
- ▶ A production plan $y = (y_1, \dots, y_n)$ is a list of the net outputs of various goods and services where y_i is net output of commodity i .

Example

- ▶ Say a government agency is devising a proposal to build a new highway.
- ▶ A production plan $y = (y_1, \dots, y_n)$ is a list of the net outputs of various goods and services where y_i is net output of commodity i .
- ▶ How could we measure the overall “size” of the plan?

Example

- ▶ Say a government agency is devising a proposal to build a new highway.
- ▶ A production plan $y = (y_1, \dots, y_n)$ is a list of the net outputs of various goods and services where y_i is net output of commodity i .
- ▶ How could we measure the overall “size” of the plan?
- ▶ What if we simply added up the net outputs of all the goods and services i.e. $\sum y_i$?

Example

- ▶ Say a government agency is devising a proposal to build a new highway.
- ▶ A production plan $y = (y_1, \dots, y_n)$ is a list of the net outputs of various goods and services where y_i is net output of commodity i .
- ▶ How could we measure the overall “size” of the plan?
- ▶ What if we simply added up the net outputs of all the goods and services i.e. $\sum y_i$?
- ▶ Since some of the net outputs are negative, it would be more appropriate to take their absolute value first.

Example

- ▶ Say a government agency is devising a proposal to build a new highway.
- ▶ A production plan $y = (y_1, \dots, y_n)$ is a list of the net outputs of various goods and services where y_i is net output of commodity i .
- ▶ How could we measure the overall “size” of the plan?
- ▶ What if we simply added up the net outputs of all the goods and services i.e. $\sum y_i$?
- ▶ Since some of the net outputs are negative, it would be more appropriate to take their absolute value first.
- ▶ Therefore one possible measure of size is

$$\|y\|_1 = \sum_{i=1}^n |y_i|$$

Example

- ▶ Another way to compensate for the negativity of some net inputs is to square the individual inputs as in

$$\|y\|_2 = \sqrt{\sum_{i=1}^n y_i^2}$$

Example

- ▶ Another way to compensate for the negativity of some net inputs is to square the individual inputs as in

$$\|y\|_2 = \sqrt{\sum_{i=1}^n y_i^2}$$

- ▶ What we probably really care about is the “output” of the project i.e. how much road will the agency produce.

Example

- ▶ Another way to compensate for the negativity of some net inputs is to square the individual inputs as in

$$\|y\|_2 = \sqrt{\sum_{i=1}^n y_i^2}$$

- ▶ What we probably really care about is the “output” of the project i.e. how much road will the agency produce.
- ▶ Can we measure the size of the plan y by the output of road, y_r ?

Example

- ▶ Another way to compensate for the negativity of some net inputs is to square the individual inputs as in

$$\|y\|_2 = \sqrt{\sum_{i=1}^n y_i^2}$$

- ▶ What we probably really care about is the “output” of the project i.e. how much road will the agency produce.
- ▶ Can we measure the size of the plan y by the output of road, y_r ?
- ▶ No: $|y_r| = 0$ does not imply $\|y\| = 0$

Example

- ▶ Another way to compensate for the negativity of some net inputs is to square the individual inputs as in

$$\|y\|_2 = \sqrt{\sum_{i=1}^n y_i^2}$$

- ▶ What we probably really care about is the “output” of the project i.e. how much road will the agency produce.
- ▶ Can we measure the size of the plan y by the output of road, y_r ?
- ▶ No: $|y_r| = 0$ does not imply $\|y\| = 0$
- ▶ It is possible for a government program to consume inputs and produce no outputs.

Example

- ▶ Another way to compensate for the negativity of some net inputs is to square the individual inputs as in

$$\|y\|_2 = \sqrt{\sum_{i=1}^n y_i^2}$$

- ▶ What we probably really care about is the “output” of the project i.e. how much road will the agency produce.
- ▶ Can we measure the size of the plan y by the output of road, y_r ?
- ▶ No: $|y_r| = 0$ does not imply $\|y\| = 0$
- ▶ It is possible for a government program to consume inputs and produce no outputs.
- ▶ A related measure that does meet the requirements of a norm uses the largest component of y as the measure of the size or the plan:

$$\|y\|_\infty = \max_{i=1}^n |y_i|$$

Euclidean Space

- ▶ The **Euclidean norm**, $\|x\|_2$ generalizes the conventional Pythagorean notion of the length of a vector in two and three-dimensional space.

Euclidean Space

- ▶ The **Euclidean norm**, $\|x\|_2$ generalizes the conventional Pythagorean notion of the length of a vector in two and three-dimensional space.
- ▶ In \mathbb{R}^2 , the Euclidean norm expresses the Pythagorean theorem that in a right triangle, the squared length of the hypotenuse is equal to the sum of the squares of the two sides:
$$\|x\|^2 = |x_1|^2 + |x_2|^2.$$

Euclidean Space

- ▶ The **Euclidean norm**, $\|x\|_2$ generalizes the conventional Pythagorean notion of the length of a vector in two and three-dimensional space.
- ▶ In \mathbb{R}^2 , the Euclidean norm expresses the Pythagorean theorem that in a right triangle, the squared length of the hypotenuse is equal to the sum of the squares of the two sides:
$$\|x\|^2 = |x_1|^2 + |x_2|^2.$$
- ▶ The space $(\mathbb{R}^n, \|\cdot\|_2)$ is referred to as a **Euclidean space**.

Banach Spaces

- ▶ As we saw in the metric spaces lecture, completeness is one of the most desirable properties of a metric space.

Banach Spaces

- ▶ As we saw in the metric spaces lecture, completeness is one of the most desirable properties of a metric space.
- ▶ Recall that a metric space is called complete if every Cauchy sequence converges in the space.

Banach Spaces

- ▶ As we saw in the metric spaces lecture, completeness is one of the most desirable properties of a metric space.
- ▶ Recall that a metric space is called complete if every Cauchy sequence converges in the space.
- ▶ A complete normed linear space is referred to as a **Banach space**.

Banach Spaces

- ▶ As we saw in the metric spaces lecture, completeness is one of the most desirable properties of a metric space.
- ▶ Recall that a metric space is called complete if every Cauchy sequence converges in the space.
- ▶ A complete normed linear space is referred to as a **Banach space**.
- ▶ Euclidean space is an example of a Banach space.

Banach Spaces

- ▶ As we saw in the metric spaces lecture, completeness is one of the most desirable properties of a metric space.
- ▶ Recall that a metric space is called complete if every Cauchy sequence converges in the space.
- ▶ A complete normed linear space is referred to as a **Banach space**.
- ▶ Euclidean space is an example of a Banach space.
- ▶ It turns out that any finite dimensional normed linear space is a Banach space.

Series

- ▶ Given a sequence x_1, x_2, x_3, \dots of elements of a normed linear space, their sum is called a **series**.

Series

- ▶ Given a sequence x_1, x_2, x_3, \dots of elements of a normed linear space, their sum is called a **series**.
- ▶ We typically deal with infinite series in game theory when we study repeated games.

Series

- ▶ Given a sequence x_1, x_2, x_3, \dots of elements of a normed linear space, their sum is called a **series**.
- ▶ We typically deal with infinite series in game theory when we study repeated games.
- ▶ For an infinite series, we can define a sequence of **partial sums** whose n th term is

$$s_n = \sum_{i=1}^n x_i$$

Series

- ▶ Given a sequence x_1, x_2, x_3, \dots of elements of a normed linear space, their sum is called a **series**.
- ▶ We typically deal with infinite series in game theory when we study repeated games.
- ▶ For an infinite series, we can define a sequence of **partial sums** whose n th term is

$$s_n = \sum_{i=1}^n x_i$$

- ▶ If s_1, s_2, \dots converges to some s , we say that the series x_1, x_2, \dots is **convergent** and we call $s = x_1 + x_2 + \dots$ the sum of the infinite series.

Example

$$\left(\frac{1}{2^k}\right)_{k=1}^{\infty} = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$$

$$s_1 = \frac{1}{2}$$

$$s_2 = \frac{1}{2} + \frac{1}{4} = \frac{3}{4}$$

$$s_3 = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} = \frac{7}{8}$$

$$s_1, s_2, s_3, \dots = \frac{1}{2}, \frac{3}{4}, \frac{7}{8}, \dots \rightarrow 1$$

$$s = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = 1$$

Divergence

- ▶ If an infinite series does not have a sum, we say that the series is **divergent**.

Divergence

- ▶ If an infinite series does not have a sum, we say that the series is **divergent**.
- ▶ *Example:* $(1)_{k=1}^{\infty} = 1 + 1 + 1 + \dots$

Divergence

- ▶ If an infinite series does not have a sum, we say that the series is **divergent**.
- ▶ $s_1, s_2, s_3, s_4, \dots = 1, 2, 3, 4, \dots$ which tends to infinity. ■

Divergence

- ▶ If an infinite series does not have a sum, we say that the series is **divergent**.
- ▶ $s_1, s_2, s_3, s_4, \dots = 1, 2, 3, 4, \dots$ which tends to infinity. ■
- ▶ When we have a convergent series, the terms in the sequence tend towards zero.

Divergence

- ▶ If an infinite series does not have a sum, we say that the series is **divergent**.
- ▶ $s_1, s_2, s_3, s_4, \dots = 1, 2, 3, 4, \dots$ which tends to infinity. ■
- ▶ When we have a convergent series, the terms in the sequence tend towards zero.
- ▶ The reverse of this, however, is not true.

Divergence

- ▶ If an infinite series does not have a sum, we say that the series is **divergent**.
- ▶ $s_1, s_2, s_3, s_4, \dots = 1, 2, 3, 4, \dots$ which tends to infinity. ■
- ▶ When we have a convergent series, the terms in the sequence tend towards zero.
- ▶ The reverse of this, however, is not true.
- ▶ A sequence that converges does not necessarily induce a sequence of partial sums that converges.

Example

- ▶ The series $(\frac{1}{k})_{k=1}^{\infty}$ is called the *harmonic series*.

Example

- ▶ The series $(\frac{1}{k})_{k=1}^{\infty}$ is called the *harmonic series*.
- ▶ The sequence $\frac{1}{n}$ converges to zero.

Example

- ▶ The series $(\frac{1}{k})_{k=1}^{\infty}$ is called the *harmonic series*.
- ▶ The sequence $\frac{1}{n}$ converges to zero.
- ▶ The sequence of partial sums of the series however diverges:

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \frac{1}{9} + \dots$$

$$> 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{4} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{16} + \dots$$

$$= 1 + \left(\frac{1}{2}\right) + \left(\frac{1}{4} + \frac{1}{4}\right) + \left(\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\right) + \left(\frac{1}{16} + \dots + \frac{1}{16}\right)$$

$$= 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \dots = \infty$$

Geometric Series

- ▶ A special type of sequence in which each term in the sequence is a constant multiple of the previous term is called a **geometric series**: $x + \delta x + \delta^2 x + \dots$

Geometric Series

- ▶ A special type of sequence in which each term in the sequence is a constant multiple of the previous term is called a **geometric series**: $x + \delta x + \delta^2 x + \dots$
- ▶ A geometric series converges if and only if $|\delta| < 1$ and converges to $\frac{x}{1-\delta}$.

Geometric Series

- ▶ A special type of sequence in which each term in the sequence is a constant multiple of the previous term is called a **geometric series**: $x + \delta x + \delta^2 x + \dots$
- ▶ A geometric series converges if and only if $|\delta| < 1$ and converges to $\frac{x}{1-\delta}$.
- ▶ *Example*: $1 + \frac{1}{2} + \frac{1}{4} + \dots = \sum_{i=1}^{\infty} \left(\frac{1}{2}\right)^{i-1} = \frac{1}{1-1/2} = 2$

Geometric Series

- ▶ A special type of sequence in which each term in the sequence is a constant multiple of the previous term is called a **geometric series**: $x + \delta x + \delta^2 x + \dots$
- ▶ A geometric series converges if and only if $|\delta| < 1$ and converges to $\frac{x}{1-\delta}$.
- ▶ *Example*: $1 + \frac{1}{2} + \frac{1}{4} + \dots = \sum_{i=1}^{\infty} (\frac{1}{2})^{i-1} = \frac{1}{1-1/2} = 2$
- ▶ Geometric series will play an important role in models where players discount the future.

Compactness in Euclidean Space

- ▶ Recall that every compact set in a metric space is closed and bounded.

Compactness in Euclidean Space

- ▶ Recall that every compact set in a metric space is closed and bounded.
- ▶ The reverse, however, is not guaranteed to hold.

Compactness in Euclidean Space

- ▶ Recall that every compact set in a metric space is closed and bounded.
- ▶ The reverse, however, is not guaranteed to hold.
- ▶ We have seen already via the Heine-Borel theorem that the two are equivalent for a subset S of (\mathbb{R}^n, d_2) .

Compactness in Euclidean Space

- ▶ Recall that every compact set in a metric space is closed and bounded.
- ▶ The reverse, however, is not guaranteed to hold.
- ▶ We have seen already via the Heine-Borel theorem that the two are equivalent for a subset S of (\mathbb{R}^n, d_2) .
- ▶ Therefore the two are also equivalent in $(\mathbb{R}^n, \|\cdot\|_2)$ i.e. in Euclidean space.

Compactness in Euclidean Space

- ▶ Recall that every compact set in a metric space is closed and bounded.
- ▶ The reverse, however, is not guaranteed to hold.
- ▶ We have seen already via the Heine-Borel theorem that the two are equivalent for a subset S of (\mathbb{R}^n, d_2) .
- ▶ Therefore the two are also equivalent in $(\mathbb{R}^n, \|\cdot\|_2)$ i.e. in Euclidean space.
- ▶ We originally took this powerful result for granted.

Compactness in Euclidean Space

- ▶ Recall that every compact set in a metric space is closed and bounded.
- ▶ The reverse, however, is not guaranteed to hold.
- ▶ We have seen already via the Heine-Borel theorem that the two are equivalent for a subset S of (\mathbb{R}^n, d_2) .
- ▶ Therefore the two are also equivalent in $(\mathbb{R}^n, \|\cdot\|_2)$ i.e. in Euclidean space.
- ▶ We originally took this powerful result for granted.
- ▶ Now we will prove it.

Limit Points

- ▶ A **limit point** (or “accumulation” point or “cluster” point) of a set A is a point a such that for all $r > 0$, there exists at least one point $c \in A \setminus \{a\}$ such that $c \in B_r(a)$.

Limit Points

- ▶ A **limit point** (or “accumulation” point or “cluster” point) of a set A is a point a such that for all $r > 0$, there exists at least one point $c \in A \setminus \{a\}$ such that $c \in B_r(a)$.
- ▶ Intuitively, a limit point is a set where no matter how little you move off of it, you intersect points of the set.

Limit Points

- ▶ A **limit point** (or “accumulation” point or “cluster” point) of a set A is a point a such that for all $r > 0$, there exists at least one point $c \in A \setminus \{a\}$ such that $c \in B_r(a)$.
- ▶ Intuitively, a limit point is a set where no matter how little you move off of it, you intersect points of the set.
- ▶ Limit points need not be in the set.

Limit Points

- ▶ A **limit point** (or “accumulation” point or “cluster” point) of a set A is a point a such that for all $r > 0$, there exists at least one point $c \in A \setminus \{a\}$ such that $c \in B_r(a)$.
- ▶ Intuitively, a limit point is a set where no matter how little you move off of it, you intersect points of the set.
- ▶ Limit points need not be in the set.
- ▶ *Example:* Every point in $[0, 1]$ is a limit point of $(0, 1)$.

Compactness in Euclidean Space

Lemma

A closed subset of a compact set is compact.

Compactness in Euclidean Space

Lemma

A closed subset of a compact set is compact.

- ▶ Let K be a closed subset of a compact set $T \subset X$ and let C_K be an open cover of K .

Compactness in Euclidean Space

Lemma

A closed subset of a compact set is compact.

- ▶ Let K be a closed subset of a compact set $T \subset X$ and let C_K be an open cover of K .
- ▶ Then $U = X \setminus K$ is an open set and $C_T = C_K \cup U$ is an open cover of T .

Compactness in Euclidean Space

Lemma

A closed subset of a compact set is compact.

- ▶ Let K be a closed subset of a compact set $T \subset X$ and let C_K be an open cover of K .
- ▶ Then $U = X \setminus K$ is an open set and $C_T = C_K \cup U$ is an open cover of T .
- ▶ Since T is compact, then C_T has a finite subcover C'_T that also covers K .

Compactness in Euclidean Space

Lemma

A closed subset of a compact set is compact.

- ▶ Let K be a closed subset of a compact set $T \subset X$ and let C_K be an open cover of K .
- ▶ Then $U = X \setminus K$ is an open set and $C_T = C_K \cup U$ is an open cover of T .
- ▶ Since T is compact, then C_T has a finite subcover C'_T that also covers K .
- ▶ Since U does not contain any point of K , the set K is already covered by $C'_K = C'_T \setminus U$ i.e. a finite subcollection of the original collection C_K .

Compactness in Euclidean Space

Lemma

A closed subset of a compact set is compact.

- ▶ Let K be a closed subset of a compact set $T \subset X$ and let C_K be an open cover of K .
- ▶ Then $U = X \setminus K$ is an open set and $C_T = C_K \cup U$ is an open cover of T .
- ▶ Since T is compact, then C_T has a finite subcover C'_T that also covers K .
- ▶ Since U does not contain any point of K , the set K is already covered by $C'_K = C'_T \setminus U$ i.e. a finite subcollection of the original collection C_K .
- ▶ It is therefore possible to extract from any open cover C_K of K a finite subcover.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- Recall the definition of compact: every open cover of S has a finite subcover.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Recall the definition of compact: every open cover of S has a finite subcover.
- ▶ We start by showing that compact implies closed.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Recall the definition of compact: every open cover of S has a finite subcover.
- ▶ We start by showing that compact implies closed.
- ▶ Let a be a limit point of S and let C be an arbitrary collection of open sets such that every open set $U \in C$ is disjoint from some neighborhood, V_U of a .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Recall the definition of compact: every open cover of S has a finite subcover.
- ▶ We start by showing that compact implies closed.
- ▶ Let a be a limit point of S and let C be an arbitrary collection of open sets such that every open set $U \in C$ is disjoint from some neighborhood, V_U of a .
- ▶ Then C fails to cover S .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Recall the definition of compact: every open cover of S has a finite subcover.
- ▶ We start by showing that compact implies closed.
- ▶ Let a be a limit point of S and let C be an arbitrary collection of open sets such that every open set $U \in C$ is disjoint from some neighborhood, V_U of a .
- ▶ Then C fails to cover S .
- ▶ Indeed $\cap V_U$ is a neighborhood W of a in \mathbb{R}^n .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Recall the definition of compact: every open cover of S has a finite subcover.
- ▶ We start by showing that compact implies closed.
- ▶ Let a be a limit point of S and let C be an arbitrary collection of open sets such that every open set $U \in C$ is disjoint from some neighborhood, V_U of a .
- ▶ Then C fails to cover S .
- ▶ Indeed $\cap V_U$ is a neighborhood W of a in \mathbb{R}^n .
- ▶ Since a is a limit point of S , W must contain a point $x \in S$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Recall the definition of compact: every open cover of S has a finite subcover.
- ▶ We start by showing that compact implies closed.
- ▶ Let a be a limit point of S and let C be an arbitrary collection of open sets such that every open set $U \in C$ is disjoint from some neighborhood, V_U of a .
- ▶ Then C fails to cover S .
- ▶ Indeed $\cap V_U$ is a neighborhood W of a in \mathbb{R}^n .
- ▶ Since a is a limit point of S , W must contain a point $x \in S$.
- ▶ This point is not covered by C since every U in C is disjoint from V_U and hence disjoint from W which contains x .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ If S is compact but not closed, then it has a limit point $a \notin S$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ If S is compact but not closed, then it has a limit point $a \notin S$.
- ▶ Consider a collection C' consisting of an open neighborhood $N(x)$ for every $x \in S$ chosen small enough not to intersect some neighborhood V_x of a .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ If S is compact but not closed, then it has a limit point $a \notin S$.
- ▶ Consider a collection C' consisting of an open neighborhood $N(x)$ for every $x \in S$ chosen small enough not to intersect some neighborhood V_x of a .
- ▶ Then C' is an open cover of S but any finite subcollection of C' has the form of C from the previous slide and thus cannot be a finite subcover of S .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ If S is compact but not closed, then it has a limit point $a \notin S$.
- ▶ Consider a collection C' consisting of an open neighborhood $N(x)$ for every $x \in S$ chosen small enough not to intersect some neighborhood V_x of a .
- ▶ Then C' is an open cover of S but any finite subcollection of C' has the form of C from the previous slide and thus cannot be a finite subcover of S .
- ▶ This contradicts the compactness of S .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ If S is compact but not closed, then it has a limit point $a \notin S$.
- ▶ Consider a collection C' consisting of an open neighborhood $N(x)$ for every $x \in S$ chosen small enough not to intersect some neighborhood V_x of a .
- ▶ Then C' is an open cover of S but any finite subcollection of C' has the form of C from the previous slide and thus cannot be a finite subcover of S .
- ▶ This contradicts the compactness of S .
- ▶ Therefore every limit point of S is in S so S is closed.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- Now we show that compact implies bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Now we show that compact implies bounded.
- ▶ Consider the open balls centered on a common point.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Now we show that compact implies bounded.
- ▶ Consider the open balls centered on a common point.
- ▶ This can cover any set, because all points in the set are some distance away from that point.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Now we show that compact implies bounded.
- ▶ Consider the open balls centered on a common point.
- ▶ This can cover any set, because all points in the set are some distance away from that point.
- ▶ Any finite subcover of this cover must be bounded because all balls in the subcover are contained in the largest open ball within that subcover.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Now we show that compact implies bounded.
- ▶ Consider the open balls centered on a common point.
- ▶ This can cover any set, because all points in the set are some distance away from that point.
- ▶ Any finite subcover of this cover must be bounded because all balls in the subcover are contained in the largest open ball within that subcover.
- ▶ Therefore any set covered by this subcover must also be bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- Finally, we show that closed and bounded implies compact.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Finally, we show that closed and bounded implies compact.
- ▶ If a set $S \in \mathbb{R}^n$ is bounded, then it can be enclosed in an n -box $T_0 = [-a, a]^n$ where $a > 0$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Finally, we show that closed and bounded implies compact.
- ▶ If a set $S \in \mathbb{R}^n$ is bounded, then it can be enclosed in an n -box $T_0 = [-a, a]^n$ where $a > 0$.
- ▶ By the lemma, if we can show that T_0 is compact, then because S is a closed subset of T_0 , that will prove that S is compact.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Finally, we show that closed and bounded implies compact.
- ▶ If a set $S \in \mathbb{R}^n$ is bounded, then it can be enclosed in an n -box $T_0 = [-a, a]^n$ where $a > 0$.
- ▶ By the lemma, if we can show that T_0 is compact, then because S is a closed subset of T_0 , that will prove that S is compact.
- ▶ Assume FSOC that that T_0 is not compact.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Finally, we show that closed and bounded implies compact.
- ▶ If a set $S \in \mathbb{R}^n$ is bounded, then it can be enclosed in an n -box $T_0 = [-a, a]^n$ where $a > 0$.
- ▶ By the lemma, if we can show that T_0 is compact, then because S is a closed subset of T_0 , that will prove that S is compact.
- ▶ Assume FSOC that that T_0 is not compact.
- ▶ Then there exists an infinite open cover C of T_0 that does not admit any finite subcover.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ By bisecting each of the sides of the T_0 box, T_0 can be broken into 2^n sub n -boxes, each of which has a diameter equal to half the diameter of T_0 .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ By bisecting each of the sides of the T_0 box, T_0 can be broken into 2^n sub n -boxes, each of which has a diameter equal to half the diameter of T_0 .
- ▶ Then at least one of the 2^n sections of T_0 must require an infinite subcover of C .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ By bisecting each of the sides of the T_0 box, T_0 can be broken into 2^n sub n -boxes, each of which has a diameter equal to half the diameter of T_0 .
- ▶ Then at least one of the 2^n sections of T_0 must require an infinite subcover of C .
- ▶ Otherwise C itself would have a subcover formed by taking the union of the finite subcovers of the sections.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ By bisecting each of the sides of the T_0 box, T_0 can be broken into 2^n sub n -boxes, each of which has a diameter equal to half the diameter of T_0 .
- ▶ Then at least one of the 2^n sections of T_0 must require an infinite subcover of C .
- ▶ Otherwise C itself would have a subcover formed by taking the union of the finite subcovers of the sections.
- ▶ Call this T_1 . The sides of T_1 can be bisected as well which yields 2^n smaller sections of T_1 , at least one of which must require an infinite subcover of C .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Continuing this process gives us a decreasing sequence of nested n -boxes: $T_0 \supset T_1 \supset T_2 \supset \dots \supset T_k \supset \dots$

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Continuing this process gives us a decreasing sequence of nested n -boxes: $T_0 \supset T_1 \supset T_2 \supset \dots \supset T_k \supset \dots$
- ▶ The side length of T_k is $\frac{2a}{2^k} \rightarrow 0$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .
- ▶ Since C covers T_0 , then it has some member $U \in C$ such that $L \in U$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .
- ▶ Since C covers T_0 , then it has some member $U \in C$ such that $L \in U$.
- ▶ Since U is open, there is an open ball $B_n(L) \subseteq U$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .
- ▶ Since C covers T_0 , then it has some member $U \in C$ such that $L \in U$.
- ▶ Since U is open, there is an open ball $B_n(L) \subseteq U$.
- ▶ For a large enough k , we have $T_k \subseteq B_n(L) \subseteq U$.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .
- ▶ Since C covers T_0 , then it has some member $U \in C$ such that $L \in U$.
- ▶ Since U is open, there is an open ball $B_n(L) \subseteq U$.
- ▶ For a large enough k , we have $T_k \subseteq B_n(L) \subseteq U$.
- ▶ But this contradicts our claim that we need an infinite number of members of C in order to cover T_k .

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .
- ▶ Since C covers T_0 , then it has some member $U \in C$ such that $L \in U$.
- ▶ Since U is open, there is an open ball $B_n(L) \subseteq U$.
- ▶ For a large enough k , we have $T_k \subseteq B_n(L) \subseteq U$.
- ▶ But this contradicts our claim that we need an infinite number of members of C in order to cover T_k .
- ▶ Therefore T_0 is compact.

Heine-Borel Theorem

Theorem

For a subset S of Euclidean space \mathbb{R}^n , S is compact if and only if it is closed and bounded.

- ▶ Define a sequence (x_k) such that each x_k is in T_k .
- ▶ The sequence is Cauchy.
- ▶ Because the sequence is Cauchy, it converges to a limit, L .
- ▶ Since each T_k is closed, $L \in T_k$ for all k .
- ▶ Since C covers T_0 , then it has some member $U \in C$ such that $L \in U$.
- ▶ Since U is open, there is an open ball $B_n(L) \subseteq U$.
- ▶ For a large enough k , we have $T_k \subseteq B_n(L) \subseteq U$.
- ▶ But this contradicts our claim that we need an infinite number of members of C in order to cover T_k .
- ▶ Therefore T_0 is compact.
- ▶ Since S is a closed subset of T_0 , S is compact.

Sequential Compactness

- ▶ Closely related to the Heine-Borel theorem is the Bolzano-Weierstrass theorem.

Sequential Compactness

- ▶ Closely related to the Heine-Borel theorem is the Bolzano-Weierstrass theorem.
- ▶ In metric spaces, a set is compact if and only if it is **sequentially compact**,

Sequential Compactness

- ▶ Closely related to the Heine-Borel theorem is the Bolzano-Weierstrass theorem.
- ▶ In metric spaces, a set is compact if and only if it is **sequentially compact**,
- ▶ A metric space is sequentially compact if every infinite sequence has a convergent subsequence.

Sequential Compactness

- ▶ Closely related to the Heine-Borel theorem is the Bolzano-Weierstrass theorem.
- ▶ In metric spaces, a set is compact if and only if it is **sequentially compact**,
- ▶ A metric space is sequentially compact if every infinite sequence has a convergent subsequence.
- ▶ The Bolzano Weierstrass theorem will allow us to use this equivalence to write a simpler proof of the Heine-Borel theorem.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.
- ▶ Divide $[0, 1]$ into two intervals, $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.
- ▶ Divide $[0, 1]$ into two intervals, $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$.
- ▶ At least one of the halves contains infinitely many terms of (x_n) .

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.
- ▶ Divide $[0, 1]$ into two intervals, $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$.
- ▶ At least one of the halves contains infinitely many terms of (x_n) .
- ▶ Denote that interval I_1 which has length $\frac{1}{2}$.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.
- ▶ Divide $[0, 1]$ into two intervals, $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$.
- ▶ At least one of the halves contains infinitely many terms of (x_n) .
- ▶ Denote that interval I_1 which has length $\frac{1}{2}$.
- ▶ Let (x_{n_k}) be the subsequence of (x_n) consisting of every term that lies in I_1 .

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.
- ▶ Divide $[0, 1]$ into two intervals, $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$.
- ▶ At least one of the halves contains infinitely many terms of (x_n) .
- ▶ Denote that interval I_1 which has length $\frac{1}{2}$.
- ▶ Let (x_{n_k}) be the subsequence of (x_n) consisting of every term that lies in I_1 .
- ▶ Now divide I_1 into two halves each of length $\frac{1}{4}$.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Let (x_n) be a bounded sequence.
- ▶ Without loss of generality assume that every term in the sequence lies in $[0, 1]$.
- ▶ Divide $[0, 1]$ into two intervals, $[0, \frac{1}{2}]$ and $[\frac{1}{2}, 1]$.
- ▶ At least one of the halves contains infinitely many terms of (x_n) .
- ▶ Denote that interval I_1 which has length $\frac{1}{2}$.
- ▶ Let (x_{n_k}) be the subsequence of (x_n) consisting of every term that lies in I_1 .
- ▶ Now divide I_1 into two halves each of length $\frac{1}{4}$.
- ▶ At least one of these contains infinitely many terms of the (x_{n_k}) and denote that half by I_2 .

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Continue this way to construct a sequence of nested intervals $I_1 \supseteq I_2 \supseteq \dots$ where the length of I_n is $(\frac{1}{2})^n$ and each interval contains an infinite number of terms of the original sequence (x_n) .

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Continue this way to construct a sequence of nested intervals $I_1 \supseteq I_2 \supseteq \dots$ where the length of I_n is $(\frac{1}{2})^n$ and each interval contains an infinite number of terms of the original sequence (x_n) .
- ▶ Now construct a subsequence (z_n) of (x_n) made up of one term for each interval I_n .

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Continue this way to construct a sequence of nested intervals $I_1 \supseteq I_2 \supseteq \dots$ where the length of I_n is $(\frac{1}{2})^n$ and each interval contains an infinite number of terms of the original sequence (x_n) .
- ▶ Now construct a subsequence (z_n) of (x_n) made up of one term for each interval I_n .
- ▶ This is a Cauchy sequence: for all $m, n > N$, $|z_m - z_n| < (\frac{1}{2})^N$.

Bolzano-Weierstrass Theorem

Theorem

Every bounded sequence in \mathbb{R}^n has a convergent subsequence.

- ▶ Continue this way to construct a sequence of nested intervals $I_1 \supseteq I_2 \supseteq \dots$ where the length of I_n is $(\frac{1}{2})^n$ and each interval contains an infinite number of terms of the original sequence (x_n) .
- ▶ Now construct a subsequence (z_n) of (x_n) made up of one term for each interval I_n .
- ▶ This is a Cauchy sequence: for all $m, n > N$, $|z_m - z_n| < (\frac{1}{2})^N$.
- ▶ Since \mathbb{R}^n is a complete metric space, (z_n) converges.

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .
- ▶ Consider a closed bounded subset S of \mathbb{R}^n .

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .
- ▶ Consider a closed bounded subset S of \mathbb{R}^n .
- ▶ Let x_n be a sequence in S .

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .
- ▶ Consider a closed bounded subset S of \mathbb{R}^n .
- ▶ Let x_n be a sequence in S .
- ▶ Because S is bounded, so is x_n .

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .
- ▶ Consider a closed bounded subset S of \mathbb{R}^n .
- ▶ Let x_n be a sequence in S .
- ▶ Because S is bounded, so is x_n .
- ▶ By BW, x_n has a convergent subsequence in \mathbb{R}^n .

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .
- ▶ Consider a closed bounded subset S of \mathbb{R}^n .
- ▶ Let x_n be a sequence in S .
- ▶ Because S is bounded, so is x_n .
- ▶ By BW, x_n has a convergent subsequence in \mathbb{R}^n .
- ▶ Since S is closed, the limit belongs to S .

Heine-Borel Again

- ▶ It is now much easier to prove that closed and bounded implies compact in \mathbb{R}^n .
- ▶ Consider a closed bounded subset S of \mathbb{R}^n .
- ▶ Let x_n be a sequence in S .
- ▶ Because S is bounded, so is x_n .
- ▶ By BW, x_n has a convergent subsequence in \mathbb{R}^n .
- ▶ Since S is closed, the limit belongs to S .
- ▶ Therefore S is sequentially compact and therefore compact.