

Bayesian modeling

Refresher on frequentist modeling

- a series of *iid* (independent and identically distributed) random variables $\mathbf{Y} = (Y_1, \dots, Y_n)$

Refresher on frequentist modeling

- a series of *iid* (independent and identically distributed) random variables $\mathbf{Y} = (Y_1, \dots, Y_n)$
- we observe a sample $\mathbf{y} = (y_1, \dots, y_n)$

Refresher on frequentist modeling

- a series of *iid* (independent and identically distributed) random variables $\mathbf{Y} = (Y_1, \dots, Y_n)$
- we observe a sample $\mathbf{y} = (y_1, \dots, y_n)$
- model their probability distribution as $f(y|\theta)$, $\theta \in \Theta$

Refresher on frequentist modeling

- a series of *iid* (independent and identically distributed) random variables $\mathbf{Y} = (Y_1, \dots, Y_n)$
- we observe a sample $\mathbf{y} = (y_1, \dots, y_n)$
- model their probability distribution as $f(y|\theta)$, $\theta \in \Theta$

This model assumes there is a “true” distribution of Y characterized by the “true” value of the parameter θ^*

$$\hat{\theta} ?$$

Historical motivating example

Laplace

What is the probability of birth of girls rather than boys ?

⇒ **observations**: births observed in Paris between 1745 and 1770
(241,945 girls & 251,527 boys)

When a child is born, is it equally likely to be a girl or a boy ?

Three building blocks

- 1 the question
- 2 the sampling model
- 3 the prior

Three building blocks

1 the question

The first step in building a model is always to identify the question you want to answer

2 the sampling model

3 the prior

Three building blocks

1 the question

The first step in building a model is always to identify the question you want to answer

2 the sampling model

Which **observations** are available to inform our response to this ?
How can they be **described**?

3 the prior

Three building blocks

1 the question

The first step in building a model is always to identify the question you want to answer

2 the sampling model

Which **observations** are available to inform our response to this ?
How can they be **described**?

3 the prior

A probability distribution on the parameters θ of the sampling model

The sampling model

\mathbf{y} : the observations available

⇒ (parametric) **probabilistic model** underlying their **generation**:

$$Y_i \stackrel{iid}{\sim} f(y|\theta)$$

The *prior* distribution

In Bayesian modeling, compared to frequentist modeling, we add a **probability distribution** on the **parameters** θ

$$\theta \sim \pi(\theta)$$

$$Y_i|\theta \stackrel{iid}{\sim} f(y|\theta)$$

θ will thus be treated like a random variable,
but which is never observed !

Back to Laplace's historical example

- 1 The question
- 2 Sampling model
- 3 *prior*

Back to Laplace's historical example

① The question

...

② Sampling model

...

③ *prior*

...