

MCMC algorithms: general principle

Approximate an integral (or another function) from a target distribution

MCMC algorithms: general principle

Approximate an integral (or another function) from a target distribution

⇒ sample a Markov chain whose stationary law is the target (such as the *posterior*) distribution, then apply the Monte Carlo method

MCMC algorithms: general principle

Approximate an integral (or another function) from a target distribution

⇒ sample a Markov chain whose stationary law is the target (such as the *posterior*) distribution, then apply the Monte Carlo method

Requires **twofold convergence** :

- 1 the Markov chain must first converge to its stationary distribution:

$$\forall X_0, X_n \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \tilde{p}$$

MCMC algorithms: general principle

Approximate an integral (or another function) from a target distribution

⇒ sample a Markov chain whose stationary law is the target (such as the *posterior*) distribution, then apply the Monte Carlo method

Requires **twofold convergence** :

- 1 the Markov chain must first converge to its stationary distribution:

$$\forall X_0, X_n \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \tilde{p}$$

- 2 then Monte Carlo convergence must also happen:

$$\frac{1}{N} \sum_{i=1}^N f(X_{n+i}) \xrightarrow[N \rightarrow +\infty]{} \mathbb{E}[f(X)]$$

MCMC algorithms: general principle

Approximate an integral (or another function) from a target distribution

⇒ sample a Markov chain whose stationary law is the target (such as the *posterior*) distribution, then apply the Monte Carlo method

Requires **twofold convergence** :

- 1 the Markov chain must first converge to its stationary distribution:

$$\forall X_0, X_n \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \tilde{p}$$

- 2 then Monte Carlo convergence must also happen:

$$\frac{1}{N} \sum_{i=1}^N f(X_{n+i}) \xrightarrow[N \rightarrow +\infty]{} \mathbb{E}[f(X)]$$

$$\overbrace{X_0 \rightarrow X_1 \rightarrow X_2 \rightarrow \cdots \rightarrow X_n}^{\text{Markov chain convergence}} \rightarrow \overbrace{X_{n+1} \rightarrow X_{n+2} \rightarrow \cdots \rightarrow X_{n+N}}^{\text{Monte Carlo sample}}$$

General framework of MCMC algorithms

MCMC algorithms uses an acceptance-rejection framework

- 1 Initialise $x^{(0)}$
- 2 For $t = 1 \dots n + N$:
 - a Propose a new candidate $y^{(t)} \sim q(y^{(t)} | x^{(t-1)})$
 - b Accept $y^{(t)}$ with probability $\alpha(x^{(t-1)}, y^{(t)})$:
 if $t > n$, "save" $x^{(t)}$ (as part of the final Monte Carlo sample)

where q is the instrumental distribution for proposing new samples
and α is the acceptance probability.

Choosing the instrumental distribution

Not absolutely optimal choice for the instrumental distribution q
proposing new samples

⇒ infinite possibilities: some better than others

Choosing the instrumental distribution

Not absolutely optimal choice for the instrumental distribution q proposing new samples

⇒ infinite possibilities: some better than others

To guaranty convergence towards the target \tilde{p} :

- the support of q has to cover the support of \tilde{p}
- q must not generate periodic values

Choosing the instrumental distribution

Not absolutely optimal choice for the instrumental distribution q proposing new samples

⇒ infinite possibilities: some better than others

To guaranty convergence towards the target \tilde{p} :

- the support of q has to cover the support of \tilde{p}
- q must not generate periodic values

NB: *ideally* q is **easy** and **fast** to **compute**