

MCMC Algorithms

Markov chain definition

Markov chain: discrete time stochastic process

Definition: a series of random variables X_0, X_1, X_2, \dots (all valued over the same state space) with the “memoryless” **Markov property**:

$$p(X_i = x | X_0 = x_0, X_1 = x_1, \dots, X_{i-1} = x_{i-1}) = p(X_i = x | X_{i-1} = x_{i-1})$$

The set E of all possible values of X_i is called the **state space**

2 parameters:

- 1 initial distribution $p(X_0)$
- 2 transition probabilities $T(x, A) = p(X_i \in A | X_{i-1} = x)$

NB: only **homogeneous** Markov chains considered here:

$$p(X_{i+1} = x | X_i = y) = p(X_i = x | X_{i-1} = y)$$

Markov chains properties

Property: a Markov chain is **irreducible** if all sets of non-zero probability can be reached from any starting point (i.e. any state is accessible from any other)

Markov chains properties

Property: a Markov chain is **irreducible** if all sets of non-zero probability can be reached from any starting point (i.e. any state is accessible from any other)

Property: a Markov chain is **recurrent** if the trajectories (X_i) pass an infinite number of times in any set of non-zero probability of the state space

Markov chains properties

Property: a Markov chain is **irreducible** if all sets of non-zero probability can be reached from any starting point (i.e. any state is accessible from any other)

Property: a Markov chain is **recurrent** if the trajectories (X_i) pass an infinite number of times in any set of non-zero probability of the state space

Property: a Markov chain is **aperiodic** if nothing induces periodic behavior of the trajectories

Stationary law & ergodic theorem

Definition: A probability distribution \tilde{p} is called **invariant law** (or **stationary law**) for a Markov chain if it verifies the following property:
if $X_i \sim \tilde{p}$, then $X_{i+j} \sim \tilde{p} \forall j \geq 1$

Remark: a Markov chain can admit several stationary laws

Stationary law & ergodic theorem

Definition: A probability distribution \tilde{p} is called **invariant law** (or **stationary law**) for a Markov chain if it verifies the following property:
if $X_i \sim \tilde{p}$, then $X_{i+j} \sim \tilde{p} \forall j \geq 1$

Remark: a Markov chain can admit several stationary laws

Ergodic theorem (infinite space): A positive irreducible and recurrent Markov chain admits a single invariant probability distribution \tilde{p} and converges towards it

Markov chain example (discrete state space)– I

Doudou (a hamster) follows a Markov chain every minute with 3 states:

S sleep

E eat

W work out

⇒ its activity in 1min only depends on its current activity

Matrix of transition probabilities:

$$P = \begin{pmatrix} X_i/X_{i+1} & S & E & W \\ S & 0.9 & 0.05 & 0.05 \\ E & 0.7 & 0 & 0.3 \\ W & 0.8 & 0 & 0.2 \end{pmatrix}$$

Markov chain example (discrete state space)– I

Doudou (a hamster) follows a Markov chain every minute with 3 states:

S sleep

E eat

W work out

⇒ its activity in 1min only depends on its current activity

Matrix of transition probabilities:

$$P = \begin{pmatrix} X_i/X_{i+1} & S & E & W \\ S & 0.9 & 0.05 & 0.05 \\ E & 0.7 & 0 & 0.3 \\ W & 0.8 & 0 & 0.2 \end{pmatrix}$$

- 1) Is the Markov chain irreducible ? recurrent ? aperiodic ?
- 2) Suppose Doudou is now asleep. What about in 2 min ? in 10 min ?
- 3) Suppose now that Doudou is working out. What about in 10 min ?

Markov chain example (discrete state space) – II

1) Is the Markov chain irreducible ? recurrent ? aperiodic ?

...


2) Suppose Doudou is now asleep. What about in 2 min ? in 10 min ?

...

3) Suppose now that Doudou is working out. What about in 10 min ?

Markov chain example (discrete state space) – II

1) Is the Markov chain irreducible ? recurrent ? aperiodic ?




2) Suppose Doudou is now asleep. What about in 2 min ? in 10 min ?

...

3) Suppose now that Doudou is working out. What about in 10 min ?

Markov chain example (discrete state space) – II

1) Is the Markov chain irreducible ? recurrent ? aperiodic ?



2) Suppose Doudou is now asleep. What about in 2 min ? in 10 min ?

$$x_0 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}^T \quad x_2 = x_0 P^2 = \begin{pmatrix} 0.885 \\ 0.045 \\ 0.070 \end{pmatrix}^T \quad x_{10} = x_2 P^8 = x_0 P^{10} = \begin{pmatrix} 0.884 \\ 0.044 \\ 0.072 \end{pmatrix}^T$$

Markov chain example (discrete state space) – II

3) Suppose now that Doudou is working out. What about in 10 min ?

...

Markov chain example (discrete state space) – II

3) Suppose now that Doudou is working out. What about in 10 min ?

$$x_0 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}^T \quad x_{10} = x_0 P^{10} = \begin{pmatrix} 0.884 \\ 0.044 \\ 0.072 \end{pmatrix}^T$$

Here, the Markov chain being aperiodic, recurrent and irreducible, there is a stationary law: $\tilde{p} = \tilde{p}P$.