

Hyper-priors & hierarchical models

Hierarchical levels:

① $\pi(\theta)$

② $f(\mathbf{y}|\theta)$

Hyper-priors & hierarchical models

Hierarchical levels:

① $\eta \sim h(\eta)$

② $\pi(\theta|\eta)$

③ $f(\mathbf{y}|\theta)$

Hyper-priors & hierarchical models

Hierarchical levels:

① $\eta \sim h(\eta)$

② $\pi(\theta|\eta)$

③ $f(\mathbf{y}|\theta)$

$$p(\theta|\mathbf{y}) = \frac{f(\mathbf{y}|\theta)\pi(\theta)}{f(\mathbf{y})} = \frac{\int f(\mathbf{y}|\theta, \eta)\pi(\theta|\eta)h(\eta)d\eta}{f(\mathbf{y})}$$

Hyper-priors & hierarchical models

Hierarchical levels:

① $\eta \sim h(\eta)$

② $\pi(\theta|\eta)$

③ $f(\mathbf{y}|\theta)$

$$p(\theta|\mathbf{y}) = \frac{f(\mathbf{y}|\theta)\pi(\theta)}{f(\mathbf{y})} = \frac{\int f(\mathbf{y}|\theta, \eta)\pi(\theta|\eta)h(\eta)d\eta}{f(\mathbf{y})} = \frac{f(\mathbf{y}|\theta)\int \pi(\theta|\eta)h(\eta)d\eta}{f(\mathbf{y})}$$

NB: 3 hierarchical levels \Leftrightarrow two levels with *prior*: $\pi(\theta) = \int \pi(\theta|\eta)h(\eta)d\eta$

Hyper-priors & hierarchical models

Hierarchical levels:

① $\eta \sim h(\eta)$

② $\pi(\theta|\eta)$

③ $f(\mathbf{y}|\theta)$

$$p(\theta|\mathbf{y}) = \frac{f(\mathbf{y}|\theta)\pi(\theta)}{f(\mathbf{y})} = \frac{\int f(\mathbf{y}|\theta, \eta)\pi(\theta|\eta)h(\eta)d\eta}{f(\mathbf{y})} = \frac{f(\mathbf{y}|\theta)\int \pi(\theta|\eta)h(\eta)d\eta}{f(\mathbf{y})}$$

NB: 3 hierarchical levels \Leftrightarrow two levels with *prior*: $\pi(\theta) = \int \pi(\theta|\eta)h(\eta)d\eta$

\Rightarrow can **ease modeling** and **elicitation** of the *prior*...

Hyperprior in the historical example

Historical example of birth sex with a Beta *prior*

⇒ two Gamma hyper-priors for α and β (conjugated):

$$\alpha \sim \text{Gamma}(4, 0.5)$$

$$\beta \sim \text{Gamma}(4, 0.5)$$

$$\theta | \alpha, \beta \sim \text{Beta}(\alpha, \beta)$$

$$Y_i | \theta \stackrel{iid}{\sim} \text{Bernoulli}(\theta)$$

Empirical Bayes

Eliciting the *prior* according to its empirical marginal distribution

⇒ estimate the *prior* from the data

- 1 hyper-parameters
- 2 estimate them through frequentist methods (e.g. MLE) by $\hat{\eta}$
- 3 plug-in estimates into the *prior*
- 4 ⇒ *posterior*: $p(\theta|\mathbf{y}, \hat{\eta})$

Empirical Bayes

Eliciting the *prior* according to its empirical marginal distribution

⇒ estimate the *prior* from the data

- ① hyper-parameters
 - ② estimate them through frequentist methods (e.g. MLE) by $\hat{\eta}$
 - ③ plug-in estimates into the *prior*
 - ④ ⇒ *posterior*: $p(\theta|\mathbf{y}, \hat{\eta})$
- Combines Bayesian and frequentist frameworks
 - Concentrated *posterior* (↘ variance) but ↗ bias (data used twice !)
 - Approximate a fully Bayesian approach

Sequential Bayes

Bayes' theorem can be used sequentially:

$$p(\theta|\mathbf{y}) \propto f(\mathbf{y}|\theta)\pi(\theta)$$

If $\mathbf{y} = (\mathbf{y}_1, \mathbf{y}_2)$, then:

$$p(\theta|\mathbf{y}) \propto f(\mathbf{y}_2|\theta)f(\mathbf{y}_1|\theta)\pi(\theta) \propto f(\mathbf{y}_2|\theta)p(\theta|\mathbf{y}_1)$$

⇒ *posterior* distribution updates as new observations are acquired/available (*online updates*)