The Multinomial Distribution or...Why You Shouldn't Play Roulette

Matt Johnson

April 5, 2016

• We have *n* independent trials.

- We have *n* independent trials.
- Each trial has exactly 1 of *m* possible outcomes.

- We have *n* independent trials.
- Each trial has exactly 1 of *m* possible outcomes.
- Each possible outcome has probability p_i , with

$$\sum_{i=1:m} p_i = 1.$$

- We have *n* independent trials.
- Each trial has exactly 1 of *m* possible outcomes.
- Each possible outcome has probability p_i , with

$$\sum_{i=1:m} p_i = 1.$$

• Describes the number of each possible outcome.



- We have *n* independent trials.
- Each trial has exactly 1 of *m* possible outcomes.
- Each possible outcome has probability p_i , with

$$\sum_{i=1:m} p_i = 1.$$

- Describes the number of each possible outcome.
- Classic example: rolling a weighted die

Roulette is a common casino game

Roulette is a common casino game (the one with the wheel).



Roulette is a common casino game (the one with the wheel).



American wheel contains 38 "pockets"

Roulette is a common casino game (the one with the wheel).



- American wheel contains 38 "pockets"
- Numbered 1-36, as well as 0 and 00

Roulette is a common casino game (the one with the wheel).



- American wheel contains 38 "pockets"
- Numbered 1-36, as well as 0 and 00
- 18 are red, 18 are black, 0 and 00 are green

Roulette is a common casino game (the one with the wheel).



- American wheel contains 38 "pockets"
- Numbered 1-36, as well as 0 and 00
- 18 are red, 18 are black, 0 and 00 are green

Common to bet simply on black or red.

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) =$$

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) = \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 11.8\%$$

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) = \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 11.8\%$$

$$P(BNB) =$$

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) = \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 11.8\%$$

$$P(BNB) = \frac{18}{38} \cdot \frac{20}{38} \cdot \frac{18}{38} \approx 11.8\%$$

Suppose you bet on black three times in a row. What's the probability that you win exactly twice?

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) = \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 11.8\%$$

$$P(BNB) = \frac{18}{38} \cdot \frac{20}{38} \cdot \frac{18}{38} \approx 11.8\%$$

P(NBB) =

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) = \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 11.8\%$$

$$P(BNB) = \frac{18}{38} \cdot \frac{20}{38} \cdot \frac{18}{38} \approx 11.8\%$$

$$P(NBB) = \frac{20}{38} \cdot \frac{18}{38} \cdot \frac{18}{38} \approx 11.8\%$$

$$P(2B) = P(BBN) + P(BNB) + P(NBB)$$

$$P(BBN) = \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 11.8\%$$

$$P(BNB) = \frac{18}{38} \cdot \frac{20}{38} \cdot \frac{18}{38} \approx 11.8\%$$

$$P(NBB) = \frac{20}{38} \cdot \frac{18}{38} \cdot \frac{18}{38} \approx 11.8\%$$

$$P(2B) = 3 \cdot \frac{18}{38} \cdot \frac{18}{38} \cdot \frac{20}{38} \approx 35.4\%$$

Perform n spins. Bet has probability p of winning. What's the probability of exactly k wins?

Perform n spins. Bet has probability p of winning. What's the probability of exactly k wins?

P([one specific order]) =

Perform n spins. Bet has probability p of winning. What's the probability of exactly k wins?

$$P([\text{one specific order}]) = p^k \cdot (1-p)^{n-k}$$

Perform n spins. Bet has probability p of winning. What's the probability of exactly k wins?

$$P([\text{one specific order}]) = p^k \cdot (1-p)^{n-k}$$

How many ways can you order k wins and n - k losses?

Perform n spins. Bet has probability p of winning. What's the probability of exactly k wins?

$$P([\text{one specific order}]) = p^k \cdot (1-p)^{n-k}$$

How many ways can you order k wins and n - k losses?

$$\binom{n}{k} = \binom{n}{n-k} = \frac{n!}{k!(n-k)!}$$

Perform n spins. Bet has probability p of winning. What's the probability of exactly k wins?

$$P([\text{one specific order}]) = p^k \cdot (1-p)^{n-k}$$

How many ways can you order k wins and n - k losses?

$$\binom{n}{k} = \binom{n}{n-k} = \frac{n!}{k!(n-k)!}$$

$$P(k \text{ wins}) = \binom{n}{k} p^k (1-p)^{n-k}$$

Let
$$X_i \sim Bern(p)$$
 for $i=1:n$, and $Y \sim Bin(n,p)$. $E[Y] = E[X_1 + \ldots X_n]$

Let
$$X_i \sim Bern(p)$$
 for $i = 1 : n$, and $Y \sim Bin(n, p)$.

$$E[Y] = E[X_1 + \dots X_n]$$

= $E[X_1] + \dots + E[X_n]$

Let
$$X_i \sim Bern(p)$$
 for $i=1:n$, and $Y \sim Bin(n,p)$.
$$E[Y] = E[X_1 + \dots X_n]$$

$$= E[X_1] + \dots + E[X_n]$$

$$= p + \dots + p$$

Let
$$X_i \sim Bern(p)$$
 for $i=1:n$, and $Y \sim Bin(n,p)$.
$$E[Y] = E[X_1 + \dots X_n]$$
$$= E[X_1] + \dots + E[X_n]$$
$$= p + \dots + p$$
$$= np$$

What about describing more than two possible outcomes?

What about describing more than two possible outcomes?

Result will be a *vector* of counts that sum to n.

What about describing more than two possible outcomes?

Result will be a *vector* of counts that sum to n.

Requires a *vector* of probabilities that sum to 1.

What about describing more than two possible outcomes?

Result will be a *vector* of counts that sum to *n*.

Requires a *vector* of probabilities that sum to 1.

Example: the numbers of reds, blacks, and greens after 100 spins.

$$n = 100, p = (\frac{18}{38}, \frac{18}{38}, \frac{2}{38}).$$

One possible result is (43, 49, 8).

The Multinomial Distribution

What about describing more than two possible outcomes?

Result will be a *vector* of counts that sum to *n*.

Requires a vector of probabilities that sum to 1.

Example: the numbers of reds, blacks, and greens after 100 spins.

$$n = 100$$
, $p = (\frac{18}{38}, \frac{18}{38}, \frac{2}{38})$.
One possible result is (43, 49, 8).

Component i is binomial with parameters n and p_i . (The number of blacks was the binomial example!)



Probability Mass Functions

Binomial pmf:

$$P(Y = k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}$$
 (1)

Multinomial pmf:

$$P(Z = (k_1, k_2, ..., k_m)) = \frac{n!}{k_1! \cdots k_m!} p_1^{k_1} \cdots p_m^{k_m}$$
 (2)

Let Z be multinomial random vector with parameters n and probability vector p. Each component is binomial, so

$$E[Z] =$$

Let Z be multinomial random vector with parameters n and probability vector p. Each component is binomial, so

$$E[Z] = (E[Z_1], \ldots, E[Z_m])$$

Let Z be multinomial random vector with parameters n and probability vector p. Each component is binomial, so

$$E[Z] = (E[Z_1], \dots, E[Z_m])$$

= (np_1, \dots, np_m)

Let Z be multinomial random vector with parameters n and probability vector p. Each component is binomial, so

$$E[Z] = (E[Z_1], \dots, E[Z_m])$$

= (np_1, \dots, np_m)

For our roulette example, if Z counts the number of RBG in 190 spins, E[Z] = (90, 90, 10).

Bet \$1 on either RED or BLACK, for 10 spins.

Bet \$1 on either RED or BLACK, for 10 spins.

n = 10 spins

RBG =

3 6

Bet RED: -4
Bet BLACK: +2

Bet \$1 on either RED or BLACK, for 100 spins.

Bet \$1 on either RED or BLACK, for 100 spins.

n = 100 spins

RBG =

52 43 5

Bet RED: +4

Bet BLACK: -14

Bet \$1 on either RED or BLACK, for 1000 spins.

Bet \$1 on either RED or BLACK, for 1000 spins.

n = 1000 spins

RBG =

460 480 60

Bet RED: -80 Bet BLACK: -40

Questions?