

Bootstrap Confidence Intervals

Jialiang Mao

March 29, 2016

Overview


- ▶ What and why?
- ▶ How?
- ▶ Always good?

Example: Chick weight




- ▶ 578 observations

Example: Chick weight



- ▶ 578 observations
- ▶ Right-skewed

Example: Chick weight



- ▶ 578 observations
- ▶ Right-skewed
- ▶ median=103

Bootstrapping



- ▶ The saying "to pull oneself up by one's bootstraps" was already in use during the 19th century as an example of an impossible task.

Bootstrapping



- ▶ The saying "to pull oneself up by one's bootstraps" was already in use during the 19th century as an example of an impossible task.
- ▶ Bootstrap as a metaphor, meaning to better oneself by one's own unaided efforts, was in use in 1922.

Construct a Bootstrap confidence interval

Suppose we are given a sample (x_1, \dots, x_n) and we want to construct a bootstrap confidence interval for a statistic T (mean, median,...).

Construct a Bootstrap confidence interval

Suppose we are given a sample (x_1, \dots, x_n) and we want to construct a bootstrap confidence interval for a statistic T (mean, median,...). Take a large number N (for example, $N = 15000$). For $1 \leq i \leq N$,

Construct a Bootstrap confidence interval

Suppose we are given a sample (x_1, \dots, x_n) and we want to construct a bootstrap confidence interval for a statistic T (mean, median,...). Take a large number N (for example, $N = 15000$). For $1 \leq i \leq N$,

1. Get a bootstrap sample $(x_1^{*(i)}, \dots, x_n^{*(i)})$ from the original sample;
 - ▶ a random sample;

Construct a Bootstrap confidence interval

Suppose we are given a sample (x_1, \dots, x_n) and we want to construct a bootstrap confidence interval for a statistic T (mean, median,...). Take a large number N (for example, $N = 15000$). For $1 \leq i \leq N$,

1. Get a bootstrap sample $(x_1^{*(i)}, \dots, x_n^{*(i)})$ from the original sample;
 - ▶ a random sample;
 - ▶ same size;


Construct a Bootstrap confidence interval

Suppose we are given a sample (x_1, \dots, x_n) and we want to construct a bootstrap confidence interval for a statistic T (mean, median,...). Take a large number N (for example, $N = 15000$). For $1 \leq i \leq N$,

1. Get a bootstrap sample $(x_1^{*(i)}, \dots, x_n^{*(i)})$ from the original sample;
 - ▶ a random sample;
 - ▶ same size;
 - ▶ **sample with replacement.**
2. Calculate the bootstrap statistic $T^{*(i)}$ with the bootstrap sample in the first step.


Then we can use the quantiles of these N bootstrap statistics to construct a bootstrap confidence interval.

Example: Chick weight



- ▶ central 95% of the bootstrap distribution

Example: Chick weight



- ▶ central 95% of the bootstrap distribution
- ▶ bootstrap confidence interval: (98.0, 110.5)

Always good?

Always good?

- ▶ your statistic T is not too weird;

Always good?

- ▶ your statistic T is not too weird;
- ▶ need a representative sample to start;

In class exercise: Simulation study

- ▶ For $n = 100$, generate $x_1, \dots, x_n \sim N(\mu, 1)$, $\mu = 1$;
- ▶ Use the CLT based method to calculate the 95% CI for μ ;
- ▶ Use the Bootstrap method ($N = 5000$) to construct a 95% CI for μ ;
- ▶ Compare these two CIs;
- ▶ Change the values of n and N , re-do the experiment.