

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . What is  $A \cap B$ ?**

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . What is  $A \cap B$ ?**

$$A \cap B = \{2, 4\}.$$

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . What is  $A \cup B$ ?**

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . What is  $A \cup B$ ?**

$$A \cup B = \{1, 2, 3, 4, 5, 6, 8, 10\}.$$

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . What is  $A - B$ ?**

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . What is  $A - B$ ?**

$$A - B = \{6, 8, 10\}.$$

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . If  $U$  is the set of all positive integers less than or equal to 10, what is  $A^c$ ?**

**Let  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{1, 2, 3, 4, 5\}$ . If  $U$  is the set of all positive integers less than or equal to 10, what is  $A^c$ ?**

$$A^c = \{1, 3, 5, 7, 9\}.$$



**What is the sample space of a process that rolls one die and then flips a coin?**

**What is the sample space of a process that rolls one die and then flips a coin?**

$\{1H, 2H, 3H, 4H, 5H, 6H, 1T, 2T, 3T, 4T, 5T, 6T\}$ .

**What is the probability that a hand of poker contains all four suits?**

We'll use the naïve definition of probability, counting the number of allowed hand divided by the total number of hands:

$$\mathbb{P}(\text{all four suits}) = \frac{\#\{\text{ways to get all four suits}\}}{\#\{\text{total hands}\}}$$

We'll use the naïve definition of probability, counting the number of allowed hand divided by the total number of hands:

$$\mathbb{P}(\text{all four suits}) = \frac{\#\{\text{ways to get all four suits}\}}{\#\{\text{total hands}\}}$$

We know the denominator is  $\binom{52}{5}$ . Let's try to figure out the numerator using the the multi-stage experiment formulation.

The easiest way to solve this is to first pick the suit that will have 2 cards in the hand. There are  $\binom{4}{1}$  ways to do this.

The easiest way to solve this is to first pick the suit that will have 2 cards in the hand. There are  $\binom{4}{1}$  ways to do this.

In the second stage we select the two cards from the chosen suit; there are  $\binom{13}{2}$  such selections.

The easiest way to solve this is to first pick the suit that will have 2 cards in the hand. There are  $\binom{4}{1}$  ways to do this.

In the second stage we select the two cards from the chosen suit; there are  $\binom{13}{2}$  such selections.

In the third stage we choose the  $\binom{13}{1} \cdot \binom{13}{1} \cdot \binom{13}{1}$  cards from the other suits.



Putting this together yields:

$$\begin{aligned}\mathbb{P}(\text{all four suits}) &= \frac{\binom{4}{1} \cdot \binom{13}{2} \cdot \binom{13}{1}^3}{\binom{52}{5}} \\ &= 0.2637455\end{aligned}$$

A fairly high percentage!