

# LECTURE 21: MONTE CARLO METHODS

STAT 598Z: INTRODUCTION TO COMPUTING FOR STATISTICS

---

Vinayak Rao

Department of Statistics, Purdue University

April 11, 2017

We want to calculate integrals/summations

- often expectations w.r.t. some probability distribution  $p(x)$

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

We want to calculate integrals/summations

- often expectations w.r.t. some probability distribution  $p(x)$

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

What is the prob. a game of patience (solitaire) is solvable?

$$P(\text{Solvable}) = \frac{1}{|\Pi|} \sum_{\Pi} \mathbb{1}(\Pi \text{ is solvable})$$

We want to calculate integrals/summations

- often expectations w.r.t. some probability distribution  $p(x)$

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

If we drop 3 points on the plane, each Gaussian distributed, what is average the area of the resulting triangle?

$$\mathbb{E}[A] = \int A(x_1, x_2, x_3)p(x_1)p(x_2)p(x_3)dx_1dx_2dx_3$$

We want to calculate integrals/summations

- often expectations w.r.t. some probability distribution  $p(x)$

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

For dataset  $(X, y)$ , what is average loss if you randomly choose a weight-vector according to some distribution (e.g. `rnorm`)?

$$\mathbb{E}_w[\mathcal{L}(X, y)] = \int (y - w^T X)^2 p(w)dw$$

We want to calculate integrals:

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

We want to calculate integrals:

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

Sampling approximation: rather than visit all points in  $\mathcal{X}$ , calculate a summation over a finite set.

$$\hat{\mu} \approx \frac{1}{N} \sum_{i=1}^N f(x_i)$$

We want to calculate integrals:

$$\mu := \mathbb{E}_p[f] = \int_{\mathcal{X}} f(x)p(x)dx$$

Sampling approximation: rather than visit all points in  $\mathcal{X}$ , calculate a summation over a finite set.

$$\hat{\mu} \approx \frac{1}{N} \sum_{i=1}^N f(x_i)$$

Monte Carlo approximation:

- Obtain point by sampling from  $p(x)$

$$x_i \sim p$$



Is this a good idea?

- Very simple
- Unbiased

Is this a good idea?

- Very simple
- Unbiased

If  $x_i \sim p$ ,

$$\mathbb{E}_p[\hat{\mu}] = \mathbb{E}_p \left[ \frac{1}{N} \sum_{i=1}^N f(x_i) \right] = \frac{1}{N} \sum_{i=1}^N \mathbb{E}_p[f] = \mu \quad \text{Unbiased estimate}$$

Is this a good idea?

- Very simple
- Unbiased

If  $x_i \sim p$ ,

$$\mathbb{E}_p[\hat{\mu}] = \mathbb{E}_p \left[ \frac{1}{N} \sum_{i=1}^N f(x) \right] = \frac{1}{N} \sum_{i=1}^N \mathbb{E}_p[f] = \mu \quad \text{Unbiased estimate}$$

$$\text{Var}_p[\hat{\mu}] = \frac{1}{N} \text{Var}_p[f], \quad \text{Error} = \text{StdDev} \propto N^{-1/2}$$

Is this a good idea?

- In low-dims, use numerical methods like quadrature
- In high-dims, numerical methods become infeasible

Is this a good idea?

- In low-dims, use numerical methods like quadrature
- In high-dims, numerical methods become infeasible  
E.g. Simpson's rule in  $d$ -dimensions, with  $N$  grid points:

$$\text{error} \propto N^{-4/d}$$

Is this a good idea?

- In low-dims, use numerical methods like quadrature
- In high-dims, numerical methods become infeasible  
E.g. Simpson's rule in  $d$ -dimensions, with  $N$  grid points:

$$\text{error} \propto N^{-4/d}$$

Monte Carlo integration:

$$\text{error} \propto N^{-1/2}$$

Is this a good idea?

- In low-dims, use numerical methods like quadrature
- In high-dims, numerical methods become infeasible  
E.g. Simpson's rule in  $d$ -dimensions, with  $N$  grid points:

$$\text{error} \propto N^{-4/d}$$

Monte Carlo integration:

$$\text{error} \propto N^{-1/2}$$

Independent of dimensionality!

# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?



# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?

No!

# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?

No!

- Instead: *pseudorandom* numbers.

# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?

No!

- Instead: *pseudorandom* numbers.
- Map a seed to a 'random-looking' sequence.

# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?

No!

- Instead: *pseudorandom* numbers.
- Map a seed to a 'random-looking' sequence.
- Downside: <http://boallan.com/random-numbers.html>

# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?

No!

- Instead: *pseudorandom* numbers.
- Map a seed to a 'random-looking' sequence.
- Downside: <http://boallan.com/random-numbers.html>
- Upside: Can use seeds for reproducibility or debugging

`set.seed(1)`

# GENERATING RANDOM VARIABLES

- The simplest useful probability distribution  $\text{Unif}(0, 1)$ .
- In theory, can be used to generate any other RV.
- Easy to generate uniform RVs on a deterministic computer?

No!

- Instead: *pseudorandom* numbers.
- Map a seed to a 'random-looking' sequence.
- Downside: <http://boallan.com/random-numbers.html>
- Upside: Can use seeds for reproducibility or debugging

`set.seed(1)`

- Careful with batch/parallel processing.

## EXAMPLE MONTE CARLO SAMPLING



$x, y$  are a pair of dice. What is  $p(x, y)$ ?

## EXAMPLE MONTE CARLO SAMPLING



$x, y$  are a pair of dice. What is  $p(x, y)$ ?

What is  $\mathbb{E}[\min(x, y)] = \sum_{i=1}^6 \sum_{j=1}^6 \min(x, y) p(x, y)$



## EXAMPLE MONTE CARLO SAMPLING



$x, y$  are a pair of dice. What is  $p(x, y)$ ?

What is  $\mathbb{E}[\min(x, y)] = \sum_{i=1}^6 \sum_{j=1}^6 \min(x, y) p(x, y)$

Don't really need Monte Carlo here (but what if we had a 100 dice?), but let's try it anyway. How?

## EXAMPLE MONTE CARLO SAMPLING



$x, y$  are a pair of dice. What is  $p(x, y)$ ?

What is  $\mathbb{E}[\min(x, y)] = \sum_{i=1}^6 \sum_{j=1}^6 \min(x, y) p(x, y)$

Don't really need Monte Carlo here (but what if we had a 100 dice?), but let's try it anyway. How?

Roll a pair of dice  $N$  times. Call the  $i$ th outcome  $(x_i, y_i)$ . Then

$$\mathbb{E}[\min(x, y)] \approx \frac{1}{N} \sum_{i=1}^N \min(x_i, y_i)$$

# GENERATING RANDOM VARIABLES

In R we generate uniform random variables via `runif`

Additional functions include

`rnorm()`, `rgamma()`, `rexp()`, `sample()` etc.

In R we generate uniform random variables via `runif`

Additional functions include

`rnorm()`, `rgamma()`, `rexp()`, `sample()` etc.

In theory, we can generate any other random variable by transforming a uniform (or any other) random variable:

$$u \sim \text{Unif}(0, 1), \quad x = f(u)$$

In practice, finding this  $f$  is too hard. Need other approaches.

## RARE EVENT SIMULATION:



Let  $X = (x_1, \dots, x_{100})$  be a hundred dice.  
What is  $p(\sum_{d=1}^{100} x_d \geq 450)$ ?

## RARE EVENT SIMULATION:



Let  $X = (x_1, \dots, x_{100})$  be a hundred dice.

What is  $p(\text{Sum}(X) \geq 450)$ ?

(where  $\text{Sum}(X) = \sum_{d=1}^{100} x_d$ )

## RARE EVENT SIMULATION:



Let  $X = (x_1, \dots, x_{100})$  be a hundred dice.  
What is  $p(\text{Sum}(X) \geq 450)$ ?  
(where  $\text{Sum}(X) = \sum_{d=1}^{100} x_d$ )

$$\begin{aligned} p(\text{Sum}(X) \geq 450) &= \sum \delta(\text{Sum}(X) \geq 450) p(X) \\ &= \mathbb{E}_p[\delta(\text{Sum}(X) \geq 450)] \end{aligned}$$

- $\delta(\cdot)$  is the indicator function
- $\delta(\text{condition}) = 1$  if *condition* is true, else 0.

# NAIVE MONTE CARLO SAMPLING

- Propose from  $p(x)$
- Calculate  $\frac{1}{N} \sum_{i=1}^N \delta(\text{Sum}(X_i))$



- Propose from  $p(x)$
- Calculate  $\frac{1}{N} \sum_{i=1}^N \delta(\text{Sum}(X_i))$

Most  $\delta(\text{Sum}(X_i))$  terms will be 0

High variance

# IMPORTANCE SAMPLING

Importance sampling: assigns importance weights to samples.

Importance sampling: assigns importance weights to samples.

Scheme:

- Draw a proposal  $x_i$  from  $q(\cdot)$
- Assign it a weight  $w_i = p(x_i)/q(x_i)$

Importance sampling: assigns importance weights to samples.

Scheme:

- Draw a proposal  $x_i$  from  $q(\cdot)$
- Assign it a weight  $w_i = p(x_i)/q(x_i)$

$$\mu = \int_{\mathcal{X}} f(x)p(x)dx \approx \frac{1}{N} \sum_{i=1}^N w_i f(x_i) := \mu_{imp}$$

Importance sampling: assigns importance weights to samples.

Scheme:

- Draw a proposal  $x_i$  from  $q(\cdot)$
- Assign it a weight  $w_i = p(x_i)/q(x_i)$

$$\mu = \int_{\mathcal{X}} f(x)p(x)dx \approx \frac{1}{N} \sum_{i=1}^N w_i f(x_i) := \mu_{imp}$$

$$\mathbb{E}[\mu_{imp}] = \frac{1}{N} \sum_{i=1}^N \mathbb{E}[w_i f(x_i)] = \int_{\mathcal{X}} \frac{p(x)}{q(x)} f(x) q(x) dx$$

# IMPORTANCE SAMPLING

Importance sampling: assigns importance weights to samples.

Scheme:

- Draw a proposal  $x_i$  from  $q(\cdot)$
- Assign it a weight  $w_i = p(x_i)/q(x_i)$

$$\mu = \int_{\mathcal{X}} f(x)p(x)dx \approx \frac{1}{N} \sum_{i=1}^N w_i f(x_i) := \mu_{imp}$$

$$\mathbb{E}[\mu_{imp}] = \frac{1}{N} \sum_{i=1}^N \mathbb{E}[w_i f(x_i)] = \int_{\mathcal{X}} \frac{p(x)}{q(x)} f(x) q(x) dx$$

When does this make sense?

Sometimes it's easier to simulate from  $q(x)$  than  $p(x)$

When does this make sense?

Sometimes it's easier to simulate from  $q(x)$  than  $p(x)$

Sometimes it's better to simulate from  $q(x)$  than  $p(x)$ !

To reduce variance. E.g. rare event simulation.



For 100 dice, what is  $p(\text{Sum} > 450)$ ? A better choice might be to bias the dice.

E.g.  $q(x_i = v) \propto v$  (for  $v \in \{1, \dots, 6\}$ )

For 100 dice, what is  $p(\text{Sum} > 450)$ ? A better choice might be to bias the dice.

E.g.  $q(x_i = v) \propto v$  (for  $v \in \{1, \dots, 6\}$ )

- Propose from  $q(x)$
- Calculate weights  $w(X_i) = p(X_i)/q(X_i)$
- Calculate  $\frac{1}{N} \sum_{i=1}^N w(X_i) \delta(\text{Sum}(X_i))$

# IMPORTANCE SAMPLING

For 100 dice, what is  $p(\text{Sum} > 450)$ ? A better choice might be to bias the dice.

E.g.  $q(x_i = v) \propto v$  (for  $v \in \{1, \dots, 6\}$ )

- Propose from  $q(x)$
- Calculate weights  $w(X_i) = p(X_i)/q(X_i)$
- Calculate  $\frac{1}{N} \sum_{i=1}^N w(X_i) \delta(\text{Sum}(X_i))$

Gives a better estimate of

$$p(\text{Sum}(X) \geq 500) = \sum \delta(\text{Sum}(X) \geq 500) p(X)$$