Asthma Control Test™ for teens 12 years and older. Know the score.

If your teen is 12 years or older have him take the test now and discuss the results with your doctor

- Step 1 Write the number of each answer in the score box provided.
- Step 2 Add up each score box for the total.
- Step 3 Take the test to the doctor to talk about your child's total score.

All of the time	(1)	Most of the time	2 Some of the time	3 A little of the time	4 None of the time	5	
2. During the p	ast 4 week	s , how often	have you had shortness	of breath?			
More than once a day	1	Once a day	2 3 to 6 times a week	3 Once or twice a week	4 Not at all	5	
			did your asthma sympto lier than usual in the mo	oms (wheezing, coughing, orning?	shortness of breath, cl	nest tightness,	
4 or more	(1)	2 or 3 nights a week	2 Once a week	3 Once or twice	4 Not at all	5	
nights a week		a week	a woon				
4 . During the p		s , how often	have you used your resc	cue inhaler or nebulizer me			
	ast 4 week				edication (such as albu	aterol)?	
4. During the p	ast 4 week	s, how often 1 or 2 times per day	have you used your resc	cue inhaler or nebulizer me 3 Once a week or less	Not		
4. During the p	ast 4 week	s, how often 1 or 2 times per day	have you used your resc 2 or 3 times per week	cue inhaler or nebulizer me 3 Once a week or less	Not		

What does it mean if my child scores 19 or less?

- If your child's score is 19 or less, it may be a sign that your child's asthma is not under control.
- Make an appointment to discuss your child's asthma score with their doctor. Ask if you should change your child's asthma treatment plan.
- Ask your child's doctor about daily long-term medications that can help control airway inflammation and constriction, the two main causes of asthma symptoms. Many children may need to treat both of these on a daily basis for the best asthma control.