Doctoral Dissertation

Tracking atmospheric chemical components in accordance with the Sustainable Development Goals (SDGs)

Phan Anh

November 14, 2023

Graduate School of Engineering Chubu University

A Doctoral Dissertation submitted to Graduate School of Engineering, Chubu University in partial fulfillment of the requirements for the degree of Doctor of ENGINEERING

Phan Anh

Thesis Committee:

Professor Hiromichi Fukui (Supervisor)

Professor Kiyoshi Honda (Chubu University) Professor Jun Izutsu (Chubu University) Professor Masao Takano (Nagoya University)

Tracking atmospheric chemical components in accordance with the Sustainable Development Goals (SDGs)*

Phan Anh

Abstract

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Keywords:

 π , astronomy, mathematics, computer, algorithm

^{*}Doctoral Dissertation, Graduate School of Information Science, Nara Institute of Science and Technology, November 14, 2023.

Contents

List of Figures			
Li	st of	Tables	1
1	Introduction		2
	1.1	Background	2
2	Bac	ekground	3
	2.1	Air pollution	3
	2.2	Greenhouse gas	3
\mathbf{A}	IR P	OLLUTIOM INDUCED BY INTERVENTION EVENTS	4
3	Ukı	caine's case study	5
	3.1	Introduction	5
	3.2	Data	6
	3.3	Business-as-usual (BAU) modelling	6
	3.4	NO2 changes induced by COVID-19 lockdown	6
	3.5	NO2 changes induced by the armed conflict	6
	3.6	Conclusion	6
4	Jap	an's case study	7
\mathbf{G}	REE	NHOUSE GAS ESTIMATION AND MONITORING	8
5	Pla	nt functional types mapping based on scared data	9
6	Glo	bal upscaled of carbon fluxes	10
7	CO	2 monitoring and net zero modelling platform	11

8	Conclusion	12
Bi	bliography	14

List of Figures

List of Tables

1 Introduction

1.1 Background

2 Background

2.1 Air pollution

2.2 Greenhouse gas

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

AIR POLLUTIOM INDUCED BY INTERVENTION EVENTS

3 Ukraine's case study

3.1 Introduction

Nitrogen dioxide (NO₂) is a key air pollutant that can have harmful effects on human health. An increase in nitrogen oxide (NOx = NO + NO2) concentrations contributes to global warming through a chemical reaction that leads to the formation of ozone (O3), a short-lived climate pollutant with a potent warming effect (Stocker et al., 2013). The lifetime of NO2 is strongly influenced by photochemical reactions and meteorological parameters (Barré et al., 2021) and varies seasonally (Dragomir et al., 2015; Kendrick et al., 2015). During winter, photochemical reaction activity is reduced, resulting in a longer lifetime of the NO2. Additionally, seasonal variations in NO2 concentration are controlled by dispersion processes which are significantly affected by changes in boundary layer height (BLH), wind speed and direction patterns due to temperature inversions in summer and winter (Barré et al., 2021; Kendrick et al., 2015). NO2 concentration levels have been widely used to evaluate decreases in emissions associated with intervention events such as the COVID-19 pandemic lockdown and impacts on the air quality due to the short lifetime of NO2 in the atmosphere (Barré et al., 2021; Cooper et al., 2022). In Europe, anthropogenic NOx emissions are mainly attributed to combustion processes in transportation, as well as energy production and distribution.

In Ukraine, coal-fired power plants (CPPs) dominantly account for 80% of total SO2 and 25% of total NOx emissions, and some have been identified as the highest-emitting CPPs in the region and in the world (Lauri and Rosa, 2021). Since the pandemic started in March 2020, and now with the ongoing armed conflict with Russia, Ukraine has faced a series of threats to the economy, human security and the environment, as well as geopolitical tensions (Pereira et al.,

2022). During the pandemic response starting in 2020, many national and local lockdown restrictions were issued to prevent the spread of the virus, causing a sharp decrease in gross domestic product growth rate, as well as industrial and energy production (Danylyshyn, 2020). In 2021, Ukraine's economy started to recover from the pandemic but the recovery was eventually upended by an armed conflict with Russia that started on February 24, 2022. The conflict has been causing a multi-pronged crisis not only in Ukraine but also in Europe, with increased prices and exacerbated inflation among the many impacts. Many facilities and extensive areas of housing and other infrastructure, including some CPPs, have been reported destroyed or damaged in Ukraine. These impacts have consequently triggered an unprecedented refugee crisis in Ukraine, clogging border crossings between Ukraine and bordering European countries (Júlia et al., 2022). The many socio-economic changes that have occurred during the pandemic and the conflict could be expected to contribute to major variability in air quality in Ukraine, including NO2 pollution levels, during the 2020–2022 period.

3.2 Data

- 3.3 Business-as-usual (BAU) modelling
- 3.4 NO2 changes induced by COVID-19 lockdown
- 3.5 NO2 changes induced by the armed conflict
- 3.6 Conclusion

4 Japan's case study

GREENHOUSE GAS ESTIMATION, FORECASTING AND MONITORING

5 Plant functional types mapping based on scared data

6 Global upscaled of carbon fluxes

7 CO2 monitoring and net zero modelling platform

8 Conclusion

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Acknowledgements

Thank you, and thank you.

Bibliography

- Jérôme Barré, Hervé Petetin, Augustin Colette, Marc Guevara, Vincent-Henri Peuch, Laurence Rouil, Richard Engelen, Antje Inness, Johannes Flemming, Carlos Pérez García-Pando, et al. Estimating lockdown-induced european no< sub> 2</sub> changes using satellite and surface observations and air quality models. Atmospheric chemistry and physics, 21(9):7373–7394, 2021.
- Matthew J Cooper, Randall V Martin, Melanie S Hammer, Pieternel F Levelt, Pepijn Veefkind, Lok N Lamsal, Nickolay A Krotkov, Jeffrey R Brook, and Chris A McLinden. Global fine-scale changes in ambient no2 during covid-19 lockdowns. *Nature*, 601(7893):380–387, 2022.
- Bohdan Danylyshyn. The peculiarities of economic crisis due to covid-19 pandemic in a developing country: case of ukraine. *Problems and Perspectives in Management*, 18(2):13–22, 2020.
- Carmelia Mariana Dragomir, Daniel-Eduard Constantin, Mirela Voiculescu, Lucian Puiu Georgescu, Alexis Merlaud, and Michel Van Roozendael. Modeling results of atmospheric dispersion of no2 in an urban area using meti–lis and comparison with coincident mobile doas measurements. *Atmospheric Pollution Research*, 6(3):503–510, 2015.
- L. Júlia, M. Dylan, R. Danielle, and B Adrian. Nearly 3 million people have left ukraine, foreshadowing a massive humanitarian crisis, 2022. URL https://www.washingtonpost.com/world/2022/02/27/ukraine-refugees-photos-videos/.
- Christine M Kendrick, Peter Koonce, and Linda A George. Diurnal and seasonal variations of no, no2 and pm2. 5 mass as a function of traffic volumes alongside an urban arterial. *Atmospheric Environment*, 122:133–141, 2015.

- M. Lauri and G Rosa. Health impacts of coal power plant emissions in ukraine, 2021. URL https://energyandcleanair.org/publication/health-impacts-of-coal-power-plant-emissions-in-ukraine/.
- Paulo Pereira, Wenwu Zhao, Lyudmyla Symochko, Miguel Inacio, Igor Bogunovic, and Damia Barcelo. The russian-ukrainian armed conflict impact will push back the sustainable development goals, 2022.
- T.F. Stocker, G.-K. Plattner D. Qin, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley. Ipcc, 2013: Climate change 2013: The physical science basis. contribution of working group i to the fifth assessment report of the intergovernmental panel on climate changes. Technical report, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.