

Lecture Notes on Information Theory and Statistics

Shubhanshu Shekhar

Contents

I	The Basics	5
1	Information measures for discrete distributions	7
1.1	A Guessing Game	7
1.2	Formal Definitions	10
1.2.1	An application: generating purely random bits	16
1.3	Properties of Information Measures	17
1.3.1	Chain Rules.	17
1.3.2	Convexity/Concavity.	20
1.3.3	Data Processing Inequality.	22
1.3.4	Fano's inequality.	23
2	Information measures for general distributions	27
2.1	Continuous Distributions	27
2.1.1	Connections to discrete measures via quantization	29
2.1.2	Properties	30
2.1.3	Applications	32
2.2	General Distributions*	36
2.2.1	Variational Definition I: Gelfand-Yaglom-Peres	38
2.2.2	Variational Definition II: Donsker-Varadhan	39
2.3	f -divergences	42
3	Compression and Gambling	47
3.1	Source Codes	47
3.2	Kraft's Inequality	48
3.3	Lower Bound and SFE coding	49
3.4	Gambling, Doubling Rate	50
3.5	Portfolio Optimization + Kelly Betting	50
4	Universal Compression and Gambling	51
4.1	Universal Compression	51
4.1.1	Redundancy-Capacity	51
4.1.2	Regret-Complexity	51
4.1.3	Mixture Method	51
4.2	Universal Portfolios	51
II	Applications	53
5	Application I: From Universal Compression to Sequential Inference	55
5.1	Testing by Betting	55
5.2	Hypothesis Testing	55
5.3	Confidence Sequences	55

Part I

The Basics

Chapter 1

Information measures for discrete distributions

In this chapter, we introduce the three main information measures (entropy, relative entropy, and mutual information) for discrete distributions with finite support. We begin this chapter with a simple guessing game, which naturally leads the definition of these three terms, and we then study some simple properties of these information measures.

Notation. As mentioned earlier, throughout this chapter, we will work with discrete distributions supported on a finite set, which we denote by \mathcal{X} . The elements of this set could be anything, and in particular, we do not assume that there exists any ordering among them. We will mostly consider log to the base 2, unless otherwise stated.

1.1 A Guessing Game

We consider a collection of simple guessing games (or a 20 questions games) that will motivate the definitions of the three main information measures: entropy, relative entropy, and mutual information.

Question 1 (Guessing Game I). *In the simplest setting, suppose there are n identical bins, and a ball is placed uniformly at random in one of the bins. Denote the position of the ball with the random variable $X \sim \text{Uniform}(\mathcal{X})$, where $\mathcal{X} = \{0, 1, \dots, n-1\}$. Suppose, we can make binary queries of the form: “Is X in the set A ?” for $A \subset \mathcal{X}$. Our objective is to design a strategy of asking a series of such questions, such that the average number of questions required to identify the bin containing the ball (i.e., the value of the random variable X) is small.*

A simple strategy could be to ask the series of questions: is $X = 0$, is $X = 1$, and so on. With this strategy, we can check that the average number of questions needed are equal to

$$L = \sum_{i=0}^{n-1} \mathbb{P}(X = i)(i + 1) = \frac{1}{n} \sum_{i=0}^{n-1} (i + 1) = \frac{1}{n} (1 + 2 + \dots + n) = \frac{n}{2}.$$

Thus, the number of yes/no questions required by this strategy (called the linear search) is $\Omega(n)$. This strategy is quite inefficient, since with every query we reduce the search space by one. As we see next, we can do significantly better.

An optimal strategy for the above problem is the *binary search*, in which the queries are specifically designed to reduce the size of search space by half with each query. In particular, consider the case of $n = 4$. Then, the binary search decision tree is shown in Figure 1.1. Assigning the values $0 \leftarrow Y$ and $1 \leftarrow N$, we see that the series of questions to ascertain a value of $X = i$ is equivalent to the binary encoding or representation of i .

$$0 \equiv (YY) \equiv (00), \quad 1 \equiv (YN) \equiv (01), \quad 2 \equiv (NY) \equiv (10), \quad 3 \equiv (NN) \equiv (11).$$




Figure 1.1: The figure shows the decision tree for the optimal strategy (i.e., binary search) for the guessing game with X drawn uniformly at random from $\mathcal{X} = \{0, 1, 2, 3\}$. Each query is chosen to ensure that the outcomes (i.e., Y or N) are equiprobable. In other words, this strategy proceeds by greedily selecting a query whose outcome is the most uncertain.

Denote the number of questions needed to verify $X = i$ by ℓ_i . Then, the binary search scheme asks $\ell_i = 2$ questions for all $i \in \mathcal{X}$ (in other words, it represents each $i \in \mathcal{X}$ with a *codeword* of length $\ell_i = 2$). Interestingly, 2 is also equal to the negative of the logarithm of the probability assigned to each value $i \in \mathcal{X}$ to the base 2; that is, $2 = \log(1/p_i) = \log(4)$. The average number of questions required by this (optimal) strategy is then equal to

$$L = 2 = \sum_{i=0}^3 p_i \ell_i = \sum_{i=0}^3 p_i \log(1/p_i).$$

Thus, the above discussion suggests that the number of binary questions needed to completely remove the uncertainty about the value of $X \sim \text{Uniform}(\mathcal{X})$ is $\log(|\mathcal{X}|)$. What is the analog of this quantity for non-uniform distribution over \mathcal{X} ? We consider this question in the next version of the guessing game.

Question 2 (Guessing game II). *Consider the same setting as Question 1 with $\mathcal{X} = \{0, 1, 2, 3\}$, but assume that X is drawn from the following distribution (instead of uniformly):*

$$P_X = (p_0, p_1, p_2, p_3) = \left(\frac{1}{8}, \frac{1}{8}, \frac{1}{4}, \frac{1}{2} \right).$$

What is the optimal sequence of binary questions to identify the true value of X ?

We will develop a strategy for this problem, motivated by the 'halving' property of the binary search scheme for the uniform case. In particular, we will take the strategy which can be summarized as follows:

make queries whose outcomes are equally likely (or in other words, are most uncertain).

Note that when X is uniformly distributed, the above strategy reduces exactly to the binary search. The decision tree of this strategy for the distribution of Question 2 is shown in Section 1.1. Unlike the previous game, the decision tree is not balanced — it asks more questions of the less likely values of i . The expected number of questions in this case is equal to

$$\begin{aligned} L &= \sum_{i=0}^3 p_i \ell_i = \frac{1}{8} \times 3 + \frac{1}{4} \times 2 + \frac{1}{4} \times 2 + \frac{1}{2} \times 1 = \frac{15}{8} \\ &= - \sum_{i=1}^n p_i \log p_i := H(P_X). \end{aligned}$$

Again the average number of yes/no questions needed to learn X is characterized by the quantity, $-\sum_{i \in \mathcal{X}} p_i \log p_i$. This functional of the probability distribution is called its *entropy*, also called its self-information. As we will see later, it is a fundamental limit on the average number of yes/no questions needed to learn the value of X (or equivalently, the average length of a binary lossless representation of all realizations of X).




Figure 1.2: The figure shows the decision tree for the optimal strategy for a non-uniform probability distribution over \mathcal{X} . The key observation is that this strategy assigns fewer questions (or a shorter codeword) to the higher probability symbol (3), and more questions to the less probable symbols, such as 0 and 1.

In both the previous games, we assumed that the player knows the true distribution of X exactly. We now consider a situation where there is a mismatch between the true and the assumed distribution of X .

Question 3. Consider the same setting as Question 2, but now assume that the true distribution (P_X) of X is not known to the player, and instead he believes that $X \sim Q_X$, where $Q_X = (1/2, 1/8, 1/8, 1/4)$. What is the effect of this distribution mismatch?

Under the assumption that the true distribution is Q_X , the optimal codeword assignment is

$$0 \equiv (Y) \equiv (0), \quad 1 \equiv (NNY) \equiv (110), \quad 2 \equiv (NNN) \equiv (111), \quad 3 \equiv (NY) \equiv (10).$$

Again, as before, in this example, the number of questions needed to ascertain that $X = i$ is equal to $\log(1/q_i)$. The average number of questions needed to learn the value of X is

$$\begin{aligned} L &= \sum_{i=0}^3 p_i \log(1/q_i) = - \sum_{i=1}^3 p_i \log(p_i) + \sum_{i=0}^3 p_i \log(p_i/q_i) \\ &= H(P_X) + D_{\text{kl}}(P_X \parallel Q_X). \end{aligned}$$

The second term in the display is called the relative entropy or KL divergence between P_X and Q_X , and it denotes the price paid by the player for using the wrong model for asking the yes/no questions (or the extra average codeword length incurred due to the ignorance of the true distribution). As we will see later, this quantity is always non-negative.

Question 4. Finally, we now consider a case of guessing another random variable Y on the set $\{a, b\}$. The joint distribution of X and Y is stated in Table 1.1.

Suppose we want to ask a series of yes/no questions to find out the true value of both X and Y . Consider two strategies:

	$X = 0$	$X = 1$	$X = 2$	$X = 3$
$Y = a$	0	1/8	1/8	0
$Y = b$	1/8	0	1/8	1/2

Table 1.1: Joint distribution P_{XY} of (X, Y) .

- Player 1 develops two independent strategies for learning X and Y
- Player 2 develops a joint strategy for learning X and Y together.

Who does better in terms of the average number of questions needed to learning both X and Y ? How much is the improvement?

From the table, we can check that Y has a distribution $P_Y = (1/4, 3/4)$ over the set $\{a, b\}$. The marginal distribution of X is the same as in the previous two questions. Hence, we expect that the average number of yes/no questions need by player 1 (denoted by L_1) is

$$L_1 = H(P_X) + H(P_Y) \approx 1.75 + 0.811 \text{ bits} = 2.561 \text{ bits}$$

For the second player, who develops a joint strategy for querying about (X, Y) , we expect the average number of yes/no questions (denoted by L_2) to be

$$L_2 = H(P_{X,Y}) = 4 \times \frac{1}{8} \times 3 + \frac{1}{2} \times 1 = 2 \text{ bits.}$$

Thus, the player who jointly considers the two random variables requires fewer questions. This is because, knowing the value of Y also provides information about the X value. For instance, if we know that $Y = a$, then we know that X cannot be 0 or 3. Exploiting this leads to the reduced number of questions needed by player 2. The amount of improvement, $L_1 - L_2$, is equal to

$$L_1 - L_2 = H(X) + H(Y) - H(X, Y) := I(X; Y).$$

The term $I(X; Y)$ is called the *mutual information* between the random variables X and Y , and it precisely quantifies the amount of information that X contains about Y (or equivalently, Y contains about X ; since $I(X; Y)$ is symmetric).

Summary. The above discussion can be summarized as follows:

- Entropy $H(X) = -\sum_i p_i \log(p_i)$ quantifies the information content of a random variable X . It is also equal to the minimum average number of yes/no questions needed to learn the value of X .
- The relative entropy is a measure of discrepancy between two distributions P_X and Q_X . It quantifies the additional (on an average) number of yes/no questions needed to learn about X , under wrong model assumptions.
- The mutual information $I(X; Y)$ is measure of dependence between X and Y . It quantifies how much information about X is contained in the random variable Y .

1.2 Formal Definitions

In this section, we formally define the three information measures, and observe some of their basic properties. The next section contains a more thorough treatment of the properties of these measures.

Definition 5. Suppose X denotes an \mathcal{X} -valued random variable with probability mass function (p.m.f.) p_X . Then, the entropy of X (actually the distribution p_X) is defined as

$$H(X) \equiv H(p_X) = \sum_{x \in \mathcal{X}} -p_X(x) \log(p_X(x)). \quad (1.1)$$




Figure 1.3: Variation of the binary entropy $h_2(p) = -p \log(p) - \bar{p} \log(\bar{p})$ with the parameter p . Note that $h_2(p)$ is zero for $p \in \{0, 1\}$, when the random variable is deterministic, and is maximized at $p = 0.5$, which corresponds to the maximum uncertainty. Also note that the qualitative behavior of the curve is similar to that of the variance $p(1 - p)$; another measure of dispersion or variability of the distribution.

For a pair of distributions (X, Y) on $\mathcal{X} \times \mathcal{X}$ with joint distribution p_{XY} , their joint entropy is defined as, following (1.1),

$$H(X, Y) \equiv H(p_{XY}) = \sum_{x \in \mathcal{X}} \sup_{y \in \mathcal{X}} -p_{XY}(x, y) \log(p_{XY}(x, y)).$$

Finally, the conditional entropy of X given Y is defined as the average (over the marginal p_X) of the entropy of $Y|X = x$. That is,

$$H(Y|X) \equiv H(p_{Y|X}|p_X) = \sum_{x \in \mathcal{X}} p_X(x) \sum_{y \in \mathcal{X}} -p_{Y|X}(y|x) \log(p_{Y|X}(y|x)).$$

Remark 6. The base of the logarithm used in defining the entropy characterizes its unit: if the base is 2, entropy is measured in *bits*, while for natural logarithm with base e , the unit is called *nats*. It is easy to verify that changing the base from a to b is achieved by the simple identity: $H_a(X) = \log_b a H_b(X)$.

Entropy is defined as a quantitative measure of the amount of ‘uncertainty’ contained in a probability distribution. In fact, there exist several results that begin by stating a set of reasonable axioms to be satisfied by a good measure of uncertainty, and then prove that the above definition is the only one that simultaneously satisfies those properties [Csiszár and Körner, 2011].

We can also intuitively see that the above definition serves as a good measure of uncertainty. For instance, suppose $\mathcal{X} = \{0, 1\}$ and X is a Bernoulli distribution with parameter p . Then, the entropy of X , also called binary entropy, and denoted by $h_2(p)$, is defined as

$$h_2(p) = -p \log(p) - \bar{p} \log \bar{p}, \quad \text{where } \bar{p} := 1 - p.$$

On plotting it, we can see that the $h_2(p)$ is zero at $p \in \{0, 1\}$, and it achieves its maximum at $p = 1/2$.

The definition of entropy leads to some immediate conclusions that we record next:

Proposition 7. *The following statements are true for \mathcal{X} -valued random variables X, Y etc.:*

- (a) $H(X) \geq 0$, for all random variables X , and its minimum value of 0 is achieved if and only if X is equal to a constant with probability 1.
- (b) The joint entropy of (X, Y) is equal to the entropy of X , plus the conditional entropy of Y given X :

$$H(X, Y) = H(X) + H(Y|X) = H(Y) + H(X|Y).$$

- (c) For any function $f : \mathcal{X} \rightarrow \mathcal{X}$, we have $H(f(X)) \leq H(X)$, with equality iff f is bijective.

Proof. (a) Since $H(X) = \sum_{x \in \mathcal{X}} p(x) \log 1/p(x)$, it is a sum of non-negative terms, which implies that $H(X) \geq 0$. To achieve the equality, note that each $p(x) \log 1/p(x)$ must be equal to zero; which implies that for all $x \in \mathcal{X}$, the value of $p(x)$ must lie in $\{0, 1\}$. Since $\sum_{x \in \mathcal{X}} p(x)$ is constrained to be equal to 1, the result follows.

(b) This follows directly by the definition of joint and conditional entropies. In particular,

$$\begin{aligned} H(X, Y) &= \sum_{x, y} -p(x, y) \log p(x, y) = - \sum_{x \in \mathcal{X}} p(x) \sum_{y \in \mathcal{Y}} p(y|x) (\log p(x) + \log p(y|x)) \\ &= - \sum_{x \in \mathcal{X}} p(x) \log p(x) \sum_{y \in \mathcal{Y}} p(y|x) - \sum_{x \in \mathcal{X}} p(x) \sum_{y \in \mathcal{Y}} p(y|x) \log p(y|x) \\ &= H(X) + H(Y|X). \end{aligned}$$

Repeating the same argument, but now writing $p(x, y) = p(y)p(x|y)$, we get the other equivalent definition.

(c) This is a simple consequence of the previous two results. In particular, we set $Y = f(X)$. Then,

$$H(Y) + H(X|Y) = H(X, Y) = H(X) + H(Y|X) = H(X),$$

where the last equality is a consequence of the fact that $Y = f(X)$; and hence $H(Y|X)$ is equal to 0. Since $H(X|Y)$ is not necessarily zero, we get the required inequality $H(Y) \leq H(X)$. In words, this means that we cannot add more uncertainty (or self-information) to a signal by applying a deterministic function. □

We now present a simple application of the law of large numbers to characterize the support of a high dimensional product distribution in terms of the entropy.

Proposition 8 (Asymptotic equipartition probability (AEP)). *Let $X_1, X_2, \dots, X_n \stackrel{i.i.d.}{\sim} P_X$, and for a fixed $\epsilon > 0$, define the ‘typical set’ $A_n(\epsilon)$ as*

$$A_n(\epsilon) = \left\{ x^n \in \mathcal{X}^n : 2^{-n(H(X)-\epsilon)} \leq \mathbb{P}(X^n) = \prod_{i=1}^n P_X(X_i) \leq 2^{-n(H(X)+\epsilon)} \right\}.$$

Then, the following statements are true:

- $\lim_{n \rightarrow \infty} \mathbb{P}(A_n(\epsilon)) = 1$.
- $|A_n(\epsilon)| \leq 2^{n(H(X)+\epsilon)}$.
- For n large enough, we have $|A_n(\epsilon)| \geq (1 - \epsilon)2^{n(H(X)-\epsilon)}$.

Thus, for large n , a small subset $A_n(\epsilon)$ of \mathcal{X}^n , consisting of roughly equiprobable sequences, contains almost all the probability.

Proof. • This is simply an application of the weak LLN to $-\log(\mathbb{P}(X^n))$.

- We prove this by noting that $\mathbb{P}(A_n(\epsilon)) \leq 1$, and thus

$$1 \geq \sum_{x^n \in A_n(\epsilon)} \mathbb{P}(x^n) \geq \sum_{x^n \in A_n(\epsilon)} 2^{-nH(X)-n\epsilon} = |A_n(\epsilon)| 2^{-nH(X)-n\epsilon}.$$

- By the definition of convergence in probability, for n large enough, we have $\mathbb{P}(A_n(\epsilon)) \geq 1 - \epsilon$. Hence, we have

$$1 - \epsilon \leq \mathbb{P}(A_n(\epsilon)) = \sum_{x^n \in A_n(\epsilon)} \mathbb{P}(x^n) \leq \sum_{x^n \in A_n(\epsilon)} 2^{-n(H(X)-\epsilon)} = |A_n(\epsilon)| 2^{-n(H(X)-\epsilon)}.$$

□




Figure 1.4: Plots of $D_{\text{kl}}(P_X \parallel P_Y)$ and $D_{\text{kl}}(P_Y \parallel P_X)$, where $X \sim \text{Bernoulli}(p)$ and $X \sim \text{Bernoulli}(1/2)$, as p varies in $[0, 1]$. The figures illustrates the locally quadratic behavior of relative entropy, and also shows that it is not symmetric.

Remark 9. The above result can be used to construct a theoretically simple, but computationally infeasible, compression scheme. The idea is simple: enumerate all elements of $x^n \in A_n(\epsilon)$, and assign them their binary representation prefixed by an additional ‘0’ as the codeword; and for all points in $A_n(\epsilon)^c$, assign a codeword starting with ‘1’, followed by the binary representation after enumerating all elements. It is easy to check that this lossless compression scheme has an average codeword length (per symbol) smaller than $H(X) + (2 + \epsilon + \log(|\mathcal{X}|))/n$.

The next, and perhaps the most important, information measure that we introduce is the relative entropy, also known as the Kullback-Leibler or KL divergence.

Definition 10 (Relative Entropy). The relative entropy between two distributions P_X and Q_Y on the same domain \mathcal{X} is defined as

$$D_{\text{kl}}(P_X \parallel Q_Y) = \sum_{x \in \mathcal{X}} p(x) \log \left(\frac{p(x)}{q(x)} \right).$$

For joint distributions P_{XY} and Q_{XY} , the conditional relative entropy is defined as

$$D_{\text{kl}}(P_{Y|X} \parallel Q_{Y|X} | P_X) = \sum_{x \in \mathcal{X}} p(x) \sum_{y \in \mathcal{Y}} p(y|x) \log \left(\frac{p(y|x)}{q(y|x)} \right) = \sum_{x \in \mathcal{X}} p(x) D_{\text{kl}}(P_{Y|X=x} \parallel Q_{Y|X=x}).$$

In other words, the conditional relative entropy is the average (over the marginal P_X) relative entropy between the conditional distributions $P_{Y|X=x}$ and $Q_{Y|X=x}$.

From the definition we can see that the relative entropy is not symmetric in its arguments. As an example, consider the two ways of computing the relative entropy between $X \sim \text{Bernoulli}(p)$ and $Y \sim \text{Bernoulli}(1/2)$, as plotted in Figure 1.4.

Furthermore, if $P \not\ll Q$, then $D_{\text{kl}}(P, Q) = \infty$. We note some other basic properties of relative entropy in our next result.

Proposition 11. (a) Let $\varphi : [0, \infty) \rightarrow \mathbb{R}$ denote the function $\varphi(x) = x \log x$. Then, for two distributions P and Q with p.m.f. p and q respectively, we have $D_{\text{kl}}(P \parallel Q) = \mathbb{E}_Q[\varphi(p(X)/q(X))]$.

(b) For any two distributions P and Q , we have $D_{\text{kl}}(P \parallel Q) \geq 0$. The equality holds if and only if $P = Q$.

(c) Let U denote the uniform distribution over (the finite set) \mathcal{X} . Then, we have

$$D_{\text{kl}}(P \parallel U) = \log(|\mathcal{X}|) - H(P).$$

Proof. (a) Let $\ell(x)$ denote $p(x)/q(x)$. Then, the relative entropy between P and Q can be written as

$$\begin{aligned} D_{\text{kl}}(P \parallel Q) &= \sum_{x \in \mathcal{X}} p(x) \log(\ell(x)) = \sum_{x \in \mathcal{X}} q(x) \ell(x) \log \ell(x) \\ &= \sum_{x \in \mathcal{X}} q(x) \varphi(x) = \mathbb{E}_Q[\varphi(\ell(X))]. \end{aligned}$$

(b) It is easy to verify that the mapping $x \mapsto \varphi(x)$ is convex. Hence, by an application of Jensen's inequality, we have

$$\begin{aligned} D_{\text{kl}}(P \parallel Q) &= \mathbb{E}_Q[\varphi(\ell(X))] \geq \varphi(\mathbb{E}_Q[\ell(X)]) = \varphi\left(\sum_{x \in \mathcal{X}} q(x) \frac{p(x)}{q(x)}\right) \\ &= \varphi(1) = 0. \end{aligned}$$

The above inequality holds with equality if and only if the function $\ell(x)$ is a constant. In other words, this is an equality iff $p(x) = q(x)$ for all $x \in \mathcal{X}$.

(c) This result follows directly by definition. In particular, we have

$$\begin{aligned} D_{\text{kl}}(P \parallel U) &= \sum_{x \in \mathcal{X}} p(x) \log(|\mathcal{X}|p(x)) = \log(|\mathcal{X}|) \sum_{x \in \mathcal{X}} p(x) + \sum_{x \in \mathcal{X}} p(x) \log p(x) \\ &= \log(|\mathcal{X}|) - H(P). \end{aligned}$$

□

An immediate corollary of the above result is that the distribution over \mathcal{X} that achieves the maximum entropy is the uniform distribution. Our next result shows that this statement is more generally true: operations that make a distribution even *slightly more uniform* lead to an increase in entropy.

Proposition 12. *Given a distribution p on \mathcal{X} , let q denote a distribution which replaces $p(x)$ and $p(x')$ with $0.5(p(x) + p(x'))$. Then, we have $H(q) \geq H(p)$.*

Proof. The proof of this result is a direct consequence of Jensen's inequality. In particular, note that due to the convexity of the function $\varphi(x) = x \log x$, we have

$$\begin{aligned} q(x) \log q(x) &= \frac{p(x) + p(x')}{2} \log \left(\frac{p(x) + p(x')}{2} \right) \leq \frac{1}{2} (p(x) \log p(x) + p(x') \log p(x')) \\ q(x') \log q(x') &= \frac{p(x) + p(x')}{2} \log \left(\frac{p(x) + p(x')}{2} \right) \leq \frac{1}{2} (p(x) \log p(x) + p(x') \log p(x')). \end{aligned}$$

On adding the two inequalities, we get

$$-q(x) \log q(x) - q(x') \log q(x') \geq -p(x) \log p(x) - p(x') \log p(x').$$

Since the two entropies differ only on the terms x and x' , the result follows. □

Finally, we introduce the third important quantity, the mutual information.

Definition 13 (Mutual Information). The mutual information between two random variables, X and Y on the domain \mathcal{X} , is defined as

$$I(X; Y) = H(X) + H(Y) - H(X, Y) = H(X) - H(X|Y) = H(Y) - H(Y|X).$$

Alternatively, it is also defined as the relative entropy between the joint distribution of (X, Y) and the product of their marginals:

$$I(X; Y) = D(P_{XY} \parallel P_X \times P_Y).$$

The conditional mutual information between X and Y given Z , is defined as

$$I(X; Y|Z) = H(X|Z) + H(Y|Z) - H(X, Y|Z).$$

It is easy to verify that the two definitions are equivalent. Interestingly, the second definition immediately implies that unlike relative entropy, the mutual information is symmetric quantity. The first definition provides an intuitive explanation of mutual information: since entropy is a measure of uncertainty, the definition suggests that the mutual information between X and Y is the *reduction in uncertainty* about X that is achieved (on average) with the knowledge of Y .

Proposition 14. (a) *All the definitions of mutual information in Definition 13 are equivalent.*

(b) *For any random variable X , we have $I(X; X) = H(X)$. Hence, entropy is also known as the self information.*

(c) *$I(X; Y) \geq 0$, and $I(X; Y) = 0$ if and only if $X \perp Y$.*

(d) *Conditioning reduces entropy: for any pair (X, Y) , we have $H(X|Y) \leq H(X)$, with equality if and only if $X \perp Y$.*

Proof. (a) The first two definitions follow from the definition of joint entropy. In particular, since $H(X, Y) = H(X) + H(Y|X)$, we have

$$H(X) + H(Y) - H(X, Y) = H(X) + H(Y) - H(X) - H(Y|X) = H(Y) - H(Y|X).$$

Similarly, using $H(X, Y) = H(Y) + H(X|Y)$, we get the other definition $I(X; Y) = H(X) - H(X|Y)$.

To get the relative entropy definition, we start with $I(X; Y) = H(X) + H(Y) - H(X, Y)$, to get

$$\begin{aligned} I(X; Y) &= - \sum_{x \in \mathcal{X}} p(x) \log p(x) - \sum_{y \in \mathcal{Y}} p(y) \log p(y) + \sum_{x \in \mathcal{X}, y \in \mathcal{Y}} p(x, y) \log p(x, y) \\ &= - \sum_{x \in \mathcal{X}, y \in \mathcal{Y}} p(x, y) \log p(x) - \sum_{x \in \mathcal{X}, y \in \mathcal{Y}} p(x, y) \log p(y) + \sum_{x \in \mathcal{X}, y \in \mathcal{Y}} p(x, y) \log p(x, y) \\ &= \sum_{x \in \mathcal{X}, y \in \mathcal{Y}} p(x, y) \log \left(\frac{p(x, y)}{p(x)p(y)} \right) \\ &= D_{\text{kl}}(P_{XY} \parallel P_X P_Y). \end{aligned}$$

(b) This follows directly from the definition: $I(X; X) = H(X) - H(X|X) = H(X)$.

(c) The nonnegativity follows from the relative entropy definition. Furthermore, equality occurs if and only if $P_{XY} = P_X \times P_Y$, which is equivalent to independence of X and Y .

(d) This is a direct consequence of the nonnegativity: $0 \leq I(X; Y) = H(X) - H(X|Y)$, which implies that $H(X|Y) \leq H(X)$ as needed. Furthermore, since $I(X; Y)$ is equal to zero iff $X \perp Y$, we have $H(X|Y) = H(X)$ iff $X \perp Y$.

□

The above results imply that mutual information $I(X; Y)$ serves as a measure of dependence between the two random variables, and its value ranges from 0 (for $X \perp Y$) to $H(X)$ (when $X = Y$).

1.2.1 An application: generating purely random bits

Even with the elementary properties we have seen so far, we can obtain non-trivial results about important practical applications, such as the task of generating purely random bits (i.e., fair coin tosses) from bent coins.

Question 15. Suppose $X \sim \text{Bernoulli}(p)$ with $p \in [0, 1]$ unknown: that is, X denotes the outcome of a bent coin. Can we use this bent coin to obtain a fair coin toss $Y \sim \text{Bernoulli}(1/2)$?

Here is a simple procedure, that can simulate a fair coin toss using the bent coin:

- Toss the bent coin twice, and let (X_1, X_2) denote the outcomes. Define the event $E = \{(X_1, X_2) = (0, 1) \text{ or } (X_1, X_2) = (1, 0)\}$.
- On observing the outcomes, if event E occurs, then define $H \equiv \{(X_1, X_2) = (1, 0)\}$ and $T \equiv \{(X_1, X_2) = (0, 1)\}$. Then, we have

$$P(H|E) = P(H \cap E)/P(E) = \frac{\mathbb{P}(X_1 = 1, X_2 = 0)}{\mathbb{P}(X_1 = 1, X_2 = 0) + \mathbb{P}(X_1 = 0, X_2 = 1)} = \frac{p(1-p)}{p(1-p) + (1-p)p} = \frac{1}{2}.$$

Similarly, we can show that $P(T|E) = 1/2$.

- If the event E does not occur, then repeat the process.

Question 16. How can we generalize this scheme to extract multiple uniform bits from n i.i.d. draws from the bent coin? What is the average number of random bits we can extract from n i.i.d. draws?

Given n draws of the bent coin, denote by X_1, \dots, X_n (and assuming that n is even), we proceed as follows:

- Set $K = 0$.
- For i in the range $\{1, 2, \dots, n/2\}$, observe the pair (X_{2i-1}, X_{2i}) , and do one of two things:
 - If $(X_{2i-1}, X_{2i}) \in \{(1, 0), (0, 1)\}$, then set Z_{K+1} equal to 1 or 0 according to the previous rule. Increment $K \leftarrow K + 1$.
 - Otherwise, discard the pair (X_{2i-1}, X_{2i}) .
- Return the fair coin tosses (Z_1, Z_2, \dots, Z_K) .

It is easy to check that for this scheme, conditioned on $K = k$, all the 2^k bits are equally likely. Furthermore, the expected number of purely random bits that we can extract from n bent coin tosses is

$$\mathbb{E}[K] = \sum_{i=1}^{n/2} \mathbb{P}((X_{2i-1}, X_{2i}) \in \{(1, 0), (0, 1)\}) = np(1-p).$$

Question 17. Can we do better?

We can obtain a method agnostic upper bound on the achievable performance of any random bit generating scheme \mathcal{A} . In particular, let \mathcal{A} denote any scheme that maps $X^n = (X_1, \dots, X_n)$ to a random bits $(Z_1, Z_2, \dots, Z_K, K)$, which satisfy the property: *conditioned on $K = k$, the vector Z^k is uniformly distributed over $\{0, 1\}^k$, for all $k \in [n]$* . We then have the following:

$$\begin{aligned} nh_2(p) &= H(X^n) \geq H(Z^K, K) = H(K) + H(Z^K|K) \\ &\geq H(Z^K|K) = \sum_{k \geq 1} \mathbb{P}(K = k) H(Z^k|K = k) \\ &= \sum_{k \geq 1} \mathbb{P}(K = k) k = \mathbb{E}[K]. \end{aligned}$$

There exist methods, such as Peres' iterated extractor [Peres, 1992], that achieves this optimal rate in the limit of large n .

1.3 Properties of Information Measures

1.3.1 Chain Rules.

We begin by establishing the chain rules for the three information measures. These results decompose the information measures for a collection of random variables into a sum of simpler terms.

Theorem 18 (Chain rule for entropy). *Consider a random vector (X_1, X_2, \dots, X_n) taking values in \mathcal{X}^n . Then, we have*

$$H(X_1, \dots, X_n) = H(X_1) + H(X_2|X_1) + \dots + H(X_i|X_1, \dots, X_{i-1}) + \dots + H(X_n|X_1, \dots, X_{n-1}).$$

For any permutation $\pi : [n] \rightarrow [n]$, we have

$$H(X_1, \dots, X_n) = H(X_{\pi(1)}) + H(X_{\pi(2)}|X_{\pi(1)}) + \dots + H(X_{\pi(n)}|X_{\pi(1)}, \dots, X_{\pi(n-1)}).$$

Since conditioning reduces entropy, we also have the following inequality:

$$H(X_1, \dots, X_n) \leq \sum_{i=1}^n H(X_i),$$

with equality only if (X_1, \dots, X_n) are independent.

Proof. The proof follows by induction:

- For $n = 2$, we already proved it directly from the definition of joint entropy.
- Suppose the statement is true for $n - 1$.
- Then, we have

$$\begin{aligned} H(X_1, \dots, X_n) &= H(X^{n-1}, X_n) \stackrel{(i)}{=} H(X^{n-1}) + H(X_n|X^{n-1}) \\ &\stackrel{(ii)}{=} \sum_{i=1}^{n-1} H(X_i|X^{i-1}) + H(X_n|X^{n-1}) = \sum_{i=1}^n H(X_i|X^{i-1}), \end{aligned}$$

where (i) follows from the $n = 2$ case, and (ii) follows from the induction hypothesis. □

Theorem 19 (Chain rule for relative entropy). *Consider two joint distribution of (X_1, \dots, X_n) , denoted by P_{X^n} , and Q_{X^n} . Then, the relative entropy between these two distributions can be decomposed as*

$$D_{kl}(P_{X^n} \parallel Q_{X^n}) = \sum_{i=1}^n D_{kl}(P_{X_i|X^{i-1}} \parallel Q_{X_i|X^{i-1}} | P_{X^{i-1}}). \quad (1.2)$$

Again, for any permutation $\pi : [n] \rightarrow [n]$, we have

$$D_{kl}(P_{X^n} \parallel Q_{X^n}) = \sum_{i=1}^n D_{kl}\left(P_{X_{\pi(i)}|X_{\pi(1)}^{\pi(i-1)}} \parallel Q_{X_{\pi(i)}|X_{\pi(1)}^{\pi(i-1)}} | P_{X_{\pi(1)}^{\pi(i-1)}}\right).$$

If $Q_{X^n} = \prod_{i=1}^n Q_{X_i}$, then, we have

$$D_{kl}(P_{X^n} \parallel Q_{X^n}) = D_{kl}\left(P_{X^n} \parallel \prod_{i=1}^n P_{X_i}\right) + \sum_{i=1}^n D_{kl}(P_{X_i} \parallel Q_{X_i}) \quad (1.3)$$

$$\stackrel{(a)}{\geq} \sum_{i=1}^n D_{kl}(P_{X_i} \parallel Q_{X_i}). \quad (1.4)$$

The equality in (a) occurs when P_{X^n} is equal to the product of its marginals.

Proof. We prove (1.2) for the special case $n = 2$, since the general case follows by induction.

$$\begin{aligned}
D_{\text{kl}}(P_{X_1 X_2} \parallel Q_{X_1 X_2}) &= \sum_{x_1, x_2} p(x_1, x_2) \log \left(\frac{p(x_1, x_2)}{q(x_1, x_2)} \right) = \sum_{x_1, x_2} p(x_1, x_2) \log \left(\frac{p(x_1)p(x_2|x_1)}{q(x_1)q(x_2|x_1)} \right) \\
&= \sum_{x_1} p(x_1) \log \left(\frac{p(x_1)}{q(x_1)} \right) + \sum_{x_1, x_2} p(x_1, x_2) \log \left(\frac{p(x_2|x_1)}{q(x_2|x_1)} \right) \\
&= D_{\text{kl}}(P_{X_1} \parallel Q_{X_1}) + \sum_{x_1} p(x_1) \sum_{x_2} p(x_2|x_1) \log \left(\frac{p(x_2|x_1)}{q(x_2|x_1)} \right) \\
&= D_{\text{kl}}(P_{X_1} \parallel Q_{X_1}) + D_{\text{kl}}(P_{X_2|X_1} \parallel Q_{X_2|X_1} | P_{X_1}).
\end{aligned}$$

Next, to prove (1.3), we note that

$$\begin{aligned}
D_{\text{kl}}(P_{X^n} \parallel Q_{X^n}) &= \sum_{x^n} p(x^n) \log \left(\frac{p(x^n)}{\prod_{i=1}^n q_i(x_i)} \right) = \sum_{x^n} p(x^n) \log \left(\frac{p(x^n)}{\prod_{i=1}^n p_i(x_i)} \frac{\prod_{i=1}^n p_i(x_i)}{\prod_{i=1}^n q_i(x_i)} \right) \\
&= \sum_{x^n} p(x^n) \log \left(\frac{p(x^n)}{\prod_{i=1}^n p_i(x_i)} \right) + \sum_{x^n} p(x^n) \log \left(\frac{\prod_{i=1}^n p_i(x_i)}{\prod_{i=1}^n q_i(x_i)} \right) \\
&= D_{\text{kl}}(P_{X^n} \parallel \prod_{i=1}^n P_{X_i}) + \sum_{i=1}^n \sum_{x_i} p_i(x_i) \log \left(\frac{p_i(x_i)}{q_i(x_i)} \right) \\
&= D_{\text{kl}}(P_{X^n} \parallel \prod_{i=1}^n P_{X_i}) + \sum_{i=1}^n D_{\text{kl}}(P_{X_i} \parallel Q_{X_i}).
\end{aligned}$$

The inequality (1.4) follows due to the nonnegativity of $D_{\text{kl}}(P_{X^n} \parallel \prod_{i=1}^n P_{X_i})$, and note that the equality occurs if and only if this term is zero. This happens iff $P_{X^n} = \prod_{i=1}^n P_{X_i}$. \square

Corollary 20. *A simple consequence of the chain rule for relative entropy is the fact that “conditioning increases divergence”. Namely, suppose $P_{XY} = P_X P_{Y|X}$, and $Q_{XY} = P_X Q_{Y|X}$. Then, we have*

$$\begin{aligned}
D_{\text{kl}}(P_{XY} \parallel Q_{XY}) &= D_{\text{kl}}(P_Y \parallel Q_Y) + D_{\text{kl}}(P_{X|Y} \parallel Q_{X|Y} | P_Y) \\
&= D_{\text{kl}}(P_X \parallel P_X) + D_{\text{kl}}(P_{Y|X} \parallel Q_{Y|X} | P_X).
\end{aligned}$$

Since $D_{\text{kl}}(P_X \parallel P_X) = 0$ and $D_{\text{kl}}(P_{X|Y} \parallel Q_{X|Y} | P_Y) \geq 0$, the above implies

$$D_{\text{kl}}(P_Y \parallel Q_Y) \leq D_{\text{kl}}(P_{Y|X} \parallel Q_{Y|X} | P_X).$$

Finally, since mutual information is an instance of relative entropy, it also satisfies an analogous chain rule.

Proposition 21 (Chain rule for mutual information).

$$I(X_1, \dots, X_n; Y) = \sum_{i=1}^n I(X_i; Y | X_1^{i-1}).$$

For any permutation $\pi : [n] \rightarrow [n]$, we have

$$I(X_1, \dots, X_n; Y) = \sum_{i=1}^n I(X_{\pi(i)}; Y | X_1^{\pi(i-1)}).$$

Proof. This is a simple consequence of the chain rule for entropy. Again, we prove the result for the case of $n = 2$, since the general result follows by induction.

$$\begin{aligned}
I(X_1, X_2; Y) &= H(X_1, X_2) - H(X_1, X_2 | Y) = H(X_1) + H(X_2 | X_1) - H(X_1 | Y) - H(X_2 | X_1, Y) \\
&= (H(X_1) - H(X_1 | Y)) + (H(X_2 | X_1) - H(X_2 | X_1, Y)) \\
&= I(X_1; Y) + I(X_2; Y | X_1).
\end{aligned}$$

\square

Application: Han's inequality and an isoperimetric inequality for the binary hypercube. The simple chain rules obtained in this section often serve as an important tool in establishing various properties of information measures. We illustrate this by proving Han's inequalities for entropy and relative entropy.

Proposition 22. *Let X_1, \dots, X_n denote n , possibly dependent, \mathcal{X} -valued random variables. For any $i \in [n]$, define $X^{(i)} = (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$. Then, we have*

$$H(X_1, \dots, X_n) \leq \frac{1}{n-1} \sum_{i=1}^n H(X^{(i)}).$$

Proof. Introduce the notations $X^i = (X_1, \dots, X_i)$ and $X_i^j = (X_i, \dots, X_j)$ for $j \geq i$. Then, we have

$$\begin{aligned} H(X^n) &\stackrel{(i)}{=} H(X^{(i)}) + H(X_i | X^{(i)}) = H(X^{(i)}) + H(X_i | X^{i-1}, X_{i+1}^n) \\ &\stackrel{(i)}{\leq} H(X^{(i)}) + H(X_i | X^{i-1}), \end{aligned}$$

where (i) uses the chain rule for entropy, and (ii) follows from the fact that conditioning reduces entropy. The above inequality is true for all values of $i \in [n]$, and hence summing them up, we get

$$nH(X^n) \leq \sum_{i=1}^n H(X^{(i)}) + \sum_{i=1}^n H(X_i | X^{i-1}) \stackrel{(iii)}{=} \sum_{i=1}^n H(X^{(i)}) + H(X^n),$$

where (iii) again follows from the chain rule for entropy. Subtracting $H(X^n)$ from both sides, and dividing by $n-1$ leads to the required result. \square

Han's inequality has several applications in combinatorics and concentration of measure. We illustrate this with a simple result about the *density* of the subgraphs of binary hypercube.

Proposition 23. *Let $V = \{-1, 1\}^n$ denote the vertices of a binary hypercube in n dimensions, and let $E(V) = \{(x, x') : x, x' \in V, d_H(x, x') = 1\}$ denote the set of its edges (represented via unordered pairs of vertices). Note that $|V| = 2^n$, and $|E(V)| = n2^{n-1} = 2^{n-1} \log |V|$. Let A denote any subset of V , and $E(A)$ denote the edges between vertices in A . Then, we have*

$$|E(A)| \leq \frac{|A|}{2} \log |A|.$$

Proof. For the given subset $A \subset V = \{-1, 1\}^n$, let $X^n = (X_1, \dots, X_n)$ be a random variable distributed uniformly over A . Then, by chain rule for entropy, we have the following for any $i \in [n]$:

$$H(X^n) - H(X^{(i)}) = H(X_i | X^{(i)}) = - \sum_{x^n \in A} p(x^n) \log p(x_i | x^{(i)}).$$

Given an $x^n \in A$, let $\bar{x}^{(i)}$ denote $(x_1, \dots, x_{i-1}, -x_i, x_{i+1}, \dots, x_n)$: the vector with the i^{th} element flipped. Then, we have

$$p(x_i | x^{(i)}) = 1 - \frac{1}{2} \mathbf{1}_{\bar{x}^{(i)} \in A}.$$

In other words, $\log(p(x_i | x^{(i)})) = \log(2) = 1$ if $\bar{x}^{(i)} \in A$, and 0 otherwise. Plugging this into the above equation, we get

$$H(X^n) - H(X^{(i)}) = \frac{1}{|A|} \sum_{x^n \in A} \mathbf{1}_{\bar{x}^{(i)} \in A},$$

which on summing over $i \in [n]$, gives us

$$\begin{aligned} \sum_{i=1}^n H(X^n) - H(X^{(i)}) &= \frac{1}{|A|} \sum_{x^n \in A} \sum_{i=1}^n \mathbf{1}_{\bar{x}^{(i)} \in A} \\ &= \frac{1}{|A|} \sum_{x^n \in A} |\{x' \in A : d_H(x, x') = 1\}| = \frac{2|E|}{|A|}. \end{aligned}$$

Finally, from an application of Han's inequality, we get

$$\frac{2|E(A)|}{|A|} = \sum_{i=1}^n H(X^n) - H(X^{(i)}) = nH(X^n) - \sum_{i=1}^n H(X^{(i)}) \leq H(X^n) = \log(|A|).$$

This completes the proof. \square

1.3.2 Convexity/Concavity.

We begin with a simple consequence of the convexity of the mapping $x \mapsto x \log x$, that will be useful in establishing some properties of information measures.

Proposition 24. *Let $\{a_i, b_i : 1 \leq i \leq m\}$ denote non-negative real numbers (not all of which are zero), and define $A_m = \sum_{i=1}^m a_i$, and $B_m = \sum_{i=1}^m b_i$. Then, we have*

$$A_m \log \left(\frac{A_m}{B_m} \right) \leq \sum_{i=1}^m a_i \log \left(\frac{a_i}{b_i} \right).$$

The equality occurs if and only if a_i/b_i is a constant for all $i \in [m]$.

Proof. Introduce the terms $\tilde{a}_i = a_i/A_m$, and $\tilde{b}_i = b_i/B_m$, and note that $(\tilde{a}_1, \dots, \tilde{a}_m)$, and $(\tilde{b}_1, \dots, \tilde{b}_m)$ represent two probability distributions on $[m]$. Denote these probability distributions by P_a and P_b respectively. Then, we have

$$\begin{aligned} \sum_{i=1}^m a_i \log(a_i/b_i) &= A_m \left(\sum_{i=1}^m \tilde{a}_i \log(\tilde{a}_i/\tilde{b}_i) + \log \left(\frac{A_m}{B_m} \right) \right) \\ &= A_m \log \left(\frac{A_m}{B_m} \right) + A_m \left(\sum_{i=1}^m \tilde{a}_i \log(\tilde{a}_i/\tilde{b}_i) \right) \\ &= A_m \log \left(\frac{A_m}{B_m} \right) + A_m D_{\text{kl}}(P_a \parallel P_b) \\ &\geq A_m \log \left(\frac{A_m}{B_m} \right). \end{aligned}$$

The inequality follows from the non-negativity of relative entropy, and the fact that $A_m \geq 0$ by assumption.

The equality occurs iff the relative entropy between P_a and P_b is zero:

$$D_{\text{kl}}(P_a \parallel P_b) = 0 \Leftrightarrow P_a = P_b \Leftrightarrow \frac{a_i}{b_i} = \text{constant, for all } i \in [m].$$

\square

As an application of the log-sum inequality, we can establish the convexity of relative entropy.

Theorem 25. *Let P_1, P_2, Q_1, Q_2 be distributions over \mathcal{X} . For any $\lambda \in [0, 1]$, define $P_\lambda = \lambda P_1 + \bar{\lambda} P_2$, and $Q_\lambda = \lambda Q_1 + \bar{\lambda} Q_2$. Then, we have*

$$D_{\text{kl}}(P_\lambda \parallel Q_\lambda) \leq \lambda D_{\text{kl}}(P_1 \parallel Q_1) + \bar{\lambda} D_{\text{kl}}(P_2 \parallel Q_2).$$

Proof. For $\lambda \in \{0, 1\}$, the result holds trivially. So we consider the case of $\lambda \in (0, 1)$.

The relative entropy between P_λ and Q_λ is equal to

$$D_{\text{kl}}(P_\lambda \parallel Q_\lambda) = \sum_{x \in \mathcal{X}} \lambda p_1(x) + \bar{\lambda} p_2(x) \log \left(\frac{\lambda p_1(x) + \bar{\lambda} p_2(x)}{\lambda q_1(x) + \bar{\lambda} q_2(x)} \right).$$

We now apply the log-sum inequality (Proposition 24) to each term in the summation. In particular, for a fixed $x \in \mathcal{X}$, define

$$a_1 = \lambda p_1(x), \quad a_2 = \bar{\lambda} p_2(x), \quad b_1 = \lambda q_1(x), \quad \text{and} \quad b_2 = \bar{\lambda} q_2(x).$$

An application of Proposition 24 implies that

$$(a_1 + a_2) \log \left(\frac{a_1 + a_2}{b_1 + b_2} \right) \leq a_1 \log(a_1/b_1) + a_2 \log(a_2/b_2) = \lambda p_1(x) \log \left(\frac{p_1(x)}{q_1(x)} \right) + \bar{\lambda} p_2(x) \log \left(\frac{p_2(x)}{q_2(x)} \right).$$

The equality above uses the fact that $\lambda \notin \{0, 1\}$. On summing this upper bound over all $x \in \mathcal{X}$, we get the required result. \square

A simple consequence of Theorem 25 is that the entropy is a concave functional of the pmf.

Corollary 26. *Suppose P_1 and P_2 are two distributions on \mathcal{X} . Then, for any $\lambda \in [0, 1]$, we have*

$$H(P_\lambda) \geq \lambda H(P_1) + \bar{\lambda} H(P_2).$$

Proof. We know that the entropy of P_1 (resp. P_2) can be defined in terms of the relative entropy between P_1 (resp. P_2) and the uniform distribution over \mathcal{X} (denoted by U):

$$H(P_1) = \log(|\mathcal{X}|) - D_{\text{kl}}(P_1 \parallel U), \quad \text{and} \quad H(P_2) = \log(|\mathcal{X}|) - D_{\text{kl}}(P_2 \parallel U).$$

This implies that for any $\lambda \in [0, 1]$, we have

$$\begin{aligned} \lambda H(P_1) + \bar{\lambda} H(P_2) &= \log(|\mathcal{X}|) - (\lambda D_{\text{kl}}(P_1 \parallel U) + \bar{\lambda} D_{\text{kl}}(P_2 \parallel U)) \\ &\leq \log(|\mathcal{X}|) - D_{\text{kl}}(P_\lambda \parallel U) \\ &= H(P_\lambda), \end{aligned}$$

where the inequality follows from the convexity of relative entropy. \square

Finally, we characterize the convexity/concavity of the mutual information.

Theorem 27. *For two random variables X and Y taking values on finite sets \mathcal{X} and \mathcal{Y} respectively,*

- *for a fixed conditional distributions $\{p_{Y|X}(\cdot|x) : x \in \mathcal{X}\}$, the mapping $p_X \rightarrow I(X; Y)$ is concave.*
- *for a fixed marginal p_X , the mapping $p_{Y|X} \rightarrow I(X; Y)$ is convex.*

Proof. For the first statement, note that if the conditional distribution (or the channel, or the Markov kernel) $p_{Y|X}$ is fixed, then p_Y is a linear function of p_X . To see this, if we think of p_X, p_Y as row vectors, and use K to represent the $|\mathcal{X}| \times |\mathcal{Y}|$ transition matrix corresponding to $p_{Y|X}$, then $p_Y = p_X K$. Now, we observe that

$$I(X; Y) = H(Y) - H(Y|X) = H(Y) - \sum_{x \in \mathcal{X}} p_X(x) H(Y|X = x).$$

Since $H(Y)$ is a concave function of p_Y , which in turn is a linear function of p_X , we conclude that $H(Y)$ is a concave function of p_X . The second term is simply a linear function of p_X , and hence their difference, $I(X; Y)$, is a concave function of p_X , with $P_{Y|X}$ fixed.

For the second result, we state the mutual information in terms of relative entropy. In particular, we have

$$I(X; Y) = D_{\text{kl}}(P_{XY} \parallel P_X \times P_Y) = D_{\text{kl}}(P_X \parallel P_X) + D_{\text{kl}}(P_{Y|X} \parallel P_Y|P_X).$$

We already know that relative entropy is convex in its arguments, and the result follows by noting that for two Markov kernels, K_1 and K_2 , and a $\lambda \in [0, 1]$, we have

$$p_Y^\lambda = \lambda(p_X K_1) + \bar{\lambda}(p_X K_2) = p_X (\lambda K_1 + \bar{\lambda} K_2).$$

In particular, we have

$$\begin{aligned} D_{\text{kl}}(K^\lambda \parallel p_Y^\lambda | P_X) &= D_{\text{kl}}(\lambda K_1 + \bar{\lambda} K_2 \parallel \lambda p_Y^1 + \bar{\lambda} p_Y^2 | P_X) \\ &\leq \lambda D_{\text{kl}}(K_1 \parallel p_Y^1 | P_X) + \bar{\lambda} D_{\text{kl}}(K_2 \parallel p_Y^2 | P_X), \end{aligned}$$

as required. \square

Remark 28. We know that by definition $I(X; Y)$ is always upper bounded by $H(X)$, which is further upper bounded by $\log(|\mathcal{X}|)$. Now, for a fixed channel (i.e., transition kernel), the above result says that $I(X; Y)$ is a non-negative, concave function of the marginal p_X . Hence, the term

$$C = \max_{p_X} I(X; Y),$$

is well-defined and is called the *channel capacity* associated with the channel $\{p_{Y|X=x} : x \in \mathcal{X}\}$.

1.3.3 Data Processing Inequality.

In this section we look at a class of inequalities called that *data processing inequalities*. Informally, these results tell us that we cannot increase the information contained in a signal (Y) about another unknown signal (X) by processing Y .

Theorem 29. Suppose $X \rightarrow Y \rightarrow Z$ for a Markov chain; that is, $X \perp Z|Y$. Then, we have

$$I(X; Z) \leq I(X; Y), \quad \text{and} \quad I(X; Z) \leq I(Y; Z).$$

Proof. We show the proof of the first inequality, since the second inequality can be proved with the same argument.

We begin by writing $I(X; Z)$ in two equivalent ways:

$$I(X; Z) = H(X) - H(X|Z) = H(Z) - H(Z|X).$$

Since conditioning reduces entropy, we have $H(X|Z) \leq H(X|Z, Y)$. Furthermore, due to the Markov property, $X \perp Z|Y$, which means that $H(X|Z, Y) = H(X|Y)$. This implies that

$$I(X; Z) = H(X) - H(X|Z) \leq H(X) - H(X|Z, Y) = H(X) - H(X|Y) = I(X; Y).$$

Note that the above relation holds with an equality iff $H(X|Z) = H(X|Z, Y)$. Or in other words, $X \rightarrow Z \rightarrow Y$ form a Markov chain. \square

An immediate corollary of the above result gives us a data processing inequality for entropy.

Corollary 30. Suppose $X \rightarrow Y \rightarrow Z$. Then, we have

$$H(X|Y) \leq H(X|Z).$$

Proof. The result follows from Theorem 29 by expanding $I(X; Y)$ and $I(X; Z)$ in terms of the entropies. That is,

$$I(X; Z) \leq I(X; Y) \Rightarrow H(X) - H(X|Z) \leq H(X) - H(X|Y) \Rightarrow H(X|Y) \leq H(X|Z),$$

as required. \square

The above result simply says that the amount of residual uncertainty about X given Y is never larger than the residual uncertainty about X given Z ; where Z is some (possibly random) transform of Y .

Finally we present a DPI for relative entropy.

Theorem 31. Consider two distributions P_X and Q_X over the alphabet \mathcal{X} , and let $K_{Y|X}$ denote a transition probability matrix. Then, with $P_Y = P_X K_{Y|X}$, and $Q_Y = Q_X K_{Y|X}$, we have

$$D_{kl}(P_X \parallel Q_X) \geq D_{kl}(P_Y \parallel Q_Y).$$

In particular, if Y is any deterministic function of X , then we have

$$D_{kl}(P_X \parallel Q_X) \geq D_{kl}(P_{f(X)} \parallel Q_{f(X)}).$$

Proof. This statement is a simple consequence of the chain rule, and nonnegativity of relative entropy. In particular, using the chain rule, we can expand the relative entropy in two ways. First, we get

$$\begin{aligned} D_{\text{kl}}(P_{XY} \parallel Q_{XY}) &= D_{\text{kl}}(P_X \parallel Q_X) + D_{\text{kl}}(Q_{Y|X} \parallel Q_{Y|X}|P_X) \\ &= D_{\text{kl}}(P_X \parallel Q_X) + D_{\text{kl}}(\mathbf{K}_{Y|X} \parallel \mathbf{K}_{Y|X}|P_X) \\ &= D_{\text{kl}}(P_X \parallel Q_X). \end{aligned} \quad (1.5)$$

Now, expanding it the other way, we get

$$\begin{aligned} D_{\text{kl}}(P_{XY} \parallel Q_{XY}) &= D_{\text{kl}}(P_Y \parallel Q_Y) + D_{\text{kl}}(Q_{X|Y} \parallel Q_{X|Y}|P_Y) \\ &\geq D_{\text{kl}}(P_Y \parallel Q_Y), \end{aligned} \quad (1.6)$$

where the inequality follows from the nonnegativity of $D_{\text{kl}}(Q_{X|Y} \parallel Q_{X|Y}|P_Y)$. Combining (1.5) and (1.6), we get the required result. \square

Application: Impossibility results in hypothesis testing. Consider a hypothesis testing problem with i.i.d. observations $X_1, X_2, \dots, X_n \in \mathcal{X}$ drawn from a distribution P_X , with mean μ_X with the null and alternative hypotheses defined as follows:

$$H_0 : \mu_X = 0.5, \quad \text{versus} \quad H_1 : \mu_X \geq 0.5 + \Delta.$$

Suppose there exists a test $\Psi : \mathcal{X}^n \rightarrow [0, 1]$, with both type-I and type-II errors controlled at a level $\alpha \in (0, 1)$. That is, the following two statements are simultaneously true:

$$p_1 := \mathbb{E}_{H_1}[\Psi(X^n)] \geq 1 - \alpha := \bar{\alpha}, \quad \text{and} \quad p_0 := \mathbb{E}_{H_0}[\Psi(X^n)] \leq \alpha, \quad (1.7)$$

for some $\alpha \in (0, 1/4]$. Then, with a straightforward application of the DPI for relative entropy, we can obtain a lower bound on the parameter Δ . In particular, suppose H_1 is true, and the true distribution P_X has mean $\mu \geq 0.5 + \Delta$. Let P_0 denote the null distribution; that is Bernoulli with mean 0.5

$$\begin{aligned} nD_{\text{kl}}(P_X \parallel P_0) &= D_{\text{kl}}(P_X^n \parallel P_0^n) \stackrel{(a)}{\geq} D_{\text{kl}}(\mathbb{E}_{P_X^n}[\Psi(X^n)] \parallel \mathbb{E}_{P_0^n}[\Psi(X^n)]) \\ &\geq D_{\text{kl}}(\bar{\alpha} \parallel \alpha) = \bar{\alpha} \log(\bar{\alpha}/\alpha) + \alpha \log(\alpha/\bar{\alpha}) \\ &= (1 - 2\alpha) \log(\bar{\alpha}/\alpha) \geq \frac{1}{2} \log\left(\frac{1}{2\alpha}\right). \end{aligned}$$

The inequality (a) follows from an application of the DPI for relative entropy. The above chain of inequality says that if there exists a test Ψ satisfying (1.7), then the null and alternatives must be separated in relative entropy by at least $\log(1/2\alpha)/2n$. In other words,

$$\begin{aligned} D^*(\Delta) &:= \inf\{D_{\text{kl}}(P_X \parallel P_0) : \mu_X \geq 1/2 + \Delta\} \geq A_n := \frac{1}{2n} \log\left(\frac{1}{2\alpha}\right) \\ \Rightarrow \Delta &\geq (D^*)^{-1}(A_n) := \min\{\Delta' > 0 : D^*(\Delta') \geq A_n\}. \end{aligned}$$

1.3.4 Fano's inequality.

Consider the Markov chain $X \rightarrow Y \rightarrow \hat{X}$. Here, X might denote some signal that is transmitted through a noisy channel, Y is the output of the noisy channel, and \hat{X} might denote an estimate of X constructed on the basis of Y by the decoder. Assume that both X and \hat{X} lie in some finite set \mathcal{X} . If $p_e = \mathbb{P}(\hat{X} \neq X)$ denotes the probability of error, then it is intuitive to expect p_e to depend on the amount of residual uncertainty about X given that we know Y . Fano's inequality is one way of formalizing this intuition.

Proposition 32. *For any decoder \hat{X} , we have the following:*

$$h_2(p_e) + p_e \log(|\mathcal{X}| - 1) \geq H(X|\hat{X}) \geq H(X|Y).$$

Remark 33. Note that it suffices to prove the first inequality, as the second follows from an application of data-processing inequality for entropy.

Proof. Introduce the Bernoulli random variable $Z = \mathbf{1}_{X \neq \hat{X}}$. Note that $\mathbb{P}(Z = 1) = \mathbb{P}(X \neq \hat{X}) = p_e$. Then, consider the following:

$$H(X, Z|\hat{X}) = H(X|\hat{X}) + H(Z|X, \hat{X}) = H(X|\hat{X}), \quad (1.8)$$

where we used the fact that Z is a deterministic function of X and \hat{X} , which implies that $H(Z|X, \hat{X}) = 0$. Now, expanding $H(X, Z|\hat{X})$ another way, we get

$$\begin{aligned} H(X, Z|\hat{X}) &= H(Z|\hat{X}) + H(X|\hat{X}, Z) \stackrel{(i)}{\leq} H(Z) + H(X|\hat{X}, Z) \\ &= H(p_e) + \mathbb{P}(Z = 1)H(X|\hat{X}, Z = 1) + \mathbb{P}(Z = 0)H(X|\hat{X}, Z = 0) \\ &\stackrel{(ii)}{=} H(p_e) + \mathbb{P}(Z = 1)H(X|\hat{X}, Z = 1) + 0 \\ &\stackrel{(iii)}{\leq} H(p_e) + p_e \log(|\mathcal{X}| - 1). \end{aligned} \quad (1.9)$$

In the above display,

(i) uses the fact that conditioning reduces entropy,

(ii) uses the fact that when $Z = 0$, then X is equal to \hat{X} ,

(iii) uses the fact that when $Z = 1$, X can take one of $|\mathcal{X}| - 1$ values in \mathcal{X} not equal to \hat{X} .

The result then follows by combining (1.8) with (1.9). \square

A simple corollary of the above inequality is often useful in statistical applications.

Corollary 34. Suppose X is uniformly distributed over \mathcal{X} . Then, we have

$$p_e \geq 1 - \frac{I(X; \hat{X}) - 1}{\log(|\mathcal{X}|)}.$$

Proof. The starting point of this result is the standard version of Fano's inequality:

$$-H(X|\hat{X}) \geq -h_2(p_e) - p_e \log(|\mathcal{X}|) \Rightarrow H(X) - H(X|\hat{X}) \geq -h_2(p_e) + (1 - p_e)H(X).$$

Noting that $h_2(p_e) \leq 1$ (bits), on rearranging the above, we get

$$p_e H(X) \geq H(X) - I(X; \hat{X}) - 1.$$

We get the required statement by dividing both sides by $H(X) = \log(|\mathcal{X}|)$. \square

The two inequalities above give us a bound on the probability of error under an exact recovery criterion: that is, $p_e = \mathbb{P}(\hat{X} \neq X)$. We now present a simple generalization for the case of approximate recovery, when the domain \mathcal{X} is endowed with a distance measure d .

Corollary 35. For a real number $t > 0$, introduce $p_t = \mathbb{P}(d(\hat{X}, X) > t)$, and define

$$N_{\max}(t) = \max_{x \in \mathcal{X}} |B(x, t)|,$$

where $B(x, t) = \{x' \in \mathcal{X} : d(x, x') \leq t\}$ denotes the closed ball of radius t around x . Suppose $X \sim \text{Uniform}(\mathcal{X})$, and $X \rightarrow Y \rightarrow \hat{X}$ form a Markov chain. Then, we have

$$p_t \geq 1 - \frac{I(X, \hat{X}) - 1}{\log\left(\frac{|\mathcal{X}|}{N_{\max}(t)}\right)}.$$

Proof. This result follows from the same general idea. We define $Z = \mathbf{1}_{d(X, \hat{X}) > t}$, and note that $Z \sim \text{Bernoulli}(p_t)$.

$$H(X, Z|\hat{X}) = H(X|\hat{X}) + H(Z|X, \hat{X}) = H(Z|\hat{X}) + H(X|Z, \hat{X}).$$

As before, $H(Z|X, \hat{X}) = 0$. Furthermore, we have

$$\begin{aligned} H(Z|\hat{X}) + H(X|Z, \hat{X}) &\leq h_2(p_t)p_t H(X|Z=1, \hat{X}) + (1-p_t)H(X|Z=0, \hat{X}) \\ &\leq h_2(p_t) + p_t \log(|\mathcal{X}|) + (1-p_t) \log N_{\max}(t) \\ &= h_2(p_t) + p_t \log \left(\frac{\log |\mathcal{X}|}{N_{\max}(t)} \right) + \log(N_{\max}(t)). \end{aligned}$$

Plugging this back we get

$$H(X|\hat{X}) \leq h_2(p_t) + p_t \log \left(\frac{\log |\mathcal{X}|}{N_{\max}(t)} \right) + \log(N_{\max}(t)),$$

which implies

$$\begin{aligned} p_t &\geq \left(H(X|\hat{X}) - h_2(p_t) - \log(N_{\max}(t)) \right) / \log \left(\frac{\log |\mathcal{X}|}{N_{\max}(t)} \right) \\ &= \left(H(X|\hat{X}) - \cancel{H(X)} + \cancel{H(X)} - h_2(p_t) - \log(N_{\max}(t)) \right) / \log \left(\frac{\log |\mathcal{X}|}{N_{\max}(t)} \right) \\ &= \left(H(X) - \log(N_{\max}(t)) - I(X; \hat{X}) - h_2(p_t) \right) / \log \left(\frac{\log |\mathcal{X}|}{N_{\max}(t)} \right) \\ &= \left(\log(|\mathcal{X}|) - \log(N_{\max}(t)) - I(X; \hat{X}) - h_2(p_t) \right) / \log \left(\frac{\log |\mathcal{X}|}{N_{\max}(t)} \right) \\ &= 1 - \frac{I(X; \hat{X}) + h_2(p_t)}{\log(|\mathcal{X}|/N_{\max}(t))} \geq 1 - \frac{I(X; \hat{X}) + 1}{\log(|\mathcal{X}|/N_{\max}(t))} \end{aligned}$$

□

Chapter 2

Information measures for general distributions

We now extend the definition of the information measures beyond discrete distributions to more general spaces. First, we consider the case of continuous distributions in \mathbb{R}^d , and introduce the analogs of entropy, relative entropy, and mutual information. We study their properties and compare them with their discrete counterparts. Next, we define relative entropy and mutual information for general probability spaces, and establish two variational definitions. Finally, we introduce a class of information measures that generalize relative entropy, called the f -divergences, and study some of their properties.

2.1 Continuous Distributions

First we focus on the case of continuous distributions on $\mathcal{X} = \mathbb{R}^d$. That is, we focus on distributions P_X which admit a density, denoted by f_X , with respect to the Lebesgue measure on \mathcal{X} . This implies that for any measurable $E \subset \mathcal{X}$, we have $P_X = \int_{\mathcal{X}} \mathbf{1}_E(x) f_X(x) dx = \int_E f_X(x) dx$. For distributions (X, Y) with joint density f_{XY} , we also assume the existence of conditional densities $f_{Y|X}$ and $f_{X|Y}$ satisfying

$$P_{XY}(E) = \int_{\mathcal{X}} f_X(x) dx \int_{\mathcal{Y}} f_{Y|X}(y) \mathbf{1}_E(x, y) dy = \int_{\mathcal{Y}} f_Y(y) dy \int_{\mathcal{X}} f_{X|Y}(x) \mathbf{1}_E(x, y) dx.$$

For such distributions, we can define the continuous analogs of entropy, relative entropy, and mutual information, as well as their conditional variants.

Definition 36 (Differential Entropy). The differential entropy of a continuous random variable X , with density f_X , is defined as

$$h(X) \equiv h(f_X) = - \int_S f_X(x) \log(f_X(x)) dx,$$

where $S = \{x \in \mathcal{X} : f(x) > 0\}$ is the support of f_X . Similarly, conditional entropy of Y given X is defined as follows (where the integrals are over the appropriate supports):

$$h(Y|X) = - \int f_X(x) dx \int f_{Y|X}(y|x) \log(f_{Y|X}(y|x)) dy.$$

As in the discrete case, we can write the joint differential entropy $h(X, Y)$ as

$$h(X, Y) = h(X) - h(Y|X) = h(Y) - h(X|Y). \quad (2.1)$$

The definition of joint entropy in (2.1) implicitly assumes that at least one of the terms is not infinite. We now evaluate the differential entropy for some common distributions.

Example 37 (Uniform distribution). Suppose $\mathcal{X} = [a, b]$, and X is a the uniformly distributed random variable over \mathcal{X} with density $f_X(x) = 1/(b-a)$ for all $x \in \mathcal{X}$. Then, the differential entropy of X is equal to

$$h(X) = \int_{\mathcal{X}} \frac{1}{b-a} \log(b-a) dx = \log(b-a).$$

Thus, $h(X) = 0$ for $(b-a) = 1$, $h(X) < 0$ for $(b-a) < 1$, and $h(X) > 0$ for $(b-a) > 1$. This is in contrast with entropy defined for discrete distributions, which is always non-negative, and equal to 0 only for Dirac distributions.

Example 38 (Multivariate Gaussian). Consider a multivariate Gaussian random variable (X) over $\mathcal{X} = \mathbb{R}^d$, with mean μ and covariance matrix K . The density of this random variable is

$$f_X(x) = \frac{1}{\sqrt{(2\pi)^d |K|}} \exp \left(-\frac{1}{2} (x - \mu)^T K^{-1} (x - \mu) \right).$$

Then, the differential entropy of X is equal to

$$\begin{aligned} h(X) &= \log \left(\sqrt{(2\pi)^d |K|} \right) + \frac{1}{2} \int_{\mathcal{X}} (x - \mu)^T K^{-1} (x - \mu) f_X(x) dx \\ &= \frac{1}{2} \log ((2\pi)^d |K|) + \frac{1}{2} \mathbb{E} [(X - \mu)^T K^{-1} (X - \mu)] \\ &= \frac{1}{2} \log ((2\pi)^d |K|) + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \mathbb{E} [(X_i - \mu_i) K_{ij}^{-1} (X_j - \mu_j)] \\ &= \frac{1}{2} \log ((2\pi)^d |K|) + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d K_{ij}^{-1} \mathbb{E} [(X_i - \mu_i)(X_j - \mu_j)] \\ &= \frac{1}{2} \log ((2\pi)^d |K|) + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d K_{ij}^{-1} K_{ij} \\ &= \frac{1}{2} \log ((2\pi)^d |K|) + \frac{1}{2} \sum_{i=1}^d (I_d)_{ii} = \frac{1}{2} \log ((2\pi)^d |K|) + \frac{d}{2} \\ &= \frac{1}{2} \log ((2\pi e)^d |K|) \text{ nats/bits.} \end{aligned}$$

A simple consequence of this is that for univariate Gaussian with mean μ , and variance σ^2 , the differential entropy is equal to $(1/2) \log(2\pi e \sigma^2)$ nats/bits.

Remark 39. Note that in both the examples considered above, the differential entropy is “translation invariant”: that is, the differential entropy of X and $X + c$ for any $c \in \mathcal{X}$ is the same. This property is true generally for differential entropy, beyond the two examples above, as we will see later.

We now introduce the definition of relative entropy for continuous distributions.

Definition 40. Suppose P_X and P_Y are two continuous distributions on \mathcal{X} , with densities f_X and f_Y respectively. Then, the relative entropy between them is defined as (with S_X denoting the support of P_X):

$$D_{\text{kl}}(P_X \parallel P_Y) = \begin{cases} \int_{S_X} f_X(x) \log \left(\frac{f_X(x)}{f_Y(x)} \right) dx, & \text{if } \{f_Y = 0\} \subset \{f_X = 0\}, \\ +\infty, & \text{otherwise.} \end{cases}$$

Similarly, let P_{XY} (with density f_{XY}) and Q_{XY} (with density g_{XY}) denote two continuous joint distributions over $\mathcal{X} \times \mathcal{Y}$. Then the conditional relative entropy between $P_{Y|X}$ and $Q_{Y|X}$ is defined as

$$D_{\text{kl}}(P_{Y|X} \parallel Q_{Y|X} | P_X) = \int f_X(x) dx \int f_{Y|X}(y|x) \log \left(\frac{f_{Y|X}(y|x)}{g_{Y|X}(y|x)} \right) dy.$$

Example 41. The relative entropy between two univariate Gaussians, $P = N(\mu_1, \sigma_1^2)$ and $Q = N(\mu_2, \sigma_2^2)$ is equal to

$$\begin{aligned} D_{\text{kl}}(P \parallel Q) &= \mathbb{E}_P [\log(p(X)/q(X))] = \mathbb{E}_P \left[\log \left(\frac{\sigma_2}{\sigma_1} \right) + \frac{1}{2} \left(\frac{(X - \mu_2)^2}{\sigma_2^2} - \frac{(X - \mu_1)^2}{\sigma_1^2} \right) \right] \\ &= \log \left(\frac{\sigma_2}{\sigma_1} \right) + \frac{1}{2} \mathbb{E}_P \left[\left(\frac{(X - \mu_2)^2}{\sigma_2^2} - \frac{(X - \mu_1)^2}{\sigma_1^2} \right) \right] \\ &= \log \left(\frac{\sigma_2}{\sigma_1} \right) + \frac{1}{2} \left(\frac{\sigma_2^2 + (\mu_1 - \mu_2)^2}{\sigma_2^2} - 1 \right) = \log \left(\frac{\sigma_2}{\sigma_1} \right) + \frac{(\mu_1 - \mu_2)^2}{2}. \end{aligned}$$

Thus, if $\sigma_1 = \sigma_2$, the relative entropy is proportional to the difference in mean squared.

Example 42 (Gaussian-Cauchy). Note that the relative entropy can be infinite even if the distributions are absolutely continuous. For example, consider the relative entropy between P and Q , when P has a Cauchy distribution with density $f(x) = 1/(\pi(x^2 + 1))$, and Q has a Gaussian distribution with density $g(x) = (1/\sqrt{2\pi}) \exp(-x^2/2)$. Clearly, $P \ll Q$ and $Q \ll P$. However,

$$D_{\text{kl}}(P \parallel Q) = \mathbb{E}_P [\log(p(X)/q(X))] \asymp -h(P) + \mathbb{E}_P[X^2] = \infty,$$

since $h(P)$ can be shown to be finite, while the second moment of Cauchy distribution is infinite.

Finally, we can now introduce the definition of mutual information for continuous random variables.

Definition 43. The mutual information between two continuous random variables X and Y is defined as

$$I(X; Y) = \int_{S_{XY}} f_{XY}(x, y) \log \left(\frac{f_{XY}(x, y)}{f_X(x)f_Y(y)} \right) dx dy,$$

where $S_{XY} = \{(x, y) \in \mathcal{X} \times \mathcal{Y} : f_{XY}(x, y) > 0\}$ is the support of the joint density of (X, Y) . It is easy to check that $I(X; Y) = h(X) - h(X|Y) = h(Y) - h(Y|X)$.

2.1.1 Connections to discrete measures via quantization

Example 44. Let U denote the random variable with uniform distribution on $[0, a]$. For any $\Delta > 0$, let U_Δ denote the discrete quantization of U : that is, U_Δ takes values in the set $\{i\Delta : 1 \leq i \leq a/\Delta\}$. Then, we have the following relation:

$$H(U_\Delta) = \log(a/\Delta) = \log(a) - \log(\Delta) = h(U) - \log(\Delta).$$

Or in other words, for any $\Delta > 0$ (such that a/Δ is an integer), we have

$$H(U_\Delta) + \log(\Delta) = h(U).$$

We now show that the same relation between $H(\cdot)$ and $h(\cdot)$ holds more generally for continuous distributions.

Definition 45. Let X denote a real-valued random variable with continuous density f_X . For any $\Delta > 0$, partition $\mathcal{X} = \mathbb{R}$ into a (countable) grid of size Δ , consisting of sets

$$E_i = [i\Delta, (i+1)\Delta), \quad \text{for } -\infty \leq i \leq \infty.$$

Then, for any i , due to the mean-value theorem, there exists an $x_i \in E_i$, such that $f_X(x_i)\Delta = \int_{E_i} f_X(x)dx := p_i$. We define the Δ -quantized version of X , denoted by X_Δ , as

$$X_\Delta = x_i, \quad \text{with probability } p_i, \quad \text{for } i \in \mathbb{Z}.$$

Theorem 46. Consider the case of $\mathcal{X} = \mathbb{R}$, and let X be a continuous distribution with (a Riemann integrable) density function f_X . Then, we have the following:

$$\lim_{\Delta \downarrow 0} H(X_\Delta) + \log(\Delta) = h(X) = h(f_X),$$

where X_Δ is the Δ -quantized version of X (Definition 45).

Proof. The proof follows directly from the definition of discrete entropy. In particular,

$$\begin{aligned}
-H(X_\Delta) &= \sum_{i \in \mathbb{Z}} p_i \log p_i = \sum_{i \in \mathbb{Z}} f_X(x_i) \Delta (\log(f_X(x_i)) + \log(\Delta)) \\
&= \sum_{i \in \mathbb{Z}} f_X(x_i) \Delta \log(f_X(x_i)) + \log(\Delta) \sum_{i \in \mathbb{Z}} p_i \\
&= \sum_{i \in \mathbb{Z}} f_X(x_i) \Delta \log(f_X(x_i)) + \log(\Delta),
\end{aligned}$$

where the last equality simply uses the fact that $\sum_{i \in \mathbb{Z}} p_i = 1$. On rearranging the above relation, we get

$$H(X_\Delta) + \log \Delta = - \sum_{i \in \mathbb{Z}} f_X(x_i) \Delta \log(f_X(x_i)).$$

Since the term on the right is the Riemann sum for the integral $-\int_{\mathbb{R}} f_X(x) \log(f_X(x)) dx$ that defines the differential entropy of X , we get the required result by taking the limit $\Delta \downarrow 0$:

$$\lim_{\Delta \downarrow 0} H(X_\Delta) + \log \Delta = h(f_X).$$

□

Remark 47. One interpretation of this above result is in terms of the number of bits needed to learn a Δ -approximate version of a continuous random variable X . That is, to approximate a continuous random variable X , with an approximation error Δ , we roughly need $\log(1/\Delta) + h(f_X)$ bits. For example, going back to our uniform example, to represent a uniform $[0, a]$ random variable with up to Δ error, we need $\log(a/\Delta)$ bits (exactly).

Remark 48. Another interpretation of the above result is via the AEP for the discrete and continuous entropies. For concreteness, let X be a uniformly distributed over $\mathcal{X} = [0, a]$ random variable, and X_Δ is its Δ -quantization. Then, we have the following two statements:

- AEP for the differential entropy says that (with probability almost 1), the n i.i.d. realizations of X are concentrated in a volume of $\approx 2^{nh(X)}$ within the space \mathcal{X}^n .
- AEP for discrete entropy says that (with probability almost 1) there are $\approx 2^{nH(X_\Delta)}$ equiprobable sequences of length n in the quantized space \mathcal{X}_Δ^n .

Each point in \mathcal{X}_Δ^n denotes a unique cube in \mathcal{X}^n , of volume Δ^n . Hence, the total volume covered by the $2^{nH(X_\Delta)}$ sequences is $\approx \Delta^n 2^{nH(X_\Delta)}$. The result above says that the two volumes are approximately equal: that is,

$$\Delta^n 2^{nH(X_\Delta)} \approx 2^{nh(X)}, \quad \text{or} \quad \log(\Delta) + H(X_\Delta) \approx h(X).$$

Unlike the entropy, the behavior of relative entropy and mutual information is stable under quantization. In fact, through informal arguments, we can see that the continuous relative entropy (and thus mutual information as well) can be seen as the limits of their quantized versions, as $\Delta \rightarrow 0$.

2.1.2 Properties

We begin by noting the nonnegativity of relative entropy for continuous distributions.

Proposition 49. For two $\mathcal{X} = \mathbb{R}^d$ -valued distributions P and Q with densities f and g respectively, we have

$$D_{kl}(P \parallel Q) \equiv D_{kl}(f \parallel g) \geq 0,$$

with equality if and only if $f = g$ almost everywhere. As consequence, we also have for any two continuous random variables X and Y :

- $I(X; Y) \geq 0$, with equality if and only if $X \perp Y$.
- $h(X|Y) \leq h(X)$, with equality if and only if $X \perp Y$.

Proof. The proof of the nonnegativity of D_{kl} follows directly from the convexity of the mapping $\varphi(x) = x \log x$. First, note that if $P \not\ll Q$, then $D_{\text{kl}}(P \parallel Q) = \infty$, and there is nothing to prove. Hence, we can assume that $P \ll Q$, and $S_f = \{x : f(x) > 0\} \subset S_g = \{x : g(x) > 0\}$. Then, we have

$$\begin{aligned} D_{\text{kl}}(f \parallel g) &= \int_{S_f} g(x) \varphi\left(\frac{f(x)}{g(x)}\right) dx = \inf_{S_g} g(x) \varphi\left(\frac{f(x)}{g(x)}\right) dx \\ &\geq \varphi\left(\int_{S_g} g(x) \frac{f(x)}{g(x)} dx\right) = \varphi(1) = 0. \end{aligned}$$

Since the \geq above follows from Jensen's inequality, it holds with equality if and only if $f(x)/g(x)$ is a constant almost surely. Since both of them are densities, the constant must be 1 and thus the equality holds when $f = g$ almost surely. \square

We now look at some properties of differential entropy.

Proposition 50. *The differential entropy of continuous random variables satisfies the following properties:*

- (a) *Chain rule:* $h(X_1, \dots, X_n) = \sum_{i=1}^n h(X_i | X^{i-1})$.
- (b) *Translation invariance:* $h(X + c) = h(X)$ for all $c \in \mathcal{X}$.
- (c) *Effect of scaling:* if $\mathcal{X} = \mathbb{R}$, then $h(aX) = h(X) + \log(|a|)$. More generally, for vector valued X , we have $h(AX) = h(X) + \log(|\det(A)|)$, where A is an invertible matrix.

Proof. (a) Follows directly from the definition.

- (b) Let $Y = X + c$. Then, we have $f_Y(x) = f_X(x - c)$, and hence we have

$$h(Y) = \int_{c+S_X} f_X(x - c) \log(1/f_X(x - c)) dx = \int_{S_X} f_X(x) \log(1/f_X(x)) dx = h(X).$$

- (c) We prove the general multivariate case. Let $Y = AX$, and by the change of variable formula, we have

$$g(y) = \frac{1}{|A|} f(A^{-1}y).$$

Then, we have the following

$$\begin{aligned} h(AX) &= h(Y) = - \int g(y) \log(g(y)) dy = - \frac{1}{|A|} \int f(A^{-1}y) (\log(f(A^{-1}y)) - \log(|A|)) dy \\ &= \log(|A|) - \frac{1}{|A|} \int f(A^{-1}y) \log(f(A^{-1}y)) dy \\ &\stackrel{(i)}{=} \log(|A|) - \int f(x) \log(f(x)) dx = h(X) + \log(|A|). \end{aligned}$$

In the equality (i) above, we used the change of variable $x = A^{-1}y$, which implies $dx = |\det(A^{-1})| dy = (1/|A|) dy$. \square

Remark 51. Note that despite the scaling operation being a bijection, the differential entropy of the resulting variable (i.e., $Y = AX$) is different from the original (X). This is unlike the discrete entropy which did not change under such operations.

We end this section with a result about a characterization of Gaussian in terms of the maximum entropy achieving distribution among the class of all distributions with finite second moment.

Proposition 52. Let $\mathcal{P}_2(\mathbb{R})$ denote the class of continuous probability distributions on \mathbb{R} with zero mean and with second moment upper bounded by $b < \infty$. Then, the distribution from \mathcal{P}_2 with the largest differential entropy is $N(0, b)$.

Proof. Without loss of generality, we assume that $b = 1$.

Suppose P be any distribution in \mathcal{P}_2 , with density f , and Let ϕ denote the density of the standard normal random variable. Then, by the nonnegativity of relative entropy, we have

$$\begin{aligned} 0 &\leq D_{\text{kl}}(f \parallel \phi) = \int_{\mathbb{R}} f(x) (\log(f(x)) - \log(\phi(x))) dx \\ &= - \int_{\mathbb{R}} f(x) \log(\phi(x)) dx + \int_{\mathbb{R}} f(x) \log(f(x)) dx \\ &= - \int_{\mathbb{R}} \phi(x) \log(\phi(x)) dx - h(f) \\ &= h(\phi) - h(f). \end{aligned}$$

This completes the proof. \square

Example 53 (Capacity of Gaussian Channel). Suppose $\mathcal{X} = \mathbb{R}$, and consider a power-limited Gaussian channel, which takes in a \mathcal{X} -valued input X , and perturbs it with an additive Gaussian noise $Z \sim N(0, \sigma_N^2)$ that is independent of X . The output of the channel is denoted by $Y = X + Z$. Suppose we have an energy constraint $\mathbb{E}[X^2] \leq \sigma_P^2$. Then, the capacity of this channel is equal to

$$C = \sup_{f_X: \mathbb{E}[X^2] \leq \sigma_P^2} I(X; Y) = \frac{1}{2} \log \left(1 + \frac{\sigma_P^2}{\sigma_N^2} \right) = \frac{1}{2} \log(1 + \text{SNR}).$$

To see why this is true, note that for any arbitrary input X , we have

$$I(X; Y) = h(Y) - h(Y|X) = h(Y) - h(X + Z|X) = h(Y) - h(Z|X) = h(Y) - h(Z),$$

where the last equality uses the fact that $Z \perp X$. Hence, we have

$$I(X; Y) = h(Y) - \frac{1}{2} \log(2\pi e \sigma_N^2) \leq \frac{1}{2} \log(2\pi e \sigma_Y^2) - \frac{1}{2} \log(2\pi e \sigma_N^2),$$

where the inequality uses Proposition 52. Since $X \perp Z$, the variance of Y (denoted by σ_Y^2) is equal to $\sigma_X^2 + \sigma_N^2$. Furthermore, under the constraint, the maximum value of σ_X^2 is equal to σ_P^2 . Thus, we have

$$I(X; Y) \leq \frac{1}{2} \log \left(\frac{\sigma_N^2 + \sigma_P^2}{\sigma_N^2} \right) = \frac{1}{2} \log \left(1 + \frac{\sigma_P^2}{\sigma_N^2} \right).$$

Since X was arbitrary, this also implies that $C \leq (1/2) \log(1 + \sigma_P^2/\sigma_N^2)$. Finally, note that the inequality holds with an equality for the input distribution $f_X = N(0, \sigma_P^2)$.

In the next section, we illustrate some interesting consequences of the simple properties of continuous information measures we obtained above.

2.1.3 Applications

We now look at some simple applications of the properties of continuous information measures.

Maximum Entropy distributions. Staying with the theme of maximum entropy distributions, we show a general method for characterizing the distributions achieving the optimal entropy under different types of constraints.

Let \mathcal{F} denote a class of densities on $\mathcal{X} = \mathbb{R}$ satisfying m integral constraints:

- $f(x) \geq 0$, for all $x \in S$, with equality outside S .

- $\int_S f(x)dx = 1$.
- $\int_S f(x)r_i(x)dx = \alpha_i$, for $1 \leq i \leq m$.

Our next result characterizes the distribution from \mathcal{F} achieving the maximum entropy.

Proposition 54. *The density $f^* \in \mathcal{F}$ achieving the maximum entropy is of the form $f^*(x) = \exp(\lambda_0 + \sum_{i=1}^m \lambda_i r_i(x))$ for all $x \in S$, for some real-valued terms $\{\lambda_i : 0 \leq i \leq m\}$, chosen to ensure f^* satisfies the constraints.*

Proof. We will use a slightly informal “guess and verify” strategy to prove this result. In particular, note that entropy is a concave functional of the densities, and the domain is convex. Hence, we can write the Lagrangian as

$$L(f) = - \int f \log f + \lambda_0 \int f + \sum_{i=1}^m \int f r_i.$$

The proper way to proceed now is to take the functional derivative of J with respect to f . Instead, we present an informal argument. Suppose we write the integrals as approximate Riemann sums. That is, we consider a grid of width Δ partitioning the domain \mathcal{X} , and write

$$L(f) \approx - \sum_{j \in \mathbb{Z}} f(x_j) \log(f(x_j)) \Delta + \lambda_0 \sum_{i \in \mathbb{Z}} f(x_j) \Delta + \sum_{i=1}^m \sum_{j \in \mathbb{Z}} f(x_j) r_i(x_j) \Delta.$$

Now, we can consider $L(f)$ as a function of the variables $\{y_j = f(x_j) : j \in \mathbb{Z}\}$. So taking the (usual) derivative of $L(f)$ with $y_j = f(x_j)$, and setting it to 0, we get

$$\frac{dL(f)}{dy_j} = \Delta \left(-\log(f(x_j)) - 1 + \lambda_0 + \sum_{i=1}^m f(x_j) r_i(x_i) \right) = 0.$$

The above equation suggests a solution of the form

$$f(x) = \exp \left(-1 + \lambda_0 + \sum_{i=1}^m \lambda_i r_i(x) \right).$$

Now, constraints give us $(m+1)$ equations in $(m+1)$ unknowns $(\lambda_i)_{i=0}^m$, and we can solve this linear system to construct a candidate solution f^* of the above form, satisfying all the constraints.

Having informally justified the form of the candidate solution f^* , we now provide an information theoretic proof of its optimality. Let g denote any element of \mathcal{F} . Then, we have

$$h(g) = - \int_S g \log g = - \int_S g \log(g/f^*) - \int_S g \log f^* = -D_{\text{kl}}(g \parallel f^*) - \int_S g \log f^*.$$

Since f^* is supported on the entire S , the above operation is valid. Due to the nonnegativity of relative entropy, the above result implies that

$$h(g) \leq - \int_S g \log f^*.$$

If the integral on the right side were with respect to the density f^* , we would have obtained the required inequality. Nevertheless, we exploit the specific form of f^* to show the required inequality:

$$\begin{aligned} - \int_S g \log f^* &= - \int_S g \left(\lambda_0 + \sum_{i=1}^m \lambda_i r_i \right) = -\lambda_0 \int_S g - \sum_{i=1}^m \lambda_i \int_S g r_i \\ &\stackrel{(i)}{=} -\lambda_0 \int_S f^* - \sum_{i=1}^m \lambda_i \int_S f^* r_i \\ &= - \int_S f^* \log f^* = h(f^*). \end{aligned}$$

The equality (i) follows from the fact that both g and f^* are elements of \mathcal{F} , and thus they satisfy the required constraints. \square

Example 55. The previous result can be used to characterize the max-entropy distribution in various settings.

- $S = [a, b]$, and no other constraints. Then, the optimal distribution has the form $f^*(x) = e^{\lambda_0}$; that is, the max-ent distribution is the uniform distribution.
- Suppose $S = \mathbb{R}$, and the constraints are $\mathbb{E}[X] = 0$ and $\mathbb{E}[X^2] = \sigma^2$. Then, the optimal density has the form $f^*(x) = \exp(\lambda_0 + \lambda_1 x + \lambda_2 x^2)$. It is easy to verify that this is a Gaussian distribution with mean 0 and variance σ^2 .
- Suppose $S = [0, \infty)$, and the only constraint is $\mathbb{E}[X] = \mu$. Then, the optimal distribution is of the form $f^*(x) = \exp(\lambda_0 + \lambda_1 x)$, which turns out to be the exponential distribution with parameter μ . That is, $f^*(x) = (1/\mu)e^{-x/\mu}$.
- Suppose $S = [0, \infty)$, and the constraints are $\mathbb{E}[X] = \alpha_1$ and $\mathbb{E}[\log X] = \alpha_2$. Then, the optimal distribution has the form $f^*(x) = \exp(\lambda_0 + \lambda_1 x + \lambda_2 \log(x)) = x^{\lambda_2} \exp(\lambda_0 + \lambda_1 x)$. This is the family of Gamma distributions.
- For $S = \mathbb{R}^d$, and the constraint that $E[X_i] = 0$ for all $i \in [d]$, and $\mathbb{E}[X_i X_j] = K_{ij}$, where $K \equiv [K_{ij}]$ is a positive semi-definite matrix, a similar argument shows that the distribution achieving the maximum entropy is $N(0, K)$.

Determinant Inequalities. The properties of continuous information measures, especially for Gaussian distributions, can be used to obtain elementary proofs of several nontrivial matrix inequalities.

Proposition 56. *The following statements are true.*

- (a) *Suppose K is an $n \times n$ positive semi-definite matrix. Then, we have*

$$|K| \equiv |\det(K)| \leq \prod_{i=1}^n K_{ii},$$

with equality if and only if K is a diagonal matrix.

- (b) *For any two psd matrices K_1 and K_2 , and any $\lambda \in [0, 1]$, we have*

$$|\lambda K_1 + \bar{\lambda} K_2| \geq |K_1|^\lambda |K_2|^{\bar{\lambda}}.$$

Or, in other words, the log-determinant function is concave on the space of positive semi definite matrices.

Proof. (a) Given the psd matrix K , consider the Gaussian random vector $(X_1, \dots, X_n) \equiv X^n \sim N(0, K)$ on the domain $\mathcal{X} = \mathbb{R}^n$. Then, the chain rule for differential entropy implies that

$$h(X^n) = \sum_{i=1}^n h(X_i | X^{i-1}) \leq \sum_{i=1}^n h(X_i).$$

Since the marginals of X^n are also univariate Gaussians, with $X_i \sim N(0, K_{ii})$, we observe that

$$\frac{1}{2} \log((2\pi e)^n |K|) \leq \sum_{i=1}^n \frac{1}{2} \log(2\pi e |K_{ii}|) = \frac{n}{2} \log(2\pi e) + \frac{1}{2} \log\left(\prod_{i=1}^n K_{ii}\right).$$

On cancelling $(n/2) \log(2\pi e)$ from both sides, and exponentiating, we get the required inequality.

Furthermore, we know that the subadditivity of differential entropy holds with equality if and only if all the X_i 's are independent; or in other words, the matrix K is equal to $\text{diag}(K_{11}, \dots, K_{nn})$.

- (b) Let $X \sim N(0, K_1)$ and $Y \sim N(0, K_2)$ denote two independent multivariate Gaussian random variables on $\mathcal{X} = \mathbb{R}^n$. For a fixed $\lambda \in [0, 1]$, introduce a Bernoulli random variable $\Lambda \sim \text{Bernoulli}(\lambda)$ independent of X and Y , and use it to define the mixture $Z = \Lambda X + \bar{\Lambda} Y$.

We first observe that the covariance of Z is equal to $K_\lambda := \lambda K_1 + \bar{\lambda} K_2$. To see this, note that since $\mathbb{E}[Z] = 0$, we have

$$\begin{aligned} \text{Cov}(Z) &= \mathbb{E}[ZZ^T] = \mathbb{P}(\Lambda = 0)\mathbb{E}[YY^T|\Lambda = 0] + \mathbb{P}(\Lambda = 1)\mathbb{E}[XX^T|\Lambda = 1] \\ &= \bar{\lambda}\mathbb{E}[YY^T] + \lambda\mathbb{E}[XX^T] \\ &= \bar{\lambda}K_2 + \lambda K_1 = K_\lambda. \end{aligned} \tag{2.2}$$

Now, we use the fact that conditioning reduces entropy, to observe

$$h(Z|\Lambda) \leq h(Z) \leq h(N(0, K_\lambda)) = \frac{1}{2} \log((2\pi e)^n |K_\lambda|).$$

Finally, we calculate the conditional entropy of Z given Λ :

$$\begin{aligned} h(Z|\Lambda) &= \lambda h(Z|\Lambda = 1) + \bar{\lambda} h(Z|\Lambda = 0) \\ &= \lambda h(X) + \bar{\lambda} h(Y) \\ &= \frac{\lambda}{2} \log((2\pi e)^n |K_1|) + \frac{\bar{\lambda}}{2} \log((2\pi e)^n |K_2|). \end{aligned} \tag{2.3}$$

Now, by combining (2.2) with (2.3), we get the required statement that

$$\lambda \log(|K_1|) + \bar{\lambda} \log(|K_2|) \leq \log(|K_\lambda|).$$

□

Estimation error. For the case of discrete distributions, Fano's inequality gave us a lower bound on the probability of error in a hypothesis testing problem in terms of the entropy of the observations. We now present two analogous results for the problem of estimation.

Proposition 57. *Consider the Markov chain $X \rightarrow Y \rightarrow \hat{X}$, where \hat{X} is an estimate of the unknown random variable X . Then, we have*

$$\mathbb{E}[(X - \hat{X}(Y))^2] \geq \frac{1}{2\pi e} 2^{h(X|Y)}.$$

In the special case where the side information is not available, the above inequality reduces to

$$\mathbb{E}[(X - \hat{X})^2] \geq \frac{1}{2\pi e} 2^{h(X)}.$$

Proof. To prove this result, we begin with the observation that

$$\begin{aligned} \mathbb{E}[(X - \hat{X}(Y))^2] &\geq \mathbb{E}[(X - E[X|Y])^2] = \mathbb{E}_Y [\mathbb{E}_{X|Y} [(X - E[X|Y])^2|Y]] \\ &= \mathbb{E}_Y [\mathbb{V}(X|Y)]. \end{aligned} \tag{2.4}$$

Now, observe that

$$\begin{aligned} h(X|Y) &= - \int_{\mathcal{Y}} f(y) dy \int_{\mathcal{X}} f(x|y) \log(f(x|y)) dx \leq \int_{\mathcal{Y}} f(y) \frac{1}{2} \log(2\pi e \mathbb{V}(X|Y = y)) \\ &\stackrel{(i)}{\leq} \frac{1}{2} \log(2\pi e \mathbb{E}_Y [\mathbb{V}(X|Y)]), \end{aligned}$$

where (i) follows from an application of Jensen's inequality, along with the concavity of $x \mapsto \log(x)$. On rearranging the above inequality, we get

$$\frac{1}{2\pi e} 2^{2h(X|Y)} \leq \mathbb{E}_Y [\mathbb{V}(X|Y)]. \tag{2.5}$$

Together, (2.4) and (2.5) imply the required result. □

Bounding the entropy of discrete distribution. We end with one final application of Proposition 52. For the case of discrete distributions with finite support, we know that the entropy is upper bounded by $\log(|\mathcal{X}|)$. However, this bound is not applicable when the support of a discrete random variable is countable. We now show how to use the max entropy continuous distribution to obtain a non-vacuous upper bound on the entropy of a discrete distribution with countable support.

Proposition 58. *Let $\mathcal{X} = \{x_i : i \in \mathbb{N}\}$ denote a countable alphabet, and let X be an \mathcal{X} -valued random variable with $\mathbb{P}(X = x_i) = p_i$ for $i \in \mathbb{N}$. Then, we have*

$$H(X) = H(p_1, p_2, \dots) \leq \frac{1}{2} \log \left((2\pi e) \times \left(\sum_{i=1}^{\infty} p_i i^2 - \left(\sum_{i=1}^{\infty} i p_i \right)^2 + \frac{1}{12} \right) \right).$$

Proof. Introduce a new random variable Y , such that $\mathbb{P}(Y = i) = p_i$. Note that $H(Y) = H(X)$, since the discrete entropy is independent of the actual values taken by the random variable. Now, let U denote a Uniform $([0, 1])$ random variable, and use it to define $Z = Y + U$. We can verify that Z is a continuous random variable with a density $f_Z(x) = p_i$, where $i = \lfloor x \rfloor$.

Now, the discrete entropy of Y is equal to

$$\begin{aligned} H(X) = H(Y) &= - \sum_{i \in \mathbb{N}} p_i \log p_i = \sum_{i \in \mathbb{N}} \int_i^{i+1} f_Z(x) \log(f_Z(x)) dx \\ &= - \int_{\mathbb{R}} f_Z(x) \log(f_Z(x)) dx \\ &= h(Z). \end{aligned}$$

Note that since X and U are independent, the variance of Z is equal to $\mathbb{V}(Y) + \mathbb{V}(U)$. Thus the differential entropy of $h(Z)$ is upper bounded by that of a zero mean Gaussian with the same variance. Hence, we have

$$\begin{aligned} H(X) = h(Z) &\leq \frac{1}{2} \log((2\pi e)\mathbb{V}(Z)), \quad \text{where} \\ \mathbb{V}(Z) = \mathbb{V}(Y) + \mathbb{V}(U) &= \sum_i i^2 p_i - \left(\sum_i i p_i \right)^2 + \frac{1}{12}. \end{aligned}$$

□

2.2 General Distributions*

In this section, we present the general definition of relative entropy. To state the definition, we need to recall the Radon-Nikodym theorem.

Fact 59. *Consider a measurable space $(\mathcal{X}, \mathcal{F})$ with two σ -finite measures P and Q . Suppose that P is absolutely continuous w.r.t. Q , denoted by $P \ll Q$, which means that $Q(E) = 0 \Rightarrow P(E) = 0$, for any $E \in \mathcal{F}$. Then, there exists a measurable function $f : \mathcal{X} \rightarrow [0, \infty)$, such that*

$$P(E) = \int_E f dQ, \quad \text{for all } E \in \mathcal{F}.$$

The (not necessarily unique) function f is called the Radon-Nikodym derivative of P w.r.t. Q , and is also denoted as $\frac{dP}{dQ}$. If $P \ll Q$ and $Q \ll P$, we have

$$\frac{dP}{dQ} = \left(\frac{dQ}{dP} \right)^{-1}.$$

Finally, if $P \ll Q$ and $Q \ll R$, then we have

$$\frac{dP}{dR} = \frac{dP}{dQ} \times \frac{dQ}{dR}.$$

Since we deal with probability measures, that are finite and hence σ -finite, the only condition required for the existence of the Radon-Nikodym derivative is the absolute continuity. Using this, we have the following general definition of relative entropy.

Definition 60. For any two distributions P and Q , defined on a common measurable space $(\mathcal{X}, \mathcal{F})$, the relative entropy between P and Q is defined as

$$D_{\text{kl}}(P \parallel Q) := \begin{cases} \mathbb{E}_Q \left[\frac{dP}{dQ}(X) \log \frac{dP}{dQ}(X) \right], & \text{if } P \ll Q, \\ +\infty, & \text{if } P \not\ll Q. \end{cases}$$

Fact 61. An equivalent definition of the relative entropy between P and Q , is

$$D_{\text{kl}}(P \parallel Q) := \begin{cases} \mathbb{E}_P \left[\log \frac{dP}{dQ}(X) \right], & \text{if } P \ll Q, \\ +\infty, & \text{if } P \not\ll Q. \end{cases}$$

See Lemma 2.4 of Polyanskiy and Wu for a proof.

Definition 62. Suppose $(\mathcal{X}, \mathcal{F})$ and $(\mathcal{Y}, \mathcal{G})$ denote two measurable spaces. Then, a Markov kernel from \mathcal{X} to \mathcal{Y} is a mapping $K : \mathcal{G} \times \mathcal{X} \rightarrow [0, 1]$, such that

- For every $x \in \mathcal{X}$, the mapping $K(\cdot, x) : \mathcal{G} \rightarrow [0, 1]$ is a probability measure on $(\mathcal{Y}, \mathcal{G})$,
- For every $E \in \mathcal{G}$, the mapping $K(E, \cdot) : \mathcal{X} \rightarrow [0, 1]$ is $(\mathcal{F}, \mathbb{B}_{[0,1]})$ -measurable.

Remark 63. The Markov kernel can be interpreted as a random mapping from \mathcal{X} to a probability measure on $(\mathcal{Y}, \mathcal{G})$. Hence, we will often denote it with $P_{Y|X}$ — that is, it defines a probability distribution of Y for every realization of X . For the special case of finite \mathcal{X} and \mathcal{Y} , the Markov kernel $K \equiv P_{Y|X}$ is simply the transition probability matrix of size $|\mathcal{X}| \times |\mathcal{Y}|$.

Fact 64 (Disintegration Theorem). *Suppose P_{XY} is a joint distribution on $\mathcal{X} \times \mathcal{Y}$, and \mathcal{Y} is standard Borel (complete separable metric space with the Borel sigma-algebra). Then, there exists a Markov kernel K , such that for any measurable $E \subset \mathcal{X} \times \mathcal{Y}$, we have*

$$P_{XY}(E) = \int_{\mathcal{X}} P_X(dx) K(E^x|x), \quad \text{for } E^x := \{y \in \mathcal{Y} : (x, y) \in E\}.$$

We can now define the general version of conditional relative entropy, using the Markov kernel.

Definition 65. Suppose X is an \mathcal{X} -valued, and Y is a \mathcal{Y} -valued random variable. Let P_{XY} and Q_{XY} denote two joint distributions of (X, Y) . Then, if \mathcal{Y} is standard Borel (or ‘nice’ in the language used by Durrett), then the conditional relative entropy between $P_{Y|X}$ and $Q_{Y|X}$ given P_X is defined as

$$D_{\text{kl}}(P_{Y|X} \parallel Q_{Y|X} | P_X) = \mathbb{E}_{P_X} [D_{\text{kl}}(P_{Y|X}(\cdot, X) \parallel Q_{Y|X}(\cdot, X))].$$

Remark 66. Having defined relative entropy, and conditional relative entropy for general distributions, we can immediately use them to define mutual information and conditional mutual information.

Remark 67. All the properties that we proved for the relative entropy of discrete or continuous random variables are also valid for the general setting considered here. These properties include the nonnegativity, convexity, and the DPI.

The main summary of this section is that we can define the relative entropy can be defined for distributions on very general observation spaces (such as on manifolds, or on function spaces). In particular, the differential entropy of a random variable X can be recovered as the relative entropy of P_X and an appropriate invariant measure, such as the Lebesgue measure for $\mathcal{X} = \mathbb{R}^d$.

2.2.1 Variational Definition I: Gelfand-Yaglom-Peres

In this section, and the next, we present two *variational definitions* of relative entropy for general probability distributions — that is, we define relative entropy as the solution of an optimization problem. There are several benefits of such a representation:

- several properties, such as convexity and lower semi-continuity, can be easily inferred,
- such representations easily allow us to get upper or lower bounds by considering specific values of the objective function being optimized.

Example 68. As a simple example, consider the ℓ_1 norm of a vector: $\|\mathbf{x}\|_1 = \sum_{i=1}^d |x_i|$. It is easy to check that it can also be defined as follows:

$$\|\mathbf{x}\|_1 = \sup_{\mathbf{y} \in \mathbb{R}^d: \|\mathbf{y}\|_\infty \leq 1} \sum_{i=1}^d x_i y_i.$$

The above definition immediately implies that the map $\mathbf{x} \mapsto \|\mathbf{x}\|_1$ is convex: since it is the supremum of linear functions. Furthermore, we also immediately observe that it is lower semi-continuous: since it is the supremum of continuous functions. Furthermore, by choosing any specific value of \mathbf{y} , we get a lower bound on the ℓ_1 norm of \mathbf{x} .

We now present the first variational definition of relative entropy, which says that for general probability spaces, the relative entropy between two distributions is equal to the supremum of the relative entropy between quantized versions of the two distributions. In other words, for most purposes, analyzing the properties of relative entropy (and derived quantities, such as mutual information) for discrete distributions with finite support is without loss of generality.

Theorem 69. Let $(\mathcal{X}, \mathcal{F})$ denote a measurable space, and let P and Q denote two probability measures on this space. Let $\mathcal{E} = \{E_1, E_2, \dots, E_n\}$ denote a finite disjoint partition of \mathcal{X} , consisting of elements of \mathcal{F} . Then, we have

$$D_{kl}(P, Q) = \sup_{\mathcal{E}} \sum_{E \in \mathcal{E}} P(E) \log \left(\frac{P(E)}{Q(E)} \right) = \sup_{\mathcal{E}} D_{kl}(P_{\mathcal{E}} \parallel Q_{\mathcal{E}}),$$

where we have used $P_{\mathcal{E}}$ and $Q_{\mathcal{E}}$ to denote the quantized versions of P and Q over the partition \mathcal{E} .

Proof. We will prove the equality by showing that both \leq and \geq simultaneously hold. The lower bound is an easy consequence of the DPI for relative entropy. In particular, let \mathcal{E} denote a partition of \mathcal{X} with m elements $\{E_1, \dots, E_m\}$. Let $f : \mathcal{X} \rightarrow [m]$ be a function defined as $f(x) = \sum_{i=1}^m i \mathbf{1}_{x \in E_i}$, and let $Y = f(X)$. Then, we have

$$D_{kl}(P \parallel Q) \stackrel{(i)}{\geq} D_{kl}(P_Y \parallel Q_Y) = D_{kl}(P_{X, \mathcal{E}} \parallel Q_{X, \mathcal{E}}),$$

where (i) follows from the DPI for relative entropy. Since \mathcal{E} above was arbitrary, we can take a supremum over all such finite partitions to get the lower bound.

Proving the other direction is more involved, and we proceed in the following steps:

Step 1: We begin by noting that without loss of generality, we can assume $P \ll Q$. Because, if $P \not\ll Q$, then both sides of the required equality are infinite. In particular, if $P \not\ll Q$, then there exists a measurable set E , such that $Q(E) = 0$ but $P(E) > 0$. Then, define $\mathcal{E} = \{E, E^c\}$, and observe that $D_{kl}(P_{\mathcal{E}} \parallel Q_{\mathcal{E}}) = \infty$.

Step 2: Since $P \ll Q$, there exists a measurable Radon-Nikodym derivative $X = dP/dQ$. Since $D_{kl}(P \parallel Q) = \mathbb{E}_Q[\varphi(X)]$, for real valued sequence $c_n \rightarrow \infty$, we have $D_{kl}(P \parallel Q) = \lim_{c \rightarrow \infty} \mathbb{E}_Q[\varphi(X) \mathbf{1}_{X \leq c}]$ by the monotone convergence theorem (MCT).

Step 3: Fix a $c > 0$ and $\epsilon > 0$, and let the integer n denote c/ϵ . Construct a partition $\mathcal{E} = \{E_0, \dots, E_n\}$, where the sets E_0 through E_n are defined as

$$E_j = \{\omega : j\epsilon \leq X(\omega) \leq (j+1)\epsilon\}, \text{ for } j = 0, \dots, n-1, \\ \text{and } E_n = \{\omega : X(\omega) \geq c\}.$$

Using this partition, define a discrete approximation of X , as

$$Y_n = \sum_{j=0}^{n-1} j\epsilon \mathbf{1}_{E_j}.$$

Let $X_c = X \mathbf{1}_{X \leq c}$, and note that $Y_n \leq X_c$ and $|Y_n - X_c| \leq \epsilon$. Now, note that for a fixed $c < \infty$, the function φ restricted to the domain $[0, c]$ is uniformly continuous. Hence, there exists a $\delta \equiv \delta(c, \epsilon)$, such that we have $|\varphi(Y_n) - \varphi(X_c)| \leq \delta$ almost surely. Furthermore, for a fixed c , the term δ goes to 0 as $\epsilon \rightarrow 0$.

The above uniform continuity result implies that

$$\mathbb{E}_Q[\varphi(X_c)] - \delta \leq \mathbb{E}_Q[\varphi(Y_n)] = \sum_{j=0}^{n-1} Q(E_j) \varphi(j\epsilon). \quad (2.6)$$

Now, we observe that

$$\mathbb{E}_Q[j\epsilon] \leq \mathbb{E}_Q[X \mathbf{1}_{E_j}] = P(E_j) \leq \mathbb{E}_Q[(j+1)\epsilon],$$

which implies that

$$j\epsilon \leq \frac{P(E_j)}{Q(E_j)} \leq (j+1)\epsilon \quad \text{or} \quad \left| \varphi\left(\frac{P(E_j)}{Q(E_j)}\right) - \varphi(j\epsilon) \right| \leq \delta.$$

Plugging this back into (2.6), we get

$$\mathbb{E}_Q[\varphi(X_c)] - \delta \leq \mathbb{E}_Q[\varphi(Y_n)] = \sum_{j=0}^{n-1} Q(E_j) \varphi\left(\frac{P(E_j)}{Q(E_j)}\right) + \delta = D_{\text{kl}}(P_{\mathcal{E}} \parallel Q_{\mathcal{E}}) + \delta.$$

To complete the proof, we take $\epsilon \rightarrow 0$ for a fixed c , and then take $c \rightarrow \infty$. □

We record an immediate consequence of the above result.

Corollary 70. *For any $\epsilon > 0$, there exists a finite partition of \mathcal{X} , such that*

$$D_{\text{kl}}(P, Q) - \epsilon \leq D_{\text{kl}}(P_{\mathcal{E}}, Q_{\mathcal{E}}).$$

2.2.2 Variational Definition II: Donsker-Varadhan

We now present a functional variational representation of relative entropy, due to Donsker and Varadhan.

Theorem 71. *Consider a measurable space $(\mathcal{X}, \mathcal{F})$, with two probability measures P and Q . Suppose $P \ll Q$, and let \mathcal{C}_Q denote the set of measurable functions $f : \mathcal{X} \rightarrow \mathbb{R}$, such that $\mathbb{E}_Q[e^f(X)] < \infty$. Then, we have*

$$D_{\text{kl}}(P \parallel Q) = \sup_{f \in \mathcal{C}_Q} \mathbb{E}_P[f(X)] - \log \left(\mathbb{E}_Q[e^{f(X)}] \right).$$

Proof. First, note that we can restrict our attention to $P \ll Q$. For otherwise, if $P \not\ll Q$, then there must exist an E such that $Q(E) = 0$, but $P(E) > 0$. Then, define a function $f_c = c \mathbf{1}_E$, and note that it lies in \mathcal{C}_Q for all values of $c \in \mathbb{R}$. Then, we have

$$\sup_{f \in \mathcal{C}_Q} \mathbb{E}_P[f(X)] - \log \mathbb{E}_Q[e^{f(X)}] \geq \sup_{c > 0} \mathbb{E}_P[f_c(X)] - \log \mathbb{E}_Q[e^{f_c(X)}] = \sup_{c > 0} cP(E) = \infty.$$

Hence, we can assume that $P \ll Q$ for the rest of the proof. As in the previous case, we prove the result in two steps to show that both the \leq and \geq hold. For any $f \in \mathcal{C}_Q$, define the *tilted* distribution Q^f , such that for any measurable E , we have

$$Q^f(E) = \mathbb{E}_Q \left[e^{f(X) - Z_f} \mathbf{1}_E(X) \right], \quad \text{where} \quad Z_f = \log \left(\mathbb{E}_Q \left[e^{f(X)} \right] \right).$$

Now, observe that $Q^f \ll Q$, and

$$\log \left(\frac{dQ^f}{dQ} \right) = f(X) - Z_f,$$

which implies that

$$\mathbb{E}_P \left[\log \left(\frac{dQ^f}{dQ} \right) \right] = \mathbb{E}_P[f(X)] - Z_f.$$

Next, we note that we also have $Q \ll Q^f$. To see why this is true, suppose there exists an E such that $Q^f(E) = 0$, but $Q(E) > 0$. Define $A_0 = \{e^{f(X) - Z_f} \geq 1\} \cap E$, and for $i \geq 1$, define $A_i = \{e^{f(X) - Z_f} \in (1/(i+1), 1/i]\} \cap E$. Then, we have $A_i \cap A_j = \emptyset$ for $i \neq j$, and furthermore, $E = \cup_{i=0}^{\infty} A_i$. Due to the countable additivity, we have $Q(E) = \sum_{i=0}^{\infty} Q(A_i)$, which means that there must exist an n , such that $Q(A_n) > 0$. This implies that

$$0 = Q^f(E) = \sum_{i=0}^{\infty} Q^f(A_i) \geq \sum_{i=0}^{\infty} \frac{Q(A_i)}{i+1} \geq \frac{Q(A_n)}{n+1} > 0,$$

which is a contradiction. Hence, we have proved that $Q \ll Q^f$, which in turn implies that $P \ll Q^f$, and hence dP/dQ^f is well defined

Now, let G denote the set $\{dP/dQ > 0\}$ and note that $\mathbb{E}_P[\log(dQ^f/dQ)] = \mathbb{E}_P[\log(dQ^f/dQ)\mathbf{1}_G]$. Introduce the probability measures \tilde{Q} and \tilde{Q}^f , such that $\tilde{Q}(E) = Q(E \cap G)$, and $\tilde{Q}^f(E) = Q^f(E \cap G)$. Then, we can write

$$\begin{aligned} \mathbb{E}_P \left[\log \left(\frac{dQ^f}{dQ} \right) \right] &= \mathbb{E}_P \left[\log \left(\frac{dQ^f}{dQ} \right) \mathbf{1}_G \right] = \mathbb{E}_P \left[\log \left(\frac{d\tilde{Q}^f}{d\tilde{Q}} \frac{d\tilde{Q}}{dP} \right) \mathbf{1}_G \right] \\ &= \mathbb{E}_P \left[\log \left(\frac{d\tilde{Q}}{dP} \right) \mathbf{1}_G \right] - \mathbb{E}_P \left[\log \left(\frac{d\tilde{Q}^f}{d\tilde{Q}} \right) \mathbf{1}_G \right] \\ &= \mathbb{E}_P \left[\log \left(\frac{dP}{dQ} \right) \right] - \mathbb{E}_P \left[\log \left(\frac{dP}{dQ^f} \right) \right] \\ &= D_{\text{kl}}(P \parallel Q) - D_{\text{kl}}(P \parallel Q^f) \leq D_{\text{kl}}(P \parallel Q). \end{aligned}$$

Since f was an arbitrary element of \mathcal{C}_Q , this completes the proof of one side of the inequality.

To show the other direction, we will rely on Theorem 69. Now consider the case when $P \ll Q$. Then, consider any partition \mathcal{E} of the domain, and define f as

$$f = \sum_{E \in \mathcal{E}} \log \left(\frac{P(E)}{Q(E)} \right) \mathbf{1}_E.$$

For this function, the objective function is

$$\begin{aligned} \mathbb{E}_P[f(X)] - \log \mathbb{E}_Q[e^{f(X)}] &= \sum_E P(E) \log \left(\frac{P(E)}{Q(E)} \right) - \log \left(\sum_{E \in \mathcal{E}} Q(E) \frac{P(E)}{Q(E)} \right) \\ &= D_{\text{kl}}(P_{\mathcal{E}} \parallel Q_{\mathcal{E}}) \end{aligned}$$

For all choices of the partition \mathcal{E} , the corresponding f is still a simple function, and hence, it also belongs to \mathcal{C}_Q . Denoting the class of simple functions by \mathcal{S} , we then obtain

$$\sup_{f \in \mathcal{C}_Q} \mathbb{E}_P[f(X)] - \log \mathbb{E}_Q[e^{f(X)}] \geq \sup_{f \in \mathcal{S}} \mathbb{E}_P[f(X)] - \log \mathbb{E}_Q[e^{f(X)}] \geq \sup_{\mathcal{E}} D_{\text{kl}}(P_{\mathcal{E}}, Q_{\mathcal{E}}).$$

The last term is precisely the definition of $D_{\text{kl}}(P \parallel Q)$ from Theorem 69. \square

Remark 72. A simple consequence of the above result is that if $D_{\text{kl}}(P \parallel Q)$ is small, then we should expect $\mathbb{E}_P[f(X)]$ to be well approximated by $\mathbb{E}_Q[f(X)]$. To see why, note that $e^x \approx 1 + x$ and $\log(1 + x) \approx x$ for small enough x . Thus choosing any function $f \in \mathcal{C}_Q$, we have

$$D_{\text{kl}}(P \parallel Q) \geq \mathbb{E}_P[f(X)] - \log \left(\mathbb{E}_Q[e^{f(X)}] \right) \approx \mathbb{E}_P[f(X)] - \log(1 + \mathbb{E}_Q[f(X)]) \approx \mathbb{E}_P[f(X)] - \mathbb{E}_Q[f(X)].$$

Thus, a small value of $D_{\text{kl}}(P \parallel Q)$ implies that the expected values of $f(X)$ under P and Q must be close to each other.

Since mutual information between (X, Y) is the relative entropy between the joint distribution and the product of marginals, we have the following corollary.

Corollary 73. For $(X, Y) \sim P_{XY}$ on $\mathcal{X} \times \mathcal{Y}$, let \mathcal{C} denote the class of functions $\{f : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R} : e^f \in L^1(P_X \times P_Y)\}$. Then, we have

$$I(X; Y) = \sup_{f \in \mathcal{C}} \mathbb{E}_{P_{XY}}[f(X, Y)] - \log \mathbb{E}_{P_X \times P_Y} \left[e^{f(X, Y)} \right].$$

Application: Generalization bound in machine learning. In statistical learning theory, we are usually given a training dataset $S = (Z_1, \dots, Z_n) \in \mathcal{Z}^n$ consisting on n i.i.d. training points drawn from a distribution P . A learning algorithm, \mathcal{A} , is a channel (or a Markov kernel) from \mathcal{Z}^n to a ‘hypothesis class’ \mathcal{H} . Given a loss function $\ell : \mathcal{H} \times \mathcal{Z} \rightarrow \mathbb{R}$, we can define the population and empirical risk of a hypothesis $h \in \mathcal{H}$ as

$$L_P(h) = \mathbb{E}_{Z \sim P}[\ell(h, Z)], \quad \text{and} \quad L_S(h) = \frac{1}{n} \sum_{Z_i \in S} \ell(h, Z_i).$$

Now, let H denote the possibly random output of a learning algorithm \mathcal{A} on a training set S . Then, the generalization error of this algorithm is the expected difference between the test and training risk of H :

$$\text{gen}(P, P_{H|S}) = \mathbb{E}[L_P(H) - L_S(H)] = \mathbb{E}[L_{S'}(H) - L_S(H)],$$

where S' is an independent copy of S . Now, we can state the following bound over the generalization error.

Proposition 74 (Xu and Raginsky, 2016). Suppose the loss function is such that $\ell(h, Z)$ is σ^2 -subGaussian for all $h \in \mathcal{H}$. Then, we have

$$\text{gen}(P, P_{H|S}) \leq \sqrt{\frac{2\sigma^2 I(S; H)}{n}}.$$

Proof. To prove this result, we note the following:

$$I(S; H) = \sup_f \mathbb{E}_{P_{SH}}[f(S, H)] - \log \mathbb{E}_{P_S \times P_H} \left[e^{f(S, H)} \right]. \quad (2.7)$$

To get the bound, we will select a specific $f(s, h) = \frac{1}{n} \sum_{z_i \in S} \ell(h, z_i)$, and note that f is (σ^2/n) -subGaussian. Furthermore, note that the generalization error is equal to

$$\text{gen}(P, P_{H|S}) = \mathbb{E}_{P_{SH}}[f(S, H)] - \mathbb{E}_{P_S \times P_H}[f(S', H')].$$

For any $\lambda \in \mathbb{R}$, plugging λf in (2.7), we get

$$\begin{aligned} I(S; H) &\geq \mathbb{E}_{P_{SH}}[\lambda f(S, H)] - \log \left(\mathbb{E}_{P_S \times P_H} \left[e^{\lambda f(S', H')} \right] \right) \\ &\geq \mathbb{E}_{P_{SH}}[\lambda f(S, H)] - \log \left(\exp \left(\frac{\lambda^2 \sigma^2}{2n} + \lambda \mathbb{E}_{P_S \times P_H}[f(S', H')] \right) \right) \\ &= \mathbb{E}_{P_{SH}}[\lambda f(S, H)] - \mathbb{E}_{P_S \times P_H}[\lambda f(S', H')] - \frac{\lambda^2 \sigma^2}{2n} \\ &= \lambda \text{gen}(P, P_{H|S}) - \frac{\lambda^2 \sigma^2}{2n}. \end{aligned}$$

On optimizing for λ , we get

$$I(S; H) \geq \sup_{\lambda} \lambda \operatorname{gen}(P, P_{H|S}) - \frac{\lambda^2 \sigma^2}{2n} = \frac{n \operatorname{gen}^2(P, P_{H|S})}{2\sigma^2}.$$

On rearranging, we get the required

$$\operatorname{gen}(P, P_{H|S}) \leq \sqrt{\frac{2\sigma^2 I(S; H)}{n}}.$$

□

Now, suppose h^* denotes the element of \mathcal{H} that achieves the minimum expected loss. How does $\mathbb{E}[L_P(H)]$ compare with $L_P(h^*)$? To analyze this, note that

$$\mathbb{E}[L_P(H)] \leq \operatorname{gen}(P, P_{H|S}) + \mathbb{E}_{P_{SH}} [L_S(H)].$$

Hence, on subtracting $L_S(h^*)$ on both sides, and using Proposition 74, we get

$$\mathbb{E}[L_P(H)] - L_P(h^*) \leq \sqrt{\frac{2\sigma^2 I(S; H)}{n}} + \mathbb{E}_{P_{SH}} [L_S(H) - L_S(h^*)].$$

Consider the following two cases:

- If $|\mathcal{H}| < \infty$, then we can upper bound $I(S; H)$ with $H(H) \leq \log(|\mathcal{H}|)$.
- More generally, suppose we consider an appropriate ϵ -covering set of \mathcal{H} , denoted by $|\mathcal{H}_\epsilon|$. Then, we can bound $I(S; H)$ with $\log(|\mathcal{H}_\epsilon|)$, which recovers the metric entropy bound.

Note that if H is selected by an empirical risk minimization (ERM) method, the second term on the RHS is upper bounded by 0.

2.3 f -divergences

Definition 75. Suppose $f : (0, \infty) \rightarrow \mathbb{R}$ is a convex function with $f(1) = 0$, and $f(0) := \lim_{x \rightarrow 0} f(x)$. Then, the f -divergence between two distributions P and Q , with $P \ll Q$ is defined as

$$D_f(P \parallel Q) := \mathbb{E}_Q \left[f \left(\frac{dP}{dQ}(X) \right) \right], \quad \text{if } P \ll Q.$$

For general P and Q , let μ denote a common dominating measure (such as $P + Q$), and $q = dQ/d\mu$ and $p = dP/d\mu$. Then, the f -divergence between P and Q is defined as

$$D_f(P \parallel Q) = \int_{q>0} q(x) f \left(\frac{p(x)}{q(x)} \right) d\mu + \mathbb{P}(q=0) \lim_{x \rightarrow \infty} x f(1/x).$$

If $\mathbb{P}(q=0) = 0$, then the second term is assumed to be 0, irrespective of $\lim_{x \rightarrow \infty} x f(1/x)$.

Remark 76. For discrete distributions, we have the following simple definition of f -divergences:

$$D_f(P \parallel Q) = \sum_{x \in \mathcal{X}} Q(x) f \left(\frac{P(x)}{Q(x)} \right).$$

Remark 77. The above definition unifies several popular statistical divergences, such as

- Relative entropy, with $f(x) = x \log x$
- Total variation, with $f(x) = \frac{1}{2}|x - 1|$

- Squared Hellinger distance, with $f(x) = (1 - \sqrt{x})^2$
- χ^2 -distance, with $f(x) = (x - 1)^2$.

We can develop several properties of f -divergences, that parallel those for relative entropy: see Polyanskiy and Wu [2023, Chapter 7] for a thorough discussion. Below, we state and prove the nonnegativity, convexity, and DPI for f -divergences.

Proposition 78. *For any convex $f : [0, \infty]$, with $f(1) = 0$, and $f(0) := \lim_{x \downarrow 0} f(x)$, we have*

- (a) $D_f(P \parallel Q) \geq 0$, for all P, Q . Furthermore, if f is strictly convex, then the equality holds only for $P = Q$.
- (b) $D_f(P_{XY} \parallel Q_{XY}) \geq D_f(P_X \parallel Q_X)$.
- (c) The mapping $(P, Q) \mapsto D_f(P \parallel Q)$ is convex.
- (d) Suppose $P_{XY} = P_X \times P_{Y|X}$ and $Q_{XY} = Q_X \times P_{Y|X}$. Then, we have the following data-processing inequality:

$$D_f(P_Y \parallel Q_Y) \leq D_f(P_X \parallel Q_X).$$

Proof. We will prove the statements under the simplifying assumption that $P_{XY} \ll Q_{XY}$.

- (a) The result follows from a direct application of Jensen's inequality. In particular, we have

$$D_f(P \parallel Q) = \mathbb{E}_Q \left[f \left(\frac{dP}{dQ}(X) \right) \right] \geq f \left(\mathbb{E}_Q \left[\frac{dP}{dQ}(X) \right] \right) = f(1) = 0.$$

If f is not affine, then the equality holds only if dP/dQ is a constant Q -almost-surely.

- (b) For simplicity, we will prove this monotonicity result for the simple case of discrete distributions.

$$\begin{aligned} D_f(P_{XY} \parallel Q_{XY}) &= \sum_{x,y} q(x,y) f \left(\frac{p(x,y)}{q(x,y)} \right) = \sum_{x \in \mathcal{X}} q(x) \sum_{y \in \mathcal{Y}} q(y|x) f \left(\frac{p(x)p(y|x)}{q(x)q(y|x)} \right) \\ &\geq \sum_{x \in \mathcal{X}} q(x) f \left(\sum_{y \in \mathcal{Y}} q(y|x) \frac{p(x)p(y|x)}{q(x)q(y|x)} \right) \\ &\geq \sum_{x \in \mathcal{X}} q(x) f \left(\frac{p(x)}{q(x)} \sum_{y \in \mathcal{Y}} p(y|x) \right) = \sum_{x \in \mathcal{X}} q(x) f \left(\frac{p(x)}{q(x)} \right) = D_f(P_X \parallel Q_X). \end{aligned}$$

- (c) The proof of this result uses the monotonicity. In particular, let P_0, P_1 and Q_0, Q_1 denote two pairs of distributions. Now, let $X \sim \text{Bernoulli}(\lambda)$ for some $\lambda \in [0, 1]$. Then, define the following joint distributions (P_{XY} and Q_{XY}):

$$P_{Y|X=0} = P_0, \quad P_{Y|X=1} = P_1, \quad \text{and} \quad Q_{Y|X=0} = Q_0, \quad Q_{Y|X=1} = Q_1.$$

Then, by the monotonicity result, we have

$$\begin{aligned} D_f(P_{XY} \parallel Q_{XY}) &= \bar{\lambda} \sum_y Q_0(y) f \left(\frac{\bar{\lambda} P_0(y)}{\bar{\lambda} P_1(y)} \right) + \lambda \sum_y Q_1(y) f \left(\frac{\lambda P_1(y)}{\lambda Q_1(y)} \right) \\ &\geq D_f(P_Y \parallel Q_Y) \\ &= D_f(\bar{\lambda} P_0 + \lambda P_1 \parallel \bar{\lambda} Q_0 + \lambda Q_1). \end{aligned}$$

(d) Again, we prove this in the simple setting of discrete distribution. By monotonicity, we know that

$$D_f(P_Y \parallel Q_Y) \leq D_f(P_{XY} \parallel Q_{XY}).$$

Note that since the conditional distributions $P_{Y|X}$ and $Q_{Y|X}$ are the same (i.e., the marginals P_X and P_Y are passed through the same channel), we have

$$\begin{aligned} D_f(P_{XY} \parallel Q_{XY}) &= \sum_{x \in \mathcal{X}} q(x) \sum_{y \in \mathcal{Y}} q(y|x) f\left(\frac{p(x)p(y|x)}{q(x)q(y|x)}\right) \\ &= \sum_{x \in \mathcal{X}} q(x) \sum_{y \in \mathcal{Y}} q(y|x) f\left(\frac{p(x)}{q(x)}\right) \\ &= \sum_{x \in \mathcal{X}} q(x) f\left(\frac{p(x)}{q(x)}\right) \sum_{y \in \mathcal{Y}} q(y|x) = D_f(P_X \parallel Q_X). \end{aligned}$$

□

As for the case of relative entropy, we can obtain the analogs of the two variational definitions for f -divergences as well.

Theorem 79. *Let P and Q be any two probability measures defined on some measurable space $(\mathcal{X}, \mathcal{F})$. For any finite disjoint partition of \mathcal{X} , denoted by $\mathcal{E} = \{E_i \in \mathcal{F} : 1 \leq i \leq n\}$, let $P_{\mathcal{E}}$ and $Q_{\mathcal{E}}$ denote the discrete distributions with pmfs $\{P(E) : E \in \mathcal{E}\}$ and $\{Q(E) : E \in \mathcal{E}\}$. Then, we have the following:*

$$D_f(P \parallel Q) = \sup_{\mathcal{E}} D_f(P_{\mathcal{E}} \parallel Q_{\mathcal{E}}) = \sup_{\mathcal{E}} \sum_{E \in \mathcal{E}} Q(E) f\left(\frac{P(E)}{Q(E)}\right).$$

The proof of this statement follows the general argument used in proving Theorem 69 for the analogous result for relative entropy. We refer the reader to Polyanskiy and Wu [2023, § 7.14] for details.

Next, we state and prove an analog of Theorem 71 for f -divergences on $\mathcal{X} = \mathbb{R}^d$.

Theorem 80. *Let $P \ll Q$ be two distributions on $\mathcal{X} = \mathbb{R}^d$, and furthermore assume that both P and Q admit densities (p , and q respectively) w.r.t. the Lebesgue measure μ . With f^* denoting the convex-conjugate of f , we have the following variational representation of $D_f(P \parallel Q)$:*

$$D_f(P \parallel Q) = \mathbb{E}_Q \left[f\left(\frac{dP}{dQ}(X)\right) \right] = \sup_{g: \mathcal{X} \rightarrow \mathbb{R}} \mathbb{E}_P[g(X)] - \mathbb{E}_Q[f^*(g(X))],$$

where g is restricted to ensure that both expectations are finite.

Proof. The proof follows directly from the definition of convex conjugates. First note that since f is convex, we have $f = (f^*)^*$. Hence, for any $u \in (0, \infty)$, we have

$$f(u) = \sup_{v \in \mathbb{R}} uv - f^*(v).$$

Thus, with $u = p(x)/q(x)$, we have

$$f\left(\frac{p(x)}{q(x)}\right) = \sup_{v \in \mathbb{R}} \frac{p(x)v}{q(x)} - f^*(v).$$

For any function g , plugging the value of $v = g(x)$ above gives us a lower bound on $f(p(x)/q(x))$. Hence, for an arbitrary function g , we have

$$D_f(P \parallel Q) \geq \sup_g \int_{\mathcal{X}} q(x) \left(g(x) \frac{p(x)}{q(x)} - f^*(g(x)) \right) dx = \sup_g \mathbb{E}_P[g(X)] - \mathbb{E}_Q[f^*(g(X))].$$

To show the other direction: fix an arbitrary $\epsilon > 0$, and define g_{ϵ} as the function with values $g(x) = v_x$; where v_x ensures $v_x p(x)/q(x) - f^*(v_x) \geq f(p(x)/q(x)) - \epsilon$. Plugging this in, we get

$$D_f(P \parallel Q) \leq \mathbb{E}_P[g_{\epsilon}(X)] - \mathbb{E}_Q[f^*(g_{\epsilon}(X))] + \epsilon.$$

□

Example 81 (χ^2 -divergence). Recall that χ^2 distance corresponds to $f(x) = (x-1)^2$. The convex conjugate of this function is equal to $f^*(u) = u^2/4 + u = (u/2 + 1)^2 - 1$. To see this, note that

$$f^*(u) = \sup_{v \in \mathbb{R}} uv - (v-1)^2 = \sup_{v \in \mathbb{R}} g(v)$$

Note that g is a concave function, and $g'(v) = u - 2(v-1)$. Hence, its optimal value is achieved at $v^* = 1 + u/2$, and is equal to

$$f^*(u) = u \left(\frac{u}{2} + 1 \right) - \frac{u^2}{4} = \frac{u^2}{4} + u = \left(\frac{u}{2} + 1 \right)^2 - 1.$$

Plugging this into the variational definition of f -divergences, we get

$$\begin{aligned} \chi^2(P \parallel Q) &= \sup_g \mathbb{E}_P[g(X)] - \mathbb{E}_Q \left[\left(\frac{g(X)}{2} + 1 \right)^2 \right] + 1 \\ &= \sup_g 2\mathbb{E}_P \left[\left(\frac{g(X)}{2} + 1 \right) \right] - \mathbb{E}_Q \left[\left(\frac{g(X)}{2} + 1 \right)^2 \right] - 1 \\ &= \sup_h 2\mathbb{E}_P[h(X)] - \mathbb{E}_Q[h^2(X)] - 1, \end{aligned}$$

where in the last inequality, we used the change of variable $h(x) \leftarrow g(x)/2 + 1$.

Note that the above variational definition is not scale-invariant, and we can replace $h(\cdot)$ with an arbitrary $\lambda h(\cdot)$, to get

$$\chi^2(P \parallel Q) = \sup_{h, \lambda} 2\lambda \mathbb{E}_P[h(X)] - \lambda^2 \mathbb{E}_Q[h^2(X)] - 1.$$

On optimizing over λ , we get the following, more interpretable, definition

$$\chi^2(P \parallel Q) = \sup_h \frac{(\mathbb{E}_P[h(X)] - \mathbb{E}_Q[h(X)])^2}{\mathbb{V}_Q(h(X))}.$$

That is, two distributions P and Q are distinguishable in χ^2 -divergence, if there exists a function h , for which the absolute difference in means under P and Q is much larger than the standard deviation of $h(X)$ under Q .

Application: Estimation error bounds. Let us consider an estimation problem in the simplest case, where the parameter set Θ is a subset of \mathbb{R} . Now, let θ denote the true parameter, $X \sim P_\theta$, and $\hat{\theta} = \hat{\theta}(X)$ is some estimate of θ based on X . That is, we have the Markov chain $\theta \rightarrow X \rightarrow \hat{\theta}$.

Proposition 82 (Hammersley-Chapman-Robbins). *In the setting described above, the expected squared error of any estimator $\hat{\theta}$ satisfies*

$$\mathbb{E}_\theta[(\theta - \hat{\theta})^2] \geq \mathbb{V}_\theta(\hat{\theta}) \geq \sup_{\theta' \neq \theta} \frac{(\mathbb{E}_\theta[\hat{\theta}(X)] - \mathbb{E}_{\theta'}[\hat{\theta}(X)])^2}{\chi^2(P_{\theta'} \parallel P_\theta)}.$$

Proof. This result follows by a simple application of the DPI for f -divergences, and the variational definition of χ^2 -divergence obtained in Example 81. In particular, let $\theta' \in \Theta$ such that $\theta' \neq \theta$. Then, with $P_X = P_{\theta'}$ and $Q_X = P_\theta$, we have

$$\chi^2(P_X \parallel Q_X) \stackrel{(i)}{\geq} \chi^2(P_{\hat{\theta}} \parallel Q_{\hat{\theta}}) \stackrel{(ii)}{\geq} \frac{(\mathbb{E}_\theta[\hat{\theta}] - \mathbb{E}_{\theta'}[\hat{\theta}])^2}{\mathbb{V}_\theta(\hat{\theta})},$$

where (i) is due to the data-processing inequality, and (ii) is due to the variational representation of χ^2 -divergence with $h(x) = x$. On rearranging, we get

$$\mathbb{V}_\theta(\hat{\theta}) \geq \frac{\left(\mathbb{E}_\theta[\hat{\theta}] - \mathbb{E}_{\theta'}[\hat{\theta}]\right)^2}{\chi^2(P_{\theta'} \parallel P_\theta)}.$$

As $\theta' \neq \theta$ is an arbitrary element of Θ , the result follows by taking a supremum over all such θ' . \square

Corollary 83 (Cramer-Rao lower bound). *Suppose that the distributions $\{P_\theta : \theta \in \Theta \subset \mathbb{R}\}$ admit densities (w.r.t. Lebesgue measure) that are twice continuously differentiable. If we restrict our attention to unbiased $\hat{\theta}$, then Proposition 82 implies the following:*

$$\mathbb{E}_\theta[(\hat{\theta} - \theta)^2] = \mathbb{V}_\theta(\hat{\theta}) \geq \frac{1}{J_F(\theta)}, \quad \text{where} \quad J_F(\theta) = \mathbb{E}_\theta \left[\left(\frac{\partial \log(p_\theta(X))}{\partial \theta} \right)^2 \right].$$

Proof. The conclusion follows from a fact about χ^2 -distance that

$$\chi^2(P_{\theta'} \parallel P_\theta) = J_F(\theta)(\theta - \theta')^2 + o((\theta - \theta')^2).$$

To see why this is true, consider the definition of χ^2 -divergence between $P_{\theta'}$ and P_θ

$$\chi^2(P_{\theta'} \parallel P_\theta) = \int_{\mathbb{R}} p_\theta(x) \left(\frac{p_{\theta'}(x)}{p_\theta(x)} - 1 \right)^2 dx = \int_{\mathbb{R}} \frac{(p_{\theta'}(x) - p_\theta(x))^2}{p_\theta(x)} dx.$$

Now, by a Taylor' expansion around $p_\theta(x)$, we have

$$p_{\theta'}(x) = p_\theta(x) + \dot{p}_\theta(x)(\theta' - \theta) + o(|\theta' - \theta|).$$

Plugging this approximation back into the definition of $\chi^2(P_{\theta'} \parallel \theta)$, we get that

$$\chi^2(P_{\theta'} \parallel P_\theta) = (\theta' - \theta)^2 \int \frac{\dot{p}_\theta^2(x)}{p_\theta(x)} dx + o((\theta' - \theta)^2).$$

\square

Chapter 3

Compression and Gambling

- Definition of a D-ary code
- Singular, Uniquely Decodable, Prefix Codes
- Kraft's inequality for prefix codes, and UD codes
- Entropy as a lower bound on the codeword length
- Relative entropy as a measure of redundancy
- Achievability: Shannon Codes, SFE codes, Huffman code
- Doubling rate in gambling
- Value of side-information in gambling
- Portfolio optimization + Proportion (or Kelly) Betting

3.1 Source Codes

Definition 84. A D -ary source code for symbols from an alphabet \mathcal{X} is a mapping $C : \mathcal{X} \rightarrow \mathcal{D}^*$, where $\mathcal{D}^* = \cup_{m=1}^{\infty} \mathcal{D}^m$. For any x , we refer to $C(x)$ as the codeword associated with x , and use $l(x)$ to denote its length. The average codeword length of an \mathcal{X} -valued random variable $X \sim P_X$, with the coding scheme C , is defined as $L_C(X) = \sum_{x \in \mathcal{X}} p_X(x) l(x)$.

For any $k \geq 1$, the k -extension of C , is the mapping from $\mathcal{X}^k \rightarrow \mathcal{D}^*$ that assigns $x^k = (x_1, \dots, x_k)$ to $C(x^k) = C(x_1)C(x_2) \dots C(x_k)$.

Definition 85. We are interested mainly in the following three classes of codes:

- We say a code C is *nonsingular*, if $x \neq x' \implies C(x) \neq C(x')$.
- We say a code C is *uniquely decodable*, if for all $k \geq 1$ the k extension of C is nonsingular.
- We say a code C is *instantaneously decodable*, if no codeword $C(x)$ is the prefix for another codeword $C(x')$.

Remark 86. Nonsingular codes cannot be applied in general to streams of symbols. Uniquely decodable codes may require waiting arbitrarily long before even one symbol can be decoded. Instantaneously decodable (or prefix) codes are self-punctuating, and hence they can be decoded on a per-symbol basis.

3.2 Kraft's Inequality

In this section, we will obtain a correspondence between probability distributions and compression schemes. In particular, we show that every prefix code (in fact, every UD code) induces a probability distribution on the alphabet.

Theorem 87. *Suppose \mathcal{X} is a countable alphabet, and C denotes any binary prefix code with lengths $(l_i)_{i \geq 1}$. Then, we have*

$$\sum_{i \geq 1} 2^{-l_i} \leq 1. \quad (3.1)$$

Conversely, for any $(l_i)_{i \geq 1}$ satisfying (3.1), we can construct a prefix code with these lengths.

Proof. For any binary codeword $\mathbf{y}_i = (y_1, y_2, \dots, y_{l_i}) \in \{0, 1\}^*$, we introduce the following mapping:

$$V(\mathbf{y}) = 0.y_1y_2 \dots y_{l_i} = \sum_{j=1}^{l_i} y_j 2^{-j}.$$

In words, V assigns \mathbf{y}_i to a real number in $[0, 1]$, whose binary representation is given by $0.\mathbf{y}_i$. Now, let $I(\mathbf{y}_i)$ denote the interval that consists of all binary sequences whose first l_i bits are equal to \mathbf{y}_i . It is easy to verify that

$$I(\mathbf{y}_i) = [V(\mathbf{y}_i), V(\mathbf{y}_i) + 2^{-l_i}).$$

Next, we observe that since C is a prefix code, no codeword is a prefix for another. This implies that

$$I(\mathbf{y}_i) \cap I(\mathbf{y}_j) = \emptyset, \quad \text{for } i \neq j.$$

Since the length of each interval $I(\mathbf{y}_i)$ is equal to 2^{-l_i} , and they are all contained in $[0, 1]$, the sum of their lengths must be upper bounded by 1, as required.

For the converse part, given $(l_i)_{i \geq 1}$ satisfying Kraft's inequality, we construct codewords as the binary representation of the lower endpoints of the following disjoint intervals:

$$I_i = \left[\sum_{j=1}^{i-1} 2^{-l_j}, \sum_{j=1}^i 2^{-l_j} \right), \quad \text{for all } i \geq 1.$$

The length of interval I_i is equal to 2^{-l_i} , which by assumption sums up to no larger than 1. □

Can we gain anything by considering uniquely decodable codes? The next result says no.

Theorem 88. *Suppose C is a uniquely decodable code over a countable alphabet \mathcal{X} , with lengths $(l_i)_{i \geq 1}$. Then, the lengths must satisfy Kraft's inequality.*

Furthermore, for any $(l_i)_{i \geq 1}$ satisfying Kraft's inequality, there exists a UD code over \mathcal{X} with those lengths.

Proof. If C is a UD code for symbols from \mathcal{X} , then it is also a UD code for any finite subset \mathcal{X}' of \mathcal{X} . Let \mathcal{X}_N denote the subset of \mathcal{X} consisting of its first N elements. Then, note that

$$\sum_{x \in \mathcal{X}} 2^{-l(x)} = \lim_{N \rightarrow \infty} \sum_{x \in \mathcal{X}_N} 2^{-l(x)}.$$

This means that it suffices to establish Kraft's inequality for an arbitrary finite N , since the bound follows by taking N to infinity.

Fix an $N < \infty$, and let $l_{\max} = \max_{x \in \mathcal{X}_N} l(x)$. Then, observe the following for an arbitrary $k \in \mathbb{N}$:

$$\begin{aligned} \left(\sum_{x \in \mathcal{X}_N} 2^{-l(x)} \right)^k &= \sum_{x_1 \in \mathcal{X}_N} \sum_{x_2 \in \mathcal{X}_N} \dots \sum_{x_k \in \mathcal{X}_N} 2^{-l(x_1)} 2^{-l(x_2)} \dots 2^{-l(x_k)} \\ &= \sum_{x^k \in \mathcal{X}_N^k} 2^{-l(x^k)} \\ &= \sum_{m=1}^{kl_{\max}} a(m) 2^{-m}, \end{aligned}$$

where $a(m)$ denotes the number of sequences in \mathcal{X}_N^m that are assigned a codeword of length $m \in \{1, 2, \dots, kl_{\max}\}$. Next, we make the observation that $a(m) \leq 2^m$. This is because the code C is assumed to be uniquely decodable, which means that each codeword in $\{0, 1\}^m$ is assigned to at most one element of \mathcal{X}_N^* . This implies that $a(m) 2^{-m} \leq 1$, and thus

$$\begin{aligned} \left(\sum_{x \in \mathcal{X}_N} 2^{-l(x)} \right)^k &\leq kl_{\max} \\ \implies \sum_{x \in \mathcal{X}_N} 2^{-l(x)} &\leq k^{1/k} l_{\max}^{1/k}. \end{aligned}$$

The above inequality is true for all values of $k \geq 1$, and hence, by taking the limit of $k \rightarrow \infty$, we get

$$\sum_{x \in \mathcal{X}_N} 2^{-l(x)} \leq \lim_{k \rightarrow \infty} (kl_{\max})^{1/k} = 1.$$

Since N was arbitrary, we can then conclude that

$$\sum_{x \in \mathcal{X}} 2^{-l(x)} = \lim_{N \rightarrow \infty} \sum_{x \in \mathcal{X}_N} 2^{-l(x)} \leq 1.$$

□

3.3 Lower Bound and SFE coding

Entropy as a lower bound on average codeword length. We now observe that for a random variable $X \sim P_X$ taking values on the alphabet \mathcal{X} , the average codeword length for any uniquely decodable code C is lower bounded by the entropy of X .

Theorem 89. *Suppose $X \sim P_X$ is a random variable taking values in a countable alphabet \mathcal{X} . If C is any uniquely decodable coding scheme for X , then, we have*

$$L_C(X) = \sum_{x \in \mathcal{X}} p(x) l(x) \geq H(X) = \sum_{x \in \mathcal{X}} p(x) \log(1/p(x)).$$

Proof. Since C is uniquely decodable, it satisfies Kraft's inequality. Thus, the term $A = \sum_{x \in \mathcal{X}} 2^{-l(x)}$ is less than or equal to 1. Define the probability distribution R over \mathcal{X} , with p.m.f. $r(x) = 2^{-l(x)}/A$. Then, we have the following:

$$\begin{aligned} L_C(X) - H(X) &= \sum_{x \in \mathcal{X}} p(x) l(x) - \sum_{x \in \mathcal{X}} p(x) \log(1/p(x)) \\ &= \sum_{x \in \mathcal{X}} p(x) \log \left(\frac{p(x)}{2^{-l(x)}} \right) \\ &= \sum_{x \in \mathcal{X}} p(x) \log \left(\frac{p(x)}{r(x)} \right) + \sum_{x \in \mathcal{X}} p(x) \log(1/A) \\ &= D_{\text{kl}}(P_X \parallel R) - \log(1/A) \geq D_{\text{kl}}(P_X \parallel R) \geq 0. \end{aligned}$$

In the last inequality we used the nonnegativity of relative entropy, while in the second-last inequality, we used the fact that $A \leq 1$ for uniquely decodable codes. \square

The proof of the above result also tells us when the optimal bound is achieved: if $A = 1$, and $P_X = R$.

Typical Set Coding. We saw an example of a lossless compression scheme which achieves close to the optimal compression rate in Remark 9. However, that scheme is mainly a theoretical construction, and is not feasible for practical compression tasks due to its exponential complexity.

Shannon Fano Elias Coding. We now present a practical compression scheme that achieves the optimal compression rate for large block lengths.

3.4 Gambling, Doubling Rate

3.5 Portfolio Optimization + Kelly Betting

Chapter 4

Universal Compression and Gambling

4.1 Universal Compression

4.1.1 Redundancy-Capacity

4.1.2 Regret-Complexity

4.1.3 Mixture Method

4.2 Universal Portfolios

Part II

Applications

Chapter 5

Application I: From Universal Compression to Sequential Inference

5.1 Testing by Betting

5.2 Hypothesis Testing

5.3 Confidence Sequences

Bibliography

- I. Csiszár and J. Körner. *Information theory: coding theorems for discrete memoryless systems*. Cambridge University Press, 2011.
- Y. Peres. Iterating von neumann’s procedure for extracting random bits. *The Annals of Statistics*, pages 590–597, 1992.
- Y. Polyanskiy and Y. Wu. *Information theory: from coding to learning*. Cambridge University Press, 2023.