Lecture 1: From Particle Systems to Gaussian Measures on Hilbert Spaces

Tien Anh Vu

16.10.2025

This lecture begins with empirical measures of independent Brownian particles, derives the macroscopic heat equation and the fluctuation heuristics leading to the Dean-Kawasaki picture, and then turns to Gaussian measures on Hilbert spaces, Wiener measure, covariance operators, and the inverse Laplacian.

1. Microscopic Particle System

Let us consider a system of independent Brownian motions, which we will denote as $B^1, B^2, B^3, \ldots$. We also take a test function f that is smooth. Specifically, we require $f \in C_b^2(\mathbb{R})$, which means that f is twice continuously differentiable and has bounded derivatives.

Definition 1 (Empirical average)**.** We can define the empirical average of this system of N particles at time t as $\eta_t^N(f)$. We write this as:

$$\eta_t^N(f) := \frac{1}{N}\sum_{i=1}^{N} f\big(B_t^i + x_0\big).$$

Different choices of f probe different features:

- if $f(x) = x$, then we are averaging particle positions (mean position);
- if $f(x) = x^2$, then we are averaging squared positions (second moment/spread);

If we want to emphasize sample-wise evaluation, we write

$$\eta_t^N(f)(\omega) := \frac{1}{N}\sum_{i=1}^{N} f\big(B_t^i(\omega) + x_0\big).$$

where x_0 represents the starting position of our particles. We can equivalently express this empirical average as an integral with respect to the empirical measure, utilizing the Dirac measure δ:

$$\eta_t^N(f) = \int f \, d\left(\frac{1}{N}\sum_{i=1}^{N} \delta_{B_t^i + x_0}\right).$$

This is shorthand for the same integral with the variable shown explicitly:

$$\eta_t^N(f) = \int_{\mathbb{R}} f(x) \, d\left(\frac{1}{N}\sum_{i=1}^{N} \delta_{B_t^i + x_0}\right)(x).$$

2. Macroscopic Limit via SLLN

Now, we want to understand the macroscopic behavior of this system as the number of particles grows very large. We apply the Strong Law of Large Numbers (SLLN). As $N \to \infty$, this empirical average converges almost surely with respect to P-measure to the expected value:

$$\eta_t^N(f) \xrightarrow[N\to\infty]{\text{a.s. w.r.t. } P} \mathbb{E}\big[f(B_t + x_0)\big].$$

Since B_t is a standard Brownian motion, a clean way to write this is

$$B_t + x_0 \sim \mathcal{N}(x_0, t), \qquad \mu_t(dx) = p(t,x)\,dx,$$

with

$$p(t,x) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{(x-x_0)^2}{2t}\right).$$

Here μ_t is the probability distribution of $B_t + x_0$ at time t, and $p(t,x)$ is the density of μ_t (equivalently, of $B_t + x_0$) with respect to Lebesgue measure. Then the expectation becomes

$$\mathbb{E}\big[f(B_t + x_0)\big] = \int_{\mathbb{R}} f(x)p(t,x)\,dx.$$

Intuitively, we are averaging the values of f over all positions x, weighted by how likely the Brownian particle is to be at x at time t.

3. Governing PDE: Heat Equation

The natural question now is to identify the governing equation for this probability density $p(t,x)$. It turns out that $p(t,x)$ is the fundamental solution of the Heat Equation. The system is:

$$\partial_t p(t,x) = \frac{1}{2}\partial_{xx} p(t,x), \qquad t > 0,\ x \in \mathbb{R},$$

coupled with the initial condition as $t \to 0$:

$$p(t,x)dx \to \delta_{x_0}(dx).$$

In this sense, this is not a classical solution at $t = 0$, because the initial condition is not a function. Rather, the equation should be understood in a weak or distributional sense with initial datum given by the Dirac measure.

4. Physical Intuition

To build a physical intuition for the heat equation, consider the relationship between the time derivative $\partial_t p$ and the spatial second derivative $\partial_{xx} p$. If you plot the density $p(t,x)$ and look at a region with negative curvature—such as a local peak—the second spatial derivative $\partial_{xx} p(t,x)$ is negative. According to the heat equation, this means the time derivative $\partial_t p(t,x)$ must also be negative. Consequently, at points of negative curvature, the "temperature" or probability density will go down, which has the effect of smoothing out the distribution over time.

At points of positive curvature, such as local minima, the second spatial derivative is positive, so the density increases there. In this way, the heat equation averages out local extrema and smooths the profile over time.

This heuristic picture explains the macroscopic heat flow. The next step is to recover the same evolution directly from the empirical particle system and to identify the stochastic fluctuations before the limit.

5. Microscopic Fluctuations and the Macroscopic Limit

5.1. Setup and Empirical Average

We have already introduced the microscopic particle system, defined the empirical average, and identified the corresponding macroscopic heat equation. We now continue by making that passage more rigorous: we analyze the fluctuations around the Law of Large Numbers using Itô's formula and show how the stochastic part disappears as the number of particles grows.

To do this, we return to the empirical measure for N independent Brownian motions B_t^i, all starting at an initial position x_0, and evaluate it against a test function f. This gives the empirical average

$$\eta_t^N(f) = \frac{1}{N}\sum_{i=1}^{N} f(B_t^i + x_0).$$

5.2. Itô Formula and the Drift-Martingale Decomposition

To observe the time evolution of this empirical measure, we compute its stochastic differential. Applying Itô's formula, the differential is expressed as:

$$d\left(\frac{1}{N}\sum_{i=1}^{N} f(B_t^i + x_0)\right) = \frac{1}{N}\sum_{i=1}^{N} df(B_t^i + x_0).$$

When we expand $df(B_t^i + x_0)$ using the standard rules of stochastic calculus, we decompose the dynamic into two fundamentally distinct components: an Itô integral and a correction term (the drift). The expansion gives us:

$$d\eta_t^N(f) = \frac{1}{N}\sum_{i=1}^{N} f'(B_t^i + x_0)\, dB_t^i + \frac{1}{N}\sum_{i=1}^{N} \frac{1}{2} f''(B_t^i + x_0)\, dt.$$

Let us analyze these two terms closely. The second term is our correction term, which can be written compactly using our empirical measure notation as $\eta_t^N\left(\frac{1}{2}f''\right)$. This term captures the expected diffusion, effectively mapping to the Laplacian operator in our emerging macroscopic PDE.

The first term is a local Martingale, which we will denote as $M_t^N(f)$. This term represents the microscopic stochastic fluctuations of the system. To understand the macroscopic limit, we must determine the asymptotic behavior of this Itô integral as $N \to \infty$. Specifically, we need to prove that

$$M_t^N(f) \xrightarrow[N\to\infty]{} 0.$$

5.3. Vanishing of the Martingale via Itô Isometry

To prove that the fluctuations vanish, we compute the expected value of the squared martingale, $\mathbb{E}[(M_t^N(f))^2]$. We do this by applying Itô isometry, which allows us to equate the expectation of a squared stochastic integral to the expectation of the standard integral of the squared integrand. Because the Brownian motions are independent, the cross-terms in the squared sum take an expectation of zero, leaving us with:

$$\mathbb{E}[(M_t^N(f))^2] = \frac{1}{N^2}\sum_{i=1}^{N} \mathbb{E}\left[\left(\int_0^t f'(B_s^i + x_0)\, dB_s^i\right)^2\right].$$

By Itô isometry, we transform the stochastic integral into a Riemann integral:

$$\mathbb{E}[(M_t^N(f))^2] = \frac{1}{N^2} \sum_{i=1}^{N} \mathbb{E}\left[\int_0^t f'(B_s^i + x_0)^2 \, ds\right].$$

Since the N Brownian motions are identically distributed, the expectation is the same for every particle i. This allows us to simplify the sum completely:

$$\mathbb{E}[(M_t^N(f))^2] = \frac{1}{N} \mathbb{E}\left[\int_0^t f'(B_s + x_0)^2 \, ds\right].$$

5.4. Quadratic Variation

We can apply the exact same logic to compute the quadratic variation of the martingale:

$$\langle M^N(f) \rangle_t = \frac{1}{N^2} \sum_{i=1}^{N} \int_0^t f'(B_s^i + x_0)^2 \, ds.$$

This expression can be rewritten by pulling the $\frac{1}{N}$ factor out and recognizing the remaining sum as the empirical measure applied to $(f')^2$. Thus, we obtain:

$$\langle M^N(f) \rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2) \, ds.$$

The critical insight here is that the quadratic variation is of order $O\left(\frac{1}{N}\right)$. Because the variance of the martingale term scales by $\frac{1}{N}$, it completely vanishes in the limit as the number of molecules N approaches infinity.

5.5. Conclusion

This demonstrates the Law of Large Numbers for our empirical measure. The stochastic noise that characterizes the microscopic gas molecules dissipates entirely, leaving only the deterministic correction term. This remaining term dictates the macroscopic description of the system, effectively deriving the governing PDE from the underlying stochastic particle dynamics.

6. Stochastic Dynamics of the Empirical Measure

The Law of Large Numbers identifies the deterministic limit, but the weak differential form of the empirical measure is what makes the fluctuation analysis systematic. We therefore rewrite the particle dynamics directly at the level of the empirical measure.

6.1. The Evolution Equation for the Empirical Measure

We start with the fundamental stochastic differential equation for the empirical measure η_t^N tested against a function f. The equation reads:

$$d\eta_t^N(f) = dM_t^N(f) + \eta_t^N\left(\frac{1}{2} f''\right) dt.$$

Here, the evolution of our empirical measure is decomposed into two parts:

- A martingale term $dM_t^N(f)$, which captures the stochastic noise of the system. As $N \to \infty$, this term vanishes and goes to 0 due to the Law of Large Numbers.
- A drift term $\eta_t^N\left(\frac{1}{2}f''\right)dt$, which represents the deterministic macroscopic evolution driven by the Laplacian (here, the second derivative $\frac{1}{2}f''$ since we are dealing with standard Brownian motion).

6.2. Expanding into Integral Form and Density Assumptions

We then rewrite this differential equation into its integral form:

$$\eta_t^N(f) = \eta_0^N(f) + \int_0^t \eta_s^N\left(\frac{1}{2}f''\right)\,ds.$$

Below this, we expand the measure into an integral over space:

$$\eta_t^N(f) = \eta_0^N(f) + \int_0^t \int \frac{1}{2}f''(x)\eta_s(x)\,dx\,ds.$$

To do this, assume the empirical measure can be represented by a density $\eta_s(x)$ with respect to the Lebesgue measure.

The discrete empirical measure for our N particles is:

$$\eta_s^N(x) = \frac{1}{N}\sum_{i=1}^{N} \delta_{B_s^i + x_0}(x).$$

This shows that the measure is comprised of Dirac measures located at the positions of the individual independent Brownian particles, B_s^i, starting from some initial position x_0.

7. Distributional Derivatives

We next arrive at a formal equivalence:

$$\eta_t^N\left(\frac{1}{2}f''\right)dt = \frac{1}{2}\,\partial_{xx}\eta_t^N(f)\,dt.$$

This is a crucial step. By using integration by parts, we can transfer the second derivative from our test function f'' directly onto the empirical measure itself, yielding the distributional derivative $\partial_{xx}\eta_t^N$. This is how the heat equation emerges in the macroscopic limit.

8. Fluctuations and the Strong Law of Large Numbers

With the weak evolution equation in hand, we can now isolate the random part that survives after centering and rescaling. Define the standardized fluctuation by

$$\psi_t^N(f) := \sqrt{N}\big(\eta_t^N(f) - \eta_t(f)\big).$$

Using the evolution equation for η_t^N together with the heat equation satisfied by η_t, one obtains

$$d\psi_t^N(f) = \sqrt{N}\,dM_t^N(f) + \psi_t^N\left(\frac{1}{2}f''\right)dt.$$

This shows that the fluctuation field is driven by a rescaled martingale term together with the same second-order drift. The martingale term contains the random deviations from the macroscopic limit. When $N \to \infty$, this martingale term vanishes at the Law of Large Numbers scale, but after rescaling by $\sqrt{N}$, it survives and becomes the fluctuation field.

We write out the explicit Itô representation for the martingale term:

$$M_t^N(f) = \frac{1}{N} \sum_{i=1}^{N} \int_0^t f'(B_s^i + x_0)\, dB_s^i.$$

Its quadratic variation is

$$\left\langle \frac{1}{N} \sum_{i=1}^{N} \int_0^t f'(B_s^i + x_0)\, dB_s^i \right\rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

The single Brownian integral has the same quadratic variation:

$$\left\langle \frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s \right\rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

Therefore, the original martingale term has fluctuation size $N^{-1/2}$, and both sides below have the same fluctuation. For this reason, one writes

$$\frac{1}{N} \sum_{i=1}^{N} \int_0^t f'(B_s^i + x_0)\, dB_s^i \simeq \frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s.$$

The integrand $\eta_s^N((f')^2)$ corresponds to the quadratic variation, while dB_s represents one Brownian motion. The $1/\sqrt{N}$ scaling is the classic Central Limit Theorem scaling characterizing the size of the fluctuations.

9. The Space-Time Fluctuation Limit

To make the fluctuation term spatially explicit, rewrite its variance structure in density form and then match it with a space-time stochastic integral:

$$\eta_t(dx) = p(t, x)\, dx.$$

Thus, in the limit one has $\eta_t(x) = p(t, x)$, where $p(t, x)$ is the transition probability density, or heat kernel.

Assuming that the empirical measure admits a density, so that $\eta_s^N(dx) = \eta_s^N(x)\, dx$, one has

$$\eta_s^N((f')^2) = \int_{\mathbb{R}} (f'(x))^2 \eta_s^N(x)\, dx.$$

Substituting this into the quadratic variation gives

$$\left\langle \frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s \right\rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

The corresponding space-time stochastic integral has the same quadratic variation:

$$\left\langle \frac{1}{\sqrt{N}} \int_0^t \int_{\mathbb{R}} f'(x) \sqrt{\eta_s^N(x)}\, d\xi(s, x) \right\rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds = \frac{1}{N} \int_0^t \int_{\mathbb{R}} (f'(x))^2 \eta_s^N(x)\, dx\, ds.$$

Thus, one writes

$$dM_t^N(f) \simeq \frac{1}{\sqrt{N}} \int_{\mathbb{R}} f'(x) \sqrt{\eta_t^N(x)}\, d\xi(t,x).$$

Thus, at a formal level, the martingale term may be represented by the space-time integral above, where the noise is written as a space-time differential $d\xi(s,x)$ or $dB(s,x)$.

10. The Central Limit Theorem for the Martingale Term

The space-time representation identifies the correct fluctuation scale. The next step is to state the limiting Gaussian law of the rescaled martingale.

Scale the martingale term $M_t^N(f)$ by $\sqrt{N}$ and examine its fluctuation limit:

$$\sqrt{N}\, M_t^N(f) \to \int_0^t \sqrt{\eta_s\big((f')^2\big)}\, dB_s.$$

The limit is a Gaussian martingale represented as an Itô integral with respect to a standard Brownian motion. For each fixed time t, the Central Limit Theorem for martingales gives the Normal law

$$\sqrt{N}\, M_t^N(f) \to \mathcal{N}\left(0, \int_0^t \eta_s\big((f')^2\big)\, ds\right).$$

Thus, the asymptotic fluctuations of the empirical measure tested against f are characterized by a Gaussian variance determined by the macroscopic density η_s and the squared gradient $(f')^2$.

11. The Approximate Stochastic Differential Equation

This limiting variance structure suggests the following approximate stochastic differential equation for the empirical measure:

$$d\eta_t^N(f) \approx \eta_t^N\left(\frac{1}{2} f''\right) dt + \frac{1}{\sqrt{N}} \sqrt{\eta_t^N((f')^2)}\, dB_t.$$

This encapsulates both the deterministic drift, namely the Laplacian term derived earlier, and the newly scaled fluctuation term driven by a single Brownian motion dB_t. The variance of the Itô integral is equal to the expectation of the integral of the squared integrand. Here,

$$\mathrm{Var}\left(\frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s\right) = \mathbb{E}\left[\left(\frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s\right)^2\right] = \frac{1}{N}\, \mathbb{E}\left[\int_0^t \eta_s^N((f')^2)\, ds\right].$$

12. Space-Time White Noise and Variance Equivalence

To pass from the single-Brownian representation to a spatial noise term, introduce space-time white noise $\xi(t,x)$ with covariance

$$\mathbb{E}(\xi(t,x)\xi(s,y)) = \delta(t-s)\delta(x-y).$$

This indicates that the noise is completely uncorrelated in both space and time. For this to work formally, we write

$$\eta_s^N(dx) = \eta_s^N(x)\, dx,$$

so that the empirical measure is represented by a spatial density.

We now show that the space-time white noise formulation matches the variance of the original particle system. First, we look at the scaled variance

$$V_t^N(f) := \frac{1}{N}\,\mathbb{E}\bigg(\int_0^t \eta_s^N((f')^2)\,ds\bigg).$$

Next, we expand the empirical measure into its explicit spatial integral:

$$= V_t^N(f) = \frac{1}{N}\,\mathbb{E}\bigg(\int_0^t \int_{\mathbb{R}} (f'(x))^2 \eta_s^N(x)\,dx\,ds\bigg).$$

Finally, applying Itô isometry in reverse with the space-time white noise, we can express this same expectation as the square of a stochastic integral:

$$= V_t^N(f) = \frac{1}{N}\,\mathbb{E}\bigg[\bigg(\int_0^t \int_{\mathbb{R}} f'(x)\sqrt{\eta_s^N(x)}\,d\xi(s,x)\bigg)^2\bigg].$$

This is the critical mathematical bridge. It proves that the macroscopic noise acting on the system can be represented by integrating $f'(x)$ against $\sqrt{\eta_s^N(x)}$ multiplied by the space-time white noise $d\xi(s,x)$.

13. The Dean-Kawasaki Equation

Combining the weak empirical evolution with the space-time representation of the martingale term leads to the formal Dean-Kawasaki equation. Recalling the earlier evolution equation

$$d\eta_t^N(f) = dM_t^N(f) + \eta_t^N\left(\frac{1}{2}f''\right)dt,$$

and using the space-time noise representation obtained earlier,

$$dM_t^N(f) \simeq \frac{1}{\sqrt{N}}\int_{\mathbb{R}} f'(x)\sqrt{\eta_t^N(x)}\,d\xi(t,x),$$

one arrives at the weak approximation

$$d\eta_t^N(f) \simeq \eta_t^N\left(\frac{1}{2}f''\right)dt + \frac{1}{\sqrt{N}}\int_{\mathbb{R}} f'(x)\sqrt{\eta_t^N(x)}\,d\xi(t,x).$$

To rewrite this in density form, formally assume that the empirical measure admits a density $\eta_t^N(x)$. Then the drift term becomes

$$\eta_t^N\left(\frac{1}{2}f''\right) = \int_{\mathbb{R}} \frac{1}{2}f''(x)\eta_t^N(x)\,dx = \int_{\mathbb{R}} f(x)\,\frac{1}{2}\partial_{xx}\eta_t^N(x)\,dx,$$

again formally by integration by parts. For the noise term, one likewise writes

$$\int_{\mathbb{R}} f'(x)\sqrt{\eta_t^N(x)}\,d\xi(t,x) = -\int_{\mathbb{R}} f(x)\,\partial_x\big(\sqrt{\eta_t^N(x)}\,\xi(t,x)\big)\,dx,$$

and the sign may be absorbed into the noise, since replacing ξ by $-\xi$ does not change its law. On the left-hand side, the weak form of the time derivative is

$$d\eta_t^N(f) = \int_{\mathbb{R}} f(x)\,\partial_t\eta_t^N(x)\,dx\,dt$$

so the weak approximation can be rewritten as

$$\int_{\mathbb{R}} f(x)\,\partial_t \eta_t^N(x)\,dx\,dt \simeq \int_{\mathbb{R}} f(x)\left[\frac{1}{2}\partial_{xx}\eta_t^N(x) + \frac{1}{\sqrt{N}}\partial_x\big(\sqrt{\eta_t^N(x)}\,\xi(t,x)\big)\right]dx\,dt.$$

Since this identity holds for all test functions f, one reads off the formal distributional equation

$$\partial_t \eta_t^N = \frac{1}{2}\partial_{xx}\eta_t^N(x) + \frac{1}{\sqrt{N}}\partial_x\big(\sqrt{\eta_t^N(x)}\,\xi(t,x)\big).$$

This equation still describes the empirical density at finite N. To pass to the fluctuation equation, center and rescale:

$$\psi_t^N = \sqrt{N}\big(\eta_t^N - \eta_t\big),$$

and expect the fluctuation field to converge to a limiting process ψ_t. In the formal limit, this leads to the Dean-Kawasaki SPDE

$$\partial_t \psi_t = \frac{1}{2}\partial_{xx}\psi_t + \partial_x\big(\sqrt{\eta_t(x)}\,\xi(t,x)\big).$$

Thus, the equation for η_t^N is the pre-limit heuristic equation, whereas the equation for ψ_t is the limiting fluctuation equation.

13.1. A Crucial Warning on Well-Posedness

The solution theory of this SPDE is largely open, since its precise mathematical formulation is unclear. In particular, it is not clear whether the asymptotic fluctuations can really be represented as the integral of a test function against a signed density, let alone a sufficiently differentiable one. This indicates that some of the assumptions made in the formal derivation are not easy to justify rigorously. Thus, while the Dean-Kawasaki equation is a useful formal model for fluctuating hydrodynamics, it remains severely ill-posed in rigorous mathematics.

14. A Linear Comparison

As a comparison point, consider the linear equation

$$dX_t = AX_t\,dt + C dW_t,$$

which is the standard form of an Ornstein-Uhlenbeck process. This provides a useful contrast between the highly non-linear, ill-posed nature of the Dean-Kawasaki SPDE and a well-understood, solvable linear stochastic differential equation.

15. Transition to Hilbert-Space Stochastic Analysis

The fluctuation equations above naturally lead to noises and covariance structures that are best understood in infinite-dimensional spaces. To make those objects precise, we now leave the particle-level description and move to probability measures on separable Hilbert spaces. The next sections introduce that framework and then return to the concrete example of Wiener measure.

16. Hilbert-Space Framework

We begin with the geometric and probabilistic setting needed for infinite-dimensional Gaussian measures.

16.1. The Setting: Hilbert Spaces and Operators

The fundamental mathematical stage for this chapter is the Hilbert space $(L^2(\mu), \langle \cdot, \cdot \rangle)$, which represents the classic L^2 space of square-integrable functions with respect to a measure μ, equipped with an inner product. A Hilbert space is a complete normed vector space. The inner product induces the norm, and completeness means that every Cauchy sequence converges within the space.

Next consider linear operators acting between spaces. Let

$$L : U \to V,$$

where U and V are Hilbert spaces. A crucial analytic property is

$$L \text{ is continuous} \quad \Longleftrightarrow \quad \sup_{\|u\|_U \leq 1} \|Lu\|_V < \infty.$$

Thus, a linear operator is continuous if and only if it is bounded. This is a bedrock principle of functional analysis.

16.2. A Canonical Example: The ℓ^2 Sequence Space

A quintessential example of an infinite-dimensional Hilbert space is the space of square-summable sequences, denoted by ℓ^2. It is defined by

$$\ell^2 = \left\{ (u_k)_{k \in \mathbb{N}} \in \mathbb{R} : \sum_{k=1}^{\infty} u_k^2 < \infty \right\}.$$

This means that one considers infinite sequences of real numbers for which the sum of their squares converges to a finite value. Its inner product is

$$\langle u, v \rangle = \sum_{k=1}^{\infty} u_k v_k.$$

This is the infinite-dimensional analogue of the standard Euclidean dot product.

A Hilbert space is called separable if there exists a countable dense subset $H_0 \subset H$. In infinite dimensions, spaces can be extremely large. Having a countable dense subset means that every element of the space can be approximated by a sequence from a countable set, which guarantees the existence of a countable orthonormal basis. This makes defining integrals and measures much more manageable.

16.3. The Core Concept: Gaussian Measures on Hilbert Spaces

We now move from deterministic functional analysis to probability theory.

Here $\mathcal{B}(H)$ denotes the Borel σ-algebra generated by the open sets of the Hilbert space.

Definition 2 (Gaussian measure)**.** A probability measure μ on $(H, \mathcal{B}(H))$ is called Gaussian if for all $h \in H$ the linear functional

$$\ell_h : H \to \mathbb{R}, \qquad g \mapsto \langle g, h \rangle_H$$

is normally distributed under μ.

This means that if g is distributed according to μ on H, then the real-valued random variable

$$\ell_h(g) = \langle g, h \rangle_H$$

has a one-dimensional normal distribution on $\mathbb{R}$. In other words, a measure on a Hilbert space is Gaussian if every one-dimensional projection is a Gaussian random variable.

By the Riesz representation theorem, any continuous linear functional can be represented as a scalar product with an element of the Hilbert space. Thus, the inner products $\langle \cdot, h \rangle$ exhaust all continuous linear observations of the space.

16.4. The Characteristic Function of the Projection ℓ_h

To formalize the Gaussian property, there exist $m(h) \in \mathbb{R}$ and $\sigma^2(h) \in \mathbb{R}_+$ such that for all $t \in \mathbb{R}$,

$$\int_H e^{it\ell_h(g)}\, \mu(dg) = e^{itm(h) - \frac{1}{2}t^2\sigma^2(h)}.$$

This is the characteristic function of the real-valued random variable $\ell_h(g) = \langle g, h \rangle_H$ when g is distributed according to μ. It is exactly the characteristic function of a one-dimensional normal distribution with mean $m(h)$ and variance $\sigma^2(h)$.

The variance term $\sigma^2(h)$ is not an arbitrary function of h. It has a quadratic dependence on h, meaning that there is a covariance operator Q such that

$$\sigma^2(h) = \langle Qh, h \rangle_H.$$

Here the map $h \mapsto Qh$ is linear, and the variance is obtained by pairing Qh again with h. This operator Q encodes how the Gaussian measure spreads in different directions of the Hilbert space.

Finally, the mean functional is linear. Indeed,

$$\int_H \ell_{h_1+h_2}(g)\, \mu(dg) = \int_H \ell_{h_1}(g)\, \mu(dg) + \int_H \ell_{h_2}(g)\, \mu(dg).$$

Because the mean of the projection ℓ_h is exactly its integral with respect to μ, one has

$$m(h_1 + h_2) = \int_H \ell_{h_1+h_2}(g)\, \mu(dg) = \int_H \ell_{h_1}(g)\, \mu(dg) + \int_H \ell_{h_2}(g)\, \mu(dg) = m(h_1) + m(h_2).$$

This page sets the stage for Gaussian noise spread across infinite dimensions, a concept that will be essential when returning to objects such as space-time white noise in stochastic partial differential equations.

17. Structural Characterization of Gaussian Measures

With the projection-based definition in place, we can now state the structural characterization of Gaussian measures on a Hilbert space.

17.1. The Characteristic Function of the Projection $\langle g, h \rangle_H$

Theorem 1 (Theorem 1.6)**.** A probability measure μ on $(H, \mathcal{B}(H))$ is Gaussian if and only if

$$\int_H e^{i\langle g,h\rangle_H}\, \mu(dg) = e^{i\langle m,h\rangle_H - \frac{1}{2}\langle Qh,h\rangle_H} \qquad \text{for all } h \in H.$$

The left-hand side is the characteristic function of the measure μ in the direction h, equivalently the characteristic function of the real-valued projection $\langle g, h\rangle_H$, and the right-hand side is the infinite-dimensional analogue of the one-dimensional Gaussian Fourier transform

$$\varphi(t) = e^{itm - \frac{1}{2}t^2\sigma^2},$$

where the mean is represented by the inner product $\langle m, h\rangle_H$ and the variance is replaced by the quadratic form $\langle Qh, h\rangle_H$. Thus, this is the concrete Hilbert-space version of the abstract projection formula introduced in Section 16.4, with $m(h) = \langle m, h\rangle_H$ and $\sigma^2(h) = \langle Qh, h\rangle_H$.

17.2. The Parameters: Mean Vector and Covariance Operator

Below the theorem, the properties of the objects m and Q determine the shape of the Gaussian measure.

First, $m \in H$ is the mean vector. Since H is a Hilbert space, the mean is itself an element of the space.

Next, the covariance operator

$$Q : H \to H$$

is linear and continuous, and it is symmetric in the sense that

$$\langle Qh_1, h_2\rangle_H = \langle h_1, Qh_2\rangle_H.$$

It is also positive semidefinite, that is,

$$\langle Qh, h\rangle_H \geq 0 \qquad \text{for all } h \in H.$$

This is required because variances must be non-negative.

17.3. The Trace Class Requirement

A crucial infinite-dimensional condition is that Q must have finite trace. The trace is defined by

$$\mathrm{tr}(Q) = \sum_{k=1}^{\infty} \langle Qe_k, e_k\rangle_H < \infty,$$

where $\{e_k\}_{k\geq 1}$ is a complete orthonormal system (CONS) of the Hilbert space. The trace is finite for one, and hence for all, complete orthonormal systems of H. Thus, the total variance of the Gaussian measure is an invariant of the operator Q and does not depend on the particular basis used to compute it. One cannot take the identity as the covariance operator in infinite dimensions. If the covariance were the identity, then all directions would carry the same variance. Geometrically, this means that small balls behave uniformly under rotations: one can rotate the same ball along different coordinate directions, and all these copies would have the same volume. In infinitely many dimensions, this would produce infinitely many disjoint balls of the same volume inside a bounded region. Then the total volume would be the same positive number added infinitely many times, which is impossible. Thus, the identity cannot be the covariance operator of a genuine Gaussian measure on an infinite-dimensional Hilbert space.

The formula version is

$$Q = \mathrm{Id}_H \quad \Longrightarrow \quad \mathrm{tr}(Q) = \mathrm{tr}(\mathrm{Id}_H) = \sum_{k=1}^{\infty} \langle e_k, e_k\rangle_H = \sum_{k=1}^{\infty} 1 = \infty.$$

This violates the finite trace condition. Another way to see this is through Gaussian mass of balls in high dimensions: for every finite R,

$$\mathcal{N}(0, \mathrm{Id}_n)\big(B_R(0)\big) \to 0 \qquad \text{as } n \to \infty.$$

Thus, if a standard normal distribution on an infinite-dimensional Hilbert space existed, it would assign zero mass to every finite-radius ball, which contradicts the fact that the whole space has mass one.

More generally, there can be no Gaussian measure $\mu = \mathcal{N}(0, Q)$ on H with bounded inverse Q^{-1}. To realize a Gaussian measure with covariance operator Id_H, one must embed H into a larger Hilbert space H_1 such that the embedding $J : H \hookrightarrow H_1$ is Hilbert-Schmidt, that is,

$$\sum_{k=1}^{\infty} \|Je_k\|_{H_1}^2 < \infty$$

for one, and hence for all, complete orthonormal systems (e_k) of H.

In this case one writes

$$\mu = \mathcal{N}(m, Q),$$

meaning that μ is the Gaussian measure with mean vector m and covariance operator Q. The operator Q is called the covariance operator of μ.

Once the trace condition is available, spectral theory can be applied to diagonalize the covariance operator.

17.4. A Spectral Fact for Trace-Class Operators

The following functional analytic fact will be used repeatedly.

Proposition 1 (Proposition 1.10)**.** Let Q be linear, continuous, symmetric, positive semidefinite, and assume that

$$\mathrm{tr}(Q) < \infty.$$

Then there exists a complete orthonormal system $(e_k)_{k\geq 1}$ of H consisting of eigenvectors of Q, that is,

$$Qe_k = \lambda_k e_k, \qquad \lambda_k \geq 0.$$

Moreover, 0 is the only accumulation point of the sequence of eigenvalues $(\lambda_k)_{k\geq 1}$, and

$$\sum_{k=1}^{\infty} \lambda_k = \mathrm{tr}(Q) < \infty.$$

This gives the canonical spectral decomposition of a positive trace-class operator and prepares the way for representing Gaussian measures through their eigenvalues and eigenvectors.

17.5. Three Fundamental Properties

Three basic identities follow from the definition. Recall that

$$\ell_h : H \to \mathbb{R}, \qquad \ell_h(x) = \langle x, h\rangle_H.$$

So ℓ_h is the linear functional associated with the vector h. It takes an element $x \in H$ and returns the scalar projection of x onto h.

That is why

$$E_\mu(\ell_h) = \int_H \ell_h(x)\,\mu(dx) = \int_H \langle x, h\rangle_H\,\mu(dx).$$

First, the expected value of a linear projection is

$$\int_H \langle x, h\rangle_H\,\mu(dx) = \langle m, h\rangle_H.$$

Second, the covariance between two linear projections is

$$\begin{aligned}
\mathrm{Cov}(\ell_{h_1}, \ell_{h_2}) &= E_\mu\big[(\ell_{h_1} - E_\mu \ell_{h_1})(\ell_{h_2} - E_\mu \ell_{h_2})\big] \\
&= \int_H \big(\langle x, h_1\rangle_H - \langle m, h_1\rangle_H\big)\big(\langle x, h_2\rangle_H - \langle m, h_2\rangle_H\big)\,\mu(dx) \\
&= \langle Qh_1, h_2\rangle_H \\
&= \langle h_1, Qh_2\rangle_H.
\end{aligned}$$

This shows how the operator Q acts in practice: the covariance between two projections is obtained by applying Q to one direction and taking the inner product with the other.

Third, the expected squared distance from the mean is

$$\int_H \|x - m\|_H^2\,\mu(dx).$$

Using Parseval's identity with an orthonormal basis $\{e_k\}$ gives

$$\int_H \|x - m\|_H^2\,\mu(dx) = \int_H \sum_{k=1}^{\infty} \langle x - m, e_k\rangle_H^2\,\mu(dx).$$

By Fubini's theorem, or by monotone convergence,

$$\int_H \|x - m\|_H^2\,\mu(dx) = \sum_{k=1}^{\infty} \int_H \langle x - m, e_k\rangle_H^2\,\mu(dx).$$

Recall the covariance identity

$$\int_H \big(\langle x, h_1\rangle_H - \langle m, h_1\rangle_H\big)\big(\langle x, h_2\rangle_H - \langle m, h_2\rangle_H\big)\,\mu(dx) = \langle Qh_1, h_2\rangle_H.$$

Applying this identity with $h_1 = h_2 = e_k$ gives

$$\int_H \langle x - m, e_k\rangle_H^2\,\mu(dx) = \langle Qe_k, e_k\rangle_H.$$

Therefore,

$$\int_H \|x - m\|_H^2\,\mu(dx) = \sum_{k=1}^{\infty} \langle Qe_k, e_k\rangle_H = \mathrm{tr}(Q).$$

This proves that the expected squared deviation of a Gaussian random variable in a Hilbert space is exactly the trace of its covariance operator. Recall Parseval's identity:

$$\|x\|_H^2 = \sum_{i=1}^{\infty} x_i^2 = \sum_{i=1}^{\infty} \langle x, e_i\rangle_H^2.$$

This summarizes the trace computation above. It says that the squared norm of an element of the Hilbert space can be decomposed into the sum of the squared projections along an orthonormal basis.

These abstract identities become concrete in the fundamental example of Wiener measure.

18. Wiener Measure

The first major example of an infinite-dimensional Gaussian measure is the Wiener measure.

18.1. The Path Mapping: Random Variables as Functions

Consider a standard one-dimensional Brownian motion $(\beta(t))_{t\in[0,T]}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The goal is to use this Brownian motion to construct a measure on a Hilbert space.

The key observation is that in infinite dimensions one maps an outcome not to a real number, but to an entire function. Define

$$\beta_\cdot : \Omega \to C([0,T]) \subseteq L^2([0,T]), \qquad \omega \mapsto (t \mapsto \beta(t,\omega)),$$

This is the sample path, or trajectory, of the Brownian motion. One often writes $\beta(\cdot)$ when the dependence on ω is suppressed. Since every continuous function on $[0,T]$ is square-integrable, one has

$$C([0,T]) \subseteq L^2([0,T]).$$

Therefore the Brownian path can be viewed as an element of the Hilbert space $L^2([0,T])$.

18.2. Measurability and the Induced Measure

For this construction to be rigorous, the mapping $\beta_\cdot$ must be measurable with respect to the Borel σ-algebra on $L^2([0,T])$. Since it is measurable, one can push the underlying probability measure forward and define a new measure on the Hilbert space by

$$\mu(A) = \mathbb{P}(\beta(\cdot) \in A), \qquad A \in \mathcal{B}(L^2([0,T])).$$

This measure μ is the Wiener measure. It gives the probability that the Brownian trajectory belongs to a measurable set of functions A.

18.3. Proving that the Mean is Zero

The next claim is that μ is Gaussian with mean 0, that is, it is centered. To prove this, one checks that the expectation of each linear projection is zero.

Let $h \in L^2([0,T])$. Then

$$\mathbb{E}_\mu(\ell_h) = \mathbb{E}(\ell_h(\beta)) = \mathbb{E}(\ell_h(\beta(\cdot))).$$

By the Riesz representation theorem, the linear functional ℓ_h can be written as an inner product in $L^2([0,T])$, so

$$\ell_h(f) = \langle f, h\rangle_{L^2([0,T])} = \int_0^T f(s)h(s)\,ds$$

for $f \in L^2([0,T])$. Applying this with $f = \beta$ in $L^2([0,T])$ gives

$$\mathbb{E}_\mu(\ell_h) = \mathbb{E}\left(\int_0^T h(s)\beta(s)\,ds\right).$$

By Fubini's theorem,

$$\mathbb{E}_\mu(\ell_h) = \int_0^T h(s)\mathbb{E}(\beta(s))\,ds.$$

Since a standard Brownian motion satisfies

$$\mathbb{E}(\beta(s)) = 0 \qquad \text{for all } s \in [0, T],$$

it follows that

$$\mathbb{E}_\mu(\ell_h) = 0.$$

Therefore the mean vector of the Wiener measure on $L^2([0, T])$ is the zero function. With the mean identified, the remaining task is to compute the covariance operator explicitly.

19. The Covariance Operator of the Wiener Measure

We now compute the second fundamental characteristic of the Wiener measure on $L^2([0, T])$, namely its covariance operator Q.

19.1. Formulating the Covariance

Since the Wiener measure is centered, one has $E_\mu(\ell_g) = E_\mu(\ell_h) = 0$, and therefore the covariance reduces to the expectation of the product:

$$\mathrm{Cov}(\ell_g, \ell_h) = \mathbb{E}(\ell_g \ell_h).$$

Substituting the Brownian path into the functionals gives

$$\mathrm{Cov}(\ell_g, \ell_h) = \mathbb{E}(\ell_g(\beta)\ell_h(\beta)).$$

19.2. Expanding into Double Integrals and Applying Fubini's Theorem

Using the Riesz representation theorem, the linear functionals can be written as L^2 inner products over the interval $[0, T]$, so

$$\mathrm{Cov}(\ell_g, \ell_h) = \mathbb{E}\left(\left(\int_0^T g(s)\beta(s)\,ds\right)\left(\int_0^T h(t)\beta(t)\,dt\right)\right).$$

Rewriting the product of integrals as a double integral and applying Fubini's theorem yields

$$\mathrm{Cov}(\ell_g, \ell_h) = \int_0^T \int_0^T g(s)h(t)\mathbb{E}(\beta(s)\beta(t))\,ds\,dt.$$

For a standard Brownian motion,

$$\mathbb{E}(\beta(s)\beta(t)) = s \wedge t = \min(s, t).$$

Substituting this identity gives

$$\mathrm{Cov}(\ell_g, \ell_h) = \int_0^T \int_0^T g(s)h(t)(s \wedge t)\,ds\,dt.$$

19.3. Isolating the Covariance Operator Q

By definition, the covariance operator is characterized by the bilinear form

$$\mathrm{Cov}(\ell_g, \ell_h) = \langle Qg, h \rangle_{L^2([0,T])}.$$

To identify Q, rewrite the double integral in the form of an L^2 inner product:

$$\mathrm{Cov}(\ell_g, \ell_h) = \int_0^T \left(\int_0^T g(s)(s \wedge t)\, ds \right) h(t)\, dt.$$

View the inner integral as a function of t and denote it by Qg, where Q is the integral operator

$$Qg(t) = \int_0^T g(s)(s \wedge t)\, ds.$$

Then, by definition of the L^2 inner product,

$$\langle Qg, h \rangle = \int_0^T \left(\int_0^T g(s)(s \wedge t)\, ds \right) h(t)\, dt,$$

which identifies Q as the covariance operator.

19.4. Splitting the Integral Operator

Because the minimum $s \wedge t$ changes its form according to whether $s \leq t$ or $s > t$, split the integral over $[0, T]$ into the intervals $[0, t]$ and $[t, T]$. On $[0, t]$, one has $s \wedge t = s$, while on $[t, T]$, one has $s \wedge t = t$. Therefore,

$$Qg(t) = \int_0^t g(s)s\, ds + t \int_t^T g(s)\, ds.$$

This is a useful explicit form of the covariance operator.

This operator is closely related to the fundamental solution of the Laplace operator. Its boundary behavior can be read off from the explicit formula.

At $t = 0$,

$$Qg(0) = 0,$$

which is a Dirichlet boundary condition.

At the other endpoint, the derivative satisfies

$$(Qg)'(T) = 0,$$

which is a Neumann boundary condition.

Differentiate the split formula:

$$(Qg)'(t) = \frac{d}{dt} \left(\int_0^t g(s)s\, ds + t \int_t^T g(s)\, ds \right).$$

Using the fundamental theorem of calculus on the first term and the product rule on the second term gives

$$(Qg)'(t) = g(t)t + \frac{d}{dt} \left(t \int_t^T g(s)\, ds \right).$$

For the second term,

$$\frac{d}{dt}\left(t\int_t^T g(s)\,ds\right) = \int_t^T g(s)\,ds + t\frac{d}{dt}\left(\int_t^T g(s)\,ds\right) = \int_t^T g(s)\,ds - tg(t).$$

Hence,

$$(Qg)'(t) = g(t)t + \int_t^T g(s)\,ds - tg(t).$$

The terms $g(t)t$ and $-tg(t)$ cancel, so

$$(Qg)'(t) = \int_t^T g(s)\,ds.$$

Evaluating at $t = T$ yields

$$(Qg)'(T) = \int_T^T g(s)\,ds = 0,$$

which proves the Neumann boundary condition. Differentiating once more gives

$$(Qg)''(t) = -g(t),$$

showing that Q is the inverse of the Laplacian in this sense.

20. The Covariance Operator as the Inverse Laplacian

The derivative computation above now identifies the covariance operator more conceptually as the inverse of the negative Laplacian.

From the previous derivation,

$$(Qg)'(t) = \int_t^T g(s)\,ds.$$

Applying the fundamental theorem of calculus,

$$\frac{d}{dt}\int_t^T g(s)\,ds = -g(t).$$

Therefore, differentiating once more gives

$$(Qg)''(t) = -g(t).$$

This leads directly to the realization that

$$Q = (-\Delta)^{-1},$$

that is, the covariance operator for the standard Wiener measure is the inverse of the negative one-dimensional Laplacian.

This inversion is not understood without boundary conditions. A differential operator such as the Laplacian cannot simply be inverted on its own, since one must specify the boundary conditions. Here one denotes the Laplace operator by Δ_x and imposes Dirichlet boundary conditions. Probabilistically, this corresponds to a fixed starting point at 0, which matches the condition

$$Qg(0) = 0.$$

Different boundary conditions lead to different Gaussian path measures. In particular, pinning both endpoints leads to the Brownian bridge, which may be viewed as a related Gaussian measure on path space.

21. The Impossibility of Lebesgue Measure in Infinite Dimensions

The covariance discussion above highlights a broader infinite-dimensional phenomenon: there is no analogue of Lebesgue measure on an infinite-dimensional space. In finite dimensions, Lebesgue measure is translation-invariant, so a set keeps the same volume under translation. In infinite dimensions, one may take a small ball and rotate it along different coordinate directions; under a Lebesgue-type measure, all of these rotated balls would have the same volume, and one could produce infinitely many such disjoint balls inside a bounded region. This would lead to

$$\sum_{i=1}^{\infty} \mathrm{Vol}(\mathrm{ball}).$$

Because one is adding the same non-zero number infinitely many times, the sum diverges. It cannot be equal to 1.

Appendix

A.1. Brownian Motion

Definition 3 (Brownian motion)**.** Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$ be a filtered probability space. A process $(W_t)_{t\geq 0}$ is a standard Brownian motion if:

- $W_0 = 0$ almost surely;
- for every $0 \leq s < t$, $W_t - W_s \sim \mathcal{N}(0, t-s)$;
- increments on disjoint time intervals are independent;
- sample paths are continuous.

In this lecture, the particle position is modeled by $X_t^i = B_t^i + x_0$, where each B^i is a Brownian motion.

A.2. Empirical Average

Definition 4 (General empirical average)**.** Let E be a measurable space, let $Y^1, \ldots, Y^N$ be E-valued random variables, and let $f : E \to \mathbb{R}$ be bounded measurable. The empirical average is the random variable

$$\eta^N(f) := \frac{1}{N} \sum_{i=1}^{N} f(Y^i).$$

For a fixed realization ω, this becomes the realized empirical average

$$\eta^N(f)(\omega) := \frac{1}{N} \sum_{i=1}^{N} f\big(Y^i(\omega)\big).$$

This quantity is the sample mean of the observable f evaluated on the N particles. Thus, when we simply say "empirical average," we usually mean the random variable $\eta^N(f)$, while $\eta^N(f)(\omega)$ denotes the realized average for one sample ω. In this lecture, we take

$$Y^i = X_t^i := B_t^i + x_0,$$

so

$$\eta_t^N(f) = \frac{1}{N}\sum_{i=1}^{N} f(B_t^i + x_0),$$

and for a fixed sample ω,

$$\eta_t^N(f)(\omega) = \frac{1}{N}\sum_{i=1}^{N} f\big(B_t^i + x_0\big)(\omega).$$

Thus $\eta_t^N(f)$ is the particle-system empirical average of f at time t.

A.3. Empirical Measure

Definition 5 (Empirical measure)**.** For $Y^1, \ldots, Y^N$ as above, the empirical measure is the random measure

$$\mu^N := \frac{1}{N}\sum_{i=1}^{N} \delta_{Y^i}.$$

For a fixed realization ω, this becomes the realized empirical measure

$$\mu^N(\omega) := \frac{1}{N}\sum_{i=1}^{N} \delta_{Y^i(\omega)}.$$

Then

$$\eta^N(f)(\omega) = \int_E f(y)\, \mu^N(\omega)(dy).$$

Thus, the empirical average is obtained by testing the empirical measure against the function f. In this lecture, at time t,

$$\mu_t^N := \frac{1}{N}\sum_{i=1}^{N} \delta_{B_t^i + x_0},$$

and for a fixed sample ω,

$$\mu_t^N(\omega) := \frac{1}{N}\sum_{i=1}^{N} \delta_{B_t^i(\omega) + x_0}.$$

Then

$$\eta_t^N(f)(\omega) = \int_{\mathbb{R}} f(x)\, \mu_t^N(\omega)(dx).$$

A.4. Strong Law of Large Numbers (SLLN)

Theorem 2 (SLLN)**.** Let $(Z_i)_{i\geq 1}$ be i.i.d. random variables with $\mathbb{E}[|Z_1|] < \infty$. Then

$$\frac{1}{N}\sum_{i=1}^{N} Z_i \xrightarrow[N\to\infty]{\text{a.s.}} \mathbb{E}[Z_1].$$

This says that the sample average converges almost surely to the true expectation.

Remark 1 (Application in this lecture). For fixed t and fixed test function f, take

$$Z_i := f(B_t^i + x_0).$$

Since $(B_t^i)_i$ are independent and identically distributed, so are $(Z_i)_i$. Therefore

$$\eta_t^N(f) = \frac{1}{N}\sum_{i=1}^{N} f(B_t^i + x_0) \xrightarrow[N\to\infty]{\text{a.s.}} \mathbb{E}[f(B_t + x_0)].$$

This is exactly the convergence used earlier in the notes.

A.5. Types of Convergence for Random Variables

Definition 6 (Almost sure convergence)**.** We say $X_n \to X$ almost surely (a.s.) if

$$\mathbb{P}\Big(\big\{\omega : \lim_{n\to\infty} X_n(\omega) = X(\omega)\big\}\Big) = 1.$$

In words: for almost every outcome ω, if we look at the sequence $X_n(\omega)$, it converges pointwise to $X(\omega)$.

Definition 7 (Convergence in probability)**.** We say $X_n \to X$ in probability if for every $\varepsilon > 0$,

$$\lim_{n\to\infty} \mathbb{P}(|X_n - X| > \varepsilon) = 0.$$

So if you fix any error tolerance ε, the chance that X_n is farther than ε from X goes to zero as n increases.

Definition 8 (Convergence in L^p)**.** For $p \geq 1$, we say $X_n \to X$ in L^p if

$$\lim_{n\to\infty} \mathbb{E}\big[|X_n - X|^p\big] = 0.$$

You can think of this as mean convergence with a p-power penalty on the error; larger p puts more weight on large deviations.

Definition 9 (Convergence in distribution)**.** We say $X_n \to X$ in distribution (or weakly) if

$$\lim_{n\to\infty} \mathbb{E}[\phi(X_n)] = \mathbb{E}[\phi(X)]$$

for every bounded continuous test function ϕ. Equivalently, if F_n and F are the distribution functions of X_n and X, then

$$\lim_{n\to\infty} F_n(x) = F(x)$$

at every continuity point x of F. Here we are not tracking sample paths directly; we are saying that the probability laws themselves converge.

Remark 2 (Standard implications). The main implication chain is

$$X_n \to X \text{ in } L^p \implies X_n \to X \text{ in probability} \implies X_n \to X \text{ in distribution.}$$

Also, almost sure convergence implies convergence in probability. When teaching this, a useful rule of thumb is: stronger notions (like a.s. or L^p) usually give weaker ones (probability, then distribution).

A.6. Expectation

Definition 10 (Expectation)**.** Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $X : \Omega \to \mathbb{R}$ be an integrable random variable (that is, $\mathbb{E}[|X|] < \infty$). Its expectation is

$$\mathbb{E}[X] := \int_\Omega X(\omega)\, \mathbb{P}(d\omega).$$

If X has distribution μ_X on $\mathbb{R}$, then equivalently

$$\mathbb{E}[X] = \int_\mathbb{R} x\, \mu_X(dx).$$

More generally, for measurable f with $\mathbb{E}[|f(X)|] < \infty$,

$$\mathbb{E}[f(X)] = \int_\mathbb{R} f(x)\, \mu_X(dx).$$

In our lecture notation, this is exactly the form

$$\mathbb{E}[f(B_t + x_0)] = \int_\mathbb{R} f(x)\, p(t, x)\, dx.$$

A.7. Distribution μ_X

Definition 11 (Distribution (probability law) of a random variable)**.** Let $X : (\Omega, \mathcal{F}, \mathbb{P}) \to \mathbb{R}$ be a random variable. Its distribution (or probability law, equivalently probability distribution) is the measure μ_X on $\mathbb{R}$ defined by

$$\mu_X(A) := \mathbb{P}(X \in A),$$

for every Borel set $A \subseteq \mathbb{R}$.

So "distribution of X," "probability law of X," "probability distribution of X," and "the probability measure induced by X" all refer to this same measure μ_X. So μ_X tells us how the values of X are distributed, without referring to the sample space directly.

Remark 3. For any bounded measurable f,

$$\int_\mathbb{R} f(x)\, \mu_X(dx) = \mathbb{E}[f(X)].$$

This is why we can rewrite expectations as integrals with respect to μ_X.

Remark 4 (Pushforward measure viewpoint). The same measure can be written as

$$\mu_X = \mathbb{P} \circ X^{-1},$$

read as "the pushforward of $\mathbb{P}$ by X." Concretely, this means

$$(\mathbb{P} \circ X^{-1})(A) = \mathbb{P}\big(X^{-1}(A)\big) = \mathbb{P}(X \in A),$$

which is exactly the definition of μ_X above.

A.8. Density

Definition 12 (Density of a random variable)**.** Let X be a real-valued random variable with law μ_X. We say that X has a density p_X if

$$\mu_X(A) = \int_A p_X(x)\,dx$$

for every Borel set $A \subseteq \mathbb{R}$. Equivalently, μ_X is absolutely continuous with respect to Lebesgue measure λ (written $\mu_X \ll \lambda$), and p_X is the Radon–Nikodym derivative:

$$p_X = \frac{d\mu_X}{d\lambda},$$

which describes how the probability mass is distributed over $\mathbb{R}$.

Remark 5. If X has density p_X, then for every bounded measurable f,

$$\mathbb{E}[f(X)] = \int_{\mathbb{R}} f(x)\,p_X(x)\,dx.$$

In our lecture, $X = B_t + x_0$ has Gaussian density

$$p(t,x) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{(x-x_0)^2}{2t}\right),$$

so

$$\mathbb{E}[f(B_t + x_0)] = \int_{\mathbb{R}} f(x)\,p(t,x)\,dx.$$

A.9. Meaning of $\delta_{x_0}(dx)$

Definition 13 (Dirac measure at x_0)**.** The symbol δ_{x_0} denotes the Dirac measure concentrated at the point x_0. It is defined by

$$\delta_{x_0}(A) = \mathbf{1}_A(x_0)$$

for every Borel set $A \subseteq \mathbb{R}$. Equivalently,

$$\delta_{x_0}(A) = \begin{cases} 1, & \text{if } x_0 \in A, \\ 0, & \text{if } x_0 \notin A. \end{cases}$$

Remark 6. Writing $\delta_{x_0}(dx)$ emphasizes that this is a measure in the variable x. Its key property is

$$\int_{\mathbb{R}} f(x)\,\delta_{x_0}(dx) = f(x_0).$$

So in our note,

$$p(t,x)\,dx \to \delta_{x_0}(dx) \quad (t \downarrow 0)$$

means the probability law with density $p(t,x)$ converges to a point mass at x_0.

A.10. Notation for η_t^N, η_t, and Their Densities

Remark 7 (Notation guide). Several closely related symbols appear throughout these notes:

$$\eta_t^N(f) = \frac{1}{N}\sum_{i=1}^{N} f(B_t^i + x_0).$$

This is the empirical average of the test function f over the N particles at time t.

The corresponding empirical measure is

$$\eta_t^N(dx) = \frac{1}{N}\sum_{i=1}^{N} \delta_{B_t^i + x_0}(dx).$$

Thus $\eta_t^N(f)$ means the empirical measure tested against f:

$$\eta_t^N(f) = \int_{\mathbb{R}} f(x)\, \eta_t^N(dx).$$

In the macroscopic limit, the deterministic limiting measure is denoted by

$$\eta_t(dx) = p(t,x)\, dx.$$

Accordingly,

$$\eta_t(f) = \int_{\mathbb{R}} f(x)\, \eta_t(dx) = \int_{\mathbb{R}} f(x)\, p(t,x)\, dx.$$

Whenever we formally write $\eta_t^N(x)$ or $\eta_t(x)$, this means that the corresponding measure is being represented by a density with respect to Lebesgue measure:

$$\eta_t^N(dx) = \eta_t^N(x)\, dx, \qquad \eta_t(dx) = \eta_t(x)\, dx.$$

In particular, for the limiting measure one has

$$\eta_t(x) = p(t,x).$$

Here η_t denotes the measure, while $\eta_t(x)$ denotes its density with respect to Lebesgue measure. Thus, $\eta_t(x) = p(t,x)$ means that the density of the limiting measure η_t is exactly the heat kernel $p(t,x)$.

A.11. Space-Time White Noise

Definition 14 (Space-time white noise)**.** Space-time white noise is a generalized Gaussian field $\xi(t,x)$ with covariance

$$\mathbb{E}[\xi(t,x)\xi(s,y)] = \delta(t-s)\delta(x-y).$$

This means that the noise is formally uncorrelated in both time and space.

Remark 8. The symbol $\xi(t,x)$ should not be understood as an ordinary function of (t,x). It is a distribution-valued object, so expressions such as $\xi(t,x)$ or $d\xi(t,x)$ are formal notations for stochastic integration against this noise.

Remark 9. Formally, one may think of $\xi(t,x)$ as the derivative of a Brownian sheet $W(t,x)$:

$$\xi(t,x) = \partial_t \partial_x W(t,x).$$

This identity is only symbolic, because the Brownian sheet is not classically differentiable in either variable.

Remark 10. Accordingly, an expression of the form

$$\int_0^t \int_{\mathbb{R}} g(s,x)\, d\xi(s,x)$$

denotes stochastic integration with respect to space-time white noise. Its quadratic variation is given formally by

$$\left\langle \int_0^t \int_{\mathbb{R}} g(s,x)\, d\xi(s,x) \right\rangle_t = \int_0^t \int_{\mathbb{R}} g(s,x)^2\, dx\, ds.$$

A.12. Brownian Sheet

Definition 15 (Brownian sheet)**.** A Brownian sheet is a two-parameter Gaussian random field

$$W(t,x), \qquad t \geq 0,\ x \geq 0,$$

with mean zero and covariance

$$\mathbb{E}[W(t,x)W(s,y)] = (t \wedge s)(x \wedge y).$$

Remark 11. It is the natural two-parameter analogue of Brownian motion. While Brownian motion is indexed by one variable, usually time, the Brownian sheet is indexed by both time and space and can be viewed heuristically as a random surface.

Remark 12. Space-time white noise can be understood formally as the mixed derivative of the Brownian sheet:

$$\xi(t,x) = \partial_t \partial_x W(t,x).$$

This identity is only symbolic, since the Brownian sheet is not classically differentiable in either variable.

A.13. Quadratic Variation Identities Used in the Fluctuation Argument

Lemma 1. For the martingale term

$$M_t^N(f) = \frac{1}{N} \sum_{i=1}^N \int_0^t f'(B_s^i + x_0)\, dB_s^i,$$

one has

$$\left\langle \frac{1}{N} \sum_{i=1}^N \int_0^t f'(B_s^i + x_0)\, dB_s^i \right\rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

Moreover,

$$\left\langle \frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s \right\rangle_t = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

Proof. For the first identity, use that the Brownian motions $B^1, \dots, B^N$ are independent. Hence the cross-variations vanish, and

$$\left\langle \frac{1}{N} \sum_{i=1}^{N} \int_0^{\cdot} f'(B_s^i + x_0)\, dB_s^i \right\rangle_t = \frac{1}{N^2} \sum_{i=1}^{N} \left\langle \int_0^{\cdot} f'(B_s^i + x_0)\, dB_s^i \right\rangle_t .$$

The quadratic variation of each Itô integral is the time integral of the squared integrand, so

$$= \frac{1}{N^2} \sum_{i=1}^{N} \int_0^t f'(B_s^i + x_0)^2\, ds.$$

Recognizing the empirical measure in the sum gives

$$= \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

For the second identity, apply the general rule

$$\left\langle \int_0^t H_s\, dB_s \right\rangle_t = \int_0^t H_s^2\, ds$$

with

$$H_s = \frac{1}{\sqrt{N}} \sqrt{\eta_s^N((f')^2)}.$$

Then

$$\left\langle \frac{1}{\sqrt{N}} \int_0^t \sqrt{\eta_s^N((f')^2)}\, dB_s \right\rangle_t = \int_0^t \frac{1}{N} \eta_s^N((f')^2)\, ds = \frac{1}{N} \int_0^t \eta_s^N((f')^2)\, ds.$$

This proves the claim. □

A.14. Random Elements Distributed According to a Measure

Definition 16 (Law of a random element)**.** Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let

$$g : \Omega \to H$$

be an H-valued random element. Its law is the probability measure μ on $(H, \mathcal{B}(H))$ defined by

$$\mu(A) = \mathbb{P}(g \in A), \qquad A \in \mathcal{B}(H).$$

Remark 13. Saying that g is distributed according to μ means exactly that the law of g is μ. In other words, for every Borel set $A \subset H$,

$$\mathbb{P}(g \in A) = \mu(A).$$

Thus μ describes how the random point g is distributed over the Hilbert space.

Remark 14. If $h \in H$ is fixed, then

$$\ell_h(g) = \langle g, h \rangle_H$$

is a real-valued random variable obtained by projecting the random element g onto the direction h. This is the object whose distribution is required to be Gaussian in the definition of a Gaussian measure on a Hilbert space.

A.15. The First Two Identities for the Gaussian Measure

Lemma 2. Let $\mu = \mathcal{N}(m, Q)$ on $(H, \mathcal{B}(H))$. Then for all $h, h_1, h_2 \in H$,

$$\int_H \langle x, h\rangle_H \, \mu(dx) = \langle m, h\rangle_H$$

and

$$\int_H \big(\langle x, h_1\rangle_H - \langle m, h_1\rangle_H\big)\big(\langle x, h_2\rangle_H - \langle m, h_2\rangle_H\big) \, \mu(dx) = \langle Qh_1, h_2\rangle_H.$$

Proof. Fix $h \in H$. By the characteristic function formula,

$$\varphi_h(t) := \int_H e^{it\langle x,h\rangle_H} \, \mu(dx) = e^{it\langle m,h\rangle_H - \frac{1}{2}t^2\langle Qh,h\rangle_H}.$$

Differentiating at $t = 0$ gives

$$\varphi_h'(0) = i \int_H \langle x, h\rangle_H \, \mu(dx) = i\langle m, h\rangle_H.$$

Therefore,

$$\int_H \langle x, h\rangle_H \, \mu(dx) = \langle m, h\rangle_H.$$

For the covariance identity, fix $h_1, h_2 \in H$ and consider the centered random variables

$$Y_j(x) := \langle x, h_j\rangle_H - \langle m, h_j\rangle_H, \qquad j = 1, 2.$$

For $s, t \in \mathbb{R}$, the characteristic function of $sY_1 + tY_2$ is

$$\int_H e^{i(sY_1(x)+tY_2(x))} \, \mu(dx) = e^{-\frac{1}{2}\langle Q(sh_1+th_2), sh_1+th_2\rangle_H}.$$

Expanding the quadratic form gives

$$\langle Q(sh_1 + th_2), sh_1 + th_2\rangle_H = s^2\langle Qh_1, h_1\rangle_H + 2st\langle Qh_1, h_2\rangle_H + t^2\langle Qh_2, h_2\rangle_H.$$

Differentiating the characteristic function once in s and once in t at $(0, 0)$ yields

$$-\int_H Y_1(x)Y_2(x) \, \mu(dx) = -\langle Qh_1, h_2\rangle_H.$$

Hence

$$\int_H \big(\langle x, h_1\rangle_H - \langle m, h_1\rangle_H\big)\big(\langle x, h_2\rangle_H - \langle m, h_2\rangle_H\big) \, \mu(dx) = \langle Qh_1, h_2\rangle_H.$$

This proves the claim. □

A.16. Gaussian Martingales

Definition 17 (Gaussian martingale)**.** A continuous martingale $(M_t)_{t\geq 0}$ is called a Gaussian martingale if, for every finite collection of times

$$0 \leq t_1 < \cdots < t_n,$$

the random vector $(M_{t_1}, \ldots, M_{t_n})$ has a multivariate Gaussian distribution.

Remark 15. Brownian motion is the standard example of a Gaussian martingale. More generally, if

$$M_t = \int_0^t H_s \, dB_s$$

with deterministic square-integrable integrand H, then M is again a Gaussian martingale and

$$\langle M \rangle_t = \int_0^t H_s^2 \, ds.$$

If the bracket is not equal to t, then M is usually not a Brownian motion, even though it is still Gaussian.

Remark 16. In the fluctuation argument of this lecture, the limit

$$M_t(f) := \int_0^t \sqrt{\eta_s((f')^2)} \, dB_s$$

is described as a Gaussian martingale. For each fixed time t, it is a centered normal random variable with variance

$$\mathrm{Var}(M_t(f)) = \int_0^t \eta_s((f')^2) \, ds,$$

and as a process its covariance structure is determined by its quadratic variation.

A.17. Characteristic Functions in Finite and Infinite Dimensions

Remark 17. If X is an $\mathbb{R}^n$-valued random variable, then its characteristic function is

$$\varphi_X(\xi) = \mathbb{E}\big[e^{i\xi \cdot X}\big], \qquad \xi \in \mathbb{R}^n.$$

For a Gaussian random vector $X \sim \mathcal{N}(m, Q)$ on $\mathbb{R}^n$, this becomes

$$\varphi_X(\xi) = e^{i\xi \cdot m - \frac{1}{2}\xi \cdot Q\xi}.$$

Remark 18. On a Hilbert space H, one uses the same idea, but the role of the finite-dimensional parameter ξ is taken by a direction $h \in H$. If μ is a probability measure on H, then the characteristic function of the one-dimensional projection

$$\ell_h(g) = \langle g, h \rangle_H$$

is

$$\int_H e^{i\langle g,h \rangle_H} \, \mu(dg).$$

If $\mu = \mathcal{N}(m, Q)$ is Gaussian on H, then

$$\int_H e^{i\langle g,h \rangle_H} \, \mu(dg) = e^{i\langle m,h \rangle_H - \frac{1}{2}\langle Qh,h \rangle_H}.$$

Thus, the infinite-dimensional formula is the Hilbert-space analogue of the finite-dimensional Gaussian characteristic function, with the scalar product replacing the Euclidean dot product.