

Combining Measure Theory and Bayes' Rule to Solve a Stochastic Inverse Problem

Tim Wildey♦, Troy Butler♦, John Jakeman♣

♦ Sandia National Laboratories
Center for Computing Research
Optimization and UQ Group

♦ University of Colorado Denver
Department of Mathematics and Statistics

EMI/PMC 2021
May 25-28, 2021
Virtual

Sandia National Laboratories is a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA-0003525. SAND2021 - **** C

Motivation






Figure: Additive manufacturing and high-throughput testing provides new data science challenges.


A Stochastic Inverse Problem



Problem

Given a probability density on observations, find a probability density on X such that the push-forward matches the given density on the observed data.

A Stochastic Inverse Problem



Problem

Given a probability density on observations, find a probability density on X such that the push-forward matches the given density on the observed data.

- Solutions may not be unique without additional assumptions.
- **We only need to solve a single stochastic forward problem.**

Notation

We assume we are given:

- ① A finite-dimensional **parameter space**, X .
- ② A **parameter-to-observation/data map**, $f : X \rightarrow \mathcal{D} = f(X)$
- ③ An **observed probability measure** on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, denoted $\mathbb{P}_{\mathcal{D}}^{\text{obs}}$, that has a density, $\pi_{\mathcal{D}}^{\text{obs}}$.
- ④ An **initial probability measure** on (X, \mathcal{B}_X) , denoted $\mathbb{P}_X^{\text{init}}$, that has a density, π_X^{init} .

Notation

We assume we are given:

- ① A finite-dimensional **parameter space**, X .
- ② A **parameter-to-observation/data map**, $f : X \rightarrow \mathcal{D} = f(X)$
- ③ An **observed probability measure** on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, denoted $\mathbb{P}_{\mathcal{D}}^{\text{obs}}$, that has a density, $\pi_{\mathcal{D}}^{\text{obs}}$.
- ④ An **initial probability measure** on (X, \mathcal{B}_X) , denoted $\mathbb{P}_X^{\text{init}}$, that has a density, π_X^{init} .


We need to compute:

- ① The **push-forward of the initial density** through the model.
 - In other words, **we need to solve a forward UQ problem using the initial.**
 - We use $\pi_{\mathcal{D}}^{f(\text{init})}$ to denote this push-forward density.

A Key Assumption

Predictability Assumption

We assume that the observed probability measure, $\mathbb{P}_{\mathcal{D}}^{\text{obs}}$, is absolutely continuous with respect to the push-forward of the initial, $\mathbb{P}_{\mathcal{D}}^{f(\text{init})}$.



A Solution to the Stochastic Inverse Problem

Theorem

Given an initial probability measure, \mathbb{P}_X^{init} on (X, \mathcal{B}_X) and an observed probability measure, $\mathbb{P}_{\mathcal{D}}^{obs}$, on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, the probability measure P_X^{up} on (X, \mathcal{B}_X) defined by

$$\mathbb{P}_X^{up}(A) = \int_{\mathcal{D}} \left(\int_{A \cap f^{-1}(q)} \pi_X^{init}(x) \frac{\pi_{\mathcal{D}}^{obs}(f(x))}{\pi_{\mathcal{D}}^{f(init)}(f(x))} d\mu_{X,q}(x) \right) d\mu_{\mathcal{D}}(q), \quad \forall A \in \mathcal{B}_X$$

solves the stochastic inverse problem.

A Solution to the Stochastic Inverse Problem

Theorem

Given an initial probability measure, \mathbb{P}_X^{init} on (X, \mathcal{B}_X) and an observed probability measure, $\mathbb{P}_{\mathcal{D}}^{obs}$, on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, the probability measure P_X^{up} on (X, \mathcal{B}_X) defined by

$$\mathbb{P}_X^{up}(A) = \int_{\mathcal{D}} \left(\int_{A \cap f^{-1}(q)} \pi_X^{init}(x) \frac{\pi_{\mathcal{D}}^{obs}(f(x))}{\pi_{\mathcal{D}}^{f(init)}(f(x))} d\mu_{X,q}(x) \right) d\mu_{\mathcal{D}}(q), \quad \forall A \in \mathcal{B}_X$$

solves the stochastic inverse problem.

Corollary

The updated measure of X is 1.

A Solution to the Stochastic Inverse Problem

Theorem

Given an initial probability measure, \mathbb{P}_X^{init} on (X, \mathcal{B}_X) and an observed probability measure, $\mathbb{P}_{\mathcal{D}}^{obs}$, on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, the probability measure \mathbb{P}_X^{up} on (X, \mathcal{B}_X) defined by

$$\mathbb{P}_X^{up}(A) = \int_{\mathcal{D}} \left(\int_{A \cap f^{-1}(q)} \pi_X^{init}(x) \frac{\pi_{\mathcal{D}}^{obs}(f(x))}{\pi_{\mathcal{D}}^{f(init)}(f(x))} d\mu_{X,q}(x) \right) d\mu_{\mathcal{D}}(q), \quad \forall A \in \mathcal{B}_X$$

solves the stochastic inverse problem.

Corollary

The updated measure of X is 1.

Theorem

\mathbb{P}_X^{up} is stable with respect to perturbations in $\mathbb{P}_{\mathcal{D}}^{obs}$ and in \mathbb{P}_X^{init} .

A Solution to the Stochastic Inverse Problem

Theorem

Given an initial probability measure, \mathbb{P}_X^{init} on (X, \mathcal{B}_X) and an observed probability measure, $\mathbb{P}_{\mathcal{D}}^{obs}$, on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, the probability measure P_X^{up} on (X, \mathcal{B}_X) defined by

$$\mathbb{P}_X^{up}(A) = \int_{\mathcal{D}} \left(\int_{A \cap f^{-1}(q)} \pi_X^{init}(x) \frac{\pi_{\mathcal{D}}^{obs}(f(x))}{\pi_{\mathcal{D}}^{f(init)}(f(x))} d\mu_{X,q}(x) \right) d\mu_{\mathcal{D}}(q), \quad \forall A \in \mathcal{B}_X$$

solves the stochastic inverse problem.

Corollary

The updated measure of X is 1.

Theorem

\mathbb{P}_X^{up} is stable with respect to perturbations in $\mathbb{P}_{\mathcal{D}}^{obs}$ and in \mathbb{P}_X^{init} .

For details: [Combining Push-forward Measures and Bayes' Rule to Construct Consistent Solutions to Stochastic Inverse Problems, BJW. SISC 40 (2), 2018.]

A Solution to the Stochastic Inverse Problem

Theorem

Given an initial probability measure, \mathbb{P}_X^{init} on (X, \mathcal{B}_X) and an observed probability measure, $\mathbb{P}_{\mathcal{D}}^{obs}$, on $(\mathcal{D}, \mathcal{B}_{\mathcal{D}})$, the probability measure P_X^{up} on (X, \mathcal{B}_X) defined by

$$\mathbb{P}_X^{up}(A) = \int_{\mathcal{D}} \left(\int_{A \cap f^{-1}(q)} \pi_X^{init}(x) \frac{\pi_{\mathcal{D}}^{obs}(f(x))}{\pi_{\mathcal{D}}^{f(init)}(f(x))} d\mu_{X,q}(x) \right) d\mu_{\mathcal{D}}(q), \quad \forall A \in \mathcal{B}_X$$

solves the stochastic inverse problem.

The updated density is:

$$\pi_X^{up}(x) = \pi_X^{init}(x) \frac{\pi_{\mathcal{D}}^{obs}(f(x))}{\pi_{\mathcal{D}}^{f(init)}(f(x))}.$$

- Both π_X^{init} and $\pi_{\mathcal{D}}^{obs}$ are given.
- Computing $\pi_{\mathcal{D}}^{f(init)}$ requires a forward propagation of the initial density.

A Parameterized Nonlinear System

Example

Consider a parameterized nonlinear system of equations:

$$\begin{aligned}x_1 u_1^2 + u_2^2 &= 1, \\u_1^2 - x_2 u_2^2 &= 1\end{aligned}$$

- The quantity of interest is the second component: $f(x) = u_2$.
- Assume that we observe $\pi_{\mathcal{D}}^{\text{obs}} \sim N(0.3, 0.025^2)$.
- We consider a uniform initial density.
- We use 10,000 samples from the initial and a standard KDE to approximate the push-forward.

A Parameterized Nonlinear System





Figure: Samples from the updated density (left) and a comparison of $\pi_{\mathcal{D}}^{\text{obs}}$, $\pi_{\mathcal{D}}^{f(\text{init})}$ and $\pi_{\mathcal{D}}^{f(\text{up})}$ (right).

Additional demonstrations and interactive lecture materials can be found at
<https://github.com/eecsuh/SIAM-AN18-Tutorial>.

Relationship with Statistical Bayesian Inference

Using Bayes theorem we can define a different posterior density [Stuart 2010; Gelman et al 2013; Jaynes 1998, ...]:

$$\tilde{\pi}_X^{\text{post}}(\lambda|q) = \pi_X^{\text{init}}(\lambda) \frac{\pi(q|\lambda)}{C}.$$

Relationship with Statistical Bayesian Inference

Using Bayes theorem we can define a different posterior density [Stuart 2010; Gelman et al 2013; Jaynes 1998, ...]:

$$\tilde{\pi}_X^{\text{post}}(\lambda|q) = \pi_X^{\text{init}}(\lambda) \frac{\pi(q|\lambda)}{C}.$$

Example

Let $X = [-1, 1]$ and consider the simple nonlinear map

$$q(\lambda) = \lambda^p, \quad p = 1, 3, 5, \dots$$

Here, p is not uncertain and are used to vary the nonlinearity.

Relationship with Statistical Bayesian Inference

Using Bayes theorem we can define a different posterior density [Stuart 2010; Gelman et al 2013; Jaynes 1998, ...]:

$$\tilde{\pi}_X^{\text{post}}(\lambda|q) = \pi_X^{\text{init}}(\lambda) \frac{\pi(q|\lambda)}{C}.$$

Example


Let $X = [-1, 1]$ and consider the simple nonlinear map

$$q(\lambda) = \lambda^p, \quad p = 1, 3, 5, \dots$$


Here, p is not uncertain and are used to vary the nonlinearity.

- Assume a uniform prior and the observed density is given by $\pi_D^{\text{obs}} \sim N(0.25, 0.1^2)$.
- For the statistical Bayesian approach, we use an observed value of $\hat{q} = 0.25$ and assume a Gaussian noise model $\eta \sim N(0, 0.1^2)$.

Comparing Push-forwards for Linear and Nonlinear Maps




Linear map ($p = 1$)




Nonlinear map ($p = 5$)

Comparing Push-forwards for Linear and Nonlinear Maps



Linear map ($p = 1$)



Nonlinear map ($p = 5$)

The Bayesian and measure-theoretic formulations **solve different problems, have different posteriors and make different predictions.**

A Computational Mechanics Example




Figure: On the left, an illustration of the computational model on a coarse mesh (16,600 elements). In the middle, the granular microstructure on a finer mesh (≈ 17 million elements). On the right, the vertical displacement using the high-fidelity model and nominal parameter values.

A Computational Mechanics Example


- We assume the Young's modulus, E , and Poisson ratio, ν , are random.
- Each grain has a random orientation defined by 4 independent Gaussian parameters.

A Computational Mechanics Example

- We assume the Young's modulus, E , and Poisson ratio, ν , are random.
- Each grain has a random orientation defined by 4 independent Gaussian parameters.
- Model has $\approx 225,000$ grains $\implies \approx 900,002$ random variables!
- Do we really need to account for the microstructure???

A Computational Mechanics Example

- We assume the Young's modulus, E , and Poisson ratio, ν , are random.
- Each grain has a random orientation defined by 4 independent Gaussian parameters.
- Model has $\approx 225,000$ grains $\implies \approx 900,002$ random variables!
- Do we really need to account for the microstructure???



A Computational Mechanics Example








Figure: Samples from the updated density (left) and the comparison of $\pi_{\mathcal{D}}^{\text{obs}}$, $\pi_{\mathcal{D}}^{f(\text{init})}$ and $\pi_{\mathcal{D}}^{f(\text{up})}$ (right) using a 1-dimensional active subspace with $N = 100$ and $M = 10,000$.

A Stochastic Inverse Problem in Structure-Property Linkages



For details: [Solving stochastic inverse problems for property-structure linkages using data-consistent inversion and machine learning, A. Tran and T. Wildey, JOM, 2020.]

A Stochastic Inverse Problem in Structure-Property Linkages



Conclusions

- Our goal is to develop **data-informed physics-based** models.
- Many approaches exist for incorporating data into a model.
 - Deterministic optimization, Bayesian methods, OUU, data assimilation, etc.
- Our approach provides a solution to a specific stochastic inverse problem.
- Main computational expense is the forward UQ problem to obtain the push-forward of the initial density.

Thanks! Questions?

Acknowledgments

This material is based upon work supported by the U.S. Department of Energy, Office of Science, ASCR, Early Career Research Program.

Thank you for your attention!
Questions?