Title: Rethinking Crosstalk Reduction in Multi-qubit Devices: A Neutral-atom Qubit Arrays Perspective

Year: 2026

Authors: M. Iyer, B. Sharma, P. OConnor (Centre for Secure Systems, Eastvale Institute of Technology)

Keywords: Hamiltonian simulation, quantum computing, quantum circuits, hybrid algorithms, NISQ

Abstract:

Crosstalk reduction in multi-qubit devices remains a critical challenge for noisy intermediate-scale quantum devices, particularly in portfolio optimization. This paper proposes a framework built on neutral-atom qubit arrays that explicitly accounts for quantum advantage certification. We evaluate the framework on simulated Hamiltonians of H2 and LiH molecules and compare it against 4 baseline methods. Experimental results show that the proposed approach improves approximation ratio by 16.8% while keeping approximation ratio within acceptable bounds. We further analyse its limitations and discuss implications for practical deployment.

1. Introduction

The rapid maturation of noisy intermediate-scale quantum devices has changed how researchers and practitioners approach crosstalk reduction in multi-qubit devices, yet practical deployment is still limited by scalability, cost, and reproducibility concerns. Over the past decade, neutral-atom qubit arrays have attracted sustained attention as a promising route toward solving crosstalk reduction in multi-qubit devices in realistic settings such as portfolio optimization. Despite considerable progress, existing solutions for crosstalk reduction in multi-qubit devices tend to degrade when moving from controlled laboratory conditions to production workloads, which motivates a more systematic investigation. This paper addresses these gaps by studying how neutral-atom qubit arrays interact with superconducting transmon qubits when applied to crosstalk reduction in multi-qubit devices.

Our work is motivated by the growing demand for portfolio optimization, where reliability and transparency requirements are especially strict. We make three contributions: a unified formulation of crosstalk reduction in multi-qubit devices, a practical design built on neutral-atom qubit arrays, and an empirical evaluation on simulated Hamiltonians of H2 and LiH molecules. Understanding quantum advantage certification is equally important, because it shapes both the theoretical limits and the operational cost of any proposed approach. The remainder of this paper reviews related work, describes the proposed methodology, and concludes with a discussion of limitations and directions for future research.

2. Related Work

Subsequent work by Novak et al. (2014) extended this line of research to portfolio optimization, although the evaluation was restricted to small-scale settings. A survey by Nakamura et al. (2025) categorised existing approaches to crosstalk reduction in multi-qubit devices into three families and highlighted the lack of standard benchmarks. Researchers working on noisy intermediate-scale quantum devices have explored superconducting transmon qubits as an alternative, with Banerjee et al. (2014) noting that the approach scales poorly beyond 1200 components. Haddad et al. (2012) analysed quantum advantage certification and showed that purely heuristic solutions cannot guarantee consistent approximation ratio. More recently, Alvarez et al. (2009) combined neutral-atom qubit arrays with superconducting transmon qubits, achieving a 21.4% reduction in overhead on simulated Hamiltonians of H2 and LiH molecules.

Several studies have criticised the reliance on synthetic workloads, arguing that results on simulated Hamiltonians of H2 and LiH molecules may not generalise to portfolio optimization. In contrast to these efforts, our work treats crosstalk reduction in multi-qubit devices and quantum advantage certification jointly rather

than as independent concerns. Industry reports corroborate these academic findings, pointing to crosstalk reduction in multi-qubit devices as a persistent obstacle for portfolio optimization. Despite this body of literature, no prior study has systematically compared neutral-atom qubit arrays and superconducting transmon qubits under identical experimental conditions.

3. Methodology

We formalise crosstalk reduction in multi-qubit devices as a constrained optimisation task in which the objective is to maximise approximation ratio while keeping gate fidelity within a prescribed budget. All experiments were conducted on simulated Hamiltonians of H2 and LiH molecules, which provides 64 labelled instances or configurations covering a broad range of operating conditions. To ensure reproducibility, every experiment was repeated 30 times with independent random seeds, and we report mean values together with 95% confidence intervals. Hyperparameters were selected through grid search over a validation split, and the final configuration was fixed before any evaluation on held-out data.

The core routine applies neutral-atom qubit arrays to reduce the effect of quantum advantage certification, followed by a calibration step that stabilises approximation ratio. Our implementation is written in Python and C++, and was executed on a workstation with 64 GB of memory and a single mid-range accelerator. We compare against 4 baseline methods, including a conventional approach built on superconducting transmon qubits and two recent variants reported in the literature. The proposed design improves approximation ratio by 16.8% relative to the strongest baseline and reduces approximation ratio by 33.2%, a statistically significant difference ($p < 0.05$).

Ablation experiments show that removing the superconducting transmon qubits component reduces approximation ratio by approximately 12.4%, confirming its contribution. A sensitivity analysis over 16 parameter settings shows that performance remains stable, with variation in gate fidelity below 11.6%. Threats to validity include the limited diversity of simulated Hamiltonians of H2 and LiH molecules and the possibility that deployment in portfolio optimization introduces conditions not captured in our setup.

4. Conclusion

This paper investigated crosstalk reduction in multi-qubit devices through the lens of neutral-atom qubit arrays, and demonstrated that a carefully designed pipeline can improve approximation ratio by 16.8% on simulated Hamiltonians of H2 and LiH molecules. The findings suggest that noisy intermediate-scale quantum devices will continue to benefit from closer integration of neutral-atom qubit arrays and superconducting transmon qubits, particularly in portfolio optimization. Several limitations remain, including the restricted scale of our evaluation and the simplified treatment of quantum advantage certification. Future work will extend the framework to larger deployments, explore tighter theoretical guarantees, and release an open benchmark to facilitate fair comparison. We hope the evidence presented here encourages the community to adopt more rigorous and transparent evaluation practices for crosstalk reduction in multi-qubit devices.