

Shortest Path with Multiples of k

Aniket Gupta (2022073)

Credits: Avi Sharma (2022119), Parth Sandeep Rastogi (2022352)

April 29, 2024

1 Problem Statement

Given a graph $G = (V, E)$, with vertices V , edges E , a source vertex s , a target vertex t , and edge weights $c(uv)$, where each weight $c(uv) > 0$ represents the cost associated with traveling from vertex u to v or from v to u , the objective is to find the shortest path from s to t such that the total cost of the path is a multiple of a given positive integer k .

2 Algorithm

The proposed algorithm is based on modifying the graph and applying Dijkstra's algorithm. Here's the pseudocode:

Algorithm 1 ShortestPathWithMultiplesOfK(G, s, t, k)

- 1: Let R be the set $\{0, 1, 2, \dots, k-1\}$
 - 2: Define a new graph $G' = (V', E')$
 - 3: where $V' = V \times R$
 - 4: $E' := \{(u, r) \rightarrow (v, (r + c(uv)) \bmod k) \mid (u, v) \in E, r \in R\}$
 - 5: Weight of each edge $(u, r) \rightarrow (v, r')$ is $c(uv)$
 - 6: Apply Dijkstra's algorithm on G' with source vertex $(s, 0)$
 - 7: **return** shortest distance from $(s, 0)$ to $(t, 0)$
-

3 Implementation

Below is the C++ implementation of the algorithm:

// Code snippet: <https://pastebin.com/Q1EU9tFc>

4 Time Complexity

The time complexity of the algorithm is $O(E' \log V') = O(mk \log(nk)) = O(m \log n)$, since k is treated as a constant factor.