

Discrete Mathematics and Logic (UE18CS205)

Unit 2 - Sets, Functions and Relations

Prof CB, Prof SVI, Prof RJ, Prof PRD, Prof SS, Prof KKV
Department of CS&E, PES University

Sets

- A **set** is an unordered collection of objects.
- Sets are discrete structures used to group objects together, often the objects having similar properties.
- The objects in a set are called the **elements** or **members** of the set.
- A set is said to **contain** its elements.
- An element is said to **belong** to the set.
- $a \in S$ denotes that “a” is an element of the set S.
- $a \notin S$ denotes that “a” is not an element of the set S.

Set roster form: All the members of the set are listed separated by commas enclosed in curly braces.

Eg: A set of natural numbers from 1 to 5.

$$S = \{1, 2, 3, 4, 5\}$$

Eg: the set of positive integers less than hundred

$$S = \{1, 2, 3, \dots, 99\}$$

Set builder form: Elements in the set are characterized by stating the property or properties they must have to be members.

Eg: the set of positive integers less than hundred

$$A = \{x \in \mathbf{Z}^+ \mid x < 100\}$$

Eg: set of rational numbers.

$$B = \{p/q \mid p \in \mathbf{Z}; q \in \mathbf{Z}, \text{ and } q \neq 0\}$$

Well known number sets:

- $\mathbf{N} = \{0, 1, 2, 3, \dots\}$, the set of natural numbers
- $\mathbf{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$, the set of integers
- $\mathbf{Z}^+ = \{1, 2, 3, 4, \dots\}$, set of positive integers
- $\mathbf{Q} = \{p/q \mid p \in \mathbf{Z} ; q \in \mathbf{Z} \text{ and } q \neq 0\}$ the set of rational numbers
- \mathbf{R} the set of real numbers
- \mathbf{C} the set of complex numbers

Equality of two sets

Two sets are equal if and only if they have the same elements.

i.e., if A and B are sets, then A and B are equal if and only if
 $\forall x (x \in A \leftrightarrow x \in B)$.

if A and B are equal sets, it can be denoted as **A = B**.

Eg: $\{a, b\} = \{a, b\}$

Eg: $\{1, 3, 5\} = \{3, 1, 5\}$





Eg: $\{1, 3, 3, 5, 5, 5, 5\} = \{1, 3, 5\}$

Eg: $\{1\} \neq \{\{1\}\}$

- Empty set / Null set
 $\{ \} = \Phi$
- Singleton set
 $\{ a \}, \{ Z^+ \}$
- $\Phi \neq \{\Phi\}$


Venn Diagram

Sets can be represented graphically using Venn diagrams.




Subsets:

- The set A is called a subset of set B if and only if every element of A is also an element of B .
- We use the notation $A \subseteq B$ to indicate that A is a subset of set B .
- $A \subseteq B$ if and only if $\forall x (x \in A \rightarrow x \in B)$ is true.



Proper Subset:

- When set A is a subset of set B but $A \neq B$, we write $A \subset B$ and say that A is a **proper subset** of B .
- For $A \subset B$ to be true it must be the case that $A \subseteq B$ and there must exist an element x of B that is not an element of A .
- That is, A is a proper subset of B if $\forall x (x \in A \rightarrow x \in B) \wedge \exists x (x \in B \wedge x \notin A)$ is true.



Theorem: For every set S ,

1. $\emptyset \subseteq S$
2. $S \subseteq S$

Proof of $\emptyset \subseteq S$:

Let S be a set.

$\emptyset \subseteq S$ iff $\forall x(x \in \emptyset \rightarrow x \in S)$.

Because the empty set contains no elements, it follows that $x \in \emptyset$ is always false.

It follows that the conditional statement $x \in \emptyset \rightarrow x \in S$ is always true, because the hypothesis is always false and a conditional statement with a false hypothesis is true.

That is $\forall x(x \in \emptyset \rightarrow x \in S)$ is true.

$\therefore \emptyset \subseteq S$

It's a vacuous proof.

Cardinality: If there are exactly n distinct elements in S where n is a non-negative integer, we say that S is a **finite set** and that n is the **cardinality** of S . The cardinality of S is denoted by $|S|$.

Eg: Set S of letters in the English alphabet. Then $|S| = 26$.

Eg: $|\emptyset| = 0$.

A set is said to be **infinite**, if it is not finite.

Eg: The set of positive integers $\{1, 2, 3, \dots\}$ is infinite.

Power Set of S , $P(S)$, is the set of all subsets of the set S .

If a set has n elements, then its power set has 2^n elements.

Eg: $P(\{0,1,2\}) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0,1\}, \{1,2\}, \{0,2\}, \{0,1,2\}\}$

The **Ordered n-tuple** (a_1, a_2, \dots, a_n) is the ordered collection that has a_1 as its first element, a_2 as its second element, \dots , a_n as its n^{th} element.

Cartesian product of A and B is

$$\mathbf{A \times B = \{(a, b) \mid a \in A \wedge b \in B\}}$$

Eg: $A = \{a, b\}$ and $B = \{1, 2, 3\}$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

$$B \times A = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$$

$$A \times B \neq B \times A$$

$$|A \times B| = |A| * |B|$$

A subset R of $A \times B$ is called a **relation** from set A to set B .

The **cartesian product** of the sets A_1, A_2, \dots, A_n , denoted by $A_1 \times A_2 \times \dots \times A_n$, is the set of ordered n -tuples (a_1, a_2, \dots, a_n) , where a_i belongs to set A_i for $i = 1, 2, \dots, n$.

That is, $A_1 \times A_2 \times \dots \times A_n =$
 $\{(a_1, a_2, \dots, a_n) \mid a_i \in A_i \text{ for } i = 1, 2, \dots, n\}.$

Truth Sets

Given Predicate P , and domain D , the truth set of P is the set of elements x in D for which $P(x)$ is true.

The truth set of $P(x)$ is denoted by $\{x \in D \mid P(x)\}$.

Q: What are the truth sets of the predicates $P(x)$, $Q(x)$, and $R(x)$, where the domain is the set of integers and $P(x)$ is “ $|x| = 1$ ”, $Q(x)$ is “ $x^2 = 2$ ”, and $R(x)$ is “ $|x| = x$ ” ?

$\forall_{x \in S} (P(x))$ denotes the universal quantification of $P(x)$ over all elements in the set S .

i.e., $\forall_{x \in S} (P(x))$ is shorthand for $\forall x (x \in S \rightarrow P(x))$.

$\exists_{x \in S} (P(x))$ denotes the existential quantification of $P(x)$ over all elements in S .

i.e., $\exists_{x \in S} (P(x))$ is shorthand for $\exists x (x \in S \wedge P(x))$.

Russell's Paradox

Let the domain be the set of all sets

$$S = \{x \mid x \notin x\}$$

Is S a member of itself?

Suppose, $S \in S$. Then, the predicate $x \notin x$ is false. Hence, S should not belong to S . It's a contradiction.

Suppose, $S \notin S$. Then, the predicate $x \notin x$ is true. Hence, S should belong to S . It's a contradiction.

Therefore, it is a paradox.

Analogy:

Predicate: I help people who can't help themselves.

Suppose I'm one of those people.

When I'm sick (i.e. I can't help myself), I'm in a paradox.


According to the predicate, I should help "me", but I can't do that because I'm sick. If I don't help myself, I'm violating the predicate.

The Big Bang Theory Scene: Leonard's car. "Play that funky music, white boy" is playing on the stereo.


- **Sheldon:** So they're requesting that the white boy play the funky music, yes?
 - **Leonard:** Yes.
- **Sheldon:** And this music we're listening to right now is funky as well?
 - **Leonard:** Sure.
- **Sheldon:** Let me ask you this. Do you think this song is the music the white boy ultimately plays?
 - **Leonard:** It could be.
- **Sheldon:** So it's like the musical equivalent of **Russell's Paradox**, the question of whether the set of all sets that don't contain themselves as members contains itself?
 - **Leonard:** Exactly.
- **Sheldon:** Well then I hate it. Music should just be fun!

Set Operations

- Union of sets A and B contains those elements in A , B or both.
 - $A \cup B = \{x \mid x \in A \vee x \in B\}$
- Intersection of sets A and B contains those elements in both A and B .
 - $A \cap B = \{x \mid x \in A \wedge x \in B\}$



$A \cup B$ is shaded.



$A \cap B$ is shaded.

Two sets are **disjoint** when their intersection is empty.

$$A \cap B = \emptyset$$

$$\text{Eg: } \{1, 2\} \cap \{3, 4\} = \emptyset$$

Q: What are the resulting sets of the following.

$$1. \{1, 2, 3\} \cap \{1, 2\} = \{1, 2\}$$

$$2. \{1, 2, 3\} \cap \{R, G, B\} = \emptyset$$

$$3. \{1, 2, 3\} \cap \emptyset = \emptyset$$

$$4. \{1, 2, 3\} \cup \{1, 4\} = \{1, 2, 3, 4\}$$

$$5. \{1, 2, 3\} \cup \{R, G, B\} = \{1, 2, 3, R, G, B\}$$

$$6. \{1, 2, 3\} \cup \emptyset = \{1, 2, 3\}$$

$$7. \{1, 2, 3\} \cup \{\} = \{1, 2, 3\}$$

Q: What are the resulting sets of the following.

1. $\{1, 2, 3\} \cap \{1, 2\} = \{1, 2\}$

2. $\{1, 2, 3\} \cap \{R, G, B\} = \emptyset$

3. $\{1, 2, 3\} \cap \emptyset = \emptyset$

4. $\{1, 2, 3\} \cup \{1, 4\} = \{1, 2, 3, 4\}$

5. $\{1, 2, 3\} \cup \{R, G, B\} = \{1, 2, 3, R, G, B\}$

6. $\{1, 2, 3\} \cup \emptyset = \{1, 2, 3\}$

7. $\{1, 2, 3\} \cup \{\} = \{1, 2, 3\}$

Two sets are **disjoint** when their intersection is empty.

$$A \cap B = \emptyset$$

Cardinality of union:

If sets A and B are disjoint, $|A \cup B| = |A| + |B|$

In general, $|A \cup B| = |A| + |B| - |A \cap B|$

Generalization of this result (of n sets) is called the **principle of inclusion-exclusion**.

Eg: $A = \{1, 2, 3\}$, $B = \{3, 4\}$

$$|A \cup B| = |A| + |B| - |A \cap B| = 3+2-1 = 4$$

$$\text{Eg: } |\{1, 2, 3\} \cup \{2, 3, 4\}| = 3+3-2 = 4$$

Difference of two sets:

Difference of two sets, $A - B$, is set containing elements in A but not in B .

$$A - B = \{ x \mid x \in A \wedge x \notin B \}$$

$$\text{Eg: } \{1, 2\} - \{3, 4\} = \{1, 2\}$$


$$\text{Eg: } \{1, 2, 3\} - \{3, 4\} = \{1, 2\}$$

$$\text{Eg: } \{1, 2, 3\} - \{1, 2, 3\} = \emptyset$$

$$\text{Eg: } \{1, 2, 3\} - \{1, 2\} = \{3\}$$

$$\text{Eg: } \{1, 2, 3\} - \{1, 2, 3\} = \{ \}$$

$$\text{Eg: } \{1, 2, 3\} - \emptyset = \{1, 2, 3\}$$



$A - B$ is shaded.

Complement of a set:

Complement, \bar{A} (A bar) or A^c , is the complement with respect to the universal set, U . That is, the difference $U - A$ is the complement of A .

$$\bar{A} = \{ x \mid x \in U \wedge x \notin A \}$$

$$\bar{A} = \{ x \mid x \notin A \}$$

Eg: $U = \{ 1, 2, 3, 4, 5, 6 \}$


$$A = \{ 1, 2, 3, 4 \}$$

$$\bar{A} = \{ 5, 6 \}$$

Eg: $U = \{ a, e, i, o, u \}$

$$A = \{ a, e, o \}$$

$$\bar{A} = \{ i, u \}$$



\bar{A} is shaded.

TABLE 1 Set Identities.

<i>Identity</i>	<i>Name</i>
$A \cup \emptyset = A$ $A \cap U = A$	Identity laws
$A \cup U = U$ $A \cap \emptyset = \emptyset$	Domination laws
$A \cup A = A$ $A \cap A = A$	Idempotent laws
$\overline{\overline{A}} = A$	Complementation law
$A \cup B = B \cup A$ $A \cap B = B \cap A$	Commutative laws
$A \cup (B \cap C) = (A \cup B) \cap C$ $A \cap (B \cup C) = (A \cap B) \cup C$	Associative laws
$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$	Distributive laws
$\overline{A \cup B} = \overline{A} \cap \overline{B}$ $\overline{A \cap B} = \overline{A} \cup \overline{B}$	De Morgan's laws
$A \cup (A \cap B) = A$ $A \cap (A \cup B) = A$	Absorption laws
$A \cup \overline{A} = U$ $A \cap \overline{A} = \emptyset$	Complement laws

Membership Tables

(Observe the similarities with the Truth Tables)

Eg: Use a **membership table** to show

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

TABLE 2 A Membership Table for the Distributive Property.

A	B	C	$B \cup C$	$A \cap (B \cup C)$	$A \cap B$	$A \cap C$	$(A \cap B) \cup (A \cap C)$
1	1	1	1	1	1	1	1
1	1	0	1	1	1	0	1
1	0	1	1	1	0	1	1
1	0	0	0	0	0	0	0
0	1	1	1	0	0	0	0
0	1	0	1	0	0	0	0
0	0	1	1	0	0	0	0
0	0	0	0	0	0	0	0

Membership Table

Prove De Morgan's law $(A \cup B)' = A' \cap B'$ using membership table.

A	B	$A \cup B$	$\overline{A \cup B}$	\overline{A}	\overline{B}	$\overline{A} \cap \overline{B}$
1	1	1	0	0	0	0
1	0	1	0	0	1	0
0	1	1	0	1	0	0
0	0	0	1	1	1	1

Q: Prove De Morgan's law $(A \cap B)' = A' \cup B'$ **without** using membership table.

Q: Use set builder notation and logical equivalences to establish the De Morgan's law $\overline{A \cap B} = \overline{A} \cup \overline{B}$.

Soln:

$\overline{A \cap B} = \{x \mid x \notin A \cap B\}$	By definition of complement
$= \{x \mid \neg(x \in (A \cap B))\}$	By definition of \notin symbol
$= \{x \mid \neg((x \in A) \wedge (x \in B))\}$	By definition of intersection
$= \{x \mid \neg(x \in A) \vee \neg(x \in B)\}$	By De Morgan's law of logic
$= \{x \mid x \notin A \vee x \notin B\}$	By definition of \notin symbol
$= \{x \mid x \in \overline{A} \vee x \in \overline{B}\}$	By definition of complement
$= \{x \mid x \in \overline{A} \cup \overline{B}\}$	By definition of union
$= \overline{A} \cup \overline{B}$	By meaning of set builder notation

Therefore, $\overline{A \cap B} = \overline{A} \cup \overline{B}$

GENERALIZED UNIONS AND INTERSECTIONS

UNION:


Union of a collection of sets contains elements that are members of at least one set in the collection.

$$A_1 \cup A_2 \cup \dots \cup A_n = \bigcup_{i=1}^n A_i$$


INTERSECTION

Intersection of a collection of sets contains elements that are members of all sets in the collection.

$$A_1 \cap A_2 \cap \dots \cap A_n = \bigcap_{i=1}^n A_i$$



(a) $A \cup B \cup C$ is shaded.



(b) $A \cap B \cap C$ is shaded.

Computer Representation of Sets:

$$U = \{ 5, 4, 3, 2, 1, 0 \} = 111111$$

$$A = \{ 2, 1, 0 \} = 000111$$

$$B = \{ 3, 2 \} = 001100$$

$$A \cup B = A \vee B$$

$$\{ 2, 1, 0 \} \cup \{ 3, 2 \} = \{ 3, 2, 1, 0 \}$$

$$\begin{array}{rcl} & 000111 & \{ 2, 1, 0 \} \\ \vee & \underline{001100} & \cup \quad \underline{\{ 3, 2 \}} \\ & 001111 & \{ 3, 2, 1, 0 \} \end{array}$$

$$A \cap B = A \wedge B$$

$$\{ 2, 1, 0 \} \cap \{ 3, 2 \} = \{ 2 \}$$

$$\begin{array}{rcl} & 000111 & \{ 2, 1, 0 \} \\ \wedge & \underline{001100} & \cap \quad \underline{\{ 3, 2 \}} \\ & 000100 & \{ 2 \} \end{array}$$


Functions



Function f from A to B is assignment of **exactly one element of B to each element of A** . $f(a) = b$ where 'b' is an element of B assigned by 'f' to the element 'a' of A .

$$f : A \rightarrow B$$

Functions also called **mappings** or **transformations**.

Eg: $g : \text{Students} \rightarrow \text{Grades}$





Domain of the function 'g' is the set of Students.

Co-domain of the function 'g' is the set of Grades.

Image of g(Goodfriend) is B.

Range of the function 'g' is {A, B, C, F}

Domain of $f : A \rightarrow B$ is A

Co-domain of $f : A \rightarrow B$ is B

Image $f(a) = b$ is b

Preimage $f(a) = b$ is a

Range of $f : A \rightarrow B$ is set of all images of elements of A

Two real-valued functions with the same domain can be added and multiplied.

Let $f_1 : A \rightarrow \mathbb{R}$ and $f_2 : A \rightarrow \mathbb{R}$

Then, $(f_1 + f_2)(x) = f_1(x) + f_2(x)$

$(f_1 \cdot f_2)(x) = f_1(x) \cdot f_2(x)$

Eg:

$f_1 : \mathbb{R} \rightarrow \mathbb{R}$

$f_2 : \mathbb{R} \rightarrow \mathbb{R}$

$f_1(x) = x^2$

$f_2(x) = x - x^2$

$(f_1 + f_2)(x) = f_1(x) + f_2(x) = x^2 + x - x^2 = x$

$(f_1 \cdot f_2)(x) = f_1(x) \cdot f_2(x) = x^2 \cdot (x - x^2) = x^3 - x^4$

Image of a subset of the domain:

$$f : A \rightarrow B$$

$$C \subseteq A$$

Image of C under f is a subset of B .

$$f(C) \subseteq B$$

$$f(C) = \{ x \mid \exists c \in C (x = f(c)) \}$$

Types of functions

- One-to-One (Injective) function
- Onto (Surjective) function
- One-to-One Correspondence (Bijective) function

One-to-One (Injective) functions:

A function is said to be One-to-one or injective, if and only if $f(a)=f(b)$ implies that $a=b$ for all a and b in the domain of f .

That is, **f is one-to-one iff $\forall a \forall b (f(a) = f(b) \rightarrow a = b)$**

That is, **f is one-to-one iff $\forall a \forall b (a \neq b \rightarrow f(a) \neq f(b))$**

Eg: $A = \{ a, b, c, d \}$,

$B = \{ 1, 2, 3, 4, 5 \}$


$g : A \rightarrow B$

$g(a) = 4,$

$g(b) = 5,$

$g(c) = 1,$

$g(d) = 3$



Let f be a function. Which of the following defines one-to-one function f ?

1. $\forall a \forall b (f(a) = f(b) \leftrightarrow a = b)$
2. $\forall a \forall b (a = b \rightarrow f(a) = f(b))$
3. $\forall a \forall b (f(a) = f(b) \rightarrow a = b)$
4. $\forall a \forall b (a \neq b \rightarrow f(a) \neq f(b))$

Increasing/Decreasing functions:

Consider a function f whose domain and codomain are subsets of the set of real numbers.

Function f is **increasing** if $f(x) \leq f(y)$ for real $x < y$

That is, $\forall x \forall y (x < y \rightarrow f(x) \leq f(y))$

Function f is **strictly increasing** if $f(x) < f(y)$ for real $x < y$.

That is, $\forall x \forall y (x < y \rightarrow f(x) < f(y))$

Strictly increasing functions must be one-to-one.

Function f is **decreasing** if $f(x) \geq f(y)$ for real $x < y$.

That is, $\forall x \forall y (x < y \rightarrow f(x) \geq f(y))$

Function f is **strictly decreasing** if $f(x) > f(y)$ for real $x < y$.

That is, $\forall x \forall y (x < y \rightarrow f(x) > f(y))$

Strictly decreasing functions must be one-to-one.

Onto (Surjective) functions:

A function f from A to B is called onto or surjective, if and only if every element $b \in B$ there is an element $a \in A$ with $f(a) = b$.

i.e., f is Onto function iff every $b \in B$ has $a \in A$ with $f(a) = b$.

In short, $\forall y \exists x (f(x) = y)$

Eg: $A = \{ a, b, c, d \}$,

$B = \{ 1, 2, 3 \}$


$G : A \rightarrow B$

$G(a) = 3$,

$G(b) = 2$,

$G(c) = 1$,

$G(d) = 3$




One-to-One Correspondence (Bijection) functions:

A function is One-to-one correspondence or bijective function if only if it is one-to-one and onto.

Eg: What kind relations are these?

1. One-to-one correspondence
2. One-to-one but not onto
3. Onto but not one-to-one
4. Neither one-to-one nor onto



Q: Diagram the following functions and mention whether they are one-to-one, onto or one-to-one correspondence:

1. $f : \{a, b, c, d\} \rightarrow \{1, 2, 3, 4\}$

$$f(a) = 1$$

$$f(b) = 2$$

$$f(c) = 3$$

$$f(d) = 4$$

1. $g : \{a, b, c, d\} \rightarrow \{1, 2, 3, 4\}$

$$g(a) = 1$$

$$g(b) = 1$$

$$g(c) = 4$$

$$g(d) = 4$$

Inverse Function:

Inverse of a function \mathbf{f} from A to B such that

$$\mathbf{f}^{-1}(b) = a \text{ when } f(a) = b.$$

Function f is **invertible** when \mathbf{f} is one-to-one correspondence (i.e. one-to-one and onto), otherwise inverse function of \mathbf{f} does not exist.

Eg: $f : \mathbf{R}^+ \rightarrow \mathbf{R}^+$

$$f(x) = x^2$$

$$f^{-1}(y) = y^{1/2}$$

$$f(3) = 3^2 = 9$$

$$f^{-1}(9) = 9^{1/2} = 3$$


Eg: $f : \mathbf{Z} \rightarrow \mathbf{Z}$

$$f(x) = x + 3$$

$$f^{-1}(y) = y - 3$$

$$f(20) = 23$$


$$f^{-1}(23) = 20$$



Composition of Functions:

Composition of function g from A to B and function f from B to C .

$$(f \circ g)(a) = f(g(a))$$



$$g: \{a, b, c\} \rightarrow \{X, Y, Z\}$$

$$g(a) = X$$

$$g(b) = Y$$

$$g(c) = Z$$

$$f: \{X, Y, Z\} \rightarrow \{1, 2, 3\}$$

$$f(X) = 1$$


$$f(Y) = 2$$

$$f(Z) = 3$$

$$(f \circ g)(a) = f(g(a)) = f(X) = 1$$

$$(f \circ g)(b) = f(g(b)) = f(Y) = 2$$

$$(f \circ g)(c) = f(g(c)) = f(Z) = 3$$



$g \circ f$ is not defined because f range, $\{1, 2, 3\}$, is not a subset of g domain, $\{a, b, c\}$

$$(g \circ f)(X) = g(f(X)) = g(1) \text{ is not defined.}$$

Eg:

$$f(x) = 5x + 7$$

$$g(x) = 3x + 2$$

$$(f \circ g)(x) = f(g(x)) = f(3x+2) = 5(3x+2) + 7 = 15x + 17$$

$$(g \circ f)(x) = g(f(x)) = g(5x+7) = 3(5x+7) + 2 = 15x + 23$$

$$(f \circ g)(x) \neq (g \circ f)(x)$$


Relations

A **binary relation** from A to B is a subset of $A \times B$.

Element a is related to b by R is denoted by aRb .

aRb denotes $(a, b) \in R$

$a \not R b$ denotes $(a, b) \notin R$



R	a	b
0	×	×
1	×	
2		×

Eg:

$A = \{0, 1, 2\}$

$B = \{a, b\}$

$A \times B = \{ (0, a), (1, a), (0, b), (1, b), (2, a), (2, b) \}$

$R = \{ (0, a), (1, a), (0, b), (2, b) \} \subseteq A \times B$

MATRIX REPRESENTATION OF RELATIONS:

2-dimensional 0-1 matrix is used for binary relations.

One row for each element of A

One column for each element of B

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$

DIRECTED GRAPH (DIGRAPH) REPRESENTATION OF RELATIONS:

Set V of vertices (nodes) representing elements of the sets.

Set E ordered pairs of elements of V called edges (arcs).

Vertex a is initial vertex and vertex b is terminal vertex of edge (a, b).

Edge (a, a) is a loop.

Relations on a Set:

Relation on the set A is a relation from A to A .



Eg: $A = \{1, 2, 3, 4\}$

$$A \times A = \{ (1,1), (1,2), (1,3), (1,4), \\ (2,1), (2,2), (2,3), (2,4), \\ (3,1), (3,2), (3,3), (3,4), \\ (4,1), (4,2), (4,3), (4,4) \}$$

$$R = \{ (a, b) \mid a \text{ divides } b \} \\ = \{ (1,1), (1,2), (1,3), (1,4), \\ (2,2), (2,4), (3,3), (4,4) \}$$

$$R_1 = \{ (a, b) \mid a \geq b \}$$


$$R_2 = \{ (a, b) \mid a = b \}$$



Eg: Set $A = \{a, b, c, d\}$

Relation $R = \{(a, d), (a, b), (b, b), (b, d), (c, b), (c, a), (d, b)\}$

	a	b	c	d
a	0	1	0	1
b	0	1	0	1
c	1	1	0	0
d	0	1	0	0



Eg: Set $A = \{1, 2, 3, 4\}$

Show the matrix and digraph representation of the relation


$R = \{(1, 1), (1, 3), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (4, 1)\}$

Eg: Set $A = \{1, 2, 3, 4\}$

Show the matrix and digraph representation of the relation

$R = \{(1, 1), (1, 3), (2, 1), (2, 3), (2, 4), (3, 1), (3, 2), (4, 1)\}$

	1	2	3	4
1	1	0	1	0
2	1	0	1	1
3	1	1	0	0
4	1	0	0	0



Relations:

- Let $A = \{1, 2, 3, 4\}$. Relation R on set A be $\{(1,1), (1,2), (2,1), (2,2), (3,4), (4,3), (3,3), (4,4)\}$.
- Let $S = \{a,b,c,d,e,f\}$. Relation R on set S be $\{(a,a),(b,b), (b,c),(c,b),(c,c),(d,d),(d,e),(d,f),(e,d),(e,e),(e,f),(f,d),(f,e),(f,f)\}$
- Relation R on the set of integers such that aRb if and only if $a = b$ or $a = -b$.
- Relation R on the set of real numbers such that aRb if and only if $a - b$ is an integer.
- Relation $R = \{(a, b) \mid a \equiv b \pmod{10}\}$.
- Relation $R = \{(a, b) \mid a \equiv b \pmod{m}\}$, where $m \in \mathbb{Z}^+ \wedge m > 1$
- Relation R on the set of strings of English letters such that aRb if and only if $\text{Length}(a) = \text{Length}(b)$.

Properties of Relations on a set:

1. Reflexive
2. Symmetric
3. Antisymmetric
4. Transitive

Reflexive Relations:

A relation R on a set A is reflexive iff

- $(a,a) \in R$ for every element $a \in A$.
- $\forall a \in A ((a, a) \in R)$.

Eg: $A = \{1, 2, 3, 4\}$

$R = \{(1, 1), (2, 2), (3, 3), (4, 4), (1, 2)\}$ is reflexive

	1	2	3	4
1	1	1	0	0
2	0	1	0	0
3	0	0	1	0
4	0	0	0	1

Symmetric Relations:

A relation R on a set A is symmetric iff

- $(b,a) \in R$ whenever $(a,b) \in R$, for all $a,b \in A$.
- $\forall a \in A \forall b \in A ((a, b) \in R \rightarrow (b, a) \in R)$
- $\forall a \in A \forall b \in A ((a, b) \notin R \rightarrow (b, a) \notin R)$

Eg: Let set $A = \{1, 2, 3, 4\}$

$R = \{(1, 2), (2, 1), (1, 4), (4, 1), (3, 3)\}$ symmetric

	1	2	3	4
1	0	1	0	1
2	1	0	0	0
3	0	0	1	0
4	1	0	0	0

Antisymmetric Relations:

A relation R on a set A is antisymmetric iff

- $a=b$ whenever $(a,b) \in R$ and $(b,a) \in R$, for all $a,b \in R$.
- $\forall a \in A \forall b \in A ((a,b) \in R \wedge (b,a) \in R \rightarrow a=b)$
- $\forall a \in A \forall b \in A ((a \neq b) \rightarrow (a,b) \notin R \vee (b,a) \notin R)$

Eg: Let set $A = \{1, 2, 3, 4\}$

$R = \{(1, 2), (3, 3), (4, 1)\}$ antisymmetric

	1	2	3	4
1	0	1	0	0
2	0	0	0	0
3	0	0	1	0
4	1	0	0	0

Symmetric? Antisymmetric?

	1	2	3
1	0	1	1
2	1	0	0
3	1	0	1

	1	2	3
1	0	1	1
2	0	0	0
3	0	1	1

	1	2	3
1	0	1	0
2	0	0	0
3	0	0	1

	1	2	3
1	0	1	1
2	0	0	0
3	1	0	1

	1	2	3
1	0	0	0
2	0	0	0
3	0	0	1

	1	2	3
1	0	0	0
2	0	0	0
3	0	0	0

Symmetric? Antisymmetric?

	1	2	3		
1	0	1	1		
2	1	0	0		
3	1	0	1	Y	N

	1	2	3		
1	0	1	1		
2	0	0	0		
3	0	1	1	N	Y

	1	2	3		
1	0	1	0		
2	0	0	0		
3	0	0	1	N	Y

	1	2	3		
1	0	1	1		
2	0	0	0		
3	1	0	1	N	N

	1	2	3		
1	0	0	0		
2	0	0	0		
3	0	0	1	Y	Y

	1	2	3		
1	0	0	0		
2	0	0	0		
3	0	0	0	Y	Y

Transitive Relations:

A relation R on a set A is transitive iff

- $(a,c) \in R$ whenever $(a,b) \in R$ and $(b,c) \in R$, for all $a,b,c \in R$.
- $\forall a \in A \forall b \in A \forall c \in A ((a,b) \in R \wedge (b,c) \in R \rightarrow (a,c) \in R)$

	1	2	3	4
1	0	1	1	1
2	0	0	1	1
3	0	0	1	0
4	0	0	1	0

	1	2	3	4
1	1	0	1	0
2	0	0	0	0
3	1	0	1	0
4	0	0	0	0

Reflexive relation R on A , if $\forall a \in A, (a, a) \in R$.









Symmetric relation R on A , if $\forall a \forall b \in A, (a, b) \in R \rightarrow (b, a) \in R$.









Antisymmetric relation R on A , if $\forall a \forall b \in A, (a, b) \in R \wedge (b, a) \in R \rightarrow a=b$.

Transitive relation R on A , if $\forall a \forall b \forall c \in A, (a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R$.

Relations on the set $A = \{1, 2, 3\}$

	<i>reflexive</i>	<i>symmetric</i>	<i>antisymmetric</i>	<i>transitive</i>
$R_0 = \{(1,1), (2,2), (3,3)\}$	Yes	Yes	Yes	Yes
$R_1 = \{(2,2), (2,3), (3,2)\}$	No	Yes	No (2,3) (3,2)	No (3,3)
$R_2 = \{(1,1), (1,2), (2,1), (2,2), (3,3)\}$	Yes	Yes	No (1,2) (2,1)	Yes
$R_3 = \{(2,3), (3,2)\}$	No	Yes	No (2,3) (3,2)	No (2,2) (3,3)
$R_4 = \{(1,2), (2,3), (1,3)\}$	No	No	Yes	Yes

Relation	Reflexive	Symmetric	Antisymmetric	Transitive
				
				
				
				
				
				
				
				

Relation	Reflexive	Symmetric	Antisymmetric	Transitive
	Y	N	Y	Y
	Y	Y	Y	Y
	N	N	Y	Y
	N	Y	N	N
	Y	Y	Y	Y
	N	N	Y	Y
	N	N	Y	N
	N	N	N	N

Q: Mention the properties (reflexive, symmetric, antisymmetric and transitive properties) of the relations.

1. Relation of $\{a, b, c\}$,

$$R = \{(a, a), (b, b), (c, c), (b, c), (c, b)\}$$

Ans: ...

2. Relation of $\{a, b, c, d\}$,

$$S = \{(a, c), (c, a), (a, d), (d, a), (d, d), (a, b), (b, a), (b, b)\}$$

Ans: ...

Q: Mention the properties (reflexive, symmetric, antisymmetric and transitive properties) of the relations.

1. Relation of $\{a, b, c\}$,

$$R = \{(a, a), (b, b), (c, c), (b, c), (c, b)\}$$

Ans: Y, Y, N, Y

2. Relation of $\{a, b, c, d\}$,

$$S = \{(a, c), (c, a), (a, d), (d, a), (d, d), (a, b), (b, a), (b, b)\}$$

Ans: N, Y, N, N

Combining Relations:

Relations from A to B are subsets of $A \times B$ and can be combined in any way two sets can be combined.

Eg: $A = \{0, 1, 2\}$, $B = \{a, b\}$

$A \times B = \{(0, a), (1, a), (0, b), (1, b), (2, a), (2, b)\}$

$R1 = \{(0, a), (0, b)\}$, $R2 = \{(0, b), (1, a), (1, b)\}$

$$R1 \cap R2 = \{(0, b)\}$$

$$R1 \cup R2 = \{(0, a), (0, b), (1, a), (1, b)\}$$

$$R1 - R2 = \{(0, a)\}$$

$$R2 - R1 = \{(1, a), (1, b)\}$$

$$R1 \oplus R2 = R1 \cup R2 - R1 \cap R2$$


$$= \{(0, a), (0, b), (1, a), (1, b)\} - \{(0, b)\}$$

$$= \{(0, a), (1, a), (1, b)\}$$

Equivalence Relations:

A relation R on a set A is called an equivalence relation if it is **reflexive**, **symmetric**, and **transitive**.

Eg: Let $S = \{a, b, c, d, e, f\}$. Relation R on set S be $\{(a, a), (b, b), (b, c), (c, b), (c, c), (d, d), (d, e), (d, f), (e, d), (e, e), (e, f), (f, d), (f, e), (f, f)\}$



	a	b	c	d	e	f
a	1					
b		1	1			
c		1	1			
d				1	1	1
e				1	1	1
f				1	1	1

Q: Let $A = \{1, 2, 3, 4\}$ and

$R = \{(1,1), (1,2), (2,1), (2,2), (3,4), (4,3), (3,3), (4,4)\}$

be a relation on A . Verify that R is an equivalence relation.

Soln:

R is reflexive since it contains $(1,1)$, $(2,2)$, $(3,3)$ and $(4,4)$.

That is, $\forall x (x,x) \in R$

R is symmetric since it contains $(1,2)$, $(2,1)$, $(3,4)$, $(4,3)$ and no (a,b) where (b,a) is not in R .

That is, $\forall x \forall y ((x,y) \in R \rightarrow (y,x) \in R)$

R is transitive since for every pair of (x,y) and (y,z) , there is (x,z) in R .

That is, $\forall x \forall y \forall z ((x,y) \in R \wedge (y,z) \in R \rightarrow (x,z) \in R)$

Therefore, R is an equivalence relation.

Q: Let R be a relation on the set of real numbers such that aRb iff $a-b$ is an integer. Prove whether R is an equivalence relation.

Soln: ...

Q: Let R be a relation on the set of real numbers such that aRb iff $a-b$ is an integer. Prove whether R is an equivalence relation.

Soln: $a-a=0$ and $0 \in \mathbb{Z}$

That is, $\forall a (aRa)$. $\therefore R$ is reflexive.

Let $a-b = k$ be an integer.

Then, $b-a = -k$, which is also an integer.

That is, if aRb , then bRa . $\therefore R$ is symmetric.

Let $a-b=k$ and $b-c=m$ where k and m are integers.

Then, $a-c = (a-b)-(c-b) = k-(-m)$, which is an integer.

That is, if aRb and bRc , then aRc . $\therefore R$ is transitive.

Because R is reflexive, symmetric and transitive,
 R is an equivalence relation.

Q: Let 'a', 'b' and 'm' are integers with $m > 1$. Show that the relation $R = \{ (a, b) \mid a \equiv b \pmod{m} \}$ is an equivalence relation on the set of integers.

Soln:

If $a \equiv b \pmod{m}$, then " $m \mid (a - b)$ " (read: m divides a-b)

...

Q: Let 'a', 'b' and 'm' are integers with $m > 1$. Show that the relation $R = \{ (a, b) \mid a \equiv b \pmod{m} \}$ is an equivalence relation on the set of integers.

Soln:

If $a \equiv b \pmod{m}$, then " $m \mid (a - b)$ " (read: m divides a-b)
 $a \equiv a \pmod{m}$ because $m \mid a - a$, which is same as $m \mid 0$.
 $\therefore R$ is reflexive.

Let $a \equiv b \pmod{m}$

i.e., $m \mid (a - b)$

$mk = a - b$, where k is an integer

$m(-k) = b - a$

i.e., $b \equiv a \pmod{m}$ because $-k$ is also an integer

$\therefore R$ is symmetric.

Q: Let 'a', 'b' and 'm' are integers with $m > 1$. Show that the relation $R = \{ (a, b) \mid a \equiv b \pmod{m} \}$ is an equivalence relation on the set of integers.

Soln:

...

Let $a \equiv b \pmod{m}$ and $b \equiv c \pmod{m}$

i.e., $mk = a-b$ and $ml = b-c$, where k and l are integers.

$$mk+ml = a-b+b-c$$

$$m(k+l) = a-c$$

i.e., $a \equiv c \pmod{m}$

$\therefore R$ is transitive.

R is reflexive, symmetric and transitive.


$\therefore R$ is an equivalence relation.

Equivalence Classes:

Let R be an equivalence relation on a set A . The set of all the elements that are related to an element 'a' of A is called equivalence class of 'a' denoted by $[a]_R$ or $[a]$ when the relation is implicit.

i.e. $[a]_R = \{s \mid (a,s) \in R\}$

Elements of $[a]_R$
are also known as
representatives of $[a]_R$.



	a	b	c	d	e	f
a	1					
b		1	1			
c		1	1			
d				1	1	1
e				1	1	1
f				1	1	1

Q: What are the equivalence classes of 0 and 1 for the congruence modulo 10?

Soln:

The equivalence class of 0 contains all integers 'a' such that $a \equiv 0 \pmod{10}$.

$$[0] = \{ \dots, -20, -10, 0, 10, 20, \dots \}$$

Similarly, the equivalence class of 1 contains all integers 'a' such that $a \equiv 1 \pmod{10}$.

$$[1] = \{ \dots, -19, -9, 1, 11, 21, \dots \}$$

Congruence classes modulo m: are the equivalence classes of the relation congruence modulo m.

$$[a]_m = \{ \dots, a-2m, a-m, \mathbf{a}, a+m, a+2m, \dots \}$$

Q: What is the equivalence class of an integer 'a' for the equivalence relation R defined by aRb iff $a = b$ or $a = -b$?

Soln:

aRb iff $a = b$ or $a = -b$ means a related itself and negative of itself.

That is, $[a]_R = \{a, -a\}$

For example, $[10]_R = \{10, -10\}$

$[-100]_R = \{100, -100\}$

$[0]_R = \{0\}$

Theorem: Let R be an equivalence relation on a set A .

These statements for elements a and b of A are equivalent:

(i) aRb (ii) $[a] = [b]$ (iii) $[a] \cap [b] \neq \emptyset$

Note:

1. Two equivalence classes are either disjoint or identical.
2. Let R be an equivalence relation on a set A and let $a, b \in A$.
 $[a] \neq [b]$ iff $[a] \cap [b] = \emptyset$.
3. For $a, b \in A$, if $b \in [a]$ then $[a] = [b]$.

Partition of a Set:

Let A be a nonempty set. Let P be a set of nonempty subsets A_1, A_2, \dots, A_n of the set A such that

$A_i \cap A_j = \emptyset$ for $i \neq j$... Mutually Exclusive

$A_1 \cup A_2 \cup \dots \cup A_n = A$... Collectively exhaustive

The set $P = \{A_1, A_2, \dots, A_n\}$ is called the partition of A .

Partial Orderings

Partial Order:

A relation R on the set S is called a partial order/ordering if it is **reflexive**, **antisymmetric** and **transitive**.

Poset (S, R) :

Relation R is a partial ordering on set S .

Eg: (\mathbb{Z}, \leq) is a Poset.

Eg: $(\mathbb{Z}^+, |)$ is a Poset.

(S, \preceq) :

(S, \preceq) is notation for poset where relation \preceq is a partial ordering on set S .

Q: Show that $(\mathbb{Z}^+, |)$ is a Poset. (It's the divisibility relation)

Soln: $a|a$ for every integer a .

$\therefore |$ is Reflexive

Whenever $a \neq b$, at least one of $a|b$ or $b|a$ is false.

$\therefore |$ is antisymmetric.

Whenever $a|b$ and $b|c$, $a|c$.

$\therefore |$ is transitive.

$\therefore (\mathbb{Z}^+, |)$ is a Poset.

Q: Show that $(P(S), \subseteq)$ is a Poset.

Soln: ...

Comparable and Incomparable:

Elements a and b are incomparable when they are elements of a poset (S, \leq) such that neither $a \leq b$ nor $b \leq a$.

Eg: In poset (\mathbb{Z}^+, \leq) , for every pair (a, b) either $a \leq b$ or $b \leq a$. For instance, $10 \leq 20$. That is, every pair (a, b) is comparable.

Eg: In poset $(\mathbb{Z}^+, |)$, 5 and 7 are incomparable because $5 \nmid 7$ is false and $7 \nmid 5$ is false. Whereas 6 and 18 are comparable because 6 divides 18.

Total Ordering (Linear Ordering):

Every pair of elements in S are comparable.



Eg: (\mathbb{Z}, \leq)

Hasse Diagram:

In a digraph of a Partial Order,

1. **remove self-loops** because we know the partial order is **reflexive**.

2. **remove direction marks** from edges because we know that the edges always point upwards as the relation is **antisymmetric**.



1. **remove transitive edges** because we know the relation is **transitive**. If there are edges (a,b) and (b,c) , remove (a,c) .

Eg: Course prerequisites

a: ICS


b: DML (requires ICS)

c: DS (requires ICS)

d: DAA (requires DS and DML)

e: DBMS (requires DS and DML)

f: AppDev (requires DAA and DBMS)



Maximal and Minimal elements:

'a' is a maximal in the poset (S, \leq) if there is no $b \in S$ such that $a < b$.

'a' is a minimal in the poset (S, \leq) if there is no $b \in S$ such that $b < a$.

Greatest and Least elements:

'a' is the greatest element of the poset (S, \leq) if $b \leq a$ for all $b \in S$.

'a' is the least element of the poset (S, \leq) if $a \leq b$ for all $b \in S$.

Upper Bound and Lower Bound elements:

If 'u' is an element of S such that $a \leq u$ for all elements $a \in A$, then u is an upper bound of A.

If 'l' is an element of S such that $l \leq a$ for all elements $a \in A$, then l is a lower bound of A.

Least Upper Bound and Greatest Lower Bound elements:

'x' is an upper bound that is less than every other upper bound of A.

'l' is a lower bound that is greater than every other lower bound of A.



Maximal elements: 12

Minimal elements: 1

Greatest element: 12

Least element: 1

**These above ones
are defined on the poset**



The following are defined on a subset of the poset.

Let the subset be $A = \{2, 3\}$

Upper bound of A: 12

Lower bound of A: 1

Least Upper bound of A: 12

Greatest Lower bound of A: 1

Eg: Poset $(\{1,2,\dots,24\}, |)$. It's a "divides" relation.

Maximal elements:

Minimal elements:

Greatest element:

Least element:

For the subset $A = \{2,3\}$,

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:


For the subset $A = \{6,10\}$

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:



Eg: Poset ($\{1,2,\dots,24\}$, $|$). It's a “divides” relation.

Maximal elements: **24, 16, 18, 20, 22, 15, 14, 21, 19, 17, 13, 23**

Minimal elements: **1**

Greatest element: **None**

Least element: **1**

For the subset $A = \{2,3\}$,

Upper bound of A: **6, 12, 18, 24**

Lower bound of A: **1**

Least Upper bound of A: **6**

Greatest Lower bound of A: **1**


For the subset $A = \{6,10\}$

Upper bound of A: **None**

Lower bound of A: **2, 1**

Least Upper bound of A: **None**

Greatest Lower bound of A: **2**



Eg: Let $S = \text{Power set of } \{a,b,c\}$. Poset (S, \subseteq) .

Maximal elements:

Minimal elements:

Greatest element:

Least element:


For the subset $A = \{\mathbf{ab}, \mathbf{b}\}$,

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:



Eg: Let $S = \text{Power set of } \{a,b,c\}$. Poset (S, \subseteq) .

Maximal elements: **abc**

Minimal elements: **ϕ**

Greatest element: **abc**

Least element: **ϕ**


For the subset $A = \{\mathbf{ab}, \mathbf{b}\}$,

Upper bound of A: **ab, abc**

Lower bound of A: **b, ϕ**

Least Upper bound of A: **ab**

Greatest Lower bound of A: **b**



Maximal elements: **f**

Minimal elements: **a**

Greatest element: **f**

Least element: **a**


For the subset $A = \{b, c\}$

Upper bound of A: **d, e, f**

Lower bound of A: **a**

Least Upper bound of A: **None**

Greatest Lower bound of A: **a**



Note: The above example demonstrates, having multiple upper bounds for a pair of elements doesn't guarantee to have a least upper bound.

Maximal elements:

Minimal elements:

Greatest element:

Least element:

For the subset $A=\{d,e,f\}$

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:


For the subset $A=\{b,d\}$

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:



Maximal elements: **g,h,i**

Minimal elements: **a,b,c,e**

Greatest element: **None**

Least element: **None**

For the subset $A=\{d,e,f\}$

Upper bound of A: **None**

Lower bound of A: **None**

Least Upper bound of A: **None**

Greatest Lower bound of A: **None**


For the subset $A=\{b,d\}$

Upper bound of A: **d,g**

Lower bound of A: **b**

Least Upper bound of A: **d**

Greatest Lower bound of A: **b**



Maximal elements:

Minimal elements:

Greatest element:

Least element:

For the subset $A=\{c,e\}$

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:


For the subset $A=\{b,i\}$

Upper bound of A:

Lower bound of A:

Least Upper bound of A:

Greatest Lower bound of A:



Maximal elements: **b,d,i,j**

Minimal elements: **a**

Greatest element: **None**

Least element: **a**

For the subset $A=\{c,e\}$

Upper bound of A: **e,f,g,h,i,j**

Lower bound of A: **c,a**

Least Upper bound of A: **e**

Greatest Lower bound of A: **c**


For the subset $A=\{b,i\}$

Upper bound of A: **None**

Lower bound of A: **a**

Least Upper bound of A: **None**



Greatest Lower bound of A: **a**



Lattice:


A partially ordered set in which every pair of elements has both a **least upper bound** and **greatest lower bound** is called a **lattice**.

Eg: Poset $(\mathbb{Z}^+, |)$





Lattice:

A partially ordered set in which every pair of elements has both a **least upper bound** and **greatest lower bound** is called a **lattice**.



If a and b are not comparable, then it is either $a \leq b$ or $b \leq a$ in the total ordering.





The solution obtained is C1, C2, C3, C4, C5

Minimal element chosen 1	5	2	4	20	12

Lemma: Every finite nonempty poset (S, \leq) has at least one minimal element.

Algorithm SourceRemoval_Toposort(V, E)

$L \leftarrow$ Empty list that will contain the sorted vertices

$S \leftarrow$ Set of all vertices with no incoming edges

while S is non-empty **do**

 remove a vertex v from S

 add v to *tail* of L

for each vertex m with an edge e from v to m **do**

 remove edge e from the graph

if m has no other incoming edges **then**

 insert m into S

return L (a topologically sorted order)

Q: Find a compatible total ordering for the poset $(\{1, 2, 4, 5, 12, 20\}, |)$.

Soln: ...

Q: Let S be Power set of $\{a,b,c\}$.
Find a compatible total ordering for the poset (S, \subseteq) .

Soln: ...

Eg: How many relations are there from a set with **m** elements to a set with **n** elements?

Ans:

Eg: How many functions are there from a set with **m** elements to a set with **n** elements?

Ans:

Eg: How many **one-to-one** functions are there from a set with **m** elements to a set with **n** elements?

Ans:

Eg: How many relations are there from a set with **m** elements to a set with **n** elements?

Ans: 2^{mn}

Eg: How many functions are there from a set with **m** elements to a set with **n** elements?

Ans: $n * n * \dots n \text{ (m times)} = n^m$

Eg: How many **one-to-one** functions are there from a set with **m** elements to a set with **n** elements?

Ans: $n * (n - 1) * (n - 2) * \dots * (n - m + 1)$, where $m \leq n$
 $= {}^n P_m$

Eg: How many **onto** functions ... ($n \leq m$)

Ans: ...

<End of Set Theory />