

# MT210 MIDTERM 1 SAMPLE 1

ILKER S. YUCE  
FEBRUARY 16, 2011

## QUESTION 1. SYSTEMS OF LINEAR EQUATIONS

Determine the values of  $k$  such that the linear system

$$\begin{array}{rcl} 9x_1 & + & kx_2 = 9 \\ kx_1 & + & x_2 = -3 \end{array}$$

is consistent.

### ANSWER

We apply row-reduction algorithm to the augmented matrix corresponding to the system given above: Assume that  $k \neq 0$ , then we get

$$\left[ \begin{array}{ccc} 9 & k & 9 \\ k & 1 & -3 \end{array} \right] \xrightarrow{(-k/9)R_1 + R_2 \rightarrow R_2} \left[ \begin{array}{ccc} 9 & k & 9 \\ 0 & 1 - \frac{k^2}{9} & -3 - k \end{array} \right].$$

By Theorem 2, we know that the system above is consistent if and only if there is no row of the form  $[0 \ 0 \ 1]$  which implies that either we must have  $1 - \frac{k^2}{9} \neq 0$  or we must have  $1 - \frac{k^2}{9} = 0$  and  $-3 - k = 0$ .

We need to examine the case  $k = 0$ . If  $k = 0$ , then we have  $9x_1 = 9$  or  $x_1 = 1$  and  $x_2 = -3$ . So, the system is consistent. Note that if  $k = -3$  the given system is still consistent. Finally, we conclude that the system above is consistent if and only if  $k \neq 3$ .

## QUESTION 2. ROW REDUCTION AND ECHELON FORMS

Determine when the augmented matrix below represents a consistent linear system.

$$\left[ \begin{array}{cccc} 1 & 0 & 2 & a \\ 2 & 1 & 5 & b \\ 1 & -1 & 1 & c \end{array} \right]$$

### ANSWER

We apply row-reduction algorithm to the augmented matrix corresponding to the system given above:

$$\left[ \begin{array}{cccc} 1 & 0 & 2 & a \\ 2 & 1 & 5 & b \\ 1 & -1 & 1 & c \end{array} \right] \xrightarrow[-1R_1+R_3 \leftrightarrow R_3]{-2R_1+R_2 \rightarrow R_2} \left[ \begin{array}{cccc} 1 & 0 & 2 & a \\ 0 & 1 & 1 & b-2a \\ 0 & -1 & -1 & c-a \end{array} \right] \xrightarrow{R_1+R_2 \rightarrow R_2} \left[ \begin{array}{cccc} 1 & 0 & 2 & a \\ 0 & 1 & 1 & b-2a \\ 0 & 0 & 0 & b-3a+c \end{array} \right].$$

By Theorem 2, we know that the system above is consistent if and only if  $b - 3a + c = 0$ .

### QUESTION 3. VECTOR EQUATIONS

Determine if  $\mathbf{b}$  is a linear combination of the vectors  $\mathbf{a}_1$ ,  $\mathbf{a}_2$ , and  $\mathbf{a}_3$  where

$$\mathbf{a}_1 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \mathbf{a}_2 = \begin{bmatrix} -2 \\ -1 \\ -1 \end{bmatrix}, \mathbf{a}_3 = \begin{bmatrix} 3 \\ -1 \\ -3 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 5 \\ -4 \\ -7 \end{bmatrix}.$$

If  $\mathbf{b}$  is a linear combination of the vectors  $\mathbf{a}_1$ ,  $\mathbf{a}_2$ , and  $\mathbf{a}_3$ , express  $\mathbf{b}$  as a linear combination of the vectors  $\mathbf{a}_1$ ,  $\mathbf{a}_2$ , and  $\mathbf{a}_3$ .

### ANSWER

We need to use the fact that  $\mathbf{b}$  is a linear combination of the vectors  $\mathbf{a}_1$ ,  $\mathbf{a}_2$ , and  $\mathbf{a}_3$  if and only if the vector equation  $x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + x_3\mathbf{a}_3 = \mathbf{b}$  has a solution. We need to reduce the augmented matrix

$$\begin{aligned} \left[ \begin{array}{cccc} 1 & -2 & 3 & 5 \\ -1 & -1 & -1 & -4 \\ 0 & -1 & -3 & -7 \end{array} \right] &\xrightarrow{R_1+R_2 \rightarrow R_2} \left[ \begin{array}{cccc} 1 & -2 & 3 & 5 \\ 0 & -3 & 2 & 1 \\ 0 & -1 & -3 & -7 \end{array} \right] \xrightarrow{\substack{-2R_3+R_1 \rightarrow R_1 \\ -3R_3+R_2 \rightarrow R_2}} \left[ \begin{array}{cccc} 1 & 0 & 9 & 19 \\ 0 & 0 & 11 & 22 \\ 0 & -1 & -3 & -7 \end{array} \right] \\ \left[ \begin{array}{cccc} 1 & 0 & 9 & 19 \\ 0 & 0 & 11 & 22 \\ 0 & -1 & -3 & -7 \end{array} \right] &\xrightarrow{\substack{R_2 \rightarrow R_3 \\ (1/11)R_3 \rightarrow R_3}} \left[ \begin{array}{cccc} 1 & 0 & 9 & 19 \\ 0 & -1 & -3 & -7 \\ 0 & 0 & 1 & 2 \end{array} \right] \xrightarrow{\substack{-9R_3+R_1 \rightarrow R_1 \\ 3R_3+R_2 \rightarrow R_2, -R_2 \rightarrow R_2}} \left[ \begin{array}{cccc} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{array} \right]. \\ \text{G.S.} &= \begin{cases} x_1 = 1 \\ x_2 = 1 \\ x_3 = 2 \end{cases} \end{aligned}$$

Finally, we see that  $1 \cdot \mathbf{a}_1 + 1 \cdot \mathbf{a}_2 + 2 \cdot \mathbf{a}_3 = \mathbf{b}$ .

#### QUESTION 4. THE MATRIX EQUATION $A\mathbf{x}=\mathbf{b}$

A. Solve the matrix equation  $A\mathbf{x} = \mathbf{b}$  where

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 0 \\ 1 & 1 & 2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

ANSWER

We need to reduce the augmented matrix

$$\left[ \begin{array}{ccc|c} 1 & 2 & 1 & 0 \\ 1 & 3 & 0 & 0 \\ 1 & 1 & 2 & 0 \end{array} \right] \xrightarrow[\begin{array}{c} -R_1+R_2 \rightarrow R_2 \\ -R_1+R_3 \rightarrow R_3 \end{array}]{\begin{array}{c} -R_1+R_2 \rightarrow R_2 \\ -R_1+R_3 \rightarrow R_3 \end{array}} \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \end{array} \right] \xrightarrow[\begin{array}{c} -2R_2+R_1 \rightarrow R_1 \\ R_2+R_3 \rightarrow R_3 \end{array}]{\begin{array}{c} R_2+R_3 \rightarrow R_3 \\ -2R_2+R_1 \rightarrow R_1 \end{array}} \left[ \begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

We see that  $x_1$  and  $x_2$  are basic variables and  $x_3$  is a free variable. We rewrite the system, i.e., we get  $x_1 + 3x_3 = 0$  and  $x_2 - x_3 = 0$  OR  $x_1 = -3x_3$  and  $x_2 = x_3$ .

$$G.S. = \begin{cases} x_1 = -3x_3 \\ x_2 = x_3 \\ x_3 \text{ is free.} \end{cases}$$

B. Is it possible to solve  $A\mathbf{x} = \mathbf{b}$  for **any given**  $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$  where  $A$  is the matrix given in part A? Explain.

ANSWER

NO. By Theorem 6,  $A\mathbf{x} = \mathbf{b}$  has a solution FOR ANY GIVEN  $\mathbf{b}$  if and only if  $A$  has 3 pivot positions. As you can see above,  $A$  has only 2 pivot positions. As a conclusion, it is not possible to solve  $A\mathbf{x} = \mathbf{b}$  for any given  $\mathbf{b}$ .

C. Describe the set of all  $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$  for which  $A\mathbf{x} = \mathbf{b}$  does have a solution.

ANSWER

We need to reduce the augmented matrix

$$\left[ \begin{array}{cccc} 1 & 2 & 1 & b_1 \\ 1 & 3 & 0 & b_2 \\ 1 & 1 & 2 & b_3 \end{array} \right] \xrightarrow[\begin{array}{c} -R_1+R_3 \rightarrow R_3 \\ -R_1+R_2 \rightarrow R_2 \end{array}]{\begin{array}{c} -R_1+R_2 \rightarrow R_2 \\ -R_1+R_3 \rightarrow R_3 \end{array}} \left[ \begin{array}{cccc} 1 & 2 & 1 & b_1 \\ 0 & 1 & -1 & b_2 - b_1 \\ 0 & -1 & 1 & b_3 - b_1 \end{array} \right] \xrightarrow[\begin{array}{c} -2R_2+R_1 \rightarrow R_1 \\ R_2+R_3 \rightarrow R_3 \end{array}]{\begin{array}{c} R_2+R_3 \rightarrow R_3 \\ -2R_2+R_1 \rightarrow R_1 \end{array}} \left[ \begin{array}{cccc} 1 & 0 & 3 & -2b_2 + 3b_1 \\ 0 & 1 & -1 & b_2 - b_1 \\ 0 & 0 & 0 & b_3 + b_2 - 2b_1 \end{array} \right]$$

By Theorem 2, the equation  $A\mathbf{x} = \mathbf{b}$  has a solution if and only if  $b_3 + b_2 - 2b_1 = 0$ .

## QUESTION 5. SOLUTION SETS OF LINEAR SYSTEMS

Consider the linear system  $A\mathbf{x} = \mathbf{b}$  where

$$A = \begin{bmatrix} 1 & -1 & -2 & -2 & -2 \\ 3 & -2 & -2 & -2 & -2 \\ -3 & 2 & 1 & 1 & -1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 3 \\ -1 \\ -1 \end{bmatrix}$$

- A. Solve the linear system.
- B. Write the general solution in parametric-vector form.
- C. Give a particular solution  $\mathbf{p}$ .
- D. Write the solution set for the homogeneous equation  $A\mathbf{x} = \mathbf{0}$ .

### ANSWER

A. We need to reduce the augmented matrix( I'll leave the details to you.)

$$\left[ \begin{array}{cccccc} 1 & -1 & -2 & -2 & -2 & 3 \\ 3 & -2 & -2 & -2 & -2 & -1 \\ -3 & 2 & 1 & 1 & -1 & -1 \end{array} \right] \sim \left[ \begin{array}{cccccc} 1 & 0 & 0 & 0 & -4 & -11 \\ 0 & 1 & 0 & 0 & -8 & -18 \\ 0 & 0 & 1 & 1 & 3 & 2 \end{array} \right]$$

We see that  $x_1$ ,  $x_2$  and  $x_3$  are basic variables and  $x_4$  and  $x_5$  are free variables. We rewrite the system, i.e., we get  $x_1 - 4x_5 = -11$ ,  $x_2 - 8x_5 = -18$ ,  $x_3 + x_4 + 3x_5 = 2$  OR  $x_1 = -11 + 4x_5$ ,  $x_2 = -18 + 8x_5$ ,  $x_3 = 2 - x_4 - 3x_5$ . Then we find that

$$G.S. = \begin{cases} x_1 = -11 + 4x_5 \\ x_2 = -18 + 8x_5 \\ x_3 = 2 - x_4 - 3x_5 \\ x_4 \text{ is free} \\ x_5 \text{ is free.} \end{cases}$$

B. The parametric vector form of the general solution set is

$$G.S. = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -11 \\ -18 \\ 2 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 0 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} 4 \\ 8 \\ -3 \\ 0 \\ 1 \end{bmatrix} : x_4, x_5 \in \mathbb{R} \right\}$$

C. The particular solution is given as  $\mathbf{p} = \begin{bmatrix} -11 \\ -18 \\ 2 \\ 0 \\ 0 \end{bmatrix}$

D. The parametric vector form of homogeneous part of the general solution set is

$$v_h = \left\{ x_4 \begin{bmatrix} 0 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} 4 \\ 8 \\ -3 \\ 0 \\ 1 \end{bmatrix} : x_4, x_5 \in \mathbb{R} \right\}$$

## QUESTION 6. LINEAR INDEPENDENCE

Determine if the following sets of vector are linearly independent. If **not**, write one vector as a linear combination of other vectors in the set.

A.)  $\left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}.$

ANSWER

This set is linearly dependent by Theorem 9, i.e., zero vector is in the set.

B.)  $\left\{ \begin{bmatrix} -5 \\ 10 \end{bmatrix}, \begin{bmatrix} -4 \\ -2 \end{bmatrix}, \begin{bmatrix} 36 \\ 12 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \end{bmatrix} \right\}$

ANSWER

This set is linearly dependent by Theorem 8, i.e., more vectors than the entries in each vectors.

C.)  $\left\{ \mathbf{v}_1 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 3 \\ 3 \end{bmatrix} \right\}$

ANSWER

This problem requires work. We need to solve the system  $x_1\mathbf{v}_1 + x_2\mathbf{v}_2 = \mathbf{0}$ :

$$\begin{bmatrix} -1 & 3 & 0 \\ 2 & 3 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \Rightarrow G.S. = \begin{cases} x_1 = 0 \\ x_2 = 0. \end{cases}$$

As a conclusion,  $\mathbf{v}_1$  and  $\mathbf{v}_2$  are linearly independent because the system  $x_1\mathbf{v}_1 + x_2\mathbf{v}_2 = \mathbf{0}$  has ONLY the trivial solution.

D.)  $\left\{ \begin{bmatrix} 1 \\ 2 \\ -4 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ -6 \end{bmatrix} \right\}$

ANSWER

This set is linearly dependent by Theorem 7, i.e., the third vector is the addition of the first two vector.

E.)  $\left\{ \mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 \\ 3 \\ -4 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -4 \\ 2 \\ -1 \end{bmatrix} \right\}$

ANSWER

This problem requires work. We need to solve the system  $x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3 = \mathbf{0}$ :

$$\begin{bmatrix} 1 & -1 & -4 \\ 0 & 3 & 2 \\ -1 & -4 & -1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow G.S. = \begin{cases} x_1 = 0 \\ x_2 = 0 \\ x_3 = 0. \end{cases}$$

As a conclusion,  $\mathbf{v}_1$ ,  $\mathbf{v}_2$  and  $\mathbf{v}_3$  are linearly independent because the system  $x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3 = \mathbf{0}$  has ONLY the trivial solution.