

**Radiographic assessment of variation in horizontal position of mental foramen in Indian population on panoramic radiographs: A retrospective study**

Dr. Vivek Bagga;<sup>1</sup>Dr.Nikhil Srivastava;<sup>2</sup>Dr. Rahul Dev Goswami;<sup>3</sup>Dr.Vijay Apparaju;<sup>4</sup>Dr.Pinhaj Ahmed Sherashiya;<sup>5</sup>Dr.Praveen Kumar.<sup>6</sup>

1. Postgraduate Student, Department of Oral and Maxillofacial Surgery, Sri Rajiv Gandhi College of Dental Sciences & Hospital, Cholanagar, RT Nagar Post, Bangalore 560032., Karnataka, INDIA
2. Postgraduate Student, Department of Oral and Maxillofacial Surgery, Sri Rajiv Gandhi College of Dental Sciences & Hospital, Cholanagar, RT Nagar Post, Bangalore 560032, Karnataka, INDIA
3. Postgraduate Student, Department of Oral Medicine and Radiology, Sri Rajiv Gandhi College of Dental Sciences & Hospital, Cholanagar, RT Nagar Post, Bangalore 560032, Karnataka, INDIA
4. Postgraduate Student, Department of Periodontology, Sri Rajiv Gandhi College of Dental Sciences & Hospital, Cholanagar, RT Nagar Post, Bangalore 560032, Karnataka, INDIA
5. Senior Oral Diagnostic Oncologist, Karnataka Cancer Society, Bengaluru-560003, Karnataka, INDIA.
6. Postgraduate Student, Department of Oral and Maxillofacial Surgery, Sri Rajiv Gandhi College of Dental Sciences & Hospital, Cholanagar, RT Nagar Post, Bangalore 560032, Karnataka, INDIA

**Running title:** Radiographic assessment of variation in horizontal position of mental foramen

**Clinical Significance:** The mental foramen (MF) is an important landmark located on the anteroinferior aspect of the body of the mandible.

**Article citation:**Dr.VivekBagga; 1Dr. Nikhil Srivastava; 2Dr. Rahul DevGoswami; 3Dr. Vijay Apparaju; 4Dr. Pinhaj AhmedSherashiya; 5Dr. Praveen Kumar. 6. Radiographic assessment of variation in horizontal position of mental foramen in Indianpopulation on panoramic radiographs: A retrospective study. JOURNAL OF ORAL MEDICINE, SURGERY, PATHOLOGY, BIOLOGY.2016; Vol 1, no 1: 32-40

**Corresponding author:**

**Dr Pinhaj Ahmed Sherashiya,**

Senior Oral Diagnostic Oncologist, Karnataka Cancer Society, Bengaluru-560003, Karnataka, INDIA

E mail add: [hi\\_pinhaz@yahoo.com](mailto:hi_pinhaz@yahoo.com)

## ABSTRACT

The mental foramen (MF) is an important landmark commonly encountered in various craniofacial, maxillofacial, plastic surgery procedures, in anesthetics, during placement of implants and even in forensics. There may be differences concerning the shape and position of the foramen. The foramen either may be completely absent or may show the presence of accessory foramina.

The mental nerve that emerges through this foramen may be damaged and cause temporary or permanent thermal, sensitive, or even tactile changes.

**Objectives:** To determine the most common horizontal position of the mental foramina in a selected population, and to compare the results with those reported for other populations.

**Materials and Methods:** Two hundred digital panoramic radiographs (DPR) were selected, and the position of mental foramen was recorded about the longitudinal axis of the nearest mandibular tooth.

**Results:** The most common position of the mental foramen was located in line with the longitudinal axis of the second premolar (54%); followed by the site between the first and second premolars (33.5%). Mental foramen was bilaterally symmetrical in 74% of the radiographs.

**Conclusion:** Studies and textbooks mention the location of the mental foramen in the Indian population as being predominantly located below the apex of the second premolar or between the apices of the first and second premolars; thereby showing agreement with our findings.

**Keywords:** Horizontal position; mental foramen; panoramic radiograph, Bangalore population

## INTRODUCTION

The mental foramen (MF) is a funnel- shaped opening, usually located closer to the anterior and inferior border of the mandible. It marks the end of the mandibular canal. The mental nerve passing through it is one of the end branches of inferior alveolar nerve.<sup>1</sup> It innervates and supplies the facial structures, such as the soft tissues of the chin, lower lip, and gingiva on the ipsilateral side of the mandible.<sup>2</sup> The foramen is invariably manipulated during various surgical procedures that are performed in the anterior region of the jaws; causing the mental nerve to be easily damaged during the administration of local anaesthetics, surgical procedures and implant placements. It is, therefore, imperative to have a method, which aids in accurate prediction of its location. Variations in the position of mental foramen are common, which may lead to perio-operative complications.

<sup>3,4</sup>According to Moiseiwitsch, individual variations could place the mental foramen anywhere below the canine, to a point between the roots of the first molar.<sup>5</sup>

The location of mental foramen has been studied in various populations. Yosue and Brooks,<sup>6,7</sup> classified the panoramic appearance of the mental foramen as:

- continuous,
- separated,
- diffuse, or
- unidentified type.<sup>7</sup>

Panoramic radiographs are the commonly used diagnostic radiographic aids, for imaging the mandible. Mental foramen is best identified in this type of radiograph. Advanced imaging techniques are expensive and have high radiation exposure.<sup>5</sup>

Literature findings, to determine the location of mental foramen on panoramic radiograph in the Indian population, are very limited. Hence, this study was conducted to document the location, symmetry, and variations in the position of the mental foramen on panoramic radiographs.

## MATERIALS AND METHODS

Digital panoramic radiographs (DPR) of randomly selected 200 patients of Indian origin; among whom 97 were males and 103 were females; were retrospectively analyzed. All patients were over 18 years of age.

Criteria for radiograph selection included:

- All radiographs must be of dentate mandible
- Free from radiolucent or radiopaque lesions.
- Radiographs that showed drifting crowding, spaced lower teeth, or previous orthodontic treatment was excluded from the study.
- High- quality images with proper angulations, contrast and reduced artefacts in the mental foramen region.
- High- quality images with proper angulations, contrast and reduced artefacts in the mental foramen region.
- Cases, where the upper premolars were missing were excluded because of the chance of an over eruption of lower premolars which can alter the location of the mental foramen.
- MF unidentified on radiograph was also excluded.

All DPRs were taken by maintaining standard criteria (tube potential: 60-90 KV, tube current: 2-15 mA, and time: 14 s). Three observers, including the principle observer, read the DPRs. The principle observer calibrated the readings. Any difference in the observations was settled with forced consensus.

The position of the MF was recorded according to a modified Haghanifar and Rokoue scale as follows (Figure. 1)


- Position 1: Situated anterior to the first premolar
- Position 2: In line with the first premolar
- Position 3: Between the first and second premolars
- Position 4: In line with the second premolar
- Position 5: Between the second premolar and mesial root of the first molar
- Position 6: In line with the mesial root of the first molar

A calibrated scale was used to identify the longitudinal axis of the nearest tooth, and its location about the MF was recorded. The position of the mental foramen on the right and left side was recorded. (Figure 1)

**Figure 1: Figure showing the variations in the position of the mental foramen on the mandible**  
**FIGURE 1**

## RESULTS

The most common location of this series was found in line with (position 4 = 54%); followed by and second premolars (position 3 = 33.50%); between second premolar and mesial root (position 5 = 10.25%); in line with the mesial (position 6 = 1.25%), in line with first premolar (position 2 = 1.9%). [Table 1]



the mental foramen in second premolar the location between first 3 = 33.50%); between of first molar (position 5 = root of first molar

**Table 1: Percentage distribution of Mental Foramen according to position**

**TABLE 1**

Position	No. of MF	%age
1	-	-
2	4	1.00
3	134	33.50
4	216	54.00
5	41	10.25
6	5	1.25
Total	400	100.00

In both males and females, position 4 (54.1% and 53.9%) was found to be the most common position of MF followed by position 3 (34% and 33%) respectively. [Table 2]

**Table 2: Percentage distribution of Mental foramen between the genders**

**TABLE 2**

Position	Male		Female	
	Frequency	%	Frequency	%
1	-	-	-	-
2	1	0.5	3	1.5
3	66	34.0	68	33.0
4	105	54.1	111	53.9
5	22	11.3	19	9.2
6	-	-	5	2.4
Total	194	100.0	206	100.0

The location of MF was bilaterally symmetrical in 74% of cases [Table 3].

**Table 3: Symmetry of location of the Mental foramen**

**TABLE 3**

Position	Symmetry		
	Male	Female	Male & Female %
1	-	-	
2	-	-	-
3	27	24	34.4%
4	42	41	56%
5	6	6	8.1%
6	-	2	1.5%
Total	75	73	100%
Total	148		74%

For the symmetrically placed MF, the most common location was position 4 (56%), followed by position 3 (34.4%).

## DISCUSSION

The mental foramen has been identified either by studying radiographs of patients or by directly measuring it on dry mandibles.

Panoramic radiographs depict the mandible as a single dimension image; thus allowing for an accurate positioning of both mental foramina. Periapical radiographs do not reveal its location if it lies outside the border of the film. Comparative analysis between right and left foramina was difficult to detect on other radiographs.<sup>4, 5, 6, 7</sup>

Predominant observations in various studies have shown that the position of the mental foramen was located along the long axis of the second premolar;<sup>8, 9, 10</sup> others have reported it to lie between the first and second premolars, with individual variations occurring occasionally.<sup>11, 12, 13</sup> However, in many studies it was found to lie between the canine and the mesial root of the first molar; and its location below the canine or first molar was either absent or very rarely present.<sup>11</sup>

In our analysis of 200 panoramic radiographs, the most common location of the foramen was in line with apex of second premolar (position 4=54%); between first and second premolar (position 3=33.51%); between second premolar and mesial root of first molar (position 5= 10.25%); in line with the mesial root of first molar (position 6=1.25%); and in line with first premolar (position 2= 1%). No mental foramen was found at position 1.

Position 4 and position 3 contributed to an overall prevalence of 87%. These locations of the MF comprised of the maximum number of cases, in our review of the literature.

There is no consensus regarding the normal position of the mental foramen in different populations. A study on Indian population, showed that these two positions (position 4 and position 3) are the most common, and was seen in 81.1% of population;<sup>4</sup> and in the Iranian population, position 3 and 4 were found in 93.2% cases, according to the study conducted by Sina Haghani et al.<sup>8</sup>

Studies conducted by authors such as Moiseiwtsch;<sup>5</sup> Rupesh et al.<sup>14</sup> Taseir Al-Khateeb et al.<sup>15</sup> and Olasoji et al.<sup>16</sup> in different ethnic populations, reported that the foramina was located precisely between the two premolar. However, studies conducted on populations, such as the Asian Indians, by Shankland,<sup>10</sup> Central regional Indians, by S Gangotri,<sup>17</sup> Malays, by Ngeow and Yuzawati,<sup>18</sup> black Zimbabweans, by Mbajorgu,<sup>19</sup> Kenyan Africans, by Mwaniki and Hassanali,<sup>20</sup> Iraqis, by Muhsen,<sup>21</sup> Saudis, by Al-Jasser and Nwoku,<sup>22</sup> and Koreans, by Kim,<sup>23</sup> have indicated that MF, is most commonly positioned in line with the second premolar (position 4).

Mental foramen was symmetrically located in the majority (74%) of our cases, but the percentage is somewhat lower as compared to other populations. It was 90.4% in Turkish, 77% in North Jordanian, 82.7% in Kurdish and Iranians with 85.7% of symmetry.<sup>17</sup> Study on the Asian Indians reported a decreased incidence of symmetry. (57%).<sup>17</sup>

There is no study, that showed 100% symmetry, in the mental foramen positioning; therefore clearly indicating that MF is not always symmetrical in every individual.

The reason for the variance in the location and symmetry of the MF in various populations can be attributed to genetic factors, diet and local factors in the growth and development of the mandible.

## CONCLUSION

Mental foramen exhibits variance in position, symmetry and size being located in various positions about the mandibular teeth. It is usually located below the mandibular second premolars or near the site between the first and second premolars with a high percentage of symmetry. Hence, it is advisable to assess this important landmark before initiating any surgical procedures in the area.

**Declaration of conflict: The authors have no conflict of interest concerning the study.**

**Source of funds: The authors have received no funds during or prior to the study.**

## REFERENCES:

1. Standring S. Gray's Anatomy. The Anatomical Basis of Clinical Practice. 39<sup>th</sup> Ed., Elsevier Churchill Livingstone. 2005; 601–02.
2. Ilayperuma I, Nanayakkara G, Palahepitiya N. Morphometric analysis of the mental foramen in adult Sri Lankan mandibles. *Int J Morphol* 2009; 27(4):1019-24.
3. White SC, Pharoah MJ. Normal radiographic anatomy. *Oral Radiology: Principles and Interpretation*. 5<sup>th</sup> ed. St Louis: Mosby; 2004: 183.
4. Fishel D, Buchner A, Hershkowitz A, Kaffe I. Roentgenologic study of the mental foramen. *Oral Surg Oral Med Oral Pathol* 1976; 41(5): 682-6.
5. Moiseiwitsch JR. Position of the mental foramen in a North American, white population. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol, Endod*.1998; 85(4): 457-60.
6. Yosue T, Brooks SL. The appearance of mental foramina on panoramic and periapical radiographs. II. Experimental evaluation. *Oral Surg, Oral Med, Oral Pathol, Endo*. 1989; 68(4): 488-92.
7. Yosue T, Brooks SL. The appearance of mental foramina on panoramic radiographs. Evaluation of patients. *Oral Surg Oral Med Oral Pathol*.1989; 68(3): 360-4.
8. Haghani S, Rokouei M. Radiographic evaluation of the mental foramen in a selected Iranian population. *Indian J Res*. 2009; 20(2):150-2.
9. Phillips JL, Weller RN, Kulid JC. The mental foramen: Part I. Size, orientation, and positional relation to the mandibular second premolar. *J Endod*.1990; 16(5):221-3.
10. Shankland WE. The position of the mental foramen in Asian Indians. *J Oral Implantol*.1994; 20(2):118-23.
11. Wang TM, Shin C, Liu JC, Kuo KJ A clinical and anatomical study of the location of the mental foramen in adult Chinese mandibles. *Acta Anat (Basel)*. 1986; 126(1):29-33.
12. Green RM. The position of the mental foramen: A comparison between the southern (Hong Kong) Chinese and other ethnic and racial groups. *Oral Surg Oral Med Oral Pathol* 1987; 63(3):287-90.
13. Al-Jasser NM, Nwoku AL. Radiographic study of the mental foramen in a selected Saudi population. *Dentomaxillofacial Radiol*.1998, 27(6):341-3.
14. Rupesh S, Winnier JJ, John SA, Joy T, Rao AP, Reddy V. Radiographic Study of the Location of Mental Foramen in a Randomly Selected Asian Indian Population on Digital Panoramic Radiographs. *J Med Sci* 2011; 11(2):90-5.

15. Al-Khateeb T, Al-Hadi Hamesha A, Ababneh KT. Position of mental foramen in northern regional Jordanian population. *Surg Radiol Anat.* 2007; 29(3):231-7.
16. Olasoji HO, Tahir A, Ekanem AU, Abubakar AA. Radiographic and anatomic locations of mental foramen in northern Nigerian adults. *Niger Postgrad Med J.* 2004; 11(3):230-3.
17. Gangotri S, Patni VM, Sathwane RS. Radiographic Determination of Position and Symmetry of Mental Foramen in Central Indian Population. *J Indian Acad Oral Med Radiol.* 2011; 23(2):101-03.
18. Ngeow WC, Yuzawati Y. The location of the mental foramen in a selected Malay population. *J Oral Sci.* 2003; 45(3):171-5.
19. Mbajjorgu EF, Mawera G, Asala SA, Zivanovic S. Position of the mental foramen in adult black Zimbabwean mandibles: a clinical anatomical study. *Central African Journal of Medicine.* 1998; 44(2):24-30.
20. Mwaniki DL, Hassanali J. Position of mandibular and mental foramina in Kenyan African mandibles. *East Afr Med J.* 1992; 69(4):210-3.
21. Salah J Muhsen. Horizontal and vertical position of the mental foramen on panoramic views of a selected Iraqi population. *J Bagh Coll Dentistry* 2011; 23(1):69-72.
22. Al Jasser NM, Nwoku AL. Radiographic study of the mental foramen in Saudi females. *Saudi Med J* 1996; 17(4): 471-74.
23. Kim IS, Kim SG, Kim YK, Kim JD. Position of the Mental Foramen in a Korean Population: A Clinical and Radiographic Study. *Implant Dent* 2006; 15(4):404–11.