



Surgical practises for Odontogenic Keratocyst/ KCOT: A survey

Dr Manoj Kumar Bhaskaran, ¹MDS; Dr Joseph Lijo, ²MDS; Dr Rachana PB, ³MDS; Dr Sumana Devadiga, ⁴MDS; Dr Susan Thomas, ⁵MDS

Keywords: Odontogenic keratocyst; keratocystic odontogenic tumor; marsupialisation; enucleation with Carnoy's solution; resection.

Professor, Department of Oral and Maxillofacial Surgery, Sri Anjaneya Dental College, Calicut, Kerala, INDIA

Associate Professor, Department of Oral and Maxillofacial Surgery, Malabar Dental College, Edappal, Kerala, INDIA

Assistant Professor, Department of Oral and Maxillofacial Surgery, KVG Sullia, Sullia, Karnataka, INDIA

Associate Professor, Department of Oral and Maxillofacial Surgery, KVG Sullia, Sullia, Karnataka, INDIA

Professor, Department of Community Dentistry, Kerala, INDIA

How to cite this article: Manoj K Bhaskaran, Joseph Lijo, Rachana PB, Sumana Devadiga and Susan Thomas. Surgical practises for Odontogenic Keratocyst/ KCOT: A survey. Journal of Oral Medicine Surgery Pathology Biology. Dec-Jul 2016; 1(2): 49-55

Address for correspondence:

Dr Manoj K Bhaskaran, Professor and Head, Department of Oral and Maxillofacial Surgery, Sri Anjaneya Dental College, Calicut, Kerala, INDIA

Running title: Surgery for Odontogenic keratocyst

Clinical significance: The various techniques for treatment of Odontogenic keratocyst has been elaborated in this article

ABSTRACT

What is the best treatment for people who have odontogenic keratocysts? The treatment of the OKC has always been challenging in view of the marked recurrence rate. Treatment is either conservative or aggressive due to a high recurrence rate from 5 to 62%.

OKC was a term coined in the mid-1950s by European oral pathologists and the lesion arises from the cell rests of the dental lamina. Multiple cysts are seen as a feature of the nevoid basal cell carcinoma (Gorlin) Syndrome (NBCCS); an autosomal dominant inherited condition.

Aims and Objectives: The aim was to study the treatment protocols for the KCOT throughout the world. The objective was to trace the treatment modalities of OKC through the generations and to find an answer to the fairly focused question of whether there could be just one management approach that would be able to treat OKC and provide a good prognosis for the patient.

Materials and Methods: Data sources; PubMed, Scopus, Cochrane, etc. were searched for articles published in English language literature during the past 10 years that reported the treatment of OKCs by the surgeons throughout the world, and whether there have been any reported changes in them since the time “OKC” has become “KCOT.” Two reviewers independently appraised articles.

Results: The treatment method for KCOT is still ambiguous and all methods have their own advantages and drawbacks. Treatment protocol is based on the size, site, lesion accessibility and tissue destruction. Maximum number of cases was seen in males (69%) and the mandible was the most affected (84.5%) site. The treatment modality ranged from marsupialisation with decompression and enucleation to simple enucleation or enucleation followed by curettage, or packing it with chemicals such as Carnoy’s solution or iodoform gauze. The most preferred method of OKC removal was enucleation (26.8%) whereas marsupialisation either alone or with enucleation (1.4%) was a hardly preferred method of treatment.

Conclusion: OKC commonly recurs following surgery. The majority of recurrence commonly occurs within the first 5 years after treatment. Because of the problematic nature of these cysts, many attempts have been made to minimize the high recurrence rate by attempting to improve the surgical techniques. It has been found in this study, that there is probably no one single treatment modality that can be considered as ideal, to treat OKCs, as all the techniques mentioned have tasted success. None of the cases have reported recurrences in the 5 years follow-up period.

INTRODUCTION

OKC is a unique and characteristic developmental odontogenic cyst that requires special attention mainly because of its locally aggressive behaviour and high recurrence rate. WHO has included this highly aggressive cyst, in the classification of odontogenic tumors in the year 2005. They introduced the term “Keratocystic Odontogenic Tumor (KCOT)” for these lesions. Multiple odontogenic cysts are found in the Nevoid Basal cell carcinoma syndrome, which is an autosomal dominant condition¹


Treatment provided for OKC depends a lot on the training, management viewpoint and expertise of the institutions in different parts of the world. ¹

MATERIALS AND METHODS

The search was based on the research question “treatment modalities for OKC” and therefore, databases from the Pubmed interface of the MEDLINE (National Library of Medicine) and other Search Engines were accumulated. The MeSH terms used were “treatment for odontogenic keratocyst” and “Keratocystic Odontogenic tumor.”

Cases that reported the OKC s but which did not include the treatment modality were excluded from the study. They were tabulated based on their age, sex, site of occurrence and the mode of treatment followed.

RESULTS



Age: Mean age **Mal- Males** **Fema- Females** **Mand- Mandible** **Max-Maxilla** **Dec- Dec-**
DecompressionEnu- Enucleation **Mar-Marupalisation** **Curr- Currrettage** **Carn-Carnoy's**
solution Rese-Resection

Column1	No of cases	Age (Years)	Male	Female	MAND	Max	Enu	Mar	Decom	Enu+ Marsu	Enu+Cur	Enu+Marsu+Decomp	Column2	Enu+Carn	Resection
Total%	71	29	69%	31%	84.50%	15.50%	26.80%	1.40%	2.80%	1.40%	8.50%	4.20%		5.60%	5.60%

TABLE 1

Table 1: Clinical parameters of OKC and the different treatment modalities³⁻⁴⁰

Total number of indexed articles selected was 40 of which the total number of cases was 71. These articles reported the features mentioned in the inclusion criteria. Cases other than OKC/ KCOT or articles that did not stress on the treatment of the OKC were excluded from the selection. Of the total number of cases 69% were males and 31% were females. The cases were predominantly found affecting the mandible (84.5%) as compared to the maxilla (11cases). The enucleation (26.8%) was the preferred mode of treatment by majority of the surgeons, enucleation with curettage (8.5%), enucleation with application of a chemical cautery such as Carnoy's solution (5.6%); resection with plate placement was reported in 1 article (1.4%), excision with disarticulation in 1 article (1.4%); one article reported a hemimandibulectomy and 1 (1.4%) case of peripheral ostectomy. Two surgeons treated their cases with decompression followed by enucleation (4.6%) of which one surgeon had performed a decompression followed by placement of the iodoform gauze. One case reported treating the cyst by marsupialisation after decompression. Decompression as the only mode of treatment was

achieved in 2.8% cases and only marsupialisation and marsupialisation with enucleation was the procedure followed in 1.4% cases.

Enucleation and cautery by Carnoy's solution was performed at a mean age of 64.5 years, enucleation alone at 56.8 years, enucleation with curettage at 27.3 years, marsupialisation at an average age of 6 years, marsupialisation with enucleation at 22 years and resection at 75 years mean. Twenty two point five % males who had 21% involvement of the mandible underwent enucleation. 32.4% males underwent enucleation with cauterisation and 42.3% cases involved the mandible. Among the males the least opted (1.4%) treatment was enucleation with curettage and most opted was enucleation with cauterisation (32.4%). In females the resection was one of the less opted methods of treatment (1.4%). In the mandible the enucleation with marsupialisation (0)

was the least preferred mode of treatment and this treatment was the least favoured even in cases of the maxillary involvement (0).

DISCUSSION

OKCs/ KCOTs are one of the most aggressive odontogenic lesions that have a high tendency to recur. It requires special care as it is one of the most commonly featured odontogenic entities. The

vast number of previously reported cases supports the theory that this is one of the aggressive lesions that penetrates the cortical plates and involves adjacent soft tissues, even sometimes extending to the skull base from the mandible or in some cases to the infratemporal fossa and to the orbit.

Column1	Column2	Column3	Column4	Column5	Column6	Column8	Column9	Column10	Column11	Column12	Column13	Column14
Clinical Para Enuc	Decomp	Marsupialisation	Enu+ Marsu	Enu+Chemical	Enu+Marsu+Decomp	Enu+Cur	Resection					
Age	56.8	0	6	22	64.5	39	41	75				
Males	16	0	0	1	23	3	2	3				
Females	3	0	1	0	12	0	4	1				
Mand	15	0	0	1	30	3	4	4				
Max	4	0	1	0	5	0	2	0				

TABLE 2

Table 2: The age, sex and site predilection of Keratocystic Odontogenic Cysts and the surgical techniques practised, based on them

In our study and from recent articles searched, a permutation and combination of the conservative and aggressive treatments such as the marsupialisation, marsupialisation with decompression and enucleation together with cauterisation with Carnoy's fixative, was found to have won favour with the surgeons. Enucleation or marsupialisation alone have also been performed as a treatment modality. Resection with bone grafting was done in only a single case. This implies that the surgeons would still prefer a simple and age old procedure like decompression and enucleation; whatever may be the clinical and histologic features; as opposed to the recommended radical and disfiguring surgery.² The incidence of male prevalence being higher than females,

Clinical Parar	Enuc	Decomp	Marsupialisation	Enu+ Marsu	Enu+Chemical	Enu+Marsu+Decomp	Enu+Cur		Resection
Age	56.8	0	6	22	64.5	39	27.3(38.5%)		
Males	16 (22.5)	0	0	1(1.4%)	23(32.4%)	3(4.2%)	2(2.8%)		3(4.2%)
Females	3(4.2%)	0	1(1.4%)	0	12(16.9%)	0	4(5.6%)		1(1.4%)
Mand	15(21.1%)	0	0	1(1.4%)	30(42.3%)	3(4.2%)	4(5.6%)		4(5.6%)
Max	4(5.6%)	0	1(1.4%)	0	5(7%)	0	2(2.8%)		0

TABLE 3

Table 3: The age, sex and site predilection of KeratocysticOdontogenicTumor and the surgical treatment practised,in %age, based on the features.

amounting to 69% of the total 71 cases and the maxilla (15.5%) being less affected than mandible (84.5%),did not have any bearing on the mode of treatment practised.

CONCLUSION

No treatment modality can be considered as the ultimate for OKC/ KCOT. In recent years, two authors, Zhang and others have proposed the treatment of KCOT by molecular methods. They have postulated the use of antagonists of SHH signalling factors that has been said to efficiently treat KCOTs. They are of the belief that the intracystic injection of an SMO protein-antagonist has a great promise as a future treatment option, as opposed to the above-mentioned conservative and aggressive surgical modalities.

Footnotes:

Declaration of conflict: No conflict of interest

Source of funds: No funds were obtained towards the article.

REFERENCES

1. Jin Fei Yeo. Treatment of odontogenic keratocysts by enucleation with adjunctive therapy has lower recurrence rates. What treatment is best for people who have odontogenic keratocysts?Evidence-Based Dentistry. 2003; 4: 53–54
2. Mamta Singh and K. C. Gupta. Surgical treatment of odontogenic keratocyst by enucleation. Contemp Clin Dent. 2010 Oct-Dec; 1(4): 263–267
3. Kiran Rao and Sudesh Kumar. The Use of Enucleation and Chemical Cauterization (Carnoy's) in the Management of Odontogenic Keratocyst of the Jaws. Indian J Otolaryngol Head Neck Surg. 2014 Jan; 66(1): 8–12.
4. Daniel Antunes FREITAS, Daniela Araújo VELOSO, Alisson Luís D'Afonseca SANTOS, Viníciu Antunes FREITASRGO. Maxillary odontogenic keratocyst: a clinical case report. Rev Gaúch Odontol, Porto Alegre, Dec 2015; 63(4): 484-488
5. Guilherme Costa Carvalho Silvaa, Edgard Carvalho Silvab, Ricardo Santiago Gomez and Tainah Couto Vieir. Odontogenic keratocyst in the maxillary sinus: Report of two cases. Oral Oncology Extra. July 2006; 42(6): 231–234

6. Ellen Cristina Gaetti Jardim, Ana Cláudia Rossi, Leonardo Perez Faveran, et al. Odontogenic Keratocyst Tumor: Report of Two Cases. *Int. J. Odontostomat.* 2013;7(1):33-38
7. Vijay Ebenezer, R. Balakrishnan and M. Siva Kumar. A Case Report on Surgical Management of Odontogenic Keratocyst . *World Journal of Medical Sciences.* 2014. 10 (2): 212-216,
8. Balasubramanian Thiagarajan Geetha Ramamoorthy. Odontogenic Keratocyst Mandible an Interesting Case report and Literature review. rhinology.in/12/1/okc.pdf
9. Ajit Auluck, MDS; Setty Suhas, MDS; Keerthilatha M. Pai. Multiple Odontogenic Keratocysts: Report of a Case. *J Can Dent Assoc* 2006; 72(7):651–6
10. Rajkumar G C, Hemalatha M, Shashikala R, Sonal P. Massive keratocystic odontogenic tumor of mandible: A case report and review of literature. *Indian J Dent Res* 2011;22:181
11. N K Suma, C Pinky, N S Venkatesh Babu, Smriti Jha. Odontogenic Keratocyst of Maxillary Premolar Region: A Case Report. *IJSS Case Reports & Reviews.* February 2015; 1(9): 38-41
12. Satish Karandikar, Kumar Nilesh, Anuj Dadhich. Large Odontogenic Keratocyst of Mandible - A Case Report. *Journal of the Indian Dental Association.* January 2011; 5(7):822-824
13. Contar et al. Keratocystic odontogenic tumor: case report. *Odonto* 2011; 19 (37): 25-32
14. Arshad F. Syndromic odontogenic keratocyst: A case report and review of literature. *J Int Soc Prevent Communit Dent.* 2016;6:84-8
15. A Chkoura, S Chbicheb, W El wady .Keratocystic Odontogenic Tumor: A Case Report and Review of the Literature. *ISPUB.COM.* 2008; 6(2): 1-6
16. K. M. Veena, Rekha Rao, H. Jagadishchandra, and Prasanna Kumar Rao. Odontogenic Keratocyst Looks Can Be Deceptive, Causing Endodontic Misdiagnosis. *Case Reports in Pathology.* Volume 2011 (2011), Article ID 159501, 3 pages
17. Belkacem Raouâa, Sioud Samèh, Touil Dorsaf, Ayachi Rahma, Selmi Jamil. Recurrence challenge in odontogenic keratocyst variants, two clinical cases. *Med Buccale Chir Buccale* 2014;20:203-208
18. Belkacem Raouâa, Sioud Samèh, Touil Dorsaf, Ayachi Rahma, Selmi Jamil. Recurrence challenge in odontogenic keratocyst variants, two clinical cases. *Med Buccale Chir Buccale* 2014;20:203-208
19. Shruti Bohra, Rohan Bhede, Hrushikesh Saraf, Minal Chaudhary, Nisu Swastika, Alka Hande. Radicular cyst or Odontogenic keratocyst? : A case report. *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS).* Jun 2014; 13(6) Ver. V: 53-57
20. Belmehdi, A., Chbicheb, S. and El Wady, W. Odontogenic Keratocyst Tumor: A Case Report and Literature Review. *Open Journal of Stomatology.*2016; 6: 171-178.
21. SES Faustino, MC Pereira, AC Rossetto and DT Oliveira. Recurrent peripheral odontogenic keratocyst: a case report. *Dentomaxillofacial Radiology.* 2008; 37: 412–41.
22. Rory O'Neill and Khadry A. Galil. Identification of an Odontogenic Keratocyst and Treatment with Guided Tissue Regeneration: Case Report. *J Can Dent Assoc* 2011;77:b6
23. Santhosh kumar S Hiremath, Atul M. Deshpande, Sanjay Byakodi, Dilip B. Magdum. Diagnostic Dilemma: A Case Report of Odontogenic Keratocyst in Lateral Periodontal Position. *International Journal of Oral & Maxillofacial Pathology;* 2011;2(3):23-26
24. Sabri C. Isler, Sabit Demircan, Taylan Can, Zerrin Cebi, Emrah Baca, Immediate Implants After Enucleation of an Odontogenic Keratocyst: An Early Return to Function, *Journal of Oral Implantology.* 2012; 38(S1):485-488.
25. Rajpal J, Arora A, Arora P, Prasad R. Deceptive periodontal pocket diagnosed later as odontogenic keratocyst. *J Interdiscip Dentistry* 2015; 5: 87-91
26. Rawson K, Kallalli BN, Telkar S, Penumatcha MR. Keratocystic odontogenic tumor of the right mandibular condyle: A rare case . *J Indian Acad Oral Med Radiol* 2014; 26: 103-6
27. Sabin Siddique, Shaji T. Varghese and Tom Mathew. Odontogenic Keratocysts of the Mandible- A case report. *wjpmr,* 2016; 2(4): 213-215
28. Garde JB, Kulkarni AU, Dadhe DP. Implant-based rehabilitation of a large mandibular odontogenic keratocyst with 7-year follow-up. *J Int Clin Dent Res Organ* 2010; 2:95-7
29. Rohan Kishor Gupta, Arun Govind Dugal, Sudhir Ramlal Pawar, Saurabh Girish Khandelwal, Apoorva Iyengar A Rare Simultaneous Occurrence of Odontogenic Keratocyst and Unicystic Ameloblastoma

- inMandible: A Case Report.. Journal of Clinical and Diagnostic Research. 2016 Aug; 10(8): ZD01-ZD04
30. Amila Brkić, Hülya Koçak-Berberoglu, Esra Eyupoğlu, Vakur Olgaç Prosthetic rehabilitation with dental implants after treatment of a keratocystic odontogenic tumor. A case report. J Clin Exp Dent. 2011; 3(Suppl1):e360-4.
 31. P.A. Dowling, P. Fleming, I.D.F. Saunders, R.J. Gorlin, S.S. Napier. Odontogenic keratocysts in a 5-year-old: Initial manifestations of nevoid basal cell carcinoma syndrome. Pediatric Dentistry. 2000; 22: 1
 32. Sarita Yanduri Veerendra Kumar B , Shyamala K , Girish Rao Orthokeratinized Odontogenic Cyst.. IJDA Archives. 02(01)
 33. Kalia V, Kaushal N, Kalra G. The syndromic multiple odontogenic keratocyst in siblings: A familial study. Ann Maxillofac Surg 2011; 1:77-82
 34. Guruprasad Y, Chauhan DS. Multiple odontogenic keratocysts in a nonsyndromic patient. J Cranio Max Dis 2012;1:36-40
 35. Yuko Tanimoto, Shouichi Miyawaki, Mikako Imai, Ryoko Takeda, and Teruko Takano-Yamamoto. Orthodontic Treatment of a Patient with an Impacted Maxillary Second Premolar and Odontogenic Keratocyst in the Maxillary Sinus. The Angle Orthodontist: November 2005; 75(6): 1077-1083.
 36. Rai S, Gauba K. Jaw cyst-Basal cell nevus-Bifid rib syndrome: A case report. J Indian Soc Pedod Prev Dent 2007; 25:137-9
 37. Molon RS, Verzola MH, Pires LC, Mascarenhas VI, da Silva RB, Cirelli JA, Barbeiro RH. Five years follow-up of a keratocyst odontogenic tumor treated by marsupialization and enucleation: A case report and literature review. Contemp Clin Dent 2015;6, Suppl S1:106-10
 38. Pushparaja Shetty, SV Srilatha. Keratocystic odontogenic tumor with mural calcification: A case report. South Asian J Cancer. 2013 Oct-Dec; 2(4): 208
 39. Jonathan Madras and Henry Lapointe. Keratocystic Odontogenic Tumour: Reclassification of the Odontogenic Keratocyst from Cyst to Tumour. JCDA. March 2008; 74(2) : 165a-h
 40. E. Ciulli, M. Rocci, R. Bollero, C. Pandolfi, L. Ottria, G. Mampieri, A. Barlattani, Jr., and P. Bollero. Maxillary cyst: description of a clinical case. Oral Implantol (Rome). 2009 Apr-Jun; 2(2): 28–33.