

# Degree Sets: Realizability and Extension Problems

Prasun Kumar

CS11S005

Advisor : Jayalal Sarma M.N.

Department of Computer Science and Engineering

Indian Institute of Technology, Madras

April 29, 2013

# Talk Plan


- Degree Set Basics
- Extension Problem for Trees
- Extension Problems for Undirected Graphs
- Tree Realizability under Multiplicity Constraints
- Degree Set Variants for Directed Graphs
- Asymmetric Directed Graph Realization of Degree Set
- Directed Tree Realization of Degree Set
- Conclusion and Future Work

# Degree Set : Definition

Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{deg(v): v \in V\}$


# Degree Set : Definition

Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{deg(v): v \in V\}$



# Degree Set : Definition


Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{\deg(v) : v \in V\}$



For this graph  
 $S = \{2, 3\}$

# Degree Set : Definition

Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{\deg(v) : v \in V\}$




For this graph

$$S = \{2, 3\}$$

$$D = (3, 3, 2, 2)$$

# Degree Set : Definition

Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{deg(v): v \in V\}$



For this graph


$$S = \{2, 3\}$$

$$D = (3, 3, 2, 2)$$

- Convention : elements in  $S$  are written in increasing order.

# Degree Set : Definition

Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{deg(v): v \in V\}$



For this graph

$$S = \{2, 3\}$$


$$D = (3, 3, 2, 2)$$

- Convention : elements in  $S$  are written in increasing order.
- $S$  is *realizable* if there is a graph respecting  $S$ .



# Degree Set : Definition

Let  $G(V, E)$  be a graph then degree set of  $G$  is  $S = \{deg(v): v \in V\}$



For this graph

$$S = \{2, 3\}$$

$$D = (3, 3, 2, 2)$$

- Convention : elements in  $S$  are written in increasing order.
- $S$  is *realizable* if there is a graph respecting  $S$ .
- Interested in *Simple graph* realizations only.

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

# Degree Set

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

- Is  $S$  realizable?

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

- Is  $S$  realizable?
- What is the minimum order possible if  $S$  is realizable?

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

- Is  $S$  realizable?
- What is the minimum order possible if  $S$  is realizable?

*If  $S$  is realizable then*

- $\Gamma(S)$  : family of simple graphs realizing  $S$

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

- Is  $S$  realizable?
- What is the minimum order possible if  $S$  is realizable?

*If  $S$  is realizable then*

- $\Gamma(S)$  : family of simple graphs realizing  $S$
- $\mu(S)$  : minimum order of a graph  $\in \Gamma(S)$

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

- Is  $S$  realizable?
- What is the minimum order possible if  $S$  is realizable?

*If  $S$  is realizable then*

- $\Gamma(S)$  : family of simple graphs realizing  $S$
- $\mu(S)$  : minimum order of a graph  $\in \Gamma(S)$

$$\mu(S) \geq a_n + 1$$

For a given set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$ , there arise two questions of interest -

- Is  $S$  realizable?
- What is the minimum order possible if  $S$  is realizable?

*If  $S$  is realizable then*

- $\Gamma(S)$  : family of simple graphs realizing  $S$
- $\mu(S)$  : minimum order of a graph  $\in \Gamma(S)$


$$\mu(S) \geq a_n + 1$$

When  $S$  - realized by a tree then we use  $\mu_T(S)$  instead of  $\mu(S)$ .




eg.  $S = \{1, 2, 3\}$


eg.  $S = \{1, 2, 3\}$



$|V| = 4$




$|V| = 5$




$|V| = 5$


eg.  $S = \{1, 2, 3\}$




$|V| = 4$



$|V| = 5$



$|V| = 5$




$|V| = 6$


$G: \mathbb{K}_4 \cup \mathbb{K}_3 \cup \mathbb{K}_2$

$|V| = 9$


eg.  $S = \{1, 2, 3\}$




$|V| = 4$



$|V| = 5$



$|V| = 5$



$|V| = 6$

$G: \mathbb{K}_4 \cup \mathbb{K}_3 \cup \mathbb{K}_2$

$|V| = 9$

$\mu(S) = 4$

# Integer Set Realization

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is always realizable and  $\mu(S) = a_n + 1$ .

# Integer Set Realization

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is always realizable and  $\mu(S) = a_n + 1$ .

Proof: by induction on  $|S| (= n)$

# Integer Set Realization

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is always realizable and  $\mu(S) = a_n + 1$ .

Proof: by induction on  $|S| (= n)$

Another **Computational Question** that we are asking

## Extension Problem

*Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = t$  ??*

# Integer Set Realization

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is always realizable and  $\mu(S) = a_n + 1$ .

Proof: by induction on  $|S| (= n)$

Another **Computational Question** that we are asking

## Extension Problem

*Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = t$  ??*

- $t < \mu(S)$  - no realization exists



# Integer Set Realization

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is always realizable and  $\mu(S) = a_n + 1$ .

Proof: by induction on  $|S| (= n)$

Another **Computational Question** that we are asking

## Extension Problem

*Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = t$  ??*

- $t < \mu(S)$  - no realization exists
- $t \geq \mu(S)$  then  $t = \mu(S) + r$  where  $r \in \mathbb{Z} \cup \{0\}$

# Integer Set Realization of Trees and TEP

## Theorem - Kapoor, Polimeni, Wall(1977)


Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree *if and only if*  $a_1 = 1$  and  $\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$

# Integer Set Realization of Trees and TEP

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree *if and only if*  $a_1 = 1$  and  $\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$

Minimum order tree construction




# Integer Set Realization of Trees and TEP

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree *if and only if*  $a_1 = 1$  and  $\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$

Minimum order tree construction



Let  $m_i$  be the *multiplicity* of  $a_i \in S$  then in this tree


- $m_1 \gg 1$  and  $m_i = 1, \forall 2 \leq i \leq n$ .

# Integer Set Realization of Trees and TEP

## Theorem - Kapoor, Polimeni, Wall(1977)

Any finite set  $S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree *if and only if*  $a_1 = 1$  and  $\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$

## Minimum order tree construction



Let  $m_j$  be the *multiplicity* of  $a_j \in S$  then in this tree

- $m_1 \gg 1$  and  $m_i = 1, \forall 2 \leq i \leq n$ .

# Tree Extension Problem(TEP)

Given a degree set  $S$  with  $a_1 = 1$ , and an integer  $r$ , check if there is a tree  $T'(V', E') \in \Gamma(S)$  such that  $|V'| = \mu_T(S) + r$ .

# Tree Extension Characterization

## Characterization

If the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is realized by a tree  $T(V, E)$  then there is another tree realization  $T' = (V', E')$  where  $|V'| = |V| + r, r \in \mathbb{Z}^+$ , *if and only if*

$$r = k_2(a_2 - 1) + k_3(a_3 - 1) + \dots + k_n(a_n - 1) \quad \forall i, k_i \in \mathbb{Z}^+ \cup \{0\}$$

# Tree Extension Characterization

## Characterization

If the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is realized by a tree  $T(V, E)$  then there is another tree realization  $T' = (V', E')$  where  $|V'| = |V| + r, r \in \mathbb{Z}^+$ , if and only if

$$r = k_2(a_2 - 1) + k_3(a_3 - 1) + \dots + k_n(a_n - 1) \quad \forall i, k_i \in \mathbb{Z}^+ \cup \{0\}$$

*Result due to Gupta, Joshi, Tripathi (2004).*

# Tree Extension Characterization

## Characterization

If the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is realized by a tree  $T(V, E)$  then there is another tree realization  $T' = (V', E')$  where  $|V'| = |V| + r, r \in \mathbb{Z}^+$ , if and only if

$$r = k_2(a_2 - 1) + k_3(a_3 - 1) + \dots + k_n(a_n - 1) \quad \forall i, k_i \in \mathbb{Z}^+ \cup \{0\}$$

*Result due to Gupta, Joshi, Tripathi (2004).*

Proof (Aliter): ( $\Rightarrow$ ) W.l.o.g. assume  $T(V, E)$  is a minimum order tree realizing  $S$  and let out of  $r$  vertices,  $k_i$  vertices are there with degree  $a_i \in S$ .



# Tree Extension Characterization

Hence

$$(\textcolor{red}{1} \sum_{i=1}^n \textcolor{blue}{a_i} - 2n + 3 + k_1, \textcolor{red}{a_2}^{1+k_2}, \dots, \textcolor{red}{a_i}^{1+k_i}, \dots, \textcolor{red}{a_n}^{1+k_n})$$

is the degree sequence of  $T'$  and  $r = \sum_{i=1}^n k_i$ .

# Tree Extension Characterization

Hence

$$(\textcolor{blue}{1}^{\sum_{i=1}^n a_i - 2n + 3 + k_1}, \textcolor{red}{a}_2^{1+k_2}, \dots, \textcolor{red}{a}_i^{1+k_i}, \dots, \textcolor{red}{a}_n^{1+k_n})$$

is the degree sequence of  $T'$  and  $r = \sum_{i=1}^n k_i$ .

lemma(AM96): For the sequence  $d = (d_1 \geq d_2 \geq \dots \geq d_n)$ , if  $\sum_{i=1}^n d_i = 2(n-1)$  then the number of pendant vertices in any tree realization of  $d$  is  $\sum_{i=1}^k (d_i - 2) + 2$  where  $k = \max\{i \mid d_i \geq 3\}$

# Tree Extension Characterization

Hence

$$(\color{blue}{1} \sum_{i=1}^n a_i - 2n + 3 + k_1, \color{red}{a_2}^{1+k_2}, \dots, \color{red}{a_i}^{1+k_i}, \dots, \color{red}{a_n}^{1+k_n})$$

is the degree sequence of  $T'$  and  $r = \sum_{i=1}^n k_i$ .

lemma(AM96): For the sequence  $d = (d_1 \geq d_2 \geq \dots \geq d_n)$ , if  $\sum_{i=1}^n d_i = 2(n-1)$  then the number of pendant vertices in any tree realization of  $d$  is  $\sum_{i=1}^k (d_i - 2) + 2$  where  $k = \max\{i \mid d_i \geq 3\}$

Since  $a_2 \geq 2$

- $\sum_{i=1}^n a_i - 2n + 3 + k_1 = \sum_{i=2}^n (k_i + 1)(a_i - 2)$

# Tree Extension Characterization

Hence

$$(\textcolor{red}{1} \sum_{i=1}^n a_i - 2n + 3 + k_1, \textcolor{red}{a}_2^{1+k_2}, \dots, \textcolor{red}{a}_i^{1+k_i}, \dots, \textcolor{red}{a}_n^{1+k_n})$$

is the degree sequence of  $T'$  and  $r = \sum_{i=1}^n k_i$ .

lemma(AM96): For the sequence  $d = (d_1 \geq d_2 \geq \dots \geq d_n)$ , if  $\sum_{i=1}^n d_i = 2(n-1)$  then the number of pendant vertices in any tree realization of  $d$  is  $\sum_{i=1}^k (d_i - 2) + 2$  where  $k = \max\{i \mid d_i \geq 3\}$

Since  $a_2 \geq 2$

- $\sum_{i=1}^n a_i - 2n + 3 + k_1 = \sum_{i=2}^n (k_i + 1)(a_i - 2)$
- $r = \sum_{i=1}^n k_i$

Solving these we get  $r = \sum_{i=2}^n k_i (a_i - 1)$ .

# Complexity Results on TEP

TREE EXTENSION PROBLEM (TEP) is equivalent to INTEGER KNAPSACK PROBLEM (IKP).

# Complexity Results on TEP

TREE EXTENSION PROBLEM (TEP) is equivalent to INTEGER KNAPSACK PROBLEM (IKP).

IKP : Given non-negative integers  $c_1, \dots, c_k$ , and a value  $d$ . Does there exist non-negative integers  $d_1, d_2, \dots, d_k$  such that  $\sum_i c_i d_i = d$ .

# Complexity Results on TEP

TREE EXTENSION PROBLEM (TEP) is equivalent to INTEGER KNAPSACK PROBLEM (IKP).

IKP : Given non-negative integers  $c_1, \dots, c_k$ , and a value  $d$ . Does there exist non-negative integers  $d_1, d_2, \dots, d_k$  such that  $\sum_i c_i d_i = d$ .

Given a degree set  $S$ , corresponding IKP instance

- $k = |S| - 1$ ,  $c_i = a_{i+1} - 1$  for all  $1 \leq i \leq k$ , and  $d = r$ .

$$\begin{array}{ccccccc} r = & k_2(a_2 - 1) + & \dots + & k_{i+1}(a_{i+1} - 1) + & \dots + & k_n(a_n - 1) \\ \updownarrow & \updownarrow & & \updownarrow & & \updownarrow \\ d = & d_1 c_1 + & \dots + & d_i c_i + & \dots + & d_k c_k \end{array}$$

# Complexity Results on TEP

TREE EXTENSION PROBLEM (TEP) is equivalent to INTEGER KNAPSACK PROBLEM (IKP).

IKP : Given non-negative integers  $c_1, \dots, c_k$ , and a value  $d$ . Does there exist non-negative integers  $d_1, d_2, \dots, d_k$  such that  $\sum_i c_i d_i = d$ .

Given a degree set  $S$ , corresponding IKP instance

- $k = |S| - 1$ ,  $c_i = a_{i+1} - 1$  for all  $1 \leq i \leq k$ , and  $d = r$ .

$$\begin{array}{ccccccc} r = & k_2(a_2 - 1) + & \dots + & k_{i+1}(a_{i+1} - 1) + & \dots + & k_n(a_n - 1) \\ \updownarrow & \updownarrow & & \updownarrow & & \updownarrow \\ d = & d_1 c_1 + & \dots + & d_i c_i + & \dots + & d_k c_k \end{array}$$

Given non-negative integers  $c_1, \dots, c_k$ , and a value  $d$  (an IKP instance), consider the degree set

- $S = \{1, c_1 + 1, \dots, c_k + 1\}$  and  $r = d$ .



# Complexity Results on TEP

TREE EXTENSION PROBLEM(TEP) is equivalent to INTEGER KNAPSACK PROBLEM(IKP).

IKP : Given non-negative integers  $c_1, \dots, c_k$ , and a value  $d$ . Does there exist non-negative integers  $d_1, d_2, \dots, d_k$  such that  $\sum_i c_i d_i = d$ .

Given a degree set  $S$ , corresponding IKP instance

- $k = |S| - 1$ ,  $c_i = a_{i+1} - 1$  for all  $1 \leq i \leq k$ , and  $d = r$ .

$$\begin{array}{ccccccc} r = & k_2(a_2 - 1) + & \dots + & k_{i+1}(a_{i+1} - 1) + & \dots + & k_n(a_n - 1) \\ \updownarrow & \updownarrow & & \updownarrow & & \updownarrow \\ d = & d_1 c_1 + & \dots + & d_i c_i + & \dots + & d_k c_k \end{array}$$

Given non-negative integers  $c_1, \dots, c_k$ , and a value  $d$  (an IKP instance), consider the degree set

- $S = \{1, c_1 + 1, \dots, c_k + 1\}$  and  $r = d$ .

(TEP is NP-Complete.)

# Unary Version of TEP

## Unary Tree Extension Problem(UTEP)

Given a tree  $T$  on  $\ell$  vertices and a string  $1^r$ , test if there is another tree  $T'$  having exactly  $\ell + r$  vertices and the degree set same as that of  $T$ .

# Unary Version of TEP

## Unary Tree Extension Problem(UTEP)

Given a tree  $T$  on  $\ell$  vertices and a string  $1^r$ , test if there is another tree  $T'$  having exactly  $\ell + r$  vertices and the degree set same as that of  $T$ .

*UTEP can be solved in log-space.*

# Unary Version of TEP

## Unary Tree Extension Problem(UTEP)

Given a tree  $T$  on  $\ell$  vertices and a string  $1^r$ , test if there is another tree  $T'$  having exactly  $\ell + r$  vertices and the degree set same as that of  $T$ .

*UTEP can be solved in log-space.*

Proof: By reducing the problem to Unary Subset Sum Problem which can be solved in log-space (Ref- Kane2010).

# Unary Version of TEP

## Unary Tree Extension Problem(UTEP)

Given a tree  $T$  on  $\ell$  vertices and a string  $1^r$ , test if there is another tree  $T'$  having exactly  $\ell + r$  vertices and the degree set same as that of  $T$ .

*UTEP can be solved in log-space.*

Proof: By reducing the problem to Unary Subset Sum Problem which can be solved in log-space (Ref- Kane2010).

## UNARY SUBSET SUM PROBLEM

Given a multiset  $A$  of  $m$  integers  $b_1, b_2, \dots, b_m$  and a value  $c$  (all inputs in unary), test if there is a subset  $A'$  of these integers such that  $\sum_{i \in A'} b_i = c$

# Reduction from UTEP to UNARY SUBSET SUM

Reduction: Given a tree  $T$  and  $r$  in unary

- write down the set  $A = \bigcup_{i=2, j=1}^{i=n, j=t_i} \{(a_i - 1)j\}$  where  $t_i = \lceil \frac{r}{a_i - 1} \rceil$  and  $r$  in unary, choose  $c = r$ .

# Reduction from UTEP to UNARY SUBSET SUM

Reduction: Given a tree  $T$  and  $r$  in unary

- write down the set  $A = \bigcup_{i=2, j=1}^{i=n, j=t_i} \{(a_i - 1)j\}$  where  $t_i = \lceil \frac{r}{a_i - 1} \rceil$  and  $r$  in unary, choose  $c = r$ .

$$r = k_2(a_2 - 1) + \cdots + k_i(a_i - 1) + \cdots + k_n(a_n - 1), \quad k_i \leq \lceil \frac{r}{a_i - 1} \rceil$$

# Reduction from UTEP to UNARY SUBSET SUM

Reduction: Given a tree  $T$  and  $r$  in unary

- write down the set  $A = \bigcup_{i=2, j=1}^{i=n, j=t_i} \{(a_i - 1)j\}$  where  $t_i = \lceil \frac{r}{a_i - 1} \rceil$  and  $r$  in unary, choose  $c = r$ .

$$r = k_2(a_2 - 1) + \cdots + k_i(a_i - 1) + \cdots + k_n(a_n - 1), \quad k_i \leq \lceil \frac{r}{a_i - 1} \rceil$$

If  $\exists A' \subseteq A$  that sums up to  $r$ , then corresponding choice of the  $j$ 's satisfies equation  $r = \sum_{i=2}^n k_i(a_i - 1)$  and for any solution  $k_i \leq t_i$  for all  $i$ .



# Reduction from UTEP to UNARY SUBSET SUM

Reduction: Given a tree  $T$  and  $r$  in unary

- write down the set  $A = \bigcup_{i=2, j=1}^{i=n, j=t_i} \{(a_i - 1)j\}$  where  $t_i = \lceil \frac{r}{a_i - 1} \rceil$  and  $r$  in unary, choose  $c = r$ .

$$r = k_2(a_2 - 1) + \cdots + k_i(a_i - 1) + \cdots + k_n(a_n - 1), \quad k_i \leq \lceil \frac{r}{a_i - 1} \rceil$$

If  $\exists A' \subseteq A$  that sums up to  $r$ , then corresponding choice of the  $j$ 's satisfies equation  $r = \sum_{i=2}^n k_i(a_i - 1)$  and for any solution  $k_i \leq t_i$  for all  $i$ .

Hence the corresponding terms  $k_i(a_i - 1)$  will appear in the set  $A$  as well. Choosing these terms in  $A'$  ensures  $\sum_{i \in A'} b_i = r = c$ .

# Parametrizations of TEP ( $r$ given in *unary*)

Two natural parametrizations of TEP (when  $r$  is given in *unary*)-

# Parametrizations of TEP ( $r$ given in *unary*)

Two natural parametrizations of TEP (when  $r$  is given in *unary*)-

- with  $|S|$  as the parameter
  - $\text{TEP} \leq_P \text{VARIETY SUBSET SUM PROBLEM}$
  - *Ref-* Fellows, Gaspers, Rosamond(2010)

# Parametrizations of TEP ( $r$ given in *unary*)

Two natural parametrizations of TEP (when  $r$  is given in *unary*)-

- with  $|S|$  as the parameter
  - $\text{TEP} \leq_P \text{VARIETY SUBSET SUM PROBLEM}$
  - *Ref-* Fellows, Gaspers, Rosamond(2010)
- with  $r$  as parameter
  - $\text{TEP} \leq_P \text{MAXIMUM KNAPSACK PROBLEM}$
  - *Ref-* Fernau(2005)

# Extension Problem for Undirected Graphs

Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = \mu(S) + r$  ?

# Extension Problem for Undirected Graphs

Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = \mu(S) + r$  ?

*Since  $|V'|$  is given  $\Rightarrow$  Degree sequence approach might be useful*

# Extension Problem for Undirected Graphs

Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = \mu(S) + r$  ?

*Since  $|V'|$  is given  $\Rightarrow$  Degree sequence approach might be useful*

Let  $D$  be the *degree sequence* of a graph  $G(V, E)$  realizing  $S$ ,  $|V| = \mu(S)$ .

# Extension Problem for Undirected Graphs

Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = \mu(S) + r$  ?

*Since  $|V'|$  is given  $\Rightarrow$  Degree sequence approach might be useful*

Let  $D$  be the *degree sequence* of a graph  $G(V, E)$  realizing  $S$ ,  $|V| = \mu(S)$ .

$$D' \longleftarrow (D, a_{i_1}, a_{i_1}, a_{i_2}, \dots, a_{i_r}) \quad \text{such that } \forall q \in [r], a_{i_q} \in S$$



# Extension Problem for Undirected Graphs

Is there a graph  $G'(V', E') \in \Gamma(S)$  such that  $|V'| = \mu(S) + r$ ?

*Since  $|V'|$  is given  $\Rightarrow$  Degree sequence approach might be useful*

Let  $D$  be the *degree sequence* of a graph  $G(V, E)$  realizing  $S$ ,  $|V| = \mu(S)$ .

$$\begin{array}{c} \textcircled{D'} \longleftarrow (\textcircled{D}, a_{i_1}, a_{i_1}, a_{i_2}, \dots, a_{i_r}) \quad \text{such that } \forall q \in [r], a_{i_q} \in S \\ \nearrow \text{graphic??} \quad \nwarrow \text{graphic} \end{array}$$

# Integer Sequence Realization

## Erdős - Gallai Theorem (1960)

The positive integer sequence  $d = (d_1 \geq d_2 \geq \dots \geq d_i \geq \dots \geq d_n)$ , where  $d_1 \leq n - 1$ , is realized by a simple graph if

- $\sum_{i=1}^n d_i$  is even, and
- $\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{d_i, k\}$  for  $1 \leq k \leq n$

# Integer Sequence Realization

## Erdős - Gallai Theorem (1960)

The positive integer sequence  $d = (d_1 \geq d_2 \geq \dots \geq d_i \geq \dots \geq d_n)$ , where  $d_1 \leq n - 1$ , is realized by a simple graph if

- $\sum_{i=1}^n d_i$  is even, and
- $\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{d_i, k\}$  for  $1 \leq k \leq n$

ZZ92 : suffices to check the inequalities for  $1 \leq k \leq m$  where

$$m = \max\{i \mid d_i \geq i, d_i \in d\}.$$

# Integer Sequence Realization

## Erdős - Gallai Theorem (1960)

The positive integer sequence  $d = (d_1 \geq d_2 \geq \cdots \geq d_i \geq \cdots \geq d_n)$ , where  $d_1 \leq n - 1$ , is realized by a simple graph if

- $\sum_{i=1}^n d_i$  is even, and
- $\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{d_i, k\}$  for  $1 \leq k \leq n$

ZZ92 : suffices to check the inequalities for  $1 \leq k \leq m$  where

$$m = \max\{i | d_i \geq i, d_i \in d\}.$$

↓

$$\begin{array}{ccccccc} \text{indices :} & \underline{1} & \underline{2} & \cdots & \underline{\mathbf{m}} & \underline{m+1} & \cdots & \underline{n} \\ d & : & \textcolor{red}{d_1} & \textcolor{red}{d_2} & \cdots & \textcolor{red}{d_m} & \textcolor{blue}{d_{m+1}} & \cdots & \textcolor{blue}{d_n} \\ & & \underbrace{\hspace{1.5cm}} & & & & \underbrace{\hspace{1.5cm}} & & \\ & & & d_i \geq j & & & & d_i < j & \end{array}$$

# Integer Sequence Realization

## Erdős - Gallai Theorem (1960)

The positive integer sequence  $d = (d_1 \geq d_2 \geq \dots \geq d_i \geq \dots \geq d_n)$ , where  $d_1 \leq n-1$ , is realized by a simple graph if

- $\sum_{i=1}^n d_i$  is even, and
- $\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{d_i, k\}$  for  $1 \leq k \leq n$

ZZ92 : suffices to check the inequalities for  $1 \leq k \leq m$  where

$$m = \max\{i \mid d_i \geq i, d_i \in d\}.$$

$$\begin{array}{ccccccc} \text{indices :} & \underline{1} & \underline{2} & \dots & \underline{m} & \underline{m+1} & \dots & \underline{n} \\ d & : & d_1 & d_2 & \dots & d_m & d_{m+1} & \dots & d_n \\ & & \underbrace{\hspace{1.5cm}}_{d_j \geq j} & & \underbrace{\hspace{1.5cm}}_{d_j < j} & & & & \end{array}$$

Idea : (can add  $d_{j'} < j'$  to get another graphic sequence  $d'$ )

# Extension Problem for Undirected Graphs

- $G(V, E)$  realizes  $S = \{a_1 < a_2 < \dots < a_n\}$ , where  $|V| = a_n + 1$ .
- Multiplicity( $m_i$ ) of different  $a_i(\in S)$ s in  $G$  are as follows

# Extension Problem for Undirected Graphs

- $G(V, E)$  realizes  $S = \{a_1 < a_2 < \dots < a_n\}$ , where  $|V| = a_n + 1$ .
- Multiplicity( $m_i$ ) of different  $a_i(\in S)$ s in  $G$  are as follows
  - for  $i \neq n, \lceil \frac{n}{2} \rceil$  :  $m_i = a_{n-i+1} - a_{n-i}$
  - for  $i = n$  :  $m_n = a_1$
  - for  $i = \lceil \frac{n}{2} \rceil$  :  $m_{\lceil \frac{n}{2} \rceil} = a_{n-\lceil \frac{n}{2} \rceil+1} - a_{n-\lceil \frac{n}{2} \rceil} + 1$

# Extension Problem for Undirected Graphs

- $G(V, E)$  realizes  $S = \{a_1 < a_2 < \dots < a_n\}$ , where  $|V| = a_n + 1$ .
- Multiplicity( $m_i$ ) of different  $a_i (\in S)$ s in  $G$  are as follows
  - for  $i \neq n, \lceil \frac{n}{2} \rceil$  :  $m_i = a_{n-i+1} - a_{n-i}$
  - for  $i = n$  :  $m_n = a_1$
  - for  $i = \lceil \frac{n}{2} \rceil$  :  $m_{\lceil \frac{n}{2} \rceil} = a_{n-\lceil \frac{n}{2} \rceil+1} - a_{n-\lceil \frac{n}{2} \rceil} + 1$

The corresponding degree sequence of  $G$  will be

$\mathbf{D} = (d_1 \geq d_2 \geq \dots \geq d_j \geq \dots \geq d_{a_n+1})$  where

$$d_j = \begin{cases} a_n & \text{if } 1 \leq j \leq a_1 \\ a_{n-i} & \text{if } 1 + a_i \leq j \leq a_{i+1}, \text{ for } 1 \leq i \leq \lfloor \frac{n}{2} \rfloor - 1 \\ a_{\lceil \frac{n}{2} \rceil} & \text{if } 1 + a_{\lfloor \frac{n}{2} \rfloor} \leq j \leq 1 + a_{\lfloor \frac{n}{2} \rfloor+1} \\ a_{n-i} & \text{if } 2 + a_i \leq j \leq 1 + a_{i+1}, \text{ for } \lfloor \frac{n}{2} \rfloor + 1 \leq i \leq n - 1 \end{cases}$$



# Extension Problem for Undirected Graphs

## Theorem

For every  $r \in \mathbb{Z}^+$ , there exists an undirected simple graph with  $\mu(S) + r$  vertices having degree set  $S = \{a_1 < a_2 < \dots < a_n\}$  except the case when  $a_1 = 1$ ,  $r$  is odd and only  $a_n$  is even.

# Extension Problem for Undirected Graphs

## Theorem

For every  $r \in \mathbb{Z}^+$ , there exists an undirected simple graph with  $\mu(S) + r$  vertices having degree set  $S = \{a_1 < a_2 < \dots < a_n\}$  except the case when  $a_1 = 1$ ,  $r$  is odd and only  $a_n$  is even.

Proof (informal): Sequence  $D$  for  $G(V, E)$  realizing  $S$  is

$$D = (a_n^{a_1} > a_{n-1}^{a_2-a_1} > \dots > a_i^{a_{n-i+1}-a_{n-i}} > \dots > a_{\lceil \frac{n}{2} \rceil}^{a_1 + \lfloor \frac{n}{2} \rfloor - a_{\lfloor \frac{n}{2} \rfloor} + 1} > \dots > a_j^{a_{n-j+1}-a_{n-j}} > \dots > a_1^{a_n-a_{n-1}})$$

# Extension Problem for Undirected Graphs

## Theorem

For every  $r \in \mathbb{Z}^+$ , there exists an undirected simple graph with  $\mu(S) + r$  vertices having degree set  $S = \{a_1 < a_2 < \dots < a_n\}$  except the case when  $a_1 = 1$ ,  $r$  is odd and only  $a_n$  is even.

Proof (informal): Sequence  $D$  for  $G(V, E)$  realizing  $S$  is

$$D = (a_n^{a_1} > a_{n-1}^{a_2 - a_1} > \dots > a_i^{a_{n-i+1} - a_{n-i}} > \dots > a_{\lceil \frac{n}{2} \rceil}^{a_1 + \lfloor \frac{n}{2} \rfloor - a_{\lfloor \frac{n}{2} \rfloor} + 1} > \dots > a_j^{a_{n-j+1} - a_{n-j}} > \dots > a_1^{a_n - a_{n-1}})$$

$n$  is even:  $D = (a_n \geq \dots \geq a_{\lceil \frac{n}{2} \rceil + 1} \geq \dots \geq a_{\lfloor \frac{n}{2} \rfloor + 1} \geq a_{\lfloor \frac{n}{2} \rfloor} \geq \dots \geq a_1)$

Here  $m = a_{\lceil \frac{n}{2} \rceil} = a_{\lfloor \frac{n}{2} \rfloor}$

$$\begin{array}{ccc} & \uparrow & \uparrow \\ & d_{a_{\lfloor \frac{n}{2} \rfloor}} & d_{a_{\lfloor \frac{n}{2} \rfloor} + 1} \end{array}$$

# Extension Problem for Undirected Graphs

## Theorem

For every  $r \in \mathbb{Z}^+$ , there exists an undirected simple graph with  $\mu(S) + r$  vertices having degree set  $S = \{a_1 < a_2 < \dots < a_n\}$  except the case when  $a_1 = 1$ ,  $r$  is odd and only  $a_n$  is even.

Proof (informal): Sequence  $D$  for  $G(V, E)$  realizing  $S$  is

$$D = (a_n^{a_1} > a_{n-1}^{a_2 - a_1} > \dots > a_i^{a_{n-i+1} - a_{n-i}} > \dots > a_{\lceil \frac{n}{2} \rceil}^{a_1 + \lfloor \frac{n}{2} \rfloor - a_{\lfloor \frac{n}{2} \rfloor} + 1} > \dots > a_j^{a_{n-j+1} - a_{n-j}} > \dots > a_1^{a_n - a_{n-1}})$$

$n$  is odd:  $D = (a_n \geq \dots \geq a_{\lceil \frac{n}{2} \rceil} \geq \dots \geq a_{\lceil \frac{n}{2} \rceil} \geq a_{\lceil \frac{n}{2} \rceil} \geq \dots \geq a_1)$

Here  $m = a_{\lceil \frac{n}{2} \rceil} = a_{\lfloor \frac{n}{2} \rfloor + 1}$

$$d_{a_{\lfloor \frac{n}{2} \rfloor + 1}} \quad d_{a_{\lfloor \frac{n}{2} \rfloor + 1} + 1}$$

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1')$

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a'_1)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1^r)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.
- Both  $r$  AND  $a_1$  is odd : Let  $a_k = \min\{a_i \in S | a_i = 2l\}$



# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1^r)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.
- Both  $r$  AND  $a_1$  is odd : Let  $a_k = \min\{a_i \in S | a_i = 2l\}$ 
  - $1 \leq k \leq \lceil \frac{n}{2} \rceil$  :  $D' = (D, a_k, a_1^{r-1})$

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1^r)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.
- Both  $r$  AND  $a_1$  is odd : Let  $a_k = \min\{a_i \in S | a_i = 2l\}$ 
  - $1 \leq k \leq \lceil \frac{n}{2} \rceil$  :  $D' = (D, a_k, a_1^{r-1})$ 
    - $d_j(\in D) = a_k \Rightarrow j \geq 1 + a_{\lfloor \frac{n}{2} \rfloor + 1} \Rightarrow j > m$ . Hence  $D'$  is graphic.

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1^r)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.
- Both  $r$  AND  $a_1$  is odd : Let  $a_k = \min\{a_i \in S | a_i = 2l\}$ 
  - $1 \leq k \leq \lceil \frac{n}{2} \rceil$  :  $D' = (D, a_k, a_1^{r-1})$ 
    - $d_j(\in D) = a_k \Rightarrow j \geq 1 + a_{\lfloor \frac{n}{2} \rfloor + 1} \Rightarrow j > m$ . Hence  $D'$  is graphic.
  - $k > \lceil \frac{n}{2} \rceil$  : do not add  $a_k$ , instead add two  $a_{\lceil \frac{n}{2} \rceil}$  and then delete  $a_k$ (first occurence) from  $D$  to get  $D'$ .

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1^r)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.
- Both  $r$  AND  $a_1$  is odd : Let  $a_k = \min\{a_i \in S | a_i = 2l\}$ 
  - $1 \leq k \leq \lceil \frac{n}{2} \rceil$  :  $D' = (D, a_k, a_1^{r-1})$ 
    - $d_j(\in D) = a_k \Rightarrow j \geq 1 + a_{\lfloor \frac{n}{2} \rfloor + 1} \Rightarrow j > m$ . Hence  $D'$  is graphic.
  - $k > \lceil \frac{n}{2} \rceil$  : do not add  $a_k$ , instead add two  $a_{\lceil \frac{n}{2} \rceil}$  and then delete  $a_k$ (first occurence) from  $D$  to get  $D'$ .
    - $D'$  is graphic if : for  $a_{n-k} + 1 \leq t \leq m(= a_{\lceil \frac{n}{2} \rceil})$ ,

$$\min\{d_{t+1}, t\} \leq 2\min\{a_{\lceil \frac{n}{2} \rceil}, t\} + a_k - d_{t+1}$$

# Extension Problem proof continued...

Consider the following exhaustive cases(except one)

- At least one of  $r$  OR  $a_1$  is even :  $D' = (D, a_1^r)$ 
  - $d_j(\in D) = a_1 \Rightarrow j > m = a_{\lceil \frac{n}{2} \rceil}$ . Hence  $D'$  is graphic.
- Both  $r$  AND  $a_1$  is odd : Let  $a_k = \min\{a_i \in S | a_i = 2l\}$ 
  - $1 \leq k \leq \lceil \frac{n}{2} \rceil$  :  $D' = (D, a_k, a_1^{r-1})$ 
    - $d_j(\in D) = a_k \Rightarrow j \geq 1 + a_{\lfloor \frac{n}{2} \rfloor + 1} \Rightarrow j > m$ . Hence  $D'$  is graphic.
  - $k > \lceil \frac{n}{2} \rceil$  : do not add  $a_k$ , instead add two  $a_{\lceil \frac{n}{2} \rceil}$  and then delete  $a_k$ (first occurence) from  $D$  to get  $D'$ .
    - $D'$  is graphic if : for  $a_{n-k} + 1 \leq t \leq m (= a_{\lceil \frac{n}{2} \rceil})$ ,

$$\min\{d_{t+1}, t\} \leq 2\min\{a_{\lceil \frac{n}{2} \rceil}, t\} + a_k - d_{t+1}$$

- condition always satisfied. Hence  $D'$  is graphic.

# Tree Realizability under Multiplicity Constraints

$S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree  $\Leftrightarrow a_1 = 1$ .

# Tree Realizability under Multiplicity Constraints

$S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree  $\Leftrightarrow a_1 = 1$ .


$$\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$$

# Tree Realizability under Multiplicity Constraints

$S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree  $\Leftrightarrow a_1 = 1$ .

$$\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$$

Minimum order tree construction






# Tree Realizability under Multiplicity Constraints

$S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree  $\Leftrightarrow a_1 = 1$ .

$$\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$$

Minimum order tree construction



Let  $m_i$  be the *multiplicity* of  $a_i \in S$  then in this tree


- $m_1 \gg 1$  and  $m_i = 1, \forall 2 \leq i \leq n$ .

# Tree Realizability under Multiplicity Constraints

$S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree  $\Leftrightarrow a_1 = 1$ .

$$\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$$

Minimum order tree construction



Let  $m_i$  be the *multiplicity* of  $a_i \in S$  then in this tree


- $m_1 \gg 1$  and  $m_i = 1, \forall 2 \leq i \leq n$ .
- degree distribution is highly skewed.

# Tree Realizability under Multiplicity Constraints

$S = \{a_1 < a_2 < \dots < a_n\}, a_i \in \mathbb{Z}^+$  is realized by a tree  $\Leftrightarrow a_1 = 1$ .

$$\mu_T(S) = \sum_{i=1}^n (a_i - 1) + 2$$

Minimum order tree construction



Let  $m_i$  be the *multiplicity* of  $a_i \in S$  then in this tree

- $m_1 \gg 1$  and  $m_i = 1, \forall 2 \leq i \leq n$ .
- degree distribution is highly skewed.

Q. Can we have another tree realization with smaller no of pendant vertices?

# Tree Realizability under Multiplicity Constraints

## Lemma

*The minimum multiplicity of pendant vertices in any tree realization for the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is  $\sum_{i=1}^n a_i - 2n + 3$*

# Tree Realizability under Multiplicity Constraints

## Lemma

*The minimum multiplicity of pendant vertices in any tree realization for the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is  $\sum_{i=1}^n a_i - 2n + 3$*

Proof: Let  $m_i$  be the multiplicity of  $a_i \in S$ . Then, the corresponding degree sequence is  $S = (1^{m_1}, a_2^{m_2}, \dots, a_n^{m_n})$ .

# Tree Realizability under Multiplicity Constraints

## Lemma

*The minimum multiplicity of pendant vertices in any tree realization for the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is  $\sum_{i=1}^n a_i - 2n + 3$*

Proof: Let  $m_i$  be the multiplicity of  $a_i \in S$ . Then, the corresponding degree sequence is  $S = (1^{m_1}, a_2^{m_2}, \dots, a_n^{m_n})$ .

Since  $a_2 \geq 2$  hence using previous lemma(AM96)

# Tree Realizability under Multiplicity Constraints

## Lemma

*The minimum multiplicity of pendant vertices in any tree realization for the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$  is  $\sum_{i=1}^n a_i - 2n + 3$*

Proof: Let  $m_i$  be the multiplicity of  $a_i \in S$ . Then, the corresponding degree sequence is  $S = (1^{m_1}, a_2^{m_2}, \dots, a_n^{m_n})$ .

Since  $a_2 \geq 2$  hence using previous lemma(AM96)

$$m_1 = 2 + (a_2 - 2)m_2 + (a_3 - 2)m_3 + \dots + (a_n - 2)m_n \quad \forall i, m_i \geq 1$$

$m_1$  will be minimum if  $m_i = 1$  for each  $i = 2, 3, \dots, n$

# Tree Realizability under Multiplicity Constraints

Minimum order tree described earlier meets exactly this requirement.

So minimum value of

$$\begin{aligned} m_1 &= 2 + (a_2 - 2) + (a_3 - 2) + \cdots + (a_n - 2) \\ &= \sum_{i=1}^n a_i - 2n + 3 \end{aligned}$$



# Tree Realizability under Multiplicity Constraints

Minimum order tree described earlier meets exactly this requirement.

So minimum value of

$$\begin{aligned} m_1 &= 2 + (a_2 - 2) + (a_3 - 2) + \cdots + (a_n - 2) \\ &= \sum_{i=1}^n a_i - 2n + 3 \end{aligned}$$

This result can be generalized in case of *Degree Multiset*.

# Degree Set Variants for Digraphs

In case of directed graphs, from degree set we mean

- *indegree set* or *outdegree set* or both

# Degree Set Variants for Digraphs

In case of directed graphs, from degree set we mean

- *indegree set* or *outdegree set* or both

Depending on the variation we call the realization  $G(V, E)$  an

## $\forall(\text{OR})$ - Realization

if  $\forall v \in V, d^+(v) \in S$  or  $d^-(v) \in S$ , and for each  $a_i \in S$  there is a vertex  $u \in V$  such that  $d^+(u) = a_i$  or  $d^-(u) = a_i$ .

# Degree Set Variants for Digraphs

In case of directed graphs, from degree set we mean

- *indegree set* or *outdegree set* or both

Depending on the variation we call the realization  $G(V, E)$  an


## $\vee(\text{OR})$ - Realization

if  $\forall v \in V, d^+(v) \in S$  or  $d^-(v) \in S$ , and for each  $a_i \in S$  there is a vertex  $u \in V$  such that  $d^+(u) = a_i$  or  $d^-(u) = a_i$ .


## $\wedge(\text{AND})$ - Realization

if  $\forall v \in V, d^+(v) \in S$  and  $d^-(v) \in S$ , and for each  $a_i \in S$  there is a vertex  $v_p$  and  $v_q \in V$  such that  $d^+(v_p) = a_i$  and  $d^-(v_q) = a_i$ .


$$S = \{2, 3\} \checkmark$$




$$S = \{2, 3\} \checkmark$$




$$S = \{2, 3\} \times$$



$$S = \{2, 3\} \checkmark$$




$$S = \{2, 3\} \times$$




- V-realized
- not  $\wedge$ -realized

$$S = \{2, 3\} \checkmark$$



- V-realized
- not  $\wedge$ -realized


$$S = \{2, 3\} \times$$



- not V-realized
- not  $\wedge$ -realized




$$S = \{2, 3\} \checkmark$$




- V-realized
- not  $\wedge$ -realized

$$S = \{1, 2\}$$



- V-realized
- $\wedge$ -realized

$$S = \{2, 3\} \times$$



- not V-realized
- not  $\wedge$ -realized

# Digraph Realization for Degree Set

A finite set  $S = \{a_1 < a_2 < \dots < a_n\}$  of non-negative integers is always realizable by a directed graph with  $a_n + 1$  vertices, under both,  $\wedge$  and  $\vee$ , notions of realizability.

# Digraph Realization for Degree Set

A finite set  $S = \{a_1 < a_2 < \dots < a_n\}$  of non-negative integers is always realizable by a directed graph with  $a_n + 1$  vertices, under both,  $\wedge$  and  $\vee$ , notions of realizability.

Proof: Construct the undirected graph  $G(V, E)$  with  $a_n + 1$  vertices for  $S$ , and then replace each undirected edge by two way symmetric edges.

# Digraph Realization for Degree Set

A finite set  $S = \{a_1 < a_2 < \dots < a_n\}$  of non-negative integers is always realizable by a directed graph with  $a_n + 1$  vertices, under both,  $\wedge$  and  $\vee$ , notions of realizability.

Proof: Construct the undirected graph  $G(V, E)$  with  $a_n + 1$  vertices for  $S$ , and then replace each undirected edge by two way symmetric edges.

- digraphs with  $a_n + 1$  vertices : *symmetric*, easy to construct

# Digraph Realization for Degree Set

A finite set  $S = \{a_1 < a_2 < \dots < a_n\}$  of non-negative integers is always realizable by a directed graph with  $a_n + 1$  vertices, under both,  $\wedge$  and  $\vee$ , notions of realizability.

Proof: Construct the undirected graph  $G(V, E)$  with  $a_n + 1$  vertices for  $S$ , and then replace each undirected edge by two way symmetric edges.

- digraphs with  $a_n + 1$  vertices : *symmetric*, easy to construct
- from realizability point of view
  - $\vee$ -realization : seems to be very relaxed

# Digraph Realization for Degree Set

A finite set  $S = \{a_1 < a_2 < \dots < a_n\}$  of non-negative integers is always realizable by a directed graph with  $a_n + 1$  vertices, under both,  $\wedge$  and  $\vee$ , notions of realizability.

Proof: Construct the undirected graph  $G(V, E)$  with  $a_n + 1$  vertices for  $S$ , and then replace each undirected edge by two way symmetric edges.

- digraphs with  $a_n + 1$  vertices : *symmetric*, easy to construct
- from realizability point of view
  - $\vee$ -realization : seems to be very relaxed

*More feasible to study an asymmetric realization under  $\wedge$ -realizability constraints.*

# Asymmetric Digraph $\wedge$ -realization

If  $\mu_A(S)$  denotes the minimum order of any asymmetric directed graph  $\wedge$ -realizing  $S = \{a_1 < a_2 < \dots < a_n\}$ ,  $n \geq 2$ ,  $a_i \in \mathbb{Z}^+$  then

$$a_1 + a_n + 1 \leq \mu_A(S) \leq a_{n-1} + a_n + 1$$

There exists some sufficient conditions which achieves the minimum order,  $(a_1 + a_n + 1)$ - vertices realization.

# Asymmetric Digraph $\wedge$ -realization

If  $\mu_A(S)$  denotes the minimum order of any asymmetric directed graph  $\wedge$ -realizing  $S = \{a_1 < a_2 < \dots < a_n\}$ ,  $n \geq 2$ ,  $a_i \in \mathbb{Z}^+$  then

$$a_1 + a_n + 1 \leq \mu_A(S) \leq a_{n-1} + a_n + 1$$

There exists some sufficient conditions which achieves the minimum order,  $(a_1 + a_n + 1)$ - vertices realization.

*Proof:* lower bound- there is least one vertex  $v$  of  $G$  such that  $d^+(v) + d^-(v) \geq a_n + a_1$  and since  $G$  is asymmetric, hence  $\mu_A(S) \geq a_1 + a_n + 1$ .



# Asymmetric Digraph $\wedge$ -realization

If  $\mu_A(S)$  denotes the minimum order of any asymmetric directed graph  $\wedge$ -realizing  $S = \{a_1 < a_2 < \dots < a_n\}$ ,  $n \geq 2$ ,  $a_i \in \mathbb{Z}^+$  then

$$a_1 + a_n + 1 \leq \mu_A(S) \leq a_{n-1} + a_n + 1$$

There exists some sufficient conditions which achieves the minimum order,  $(a_1 + a_n + 1)$ - vertices realization.

*Proof:* lower bound- there is least one vertex  $v$  of  $G$  such that  $d^+(v) + d^-(v) \geq a_n + a_1$  and since  $G$  is asymmetric, hence  $\mu_A(S) \geq a_1 + a_n + 1$ .

upper bound- By giving an inductive construction for  $G$ .

# Asymmetric Digraph $\wedge$ -realization

If  $\mu_A(S)$  denotes the minimum order of any asymmetric directed graph  $\wedge$ -realizing  $S = \{a_1 < a_2 < \dots < a_n\}$ ,  $n \geq 2$ ,  $a_i \in \mathbb{Z}^+$  then

$$a_1 + a_n + 1 \leq \mu_A(S) \leq a_{n-1} + a_n + 1$$

There exists some sufficient conditions which achieves the minimum order,  $(a_1 + a_n + 1)$ - vertices realization.

*Proof:* lower bound- there is least one vertex  $v$  of  $G$  such that  $d^+(v) + d^-(v) \geq a_n + a_1$  and since  $G$  is asymmetric, hence  $\mu_A(S) \geq a_1 + a_n + 1$ .

upper bound- By giving an inductive construction for  $G$ .

Extension : Always possible if  $r \geq a_n + a_{n-1} + 1$

# Minimum Order $\vee$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\vee$ -realizes the degree set  $S$ , is  $\sum_{i=1}^n (a_i - 1) + 2$ .*

# Minimum Order $\vee$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\vee$ -realizes the degree set  $S$ , is  $\sum_{i=1}^n (a_i - 1) + 2$ .*

*Proof: upper bound-  $\mu_{\vee}(S) \leq \mu(S) = \sum_{i=1}^n (a_i - 1) + 2$*

# Minimum Order $\vee$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\vee$ -realizes the degree set  $S$ , is  $\sum_{i=1}^n (a_i - 1) + 2$ .*

*Proof: upper bound-  $\mu_{\vee}(S) \leq \mu(S) = \sum_{i=1}^n (a_i - 1) + 2$*

Get the bipartite graph corresponding to minimum order undirected tree for  $S$ , assign same directions to all edges going across the partition.

# Minimum Order $\vee$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\vee$ -realizes the degree set  $S$ , is  $\sum_{i=1}^n (a_i - 1) + 2$ .*

*Proof: upper bound-  $\mu_{\vee}(S) \leq \mu(S) = \sum_{i=1}^n (a_i - 1) + 2$*

Get the bipartite graph corresponding to minimum order undirected tree for  $S$ , assign same directions to all edges going across the partition.

lower bound-  $\mu_{\vee}(S) \geq \sum_{i=1}^n (a_i - 1) + 2$

# Minimum Order $\vee$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{1 = a_1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\vee$ -realizes the degree set  $S$ , is  $\sum_{i=1}^n (a_i - 1) + 2$ .*

*Proof: upper bound-  $\mu_{\vee}(S) \leq \mu(S) = \sum_{i=1}^n (a_i - 1) + 2$*

Get the bipartite graph corresponding to minimum order undirected tree for  $S$ , assign same directions to all edges going across the partition.

lower bound-  $\mu_{\vee}(S) \geq \sum_{i=1}^n (a_i - 1) + 2$

For each  $i$ ,  $a_i \in S$  will appear as  $(a_i, b_j)$  or  $(b_k, a_i)$  at least once, where  $b_j, b_k \in \mathbb{Z}^+ \cup \{0\}$ . Thus

$$\sum_{v \in V} (d^-(v) + d^+(v)) = 2|E| = 2(V - 1) \geq \sum_{i=1}^n a_i + (V - n)$$

This implies the lower bound  $|V| \geq 2 + \sum_{i=1}^n (a_i - 1)$ .

# Minimum Order $\wedge$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{0 < 1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\wedge$ -realizes the degree set  $s$ , is  $2(\sum_{i=1}^n (a_i - 1)) + 2$ .*



# Minimum Order $\wedge$ - Realizability of Directed Trees

## Theorem

*For the degree set  $S = \{0 < 1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\wedge$ -realizes the degree set  $s$ , is  $2(\sum_{i=1}^n (a_i - 1)) + 2$ .*


*Proof:* Each  $a_i \in S$  will appear at least twice so we get minimum order tree for multiset  $\{1, a_2^2, a_3^2, \dots, a_n^2\}$  with all edges being assigned directions in alternate way.

# Minimum Order $\wedge$ - Realizability of Directed Trees

## Theorem

For the degree set  $S = \{0 < 1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\wedge$ -realizes the degree set  $s$ , is  $2(\sum_{i=1}^n (a_i - 1)) + 2$ .

Proof: Each  $a_i \in S$  will appear at least twice so we get minimum order tree for multiset  $\{1, a_2^2, a_3^2, \dots, a_n^2\}$  with all edges being assigned directions in alternate way.




# Minimum Order $\wedge$ - Realizability of Directed Trees

## Theorem

For the degree set  $S = \{0 < 1 < a_2 < \dots < a_n\}$ , the minimum order of a directed tree which  $\wedge$ -realizes the degree set  $s$ , is  $2(\sum_{i=1}^n (a_i - 1)) + 2$ .

Proof: Each  $a_i \in S$  will appear at least twice so we get minimum order tree for multiset  $\{1, a_2^2, a_3^2, \dots, a_n^2\}$  with all edges being assigned directions in alternate way.



# Tree Extension Problem for Directed Trees

- for undirected trees, TEP is NP-Complete.

# Tree Extension Problem for Directed Trees

- for undirected trees, TEP is NP-Complete.
- for directed trees, the task becomes easier, i.e.


For every non-negative integer  $r$ , there are directed trees with  $\mu_T(S) + r$  vertices  $\wedge$ -realizing and  $\vee$ -realizing the given degree set  $S$ .

# Tree Extension Problem for Directed Trees

- for undirected trees, TEP is NP-Complete.
- for directed trees, the task becomes easier, i.e.


For every non-negative integer  $r$ , there are directed trees with  $\mu_T(S) + r$  vertices  $\wedge$ -realizing and  $\vee$ -realizing the given degree set  $S$ .

Minimum order Directed tree  $\wedge$ -realizing ( $\vee$ -realizing) the degree set  $S = \{0 < 1 < a_2 < \dots < a_n\}$




# Directed Tree Extension

Every vertex to be added is added to the pendant vertex by an incoming or outgoing edge depending on whether the pendant vertex is a sink or source vertex respectively.



# Directed Tree Extension


Every vertex to be added is added to the pendant vertex by an incoming or outgoing edge depending on whether the pendant vertex is a sink or source vertex respectively.





# Directed Tree Extension

Every vertex to be added is added to the pendant vertex by an incoming or outgoing edge depending on whether the pendant vertex is a sink or source vertex respectively.



# Conclusion and Future Work

- Conclusion

# Conclusion and Future Work

- Conclusion
  - Extension problem for General Graphs (undirected) and Trees

# Conclusion and Future Work

- Conclusion
  - Extension problem for General Graphs (undirected) and Trees
  - Complexity aspects of Tree Extension Problem

# Conclusion and Future Work

- Conclusion
  - Extension problem for General Graphs (undirected) and Trees
  - Complexity aspects of Tree Extension Problem
  - Multiplicity lower bound for pendant vertices in Tree Realization

# Conclusion and Future Work

- Conclusion

- Extension problem for General Graphs (undirected) and Trees
- Complexity aspects of Tree Extension Problem
- Multiplicity lower bound for pendant vertices in Tree Realization
- Asymmetric Digraph  $\wedge$ -Realization

- Conclusion

- Extension problem for General Graphs (undirected) and Trees
- Complexity aspects of Tree Extension Problem
- Multiplicity lower bound for pendant vertices in Tree Realization
- Asymmetric Digraph  $\wedge$ -Realization
- Lower bound on order for Directed Trees under  $\wedge(\vee)$ -Realizability

- Conclusion

- Extension problem for General Graphs (undirected) and Trees
- Complexity aspects of Tree Extension Problem
- Multiplicity lower bound for pendant vertices in Tree Realization
- Asymmetric Digraph  $\wedge$ -Realization
- Lower bound on order for Directed Trees under  $\wedge(\vee)$ -Realizability
- Extension problem for Asymmetric Graphs and Directed Trees



- Conclusion
  - Extension problem for General Graphs (undirected) and Trees
  - Complexity aspects of Tree Extension Problem
  - Multiplicity lower bound for pendant vertices in Tree Realization
  - Asymmetric Digraph  $\wedge$ -Realization
  - Lower bound on order for Directed Trees under  $\wedge(\vee)$ -Realizability
  - Extension problem for Asymmetric Graphs and Directed Trees
- Future Work

# Conclusion and Future Work

- Conclusion

- Extension problem for General Graphs (undirected) and Trees
- Complexity aspects of Tree Extension Problem
- Multiplicity lower bound for pendant vertices in Tree Realization
- Asymmetric Digraph  $\wedge$ -Realization
- Lower bound on order for Directed Trees under  $\wedge(\vee)$ -Realizability
- Extension problem for Asymmetric Graphs and Directed Trees

- Future Work

- Bridging the gap between bounds for Asymmetric Graphs

# Conclusion and Future Work

- Conclusion

- Extension problem for General Graphs (undirected) and Trees
- Complexity aspects of Tree Extension Problem
- Multiplicity lower bound for pendant vertices in Tree Realization
- Asymmetric Digraph  $\wedge$ -Realization
- Lower bound on order for Directed Trees under  $\wedge(\vee)$ -Realizability
- Extension problem for Asymmetric Graphs and Directed Trees

- Future Work

- Bridging the gap between bounds for Asymmetric Graphs
- Minimum order graph realization for degree multiset

Questions..??

Thank You.

# Asymmetric Digraph $\wedge$ -realization

For  $D = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)

- In every realization  $G_1(V_1, E_1)$  of  $D$ ,  $\forall v \in V_1$   $d^-(v) + d^+(v) = 2a_1$  and since  $G$  is asymmetric, hence  $\mu_A(S) \geq 2a_1 + 1$ .

# Asymmetric Digraph $\wedge$ -realization

For  $D = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)

- In every realization  $G_1(V_1, E_1)$  of  $D$ ,  $\forall v \in V_1$   $d^-(v) + d^+(v) = 2a_1$  and since  $G$  is asymmetric, hence  $\mu_A(S) \geq 2a_1 + 1$ .
- To prove the upper bound we construct an asymmetric graph  $G_1$  with  $2a_1 + 1$  vertices.

# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)


$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a + 1 \text{ and } i + 1 \leq j \leq i + a\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)


$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a_1 + 1 \text{ and } i + 1 \leq j \leq i + a_1\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)


$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a_1 + 1 \text{ and } i + 1 \leq j \leq i + a_1\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)


$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a_1 + 1 \text{ and } i + 1 \leq j \leq i + a_1\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)


$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a_1 + 1 \text{ and } i + 1 \leq j \leq i + a_1\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)


$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a_1 + 1 \text{ and } i + 1 \leq j \leq i + a_1\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

For  $S = \{a_1\}$ ,  $\mu_A(S) = 2a_1 + 1$  (Chartrand *et al*, 1976)

$E_1 = \{(v_i, v_j) | 1 \leq i \leq 2a_1 + 1 \text{ and } i + 1 \leq j \leq i + a_1\}$ , where subscripts are modulo  $2a_1 + 1$



# Asymmetric Digraph $\wedge$ -realization

Divide the vertices of  $G_1$  in 3 components -  $C_x, C_y, C_z$

# Asymmetric Digraph $\wedge$ -realization


Divide the vertices of  $G_1$  in 3 components -  $C_x, C_y, C_z$

$C_x$




$$d^-(v) = d^+(v) = a_1$$

$C_y$



$$d^-(v) = d^+(v) = a_1$$

$C_z$



$$d^-(v) = d^+(v) = a_1$$




# Asymmetric Digraph $\wedge$ -realization

Divide the vertices of  $G_1$  in 3 components -  $C_x, C_y, C_z$


$C_x$



$C_y$



$C_z$



$$d^-(v) = d^+(v) = a_1$$

$$d^-(v) = d^+(v) = a_1$$

$$d^-(v) = d^+(v) = a_1$$

Now add one component  $C_1$  containing  $(a_2 - a_1)$  isolated vertices and the edge set

$$E = \{(v_x, v_1) | v_x \in C_x \wedge v_1 \in C_1\} \cup \{(v_1, v_y) | v_1 \in C_1 \wedge v_y \in C_y\}$$

to  $G_1$  to get  $G_2$ .

# Asymmetric Digraph $\wedge$ -realization

Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below

$C_x$

$$1 \ 2 \cdots a_1$$

$$d^-(v) = d^+(v) = a_1$$

$C_y$

$$(a_1 + 1) (a_1 + 2) \cdots 2a_1$$

$$d^-(v) = d^+(v) = a_1$$

$C_z$

$$2a_1 + 1$$

$$d^-(v) = d^+(v) = a_1$$

# Asymmetric Digraph $\wedge$ -realization

Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below

$C_x$

$$1 \ 2 \cdots a_1$$

$$d^-(v) = d^+(v) = a_1$$

$C_y$

$$(a_1 + 1) (a_1 + 2) \cdots 2a_1$$

$$d^-(v) = d^+(v) = a_1$$

$C_z$

$$2a_1 + 1$$

$$d^-(v) = d^+(v) = a_1$$

$$1 \ 2 \cdots (a_2 - a_1)$$

$$d^-(v) = d^+(v) = 0$$

# Asymmetric Digraph $\wedge$ -realization


Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below

$C_x$




$$d^-(v) = a_1, d^+(v) = a_2$$

$C_y$




$$d^-(v) = d^+(v) = a_1$$

$C_z$



$$d^-(v) = d^+(v) =$$

$$a_1$$




$$d^-(v) = a_1, d^+(v) = 0$$

Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below

$C_x$

$C_y$

$C_z$




Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below


$C_x$

$C_y$

$C_z$




Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below



$|V| = a_2 + a_1 + 1$ , no of components = 4

Hypothesis : Consider there exists an asymmetric directed graph  $G_{n_0}$  with degree set  $\{a_1 < a_2 < \dots < a_{n_0}\}$ ,  $n_0 < n$  with  $a_{n_0} + a_{n_0-1} + 1$  vertices and  $2n_0$  components.

Base Case(for  $|S| = 2$ ):  $G_2$  for  $\{a_1 < a_2\}$  is constructed from  $G_1$  as below



$|V| = a_2 + a_1 + 1$ , no of components = 4

Hypothesis : Consider there exists an asymmetric directed graph  $G_{n_0}$  with degree set  $\{a_1 < a_2 < \dots < a_{n_0}\}$ ,  $n_0 < n$  with  $a_{n_0} + a_{n_0-1} + 1$  vertices and  $2n_0$  components.

Need to construct  $G_{n_0+1}$  from  $G_{n_0}$ .



# Asymmetric Digraph $\wedge$ -realization

$G_{n_0}$  with degree set  $\{a_1 < a_2 < \dots < a_{n_0}\}$ ,  $n_0 < n$  and  $a_{n_0} + a_{n_0-1} + 1$  vertices has the following  $2n_0$  components :

- $C_{n_0-1}$  :  $|V| = a_{n_0} - a_{n_0-1}$ ,  $d^-(v) = d^+(v) = a_1$
- $C_i (\forall 1 \leq i \leq n_0 - 2)$  :  $|V_i| = a_{i+1} - a_i$ ,  $d^-(v) = a_{n_0-1-i}$ ,  $d^+(v) = a_1$
- $C'_i (\forall 1 \leq i' \leq n_0 - 2)$  :  $|V_i| = a_{i+1} - a_i$ ,  $d^-(v) = a_1$ ,  $d^+(v) = a_{n_0-1-i}$

# Asymmetric Digraph $\wedge$ -realization

$G_{n_0}$  with degree set  $\{a_1 < a_2 < \dots < a_{n_0}\}$ ,  $n_0 < n$  and  $a_{n_0} + a_{n_0-1} + 1$  vertices has the following  $2n_0$  components :

- $C_{n_0-1} : |V| = a_{n_0} - a_{n_0-1}, d^-(v) = d^+(v) = a_1$
- $C_i (\forall 1 \leq i \leq n_0 - 2) : |V_i| = a_{i+1} - a_i, d^-(v) = a_{n_0-1-i}, d^+(v) = a_1$
- $C'_i (\forall 1 \leq i' \leq n_0 - 2) : |V_i| = a_{i+1} - a_i, d^-(v) = a_1, d^+(v) = a_{n_0-1-i}$
- Components from base graph  $G_1$ 
  - $C_x : |V| = a_1, d^-(v) = a_{n_0-1}, d^+(v) = a_{n_0}$
  - $C_y : |V| = a_1, d^-(v) = a_{n_0}, d^+(v) = a_{n_0-1}$
  - $C_z : |V| = 1, d^-(v) = d^+(v) = a_1$

$$C_{n_o-1}$$

$$|V| = a_{n_0} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = a_1$$

$$C_1 \dots C_i \dots C_{n_o-2}$$

$$|V| = a_{i+1} - a_i, \forall 1 \leq i \leq n_o - 2$$

$$d^-(v) = a_{n_o-1-i}, d^+(v) = a_1$$

$$C'_1 \dots C'_i \dots C'_{n_o-2}$$

$$|V| = a_{i+1} - a_i, \forall 1 \leq i' \leq n_o - 2$$

$$d^-(v) = a_1, d^+(v) = a_{n_o-1-i}$$

$C_{n_o-1}$ 

$$|V| = a_{n_0} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = a_1$$

 $C_1 \dots C_i \dots C_{n_o-2}$ 

$$|V| = a_{i+1} - a_i, \forall 1 \leq i \leq n_o - 2$$

$$d^-(v) = a_{n_o-1-i}, d^+(v) = a_1$$

 $C_z$ 

$$|V| = 1$$

$$d^-(v) = d^+(v) = a_1$$

 $C'_1 \dots C'_i \dots C'_{n_o-2}$ 

$$|V| = a_{i+1} - a_i, \forall 1 \leq i' \leq n_o - 2$$

$$d^-(v) = a_1, d^+(v) = a_{n_o-1-i}$$

 $C_x$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o-1}, d^+(v) = a_{n_o}$$

 $C_y$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o}, d^+(v) = a_{n_o-1}$$

# Asymmetric Digraph $\wedge$ -realization

To obtain  $G_{n_0+1}$  from  $G_{n_0}$ , we add two new components -  
 $C_{n_0}(|V| = a_{n_0+1} - a_{n_0}), C'_{n_0-1}(|V| = a_{n_0} - a_{n_0-1})$

$C_{n_o-1}$ 

$$|V| = a_{n_o} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = a_1$$

 $C_1 \dots C_i \dots C_{n_o-2}$ 

$$|V| = a_{i+1} - a_i, \forall 1 \leq i \leq n_o - 2$$

$$d^-(v) = a_{n_o-1-i}, d^+(v) = a_1$$

 $C_z$ 

$$|V| = 1$$

$$d^-(v) = d^+(v) = a_1$$

 $C'_1 \dots C'_i \dots C'_{n_o-2}$ 

$$|V| = a_{i+1} - a_i, \forall 1 \leq i' \leq n_o - 2$$

$$d^-(v) = a_1, d^+(v) = a_{n_o-1-i}$$

 $C_x$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o-1}, d^+(v) = a_{n_o}$$

 $C_y$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o}, d^+(v) = a_{n_o-1}$$

 $C_{n_o}$ 

$$|V| = a_{n_o+1} - a_{n_o}$$

$$d^-(v) = d^+(v) = 0$$

 $C'_{n_o-1}$ 

$$|V| = a_{n_o} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = 0$$

# Asymmetric Digraph $\wedge$ -realization

To obtain  $G_{n_0+1}$  from  $G_{n_0}$ , we add two new components -

$C_{n_0}(|V| = a_{n_0+1} - a_{n_0})$ ,  $C'_{n_0-1}(|V| = a_{n_0} - a_{n_0-1})$  and the edge set

$E = E_1 \cup E_2 \cup E_3$ , where

$$\bullet E_1 = \{(v_x, v_{n_0}) \mid v_x \in C_x \wedge v_{n_0} \in C_{n_0}\} \cup \{(v_{n_0}, v_y) \mid v_{n_0} \in C_{n_0} \wedge v_y \in C_y\}$$

$C_{n_o-1}$ 

$$|V| = a_{n_o} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = a_1$$

 $C_i$ 

$$|V| = a_{i+1} - a_i$$

$$d^-(v) = a_{n_o-1-i}, d^+(v) = a_1$$

 $C_z$ 

$$|V| = 1$$

$$d^-(v) = d^+(v) = a_1$$

 $C'_{n_o-1-i}$ 

$$|V| = a_{n_o-i} - a_{n_o-1-i}$$

$$d^-(v) = a_1, d^+(v) = a_i$$

 $C_x$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o-1}, d^+(v) = a_{n_o+1}$$

 $C_y$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o+1}, d^+(v) = a_{n_o-1}$$

 $C_{n_o}$ 

$$|V| = a_{n_o+1} - a_{n_o}$$

$$d^-(v) = d^+(v) = a_1$$

 $C'_{n_o-1}$ 

$$|V| = a_{n_o} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = 0$$



# Asymmetric Digraph $\wedge$ -realization

To obtain  $G_{n_0+1}$  from  $G_{n_0}$ , we add two new components -

$C_{n_0}(|V| = a_{n_0+1} - a_{n_0})$ ,  $C'_{n_0-1}(|V| = a_{n_0} - a_{n_0-1})$  and the edge set  $E = E_1 \cup E_2 \cup E_3$ , where

- $E_1 = \{(v_x, v_{n_0}) | v_x \in C_x \wedge v_{n_0} \in C_{n_0}\} \cup \{(v_{n_0}, v_y) | v_{n_0} \in C_{n_0} \wedge v_y \in C_y\}$
- $E_2 = \{(v_y, v_{n_0-1}) | v_y \in C_y \wedge v_{n_0-1} \in C'_{n_0-1}\} \cup \{(v_{n_0-1}, v_x) | v_{n_0-1} \in C'_{n_0-1} \wedge v_x \in C_x\}$

$C_{n_o-1}$ 

$$|V| = a_{n_o} - a_{n_o-1}$$

$$d^-(v) = d^+(v) = a_1$$

 $C_i$ 

$$|V| = a_{i+1} - a_i$$

$$d^-(v) = a_{n_o-1-i}, d^+(v) = a_1$$

 $C_z$ 

$$|V| = 1$$

$$d^-(v) = d^+(v) = a_1$$

 $C'_{n_o-1-i}$ 

$$|V| = a_{n_o-i} - a_{n_o-1-i}$$

$$d^-(v) = a_1, d^+(v) = a_i$$

 $C_x$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o}, d^+(v) = a_{n_o+1}$$

 $C_y$ 

$$|V| = a_1$$

$$d^-(v) = a_{n_o+1}, d^+(v) = a_{n_o}$$

 $C_{n_o}$ 

$$|V| = a_{n_o+1} - a_{n_o}$$

$$d^-(v) = d^+(v) = a_1$$

 $C'_{n_o-1}$ 

$$|V| = a_{n_o} - a_{n_o-1}$$


$$d^-(v) = d^+(v) = a_1$$

# Asymmetric Digraph $\wedge$ -realization

To obtain  $G_{n_0+1}$  from  $G_{n_0}$ , we add two new components -

$C_{n_0}(|V| = a_{n_0+1} - a_{n_0})$ ,  $C'_{n_0-1}(|V| = a_{n_0} - a_{n_0-1})$  and the edge set  $E = E_1 \cup E_2 \cup E_3$ , where

- $E_1 = \{(v_x, v_{n_0}) | v_x \in C_x \wedge v_{n_0} \in C_{n_0}\} \cup \{(v_{n_0}, v_y) | v_{n_0} \in C_{n_0} \wedge v_y \in C_y\}$
- $E_2 = \{(v_y, v_{n_0-1}) | v_y \in C_y \wedge v_{n_0-1} \in C'_{n_0-1}\} \cup \{(v_{n_0-1}, v_x) | v_{n_0-1} \in C'_{n_0-1} \wedge v_x \in C_x\}$
- $E_3 = \{(v_i, v'_i) | v_i \in C_i \wedge v'_i \in C'_{n_0-1-i}\}$ , where  $i \in \{1, 2, \dots, n_0 - 2\}$



# Asymmetric Digraph $\wedge$ -realization

To obtain  $G_{n_0+1}$  from  $G_{n_0}$ , we add two new components -

$C_{n_0}(|V| = a_{n_0+1} - a_{n_0})$ ,  $C'_{n_0-1}(|V| = a_{n_0} - a_{n_0-1})$  and the edge set  $E = E_1 \cup E_2 \cup E_3$ , where

- $E_1 = \{(v_x, v_{n_0}) | v_x \in C_x \wedge v_{n_0} \in C_{n_0}\} \cup \{(v_{n_0}, v_y) | v_{n_0} \in C_{n_0} \wedge v_y \in C_y\}$
- $E_2 = \{(v_y, v_{n_0-1}) | v_y \in C_y \wedge v_{n_0-1} \in C'_{n_0-1}\} \cup \{(v_{n_0-1}, v_x) | v_{n_0-1} \in C'_{n_0-1} \wedge v_x \in C_x\}$
- $E_3 = \{(v_i, v'_i) | v_i \in C_i \wedge v'_i \in C'_{n_0-1-i}\}$ , where  $i \in \{1, 2, \dots, n_0 - 2\}$

$G_{n_0+1}$  resembles  $G_{n_0}$  if  $n_0$  is replaced with  $n_0 + 1$ .

Hence,  $\mu_A(S) \leq a_n + a_{n-1} + 1$

# Asymmetric Digraph $\wedge$ -realization

To obtain  $G_{n_0+1}$  from  $G_{n_0}$ , we add two new components -

$C_{n_0}(|V| = a_{n_0+1} - a_{n_0})$ ,  $C'_{n_0-1}(|V| = a_{n_0} - a_{n_0-1})$  and the edge set  $E = E_1 \cup E_2 \cup E_3$ , where

- $E_1 = \{(v_x, v_{n_0}) | v_x \in C_x \wedge v_{n_0} \in C_{n_0}\} \cup \{(v_{n_0}, v_y) | v_{n_0} \in C_{n_0} \wedge v_y \in C_y\}$
- $E_2 = \{(v_y, v_{n_0-1}) | v_y \in C_y \wedge v_{n_0-1} \in C'_{n_0-1}\} \cup \{(v_{n_0-1}, v_x) | v_{n_0-1} \in C'_{n_0-1} \wedge v_x \in C_x\}$
- $E_3 = \{(v_i, v'_i) | v_i \in C_i \wedge v'_i \in C'_{n_0-1-i}\}$ , where  $i \in \{1, 2, \dots, n_0 - 2\}$

$G_{n_0+1}$  resembles  $G_{n_0}$  if  $n_0$  is replaced with  $n_0 + 1$ .

$$\text{Hence, } \mu_A(S) \leq a_n + a_{n-1} + 1$$

Extension : Always possible if  $r \geq a_n + a_{n-1} + 1$ , by constructing  $G_1$  for  $\{a_1\}$  with  $(2a_1 + 1) + r - (a_n + a_{n-1} + 1)$  vertices and then construct the graph using same inductive approach.