Data Set Information:

In [Cortez and Morais, 2007], the output 'area' was first transformed with a ln(x+1) function. Then, several Data Mining methods were applied. After fitting the models, the outputs were post-processed with the inverse of the ln(x+1) transform. Four different input setups were used. The experiments were conducted using a 10-fold (cross-validation) x 30 runs. Two regression metrics were measured: MAD and RMSE. A Gaussian support vector machine (SVM) fed with only 4 direct weather conditions (temp, RH, wind and rain) obtained the best MAD value: 12.71 +- 0.01 (mean and confidence interval within 95% using a t-student distribution). The best RMSE was attained by the naive mean predictor. An analysis to the regression error curve (REC) shows that the SVM model predicts more examples within a lower dmitted error. In effect, the SVM model predicts better small fires, which are the majority.

Attribute Information:

```
1. X - x-axis spatial coordinate within the Montesinho park map: 1 to 9
```

2. Y - y-axis spatial coordinate within the Montesinho park map: 2 to 9

3. month - month of the year: 'jan' to 'dec'

Note: We renamed months attribute starting with "1" in order in this dataset.

(For Example: January=1, February=2... vs.)

4. day - day of the week: 'mon' to 'sun'

Note: We renamed days attribute starting with "1" in order in this dataset.

(For Example: Monday=1, Tuesday=2... vs.)

5. FFMC - FFMC index from the FWI system: 18.7 to 96.20

6. DMC - DMC index from the FWI system: 1.1 to 291.3

7. DC - DC index from the FWI system: 7.9 to 860.6

8. ISI - ISI index from the FWI system: 0.0 to 56.10

9. temp - temperature in Celsius degrees: 2.2 to 33.30

10. RH - relative humidity in %: 15.0 to 100

11. wind - wind speed in km/h: 0.40 to 9.40

12. rain - outside rain in mm/m2 : 0.0 to 6.4

13. area - the burned area of the forest (in ha): 0.00 to 1090.84

(this output variable is very skewed towards 0.0, thus it may make

sense to model with the logarithm transform).

Relevant Papers:

P. Cortez and A. Morais. A Data Mining Approach to Predict Forest Fires using Meteorological Data. In J. Neves, M. F. Santos and J. Machado Eds., New Trends in Artificial Intelligence, Proceedings of the 13th EPIA 2007 - Portuguese Conference on Artificial Intelligence, December, Guimarães, Portugal, pp. 512-523, 2007. APPIA, ISBN-13 978-989-95618-0-9. Available at: [Web Link]

Citation Request:

This dataset is public available for research. The details are described in [Cortez and Morais, 2007].

Please include this citation if you plan to use this database:

[Cortez and Morais, 2007] P. Cortez and A. Morais. A Data Mining Approach to Predict Forest Fires using Meteorological Data. In J. Neves, M. F. Santos and J. Machado Eds., New Trends in Artificial Intelligence, Proceedings of the 13th EPIA 2007 - Portuguese Conference on Artificial Intelligence, December, Guimarães, Portugal, pp. 512-523, 2007. APPIA, ISBN-13 978-989-95618-0-9. Available at: [Web Link]

Anıl Durmaz 181805080

Muhammet Furkan Yılmaz 181805069

Ferhat Kamalı 171805056