

11. Chest Pain (Acute Coronary Syndrome) Algorithm


This clinical pathway is intended to supplement, rather than substitute for, professional judgment and may be changed depending upon patient's individual needs. Failure to comply with this pathway does not represent a breach of the standard of care.

Chest Discomfort Suggestive of Ischemia
(includes **anginal equivalents** (atypical symptoms) like exertional pain in the ear, jaw, neck, shoulder, arm, back, or epigastric area; exertional dyspnoea; nausea and vomiting; diaphoresis; and fatigue.

- Monitor, support ABCs in the **Resuscitation Room (ER)**. Be prepared to provide CPR, Defibrillation and **?Thrombolysis/Fibrinolysis**
- **Obtain/review 12-lead ECG within 10 minutes of arrival to ED**
 - Do a **V4R** if ST elevation in lead V1 with simultaneous ST depression in V2 -? **Right sided STEMI**
 - Do **V7 - V9** if ST depressions ≥ 1 mm with upright T-waves in ≥ 2 contiguous anterior precordial leads (V1 to V3) -? **Posterior STEMI**
 - If there is **ST elevation in aVR ≥ 1 mm and aVR \geq V1** with widespread horizontal ST depression, most prominent in leads I, II and V4-6 – **consult an Interventional Cardiologist** immediately for PCI (**Left main coronary artery occlusion/Proximal LAD lesion/Severe sub endocardial ischaemia, nonlocalized**)
 - Sinus Tachycardia, T wave inversion in III & V1, V3 or (S1, Q3, T3) pattern -? See **15. Pulmonary Embolism Algorithm**
- Check vital signs (BP, PR, RR, SPO₂, T°C, **RBS**)
- Start Oxygen **IF** SPO₂ < 90% or if patient is dyspnoeic. Maintain SPO₂ \geq 90%
- Perform brief, targeted history, physical exam – **Indicate time of symptoms onset**

- **Consider other life threatening causes of chest pain** (pulmonary embolus, cardiac tamponade, aortic dissection, tension pneumothorax, oesophageal rupture)
- **Review initial 12-lead ECG**

Sequence of ECG changes seen during evolution of myocardial infarction



ST Elevation	MI Description	Coronaries affected
V2 – V5	Anterior	LAD
V1 – V2	Septal	Septal LAD
II, III, aVF	Inferior	RCx (20%) or RCA (80%)
V1 – V4	Anterolateral	
V3 – V6, I, aVL	Anteroseptal	
I, aVL, V5, V6	Lateral	LCx
V7, V8, V9	Posterior	RCx
V1, V4R	RV	RCA

ST elevation
ST-Elevation MI (STEMI)

See
12. STEMI Algorithm

ST depression > 0.5mm or dynamic T-wave inversion ≥ 2 mm; strongly suspicious for ischemia
High-Risk Unstable Angina/Non-ST-Elevation MI (UA/NSTEMI)

See
13. NSTEMI/UA Algorithm

Normal or Non-diagnostic changes in ST segment or T wave
Intermediate/Low Risk UA