

Q.6	(i) A bag contains 6 white, 8 red and 5 black balls. A ball is drawn at random. Find the probability that it is white. (ii) For any two events A and B, given that $P(A) = 0.4$, $P(B) = 0.7$ and $P(A \cap B) = 0.3$. Then find $P(A \cup B)$.	2
Q.7	(i) A card is drawn from a well shuffled pack of 52 cards. Find the probability that it is either a spade or a king. (ii) If A and B are two events, where $P(A) = \frac{1}{2}$, $P(B) = \frac{1}{3}$ and $P(A \cap B) = \frac{1}{4}$. Then calculate (i) $P(A B)$ (ii) $P(B A)$ (iii) $P(A \cup B)$ (iv) $P(A \cap B)$, when A and B are independent.	2
Q.8	(i) Probability that A speaks the truth is $\frac{3}{5}$ and B speaks the truth is $\frac{5}{8}$. In what percentage of cases they are likely to contradict each other? (ii) 5 man out of 100 and 25 women out of 1000 are colour blind. A colour blind person is chosen at random. What is the probability of his being a male? (Assume that male and female are in equal proportion).	2
Q.9	(i) Eight coins are thrown simultaneously. Find the probability of obtaining at least six heads. (ii) Assuming the probability of male birth as $\frac{1}{2}$, find the probability that a family of three children will have: (i) at least one girl, (ii) two boys and a girl, (iii) at the most two girls.	2
Q.10	(i) If 2% of the electric bulbs manufactured by a certain company are defective, find the probability that in a sample of 200 bulbs (i) less than two bulbs are defective (ii) more than two bulbs are defectives. ($e^{-4} = 0.0183$) (ii) Suppose that a manufactured product has 2 defects per unit of product inspected. Using Poisson's distribution, calculate the probability of finding a product without any defect, 3 defects and 4 defects. ($e^{-2} = 0.135$)	2