



# Team Contest Reference

## Team:

System.out.println(42);

Roland Haase  
Thore Tiemann  
Marcel Wienöbst

## Contents

<b>1 Algorithms</b>	<b>2</b>
1.1 Binary Search . . . . .	2
1.2 Held Karp . . . . .	2
1.3 LongestIncreasingSubsequence . . . . .	2
1.4 LongestIncreasingSubsequence . . . . .	3
1.5 NextPermutation . . . . .	3
<b>2 Graphs</b>	<b>3</b>
2.1 BellmanFord . . . . .	3
2.2 BipartiteGraphCheck . . . . .	3
2.3 Dijkstra . . . . .	4
2.4 EdmondsKarp . . . . .	4
2.5 FenwickTree . . . . .	4
2.6 FloydWarshall . . . . .	5
2.7 BFS AdjMtrx Iterativ . . . . .	5
2.8 Topologische Sortierung . . . . .	5
<b>3 Math</b>	<b>5</b>
3.1 GrahamScan . . . . .	5
<b>4 Java Knowhow</b>	<b>7</b>
4.1 System.out.printf() und String.format() . . . . .	7
4.2 Speed up IO . . . . .	7

$n$	Runtime $100 \cdot 10^6$ in 3s
[10, 11]	$\mathcal{O}(n!)$
$< 22$	$\mathcal{O}(n2^n)$
$\leq 100$	$\mathcal{O}(n^4)$
$\leq 400$	$\mathcal{O}(n^3)$
$\leq 2.000$	$\mathcal{O}(n^2 \log n)$
$\leq 10.000$	$\mathcal{O}(n^2)$
$\leq 1.000.000$	$\mathcal{O}(n \log n)$
$\leq 100.000.000$	$\mathcal{O}(n)$

byte (8 Bit, signed): -128 ... 127

short (16 Bit, signed): -32.768 ... 32.767

integer (32 Bit, signed): -2.147.483.648 ... 2.147.483.647

long (64 Bit, signed):  $-2^{63} \dots 2^{63} - 1$

**MD5:** cat <string> | tr -d [:space:] | md5sum

# 1 Algorithms

## 1.1 Binary Search

Binary searches for an element in a sorted array.

```

1 public static boolean BinarySearch(int[] array, int N,
2     int a) {
3     int lo = 0;
4     int hi = N-1;
5     while(lo <= hi) {
6         int mid = (int) ((lo + hi) / 2.0) + 0.6);
7         if(array[mid] < a) {
8             lo = mid+1;
9         } else {
10            hi = mid-1;
11        }
12    }
13    if(lo < N && array[lo] == a) {
14        return true;
15    } else {
16        return false;
17    }

```

MD5: bb87f09a50f05e688706641c26759706 |  $\mathcal{O}(\log n)$

## 1.2 Held Karp

Algorithm for TSP

```

1 public static int[] tsp(int[][] graph) {
2     int n = graph.length;
3     if(n == 1) return new int[] {0};
4     //C stores the shortest distance to node of the
5     //second dimension
6     //first dimension is the bitstring of included
7     //nodes on the way
8     int[][] C = new int[1<<n][n];
9     int[][] p = new int[1<<n][n];
10    //initialize
11    for(int k = 1; k < n; k++) {
12        C[1<<k][k] = graph[0][k];
13    }
14    for(int s = 2; s < n; s++) {
15        for(int S = 1; S < (1<<n); S++) {
16            if(Integer.bitCount(S) != s || (S&1) == 1)
17                continue;
18            for(int k = 1; k < n; k++) {
19                if((S & (1 << k)) == 0)
20                    continue;
21                //Smk is the set of nodes without k
22                int Smk = S ^ (1<<k);
23                int min = Integer.MAX_VALUE;
24                int minprev = 0;
25                for(int m=1; m<n; m++) {
26                    if((Smk & (1<<m)) == 0)
27                        continue;
28                    //distance to m with the nodes in Smk +
29                    //connection from m to k
30                    int tmp = C[Smk][m] + graph[m][k];
31                    if(tmp < min) {
32                        min = tmp;
33                        minprev = m;

```

```

34        }
35        C[S][k] = min;
36        p[S][k] = minprev;
37    }
38    }
39    }
40    //find shortest tour length
41    int min = Integer.MAX_VALUE;
42    int minprev = -1;
43    for(int k = 1; k < n; k++) {
44        //Set of all nodes except for the first + cost
45        //from 0 to k
46        int tmp = C[(1<<n) - 2][k] + graph[0][k];
47        if(tmp < min) {
48            min = tmp;
49            minprev = k;
50        }
51    }
52    //Note that the tour has not been tested yet, only
53    //the correctness of the min-tour-value
54    //backtrack tour
55    int[] tour = new int[n+1];
56    tour[n] = 0;
57    tour[n-1] = minprev;
58    int bits = (1<<n)-2;
59    for(int k = n-2; k>0; k--) {
60        tour[k] = p[bits][tour[k+1]];
61        bits = bits ^ (1<<tour[k+1]);
62    }
63    tour[0] = 0;
64    return tour;
65 }

```

MD5: 233d98980b1f4dae50ac892d7112dafb |  $\mathcal{O}(2^n n^2)$

## 1.3 LongestIncreasingSubsequence

Computes the longest increasing subsequence and is easy to be adapted.

```

1 //This has not been tested yet (adapted from tested C
2 ++ Murcia Code)
3 public static int longestInc(int[] array, int N) {
4     int[] m = new int[N];
5     for (int i = N - 1; i >= 0; i--) {
6         m[i] = 1;
7         for (int j = i + 1; j < N; j++) {
8             if (array[j] > array[i]) {
9                 if (m[i] < m[j] + 1) {
10                    m[i] = m[j] + 1;
11                }
12            }
13        }
14    }
15    int longest = 0;
16    for (int i = 0; i < N; i++) {
17        if (m[i] > longest) {
18            longest = m[i];
19        }
20    }
21    return longest;

```

MD5: 7ee618a580f2736226054b5e106d5635 |  $\mathcal{O}(n^2)$

## 1.4 LongestIncreasingSubsequence

Computes the longest increasing subsequence using binary search.

```

1 public static int[] LongestIncreasingSubsequencenlogn(
2     int[] a, int[] p) {
3     int[] m = new int[a.length+1];
4     int l = 0;
5     for(int i = 0; i < a.length; i++) {
6         int lo = 1;
7         int hi = l;
8         while(lo <= hi) {
9             int mid = (int) (((lo + hi) / 2.0) + 0.6);
10            if(a[m[mid]] < a[i]) {
11                lo = mid+1;
12            } else {
13                hi = mid-1;
14            }
15            int newL = lo;
16            p[i] = m[newL-1];
17            m[newL] = i;
18            if(newL > l) {
19                l = newL;
20            }
21        }
22        int[] s = new int[l];
23        int k = m[l];
24        for(int i = l-1; i >= 0; i--) {
25            s[i] = a[k];
26            k = p[k];
27        }
28        return s;
29    }

```

MD5: e4b7591a2e204809f3e105521a616f70 |  $\mathcal{O}(n * \log n)$

## 1.5 NextPermutation

n Returns true if there is another permutation. Can also be used to compute the nextPermutation of an array.

```

1 public static boolean nextPermutation(char[] a) {
2     int i = a.length - 1;
3     while(i > 0 && a[i-1] >= a[i]) {
4         i--;
5     }
6     if(i <= 0) {
7         return false;
8     }
9     int j = a.length - 1;
10    while (a[j] <= a[i-1]) {
11        j--;
12    }
13    char tmp = a[i - 1];
14    a[i - 1] = a[j];
15    a[j] = tmp;
16
17    j = a.length - 1;
18    while(i < j) {
19        tmp = a[i];
20        a[i] = a[j];
21        a[j] = tmp;
22        i++;
23        j--;
24    }
25    return true;

```

26 }

MD5: ca6266722db16f2dc8eae5a6cc5fcacf |  $\mathcal{O}(?)$

## 2 Graphs

### 2.1 BellmanFord

DESCRIPTION MISSING

```

1 public static boolean bellmanFord(Vertex[] vertices) {
2     //source is 0
3     vertices[0].mindistance = 0;
4     //calc distances
5     for(int i = 0; i < vertices.length-1; i++) {
6         for(int j = 0; j < vertices.length; j++) {
7             for(Edge e: vertices[j].adjacencies) {
8                 if(vertices[j].mindistance != Integer.
9                     MAX_VALUE
10                    && e.target.mindistance > vertices[j].
11                        mindistance + e.distance) {
12                    e.target.mindistance = vertices[j].
13                        mindistance + e.distance;
14                }
15            }
16        }
17    }
18    //check for negative-length cycle
19    for(int i = 0; i < vertices.length; i++) {
20        for(Edge e: vertices[i].adjacencies) {
21            if(vertices[i].mindistance != Integer.
22                MAX_VALUE && e.target.mindistance >
23                vertices[i].mindistance + e.distance) {
24                return true;
25            }
26        }
27    }
28    return false;
29 }

```

MD5: 36561a7913a81baf7b7c79b606683819 |  $\mathcal{O}(???)$

### 2.2 BipartiteGraphCheck

DESCRIPTION MISSING

```

1 public static boolean bipartiteGraphCheck(ArrayList<
2     ArrayList<Integer>> graph, int N) {
3     int[] color = new int[N];
4     for(int i = 0; i < N; i++) color[i] = -1;
5     color[0] = 1;
6     Queue<Integer> q = new LinkedList<Integer>();
7     q.add(0);
8     while(!q.isEmpty()) {
9         int u = q.poll();
10        for(int i : graph.get(u)) {
11            if(color[i] == -1) {
12                color[i] = 1-color[u];
13                q.add(i);
14            } else if(color[u] == color[i]) {
15                return false;
16            }
17        }
18    }
19    return true;

```

```
19 }
MD5: 5cb4622cf75e4ea5ffae51b0b48abf2b | O(???)
```

## 2.3 Dijkstra

Finds the shortest paths from one vertex to every other vertex in the graph (SSSP).

For negative weights, add  $|\min|+1$  to each edge, later subtract from result.

To get a different shortest path when edges are ints, add an  $\epsilon = \frac{1}{k+1}$  on each edge of the shortest path of length  $k$ , run again.

Input: A source vertex  $s$  and an adjacency list  $G$ .

Output: Modified adj. list with distances from  $s$  and predecessor vertices set.

```
1 public static void dijkstra(Vertex[] vertices, int src
2 ) {
3     vertices[src].mindistance = 0;
4     PriorityQueue<Vertex> queue = new PriorityQueue<
5         Vertex>();
6     queue.add(vertices[src]);
7     while(!queue.isEmpty()) {
8         Vertex u = queue.poll();
9         if(u.visited)
10            continue;
11         u.visited = true;
12         for(Edge e : u.adjacencies) {
13             Vertex v = e.target;
14             if(v.mindistance > u.mindistance + e.distance
15                 ) {
16                 v.mindistance = u.mindistance + e.distance
17                 ;
18                 queue.add(v);
19             }
20         }
21     }
22 }
23 class Vertex implements Comparable<Vertex> {
24     public int id;
25     public int mindistance = Integer.MAX_VALUE;
26     public LinkedList<Edge> adjacencies = new
27         LinkedList<Edge>();
28     public boolean visited = false;
29     public int compareTo(Vertex other) {
30         return Integer.compare(this.mindistance, other.
31             mindistance);
32     }
33 }
34 class Edge {
35     public Vertex target;
36     public int distance;
37
38     public Edge (Vertex target, int distance) {
39         this.target = target;
40         this.distance = distance;
41     }
42 }
```

MD5: d6882162849418a2541cfc7f6c3ddc58 |  $O(|E| \log |V|)$

## 2.4 EdmondsKarp

Finds the greatest flow in a graph.

```
public static boolean BFS(int[][] graph, int s, int t,
    int[] parent) {
    int N = graph.length;
    boolean[] visited = new boolean[N];
    for(int i = 0; i < N; i++) {
        visited[i] = false;
    }
    Queue<Integer> queue = new LinkedList<Integer>();
    queue.add(s);
    visited[s] = true;
    parent[s] = -1;
    while(!queue.isEmpty()) {
        int u = queue.poll();
        if(u == t) return true;
        for(int v = 0; v < N; v++) {
            if(visited[v] == false && graph[u][v] > 0) {
                queue.add(v);
                parent[v] = u;
                visited[v] = true;
            }
        }
    }
    return (visited[t]);
}

public static int fordFulkerson(int[][] graph, int s,
    int t) {
    int N = graph.length;
    int[][] rgraph = new int[graph.length][graph.length
    ];
    for(int u = 0; u < graph.length; u++) {
        for(int v = 0; v < graph.length; v++) {
            rgraph[u][v] = graph[u][v];
        }
    }
    int[] parent = new int[N];
    int maxflow = 0;
    while(BFS(rgraph, s, t, parent)) {
        int pathflow = Integer.MAX_VALUE;
        for(int v = t; v != s; v = parent[v]) {
            int u = parent[v];
            pathflow = Math.min(pathflow, rgraph[u][v]);
        }
        for(int v = t; v != s; v = parent[v]) {
            int u = parent[v];
            rgraph[u][v] -= pathflow;
            rgraph[v][u] += pathflow;
        }
        maxflow += pathflow;
    }
    return maxflow;
}
```

MD5: 8d85785d45794f20303d9b9f920e80dd |  $O(???)$

## 2.5 FenwickTree

Can be used for computing prefix sums.

```
int[] fwktree = new int[m + n + 1];
public static int read(int index, int[] fenwickTree) {
    int sum = 0;
    while (index > 0) {
        sum += fenwickTree[index];
        index -= (index & -index);
    }
}
```

```

8   return sum;
9 }
10 public static int[] update(int index, int addValue,
11   int[] fenwickTree) {
12   while (index <= fenwickTree.length - 1) {
13     fenwickTree[index] += addValue;
14     index += (index & -index);
15   }
16   return fenwickTree;

```

MD5: 97fd176a403e68cb76a82196191d5f19 |  $\mathcal{O}(\log n)$

## 2.6 FloydWarshall

Finds all shortest paths. Paths in array next, distances in ans.

```

1 public static void floydWarshall(int[][] graph, int
2   [][] next, int[][] ans) {
3   for(int i = 0; i < ans.length; i++) {
4     for(int j = 0; j < ans.length; j++) {
5       ans[i][j] = graph[i][j];
6     }
7   }
8   for (int k = 0; k < ans.length; k++) {
9     for (int i = 0; i < ans.length; i++) {
10      for (int j = 0; j < ans.length; j++) {
11        if (ans[i][k] + ans[k][j] < ans[i][j]
12          && ans[i][k] < Integer.MAX_VALUE && ans[k]
13            [j] < Integer.MAX_VALUE) {
14          ans[i][j] = ans[i][k] + ans[k][j];
15          next[i][j] = next[i][k];
16        }
17      }
18    }

```

MD5: 4faf8c41a9070f106e68864cc131706d |  $\mathcal{O}(n^3)$

## 2.7 BFS AdjMtrx Iterativ

Iterative BFS on adjacency matrix. Returns true or false, depending on whether there is a connection between s and g or not.

```

1 public static boolean BFSWithoutPathForAdjMatr(int s,
2   int g, int[][] graph) {
3   //s being the start and g the goal
4   boolean[] visited = new boolean[graph.length];
5   for(int i = 0; i < visited.length; i++)
6     visited[i] = false;
7   Queue<Integer> queue = new LinkedList<Integer>();
8   queue.add(s);
9   visited[s] = true;
10  while(!queue.isEmpty()) {
11    int node = queue.poll();
12    if(node == g)
13      return true;
14    for(int i = 0; i < graph.length; i++) {
15      if(graph[node][i] > 0 && !visited[i]) {
16        queue.add(i);
17        visited[i] = true;
18      }
19    }
20  }

```

return false;

```

21 }

```

MD5: 754e7dfa0a691a2511464e16104b8880 |  $\mathcal{O}(V + E?)$

## 2.8 Topologische Sortierung

Sorts a graph (represented as adjMtrx) topologically

```

1 // l enthaelt alle Knoten topologisch sortiert (Start:
2   0, Ende= n)
3 int[] l = new int[n];
4 int idx = 0;
5 // s enthaelt alle Knoten, die keine eingehende Kante
6   haben
7 ArrayList<Integer> s = new ArrayList<Integer>();
8 // initialisiere s
9 for (int i = 0; i < n; i++) {
10  if (edgesIn[i] == 0) {
11    s.add(i);
12  }
13 }
14 // Algo Beginn
15 while (!s.isEmpty()) {
16   int node = s.remove(0);
17   l[idx++] = node;
18   for (int i = 0; i < n; i++) {
19     if (adjMtrx[node][i] > 0) {
20       adjMtrx[node][i] = false;
21       edgesIn[i] -= 1;
22       if (edgesIn[i] == 0) {
23         s.add(i);
24       }
25     }

```

MD5: 01974f4bab4e48916ecdc48531a79c84 |  $\mathcal{O}(O(V + E))$

## 3 Math

### 3.1 GrahamScan

GrahamScan finds convex hull. Still has collinear point problematic at the last diagonal.

```

1 public static int ccw(Point src, Point q1, Point q2) {
2   return (q1.x - src.x) * (q2.y - src.y) - (q2.x -
3     src.x) * (q1.y - src.y);
4 }
5 public static boolean isColl(Point a, Point b, Point c) {
6   if((b.y - a.y) * (c.x - b.x) == (c.y - b.y) * (b.x
7     - a.x)) {
8     return true;
9   } else {
10    return false;
11  }
12 }
13 public static double calcDist(Point src, Point target) {
14   return Math.sqrt((src.x + target.x) * (src.x +
15     target.x) + (src.y + target.y) * (src.y +
16     target.y));

```

```

15 }
16
17 //Expects a array sorted with PolarComp as Comparator
18 //IMPORTANT! before sorting put lowest, and if two are
19 //the same leftmost, element at position 0 in array
20 public static void grahamScan(Point[] points) {
21     int m = 1;
22     for(int i = 2; i < points.length; i++) {
23         while(ccw(points[m-1], points[m], points[i]) <
24             0) {
25             if(m > 1) m--;
26             else if(i == points.length) break;
27             else i++;
28         }
29         m++;
30         Point tmp = points[i];
31         points[i] = points[m];
32         points[m] = tmp;
33     }
34 }
35
36 class Point {
37     int x;
38     int y;
39     public Point(int x, int y) {
40         this.x = x;
41         this.y = y;
42     }
43 }
44
45 class PolarComp implements Comparator<Point> {
46     Point src;
47
48     public PolarComp(Point source) {
49         src = source;
50     }
51
52     public double calcDist(Point q1, Point q2) {
53         return Math.sqrt((q1.x - q2.x) * (q1.x - q2.x) +
54             (q1.y - q2.y) * (q1.y - q2.y));
55     }
56
57     public int ccw(Point q1, Point q2) {
58         return (q1.x - src.x) * (q2.y - src.y) - (q2.x -
59             src.x) * (q1.y - src.y);
60     }
61
62     public int compare(Point q1, Point q2) {
63         int res = ccw(q1, q2);
64         double dist1 = calcDist(src, q1);
65         double dist2 = calcDist(src, q2);
66         if(res > 0) return -1;
67         else if(res < 0) return 1;
68         else if(res == 0 && dist1 < dist2) return 1;
69         else if(res == 0 && dist1 > dist2) return -1;
70         else return 0;
71     }
72 }

```

## 4 Java Knowhow

### 4.1 System.out.printf() und String.format()

**Syntax:** %[flags][width][.precision][conv]

**flags:**

- left-justify (default: right)
- + always output number sign
- 0 zero-pad numbers
- (space) space instead of minus for pos. numbers
- , group triplets of digits with ,

**width** specifies output width

**precision** is for floating point precision

**conv:**

- d byte, short, int, long
- f float, double
- c char (use C for uppercase)
- s String (use S for all uppercase)

### 4.2 Speed up IO

```
Use BufferedReader br = new BufferedReader(new InputStreamReader(System.in));
```

```
Use Double.parseDouble(Scanner.next())
```

## Theoretical Computer Science Cheat Sheet

Definitions		Series	
$f(n) = O(g(n))$	iff $\exists$ positive $c, n_0$ such that $0 \leq f(n) \leq cg(n) \forall n \geq n_0$ .	$\sum_{i=1}^n i = \frac{n(n+1)}{2}, \quad \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}, \quad \sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}.$	
$f(n) = \Omega(g(n))$	iff $\exists$ positive $c, n_0$ such that $f(n) \geq cg(n) \geq 0 \forall n \geq n_0$ .	In general:	
$f(n) = \Theta(g(n))$	iff $f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$ .	$\sum_{i=1}^n i^m = \frac{1}{m+1} \left[ (n+1)^{m+1} - 1 - \sum_{i=1}^n ((i+1)^{m+1} - i^{m+1} - (m+1)i^m) \right]$	
$f(n) = o(g(n))$	iff $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$ .	$\sum_{i=1}^{n-1} i^m = \frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} B_k n^{m+1-k}.$	
$\lim_{n \rightarrow \infty} a_n = a$	iff $\forall \epsilon > 0, \exists n_0$ such that $ a_n - a  < \epsilon, \forall n \geq n_0$ .	Geometric series:	
$\sup S$	least $b \in$ such that $b \geq s, \forall s \in S$ .	$\sum_{i=0}^{\infty} c^i = \frac{c^{n+1} - 1}{c - 1}, \quad c \neq 1, \quad \sum_{i=0}^{\infty} c^i = \frac{1}{1 - c}, \quad \sum_{i=1}^{\infty} c^i = \frac{c}{1 - c}, \quad  c  < 1,$	
$\inf S$	greatest $b \in$ such that $b \leq s, \forall s \in S$ .	$\sum_{i=0}^{\infty} ic^i = \frac{nc^{n+2} - (n+1)c^{n+1} + c}{(c-1)^2}, \quad c \neq 1, \quad \sum_{i=0}^{\infty} ic^i = \frac{c}{(1-c)^2}, \quad  c  < 1.$	
$\liminf_{n \rightarrow \infty} a_n$	$\lim_{n \rightarrow \infty} \inf \{a_i \mid i \geq n, i \in \mathbb{N}\}.$	Harmonic series:	
$\limsup_{n \rightarrow \infty} a_n$	$\lim_{n \rightarrow \infty} \sup \{a_i \mid i \geq n, i \in \mathbb{N}\}.$	$H_n = \sum_{i=1}^n \frac{1}{i}, \quad \sum_{i=1}^n iH_i = \frac{n(n+1)}{2}H_n - \frac{n(n-1)}{4}.$	
$\binom{n}{k}$	Combinations: Size $k$ sub-sets of a size $n$ set.	$\sum_{i=1}^n H_i = (n+1)H_n - n, \quad \sum_{i=1}^n \binom{i}{m} H_i = \binom{n+1}{m+1} \left( H_{n+1} - \frac{1}{m+1} \right).$	
$[n]_k$	Stirling numbers (1st kind): Arrangements of an $n$ element set into $k$ cycles.	1. $\binom{n}{k} = \frac{n!}{(n-k)!k!}, \quad 2. \sum_{k=0}^n \binom{n}{k} = 2^n, \quad 3. \binom{n}{k} = \binom{n}{n-k},$	
$\{n\}_k$	Stirling numbers (2nd kind): Partitions of an $n$ element set into $k$ non-empty sets.	4. $\binom{n}{k} = \frac{n}{k} \binom{n-1}{k-1}, \quad 5. \binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1},$	
$\langle n \rangle_k$	1st order Eulerian numbers: Permutations $\pi_1 \pi_2 \dots \pi_n$ on $\{1, 2, \dots, n\}$ with $k$ ascents.	6. $\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}, \quad 7. \sum_{k=0}^n \binom{r+k}{k} = \binom{r+n+1}{n},$	
$\langle\langle n \rangle\rangle_k$	2nd order Eulerian numbers.	8. $\sum_{k=0}^n \binom{k}{m} = \binom{n+1}{m+1}, \quad 9. \sum_{k=0}^n \binom{r}{k} \binom{s}{n-k} = \binom{r+s}{n},$	
$C_n$	Catalan Numbers: Binary trees with $n+1$ vertices.	10. $\binom{n}{k} = (-1)^k \binom{k-n-1}{k}, \quad 11. \left\{ \begin{matrix} n \\ 1 \end{matrix} \right\} = \left\{ \begin{matrix} n \\ n \end{matrix} \right\} = 1,$	
14. $\left[ \begin{matrix} n \\ 1 \end{matrix} \right] = (n-1)!,$	15. $\left[ \begin{matrix} n \\ 2 \end{matrix} \right] = (n-1)!H_{n-1},$	16. $\left[ \begin{matrix} n \\ n \end{matrix} \right] = 1,$	17. $\left[ \begin{matrix} n \\ k \end{matrix} \right] \geq \left\{ \begin{matrix} n \\ k \end{matrix} \right\},$
18. $\left[ \begin{matrix} n \\ k \end{matrix} \right] = (n-1) \left[ \begin{matrix} n-1 \\ k \end{matrix} \right] + \left[ \begin{matrix} n-1 \\ k-1 \end{matrix} \right],$	19. $\left\{ \begin{matrix} n \\ n-1 \end{matrix} \right\} = \left[ \begin{matrix} n \\ n-1 \end{matrix} \right] = \binom{n}{2},$	20. $\sum_{k=0}^n \left[ \begin{matrix} n \\ k \end{matrix} \right] = n!,$	21. $C_n = \frac{1}{n+1} \binom{2n}{n},$
22. $\left\langle \begin{matrix} n \\ 0 \end{matrix} \right\rangle = \left\langle \begin{matrix} n \\ n-1 \end{matrix} \right\rangle = 1,$	23. $\left\langle \begin{matrix} n \\ k \end{matrix} \right\rangle = \left\langle \begin{matrix} n \\ n-1-k \end{matrix} \right\rangle,$	24. $\left\langle \begin{matrix} n \\ k \end{matrix} \right\rangle = (k+1) \left\langle \begin{matrix} n-1 \\ k \end{matrix} \right\rangle + (n-k) \left\langle \begin{matrix} n-1 \\ k-1 \end{matrix} \right\rangle,$	
25. $\left\langle \begin{matrix} 0 \\ k \end{matrix} \right\rangle = \begin{cases} 1 & \text{if } k=0, \\ 0 & \text{otherwise} \end{cases}$	26. $\left\langle \begin{matrix} n \\ 1 \end{matrix} \right\rangle = 2^n - n - 1,$	27. $\left\langle \begin{matrix} n \\ 2 \end{matrix} \right\rangle = 3^n - (n+1)2^n + \binom{n+1}{2},$	
28. $x^n = \sum_{k=0}^n \left\langle \begin{matrix} n \\ k \end{matrix} \right\rangle \binom{x+k}{n},$	29. $\left\langle \begin{matrix} n \\ m \end{matrix} \right\rangle = \sum_{k=0}^m \binom{n+1}{k} (m+1-k)^n (-1)^k,$	30. $m! \left\{ \begin{matrix} n \\ m \end{matrix} \right\} = \sum_{k=0}^n \left\langle \begin{matrix} n \\ k \end{matrix} \right\rangle \binom{k}{n-m},$	
31. $\left\langle \begin{matrix} n \\ m \end{matrix} \right\rangle = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \binom{n-k}{m} (-1)^{n-k-m} k!,$	32. $\langle\langle \begin{matrix} n \\ 0 \end{matrix} \rangle\rangle = 1,$	33. $\langle\langle \begin{matrix} n \\ n \end{matrix} \rangle\rangle = 0 \quad \text{for } n \neq 0,$	
34. $\langle\langle \begin{matrix} n \\ k \end{matrix} \rangle\rangle = (k+1) \langle\langle \begin{matrix} n-1 \\ k \end{matrix} \rangle\rangle + (2n-1-k) \langle\langle \begin{matrix} n-1 \\ k-1 \end{matrix} \rangle\rangle,$	35. $\sum_{k=0}^n \langle\langle \begin{matrix} n \\ k \end{matrix} \rangle\rangle = \frac{(2n)^n}{2^n},$	36. $\left\{ \begin{matrix} x \\ x-n \end{matrix} \right\} = \sum_{k=0}^n \langle\langle \begin{matrix} n \\ k \end{matrix} \rangle\rangle \binom{x+n-1-k}{2n},$	37. $\left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\} = \sum_k \binom{n}{k} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} = \sum_{k=0}^n \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (m+1)^{n-k},$



## Theoretical Computer Science Cheat Sheet

## Identities Cont.

$$\begin{aligned}
38. \quad \begin{bmatrix} n+1 \\ m+1 \end{bmatrix} &= \sum_k \begin{bmatrix} n \\ k \end{bmatrix} \begin{bmatrix} k \\ m \end{bmatrix} = \sum_{k=0}^n \begin{bmatrix} k \\ m \end{bmatrix} n^{\overline{n-k}} = n! \sum_{k=0}^n \frac{1}{k!} \begin{bmatrix} k \\ m \end{bmatrix}, & 39. \quad \begin{bmatrix} x \\ x-n \end{bmatrix} &= \sum_{k=0}^n \left\langle \begin{bmatrix} n \\ k \end{bmatrix} \right\rangle \begin{bmatrix} x+k \\ 2n \end{bmatrix}, \\
40. \quad \left\{ \begin{bmatrix} n \\ m \end{bmatrix} \right\} &= \sum_k \begin{bmatrix} n \\ k \end{bmatrix} \left\{ \begin{bmatrix} k+1 \\ m+1 \end{bmatrix} \right\} (-1)^{n-k}, & 41. \quad \begin{bmatrix} n \\ m \end{bmatrix} &= \sum_k \begin{bmatrix} n+1 \\ k+1 \end{bmatrix} \begin{bmatrix} k \\ m \end{bmatrix} (-1)^{m-k}, \\
42. \quad \left\{ \begin{bmatrix} m+n+1 \\ m \end{bmatrix} \right\} &= \sum_{k=0}^m k \left\{ \begin{bmatrix} n+k \\ k \end{bmatrix} \right\}, & 43. \quad \begin{bmatrix} m+n+1 \\ m \end{bmatrix} &= \sum_{k=0}^m k(n+k) \begin{bmatrix} n+k \\ k \end{bmatrix}, \\
44. \quad \begin{bmatrix} n \\ m \end{bmatrix} &= \sum_k \left\{ \begin{bmatrix} n+1 \\ k+1 \end{bmatrix} \right\} \begin{bmatrix} k \\ m \end{bmatrix} (-1)^{m-k}, & 45. \quad (n-m)! \begin{bmatrix} n \\ m \end{bmatrix} &= \sum_k \begin{bmatrix} n+1 \\ k+1 \end{bmatrix} \left\{ \begin{bmatrix} k \\ m \end{bmatrix} \right\} (-1)^{m-k}, \quad \text{for } n \geq m, \\
46. \quad \left\{ \begin{bmatrix} n \\ n-m \end{bmatrix} \right\} &= \sum_k \begin{bmatrix} m-n \\ m+k \end{bmatrix} \begin{bmatrix} m+n \\ n+k \end{bmatrix} \begin{bmatrix} m+k \\ k \end{bmatrix}, & 47. \quad \begin{bmatrix} n \\ n-m \end{bmatrix} &= \sum_k \begin{bmatrix} m-n \\ m+k \end{bmatrix} \begin{bmatrix} m+n \\ n+k \end{bmatrix} \left\{ \begin{bmatrix} m+k \\ k \end{bmatrix} \right\}, \\
48. \quad \left\{ \begin{bmatrix} n \\ \ell+m \end{bmatrix} \right\} \begin{bmatrix} \ell+m \\ \ell \end{bmatrix} &= \sum_k \left\{ \begin{bmatrix} k \\ \ell \end{bmatrix} \right\} \left\{ \begin{bmatrix} n-k \\ m \end{bmatrix} \right\} \begin{bmatrix} n \\ k \end{bmatrix}, & 49. \quad \begin{bmatrix} n \\ \ell+m \end{bmatrix} \begin{bmatrix} \ell+m \\ \ell \end{bmatrix} &= \sum_k \begin{bmatrix} k \\ \ell \end{bmatrix} \begin{bmatrix} n-k \\ m \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix}.
\end{aligned}$$

## Trees

Every tree with  $n$  vertices has  $n-1$  edges.

Kraft inequality: If the depths of the leaves of a binary tree are  $d_1, \dots, d_n$ :

$$\sum_{i=1}^n 2^{-d_i} \leq 1,$$

and equality holds only if every internal node has 2 sons.

## Recurrences

Master method:

$$T(n) = aT(n/b) + f(n), \quad a \geq 1, b > 1$$

If  $\exists \epsilon > 0$  such that  $f(n) = O(n^{\log_b a - \epsilon})$  then

$$T(n) = \Theta(n^{\log_b a}).$$

If  $f(n) = \Theta(n^{\log_b a})$  then

$$T(n) = \Theta(n^{\log_b a} \log_2 n).$$

If  $\exists \epsilon > 0$  such that  $f(n) = \Omega(n^{\log_b a + \epsilon})$ , and  $\exists c < 1$  such that  $af(n/b) \leq cf(n)$  for large  $n$ , then

$$T(n) = \Theta(f(n)).$$

Substitution (example): Consider the following recurrence

$$T_{i+1} = 2^{2^i} \cdot T_i^2, \quad T_1 = 2.$$

Note that  $T_i$  is always a power of two.

Let  $t_i = \log_2 T_i$ . Then we have

$$t_{i+1} = 2^i + 2t_i, \quad t_1 = 1.$$

Let  $u_i = t_i/2^i$ . Dividing both sides of the previous equation by  $2^{i+1}$  we get

$$\frac{t_{i+1}}{2^{i+1}} = \frac{2^i}{2^{i+1}} + \frac{t_i}{2^i}.$$

Substituting we find

$$u_{i+1} = \frac{1}{2} + u_i, \quad u_1 = \frac{1}{2},$$

which is simply  $u_i = i/2$ . So we find that  $T_i$  has the closed form  $T_i = 2^{i2^{i-1}}$ .

Summing factors (example): Consider the following recurrence

$$T(n) = 3T(n/2) + n, \quad T(1) = 1.$$

Rewrite so that all terms involving  $T$  are on the left side

$$T(n) - 3T(n/2) = n.$$

Now expand the recurrence, and choose a factor which makes the left side “telescope”

$$1(T(n) - 3T(n/2) = n)$$

$$3(T(n/2) - 3T(n/4) = n/2)$$

$$\vdots \quad \vdots \quad \vdots$$

$$3^{\log_2 n-1}(T(2) - 3T(1) = 2)$$

Let  $m = \log_2 n$ . Summing the left side we get  $T(n) - 3^m T(1) = T(n) - 3^m = T(n) - n^k$  where  $k = \log_2 3 \approx 1.58496$ .

Summing the right side we get

$$\sum_{i=0}^{m-1} \frac{n}{2^i} 3^i = n \sum_{i=0}^{m-1} \left(\frac{3}{2}\right)^i.$$

Let  $c = \frac{3}{2}$ . Then we have

$$n \sum_{i=0}^{m-1} c^i = n \left( \frac{c^m - 1}{c - 1} \right)$$

$$= 2n(c^{\log_2 n} - 1)$$

$$= 2n(c^{(k-1)\log_2 n} - 1)$$

$$= 2n^k - 2n,$$

and so  $T(n) = 3n^k - 2n$ . Full history recurrences can often be changed to limited history ones (example): Consider

$$T_i = 1 + \sum_{j=0}^{i-1} T_j, \quad T_0 = 1.$$

Note that

$$T_{i+1} = 1 + \sum_{j=0}^i T_j.$$

Subtracting we find

$$\begin{aligned}
T_{i+1} - T_i &= 1 + \sum_{j=0}^i T_j - 1 - \sum_{j=0}^{i-1} T_j \\
&= T_i.
\end{aligned}$$

And so  $T_{i+1} = 2T_i = 2^{i+1}$ .

Generating functions:

1. Multiply both sides of the equation by  $x^i$ .
2. Sum both sides over all  $i$  for which the equation is valid.
3. Choose a generating function  $G(x)$ . Usually  $G(x) = \sum_{i=0}^{\infty} x^i g_i$ .
3. Rewrite the equation in terms of the generating function  $G(x)$ .
4. Solve for  $G(x)$ .
5. The coefficient of  $x^i$  in  $G(x)$  is  $g_i$ .

Example:

$$g_{i+1} = 2g_i + 1, \quad g_0 = 0.$$

Multiply and sum:

$$\sum_{i \geq 0} g_{i+1} x^i = \sum_{i \geq 0} 2g_i x^i + \sum_{i \geq 0} x^i.$$

We choose  $G(x) = \sum_{i \geq 0} x^i g_i$ . Rewrite in terms of  $G(x)$ :

$$\frac{G(x) - g_0}{x} = 2G(x) + \sum_{i \geq 0} x^i.$$

Simplify:

$$\frac{G(x)}{x} = 2G(x) + \frac{1}{1-x}.$$

Solve for  $G(x)$ :

$$G(x) = \frac{x}{(1-x)(1-2x)}.$$

Expand this using partial fractions:

$$\begin{aligned}
G(x) &= x \left( \frac{2}{1-2x} - \frac{1}{1-x} \right) \\
&= x \left( 2 \sum_{i \geq 0} 2^i x^i - \sum_{i \geq 0} x^i \right) \\
&= \sum_{i \geq 0} (2^{i+1} - 1) x^{i+1}.
\end{aligned}$$



So  $g_i = 2^i - 1$ .

## Theoretical Computer Science Cheat Sheet

$$\pi \approx 3.14159, \quad e \approx 2.71828, \quad \gamma \approx 0.57721, \quad \phi = \frac{1+\sqrt{5}}{2} \approx 1.61803, \quad \hat{\phi} = \frac{1-\sqrt{5}}{2} \approx -.61803$$

$i$	$2^i$	$p_i$	General	Probability
1	2	2	Bernoulli Numbers ( $B_i = 0$ , odd $i \neq 1$ ):	Continuous distributions: If
2	4	3	$B_0 = 1, B_1 = -\frac{1}{2}, B_2 = \frac{1}{6}, B_4 = -\frac{1}{30},$	$\Pr[a < X < b] = \int_a^b p(x) dx,$
3	8	5	$B_6 = \frac{1}{42}, B_8 = -\frac{1}{30}, B_{10} = \frac{5}{66}.$	then $p$ is the probability density function of $X$ . If
4	16	7	Change of base, quadratic formula:	$\Pr[X < a] = P(a),$
5	32	11	$\log_b x = \frac{\log_a x}{\log_a b}, \quad \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$	then $P$ is the distribution function of $X$ . If $P$ and $p$ both exist then
6	64	13	Euler's number $e$ :	$P(a) = \int_{-\infty}^a p(x) dx.$
7	128	17	$e = 1 + \frac{1}{2} + \frac{1}{6} + \frac{1}{24} + \frac{1}{120} + \dots$	Expectation: If $X$ is discrete
8	256	19	$\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x.$	$E[g(X)] = \sum_x g(x) \Pr[X = x].$
9	512	23	$\left(1 + \frac{1}{n}\right)^n < e < \left(1 + \frac{1}{n}\right)^{n+1}.$	If $X$ continuous then
10	1,024	29	$\left(1 + \frac{1}{n}\right)^n = e - \frac{e}{2n} + \frac{11e}{24n^2} - O\left(\frac{1}{n^3}\right).$	$E[g(X)] = \int_{-\infty}^{\infty} g(x)p(x) dx = \int_{-\infty}^{\infty} g(x) dP(x).$
11	2,048	31	Harmonic numbers:	Variance, standard deviation:
12	4,096	37	$1, \frac{3}{2}, \frac{11}{6}, \frac{25}{12}, \frac{137}{60}, \frac{49}{20}, \frac{363}{140}, \frac{761}{280}, \frac{7129}{2520}, \dots$	$\text{VAR}[X] = E[X^2] - E[X]^2,$
13	8,192	41	$\ln n < H_n < \ln n + 1,$	$\sigma = \sqrt{\text{VAR}[X]}.$
14	16,384	43	$H_n = \ln n + \gamma + O\left(\frac{1}{n}\right).$	For events $A$ and $B$ :
15	32,768	47	Factorial, Stirling's approximation:	$\Pr[A \vee B] = \Pr[A] + \Pr[B] - \Pr[A \wedge B]$
16	65,536	53	$1, 2, 6, 24, 120, 720, 5040, 40320, 362880, \dots$	$\Pr[A \wedge B] = \Pr[A] \cdot \Pr[B],$
17	131,072	59	$n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + \Theta\left(\frac{1}{n}\right)\right).$	iff $A$ and $B$ are independent.
18	262,144	61	Ackermann's function and inverse:	$\Pr[A B] = \frac{\Pr[A \wedge B]}{\Pr[B]}$
19	524,288	67	$a(i, j) = \begin{cases} 2^j & i = 1 \\ a(i-1, 2) & j = 1 \\ a(i-1, a(i, j-1)) & i, j \geq 2 \end{cases}$	For random variables $X$ and $Y$ :
20	1,048,576	71	$\alpha(i) = \min\{j \mid a(j, j) \geq i\}.$	$E[X \cdot Y] = E[X] \cdot E[Y],$
21	2,097,152	73	Binomial distribution:	if $X$ and $Y$ are independent.
22	4,194,304	79	$\Pr[X = k] = \binom{n}{k} p^k q^{n-k}, \quad q = 1 - p,$	$E[X + Y] = E[X] + E[Y],$
23	8,388,608	83	$E[X] = \sum_{k=1}^n k \binom{n}{k} p^k q^{n-k} = np.$	$E[cX] = c E[X].$
24	16,777,216	89	Poisson distribution:	Bayes' theorem:
25	33,554,432	97	$\Pr[X = k] = \frac{e^{-\lambda} \lambda^k}{k!}, \quad E[X] = \lambda.$	$\Pr[A_i B] = \frac{\Pr[B A_i] \Pr[A_i]}{\sum_{j=1}^n \Pr[A_j] \Pr[B A_j]}.$
26	67,108,864	101	Normal (Gaussian) distribution:	Inclusion-exclusion:
27	134,217,728	103	$p(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}, \quad E[X] = \mu.$	$\Pr\left[\bigvee_{i=1}^n X_i\right] = \sum_{i=1}^n \Pr[X_i] +$
28	268,435,456	107	The "coupon collector": We are given a random coupon each day, and there are $n$ different types of coupons. The distribution of coupons is uniform. The expected number of days to pass before we to collect all $n$ types is	$\sum_{k=2}^n (-1)^{k+1} \sum_{i_1 < \dots < i_k} \Pr\left[\bigwedge_{j=1}^k X_{i_j}\right].$
29	536,870,912	109	$nH_n.$	Moment inequalities:
30	1,073,741,824	113		$\Pr[ X  \geq \lambda E[X]] \leq \frac{1}{\lambda},$
31	2,147,483,648	127		$\Pr[ X - E[X]  \geq \lambda \cdot \sigma] \leq \frac{1}{\lambda^2}.$
32	4,294,967,296	131		Geometric distribution:
Pascal's Triangle				$\Pr[X = k] = pq^{k-1}, \quad q = 1 - p,$
1				$E[X] = \sum_{k=1}^{\infty} kpq^{k-1} = \frac{1}{p}.$
1 1				
1 2 1				
1 3 3 1				
1 4 6 4 1				
1 5 10 10 5 1				
1 6 15 20 15 6 1				
1 7 21 35 35 21 7 1				
1 8 28 56 70 56 28 8 1				
1 9 36 84 126 126 84 36 9 1				
1 10 45 120 210 252 210 120 45 10 1				

## Theoretical Computer Science Cheat Sheet

Trigonometry	Matrices	More Trig.																								
<div></div> <p>Pythagorean theorem: <math>C^2 = A^2 + B^2</math>.</p> <p>Definitions:</p> $\sin a = A/C, \quad \cos a = B/C,$ $\csc a = C/A, \quad \sec a = C/B,$ $\tan a = \frac{\sin a}{\cos a} = \frac{A}{B}, \quad \cot a = \frac{\cos a}{\sin a} = \frac{B}{A}.$ <p>Area, radius of inscribed circle:</p> $\frac{1}{2}AB, \quad \frac{AB}{A+B+C}.$ <p>Identities:</p> $\sin x = \frac{1}{\csc x}, \quad \cos x = \frac{1}{\sec x},$ $\tan x = \frac{1}{\cot x}, \quad \sin^2 x + \cos^2 x = 1,$ $1 + \tan^2 x = \sec^2 x, \quad 1 + \cot^2 x = \csc^2 x,$ $\sin x = \cos\left(\frac{\pi}{2} - x\right), \quad \sin x = \sin(\pi - x),$ $\cos x = -\cos(\pi - x), \quad \tan x = \cot\left(\frac{\pi}{2} - x\right),$ $\cot x = -\cot(\pi - x), \quad \csc x = \cot \frac{x}{2} - \cot x,$ $\sin(x \pm y) = \sin x \cos y \pm \cos x \sin y,$ $\cos(x \pm y) = \cos x \cos y \mp \sin x \sin y,$ $\tan(x \pm y) = \frac{\tan x \pm \tan y}{1 \mp \tan x \tan y},$ $\cot(x \pm y) = \frac{\cot x \cot y \mp 1}{\cot x \pm \cot y},$ $\sin 2x = 2 \sin x \cos x, \quad \sin 2x = \frac{2 \tan x}{1 + \tan^2 x},$ $\cos 2x = \cos^2 x - \sin^2 x, \quad \cos 2x = 2 \cos^2 x - 1,$ $\cos 2x = 1 - 2 \sin^2 x, \quad \cos 2x = \frac{1 - \tan^2 x}{1 + \tan^2 x},$ $\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}, \quad \cot 2x = \frac{\cot^2 x - 1}{2 \cot x},$ $\sin(x + y) \sin(x - y) = \sin^2 x - \sin^2 y,$ $\cos(x + y) \cos(x - y) = \cos^2 x - \sin^2 y.$ <p>Euler's equation:</p> $e^{ix} = \cos x + i \sin x, \quad e^{i\pi} = -1.$	<p>Multiplication:</p> $C = A \cdot B, \quad c_{i,j} = \sum_{k=1}^n a_{i,k} b_{k,j}.$ <p>Determinants: <math>\det A \neq 0</math> iff <math>A</math> is non-singular.</p> $\det A \cdot B = \det A \cdot \det B,$ $\det A = \sum_{\pi} \prod_{i=1}^n \text{sign}(\pi) a_{i,\pi(i)}.$ <p><math>2 \times 2</math> and <math>3 \times 3</math> determinant:</p> $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc,$ $\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = g \begin{vmatrix} a & b \\ d & e \end{vmatrix} - h \begin{vmatrix} a & c \\ d & f \end{vmatrix} + i \begin{vmatrix} a & b \\ d & e \end{vmatrix}$ $= aei + bfg + cdh - ceg - fha - ibd.$ <p>Permanents:</p> $\text{perm } A = \sum_{\pi} \prod_{i=1}^n a_{i,\pi(i)}.$	<div></div> <p>Law of cosines:</p> $c^2 = a^2 + b^2 - 2ab \cos C.$ <p>Area:</p> $A = \frac{1}{2}hc,$ $= \frac{1}{2}ab \sin C,$ $= \frac{c^2 \sin A \sin B}{2 \sin C}.$ <p>Heron's formula:</p> $A = \sqrt{s \cdot s_a \cdot s_b \cdot s_c},$ $s = \frac{1}{2}(a + b + c),$ $s_a = s - a,$ $s_b = s - b,$ $s_c = s - c.$ <p>More identities:</p> $\sin \frac{x}{2} = \sqrt{\frac{1 - \cos x}{2}},$ $\cos \frac{x}{2} = \sqrt{\frac{1 + \cos x}{2}},$ $\tan \frac{x}{2} = \sqrt{\frac{1 - \cos x}{1 + \cos x}},$ $= \frac{1 - \cos x}{\sin x},$ $= \frac{\sin x}{1 + \cos x},$ $\cot \frac{x}{2} = \sqrt{\frac{1 + \cos x}{1 - \cos x}},$ $= \frac{1 + \cos x}{\sin x},$ $= \frac{\sin x}{1 - \cos x},$ $\sin x = \frac{e^{ix} - e^{-ix}}{2i},$ $\cos x = \frac{e^{ix} + e^{-ix}}{2},$ $\tan x = -i \frac{e^{ix} - e^{-ix}}{e^{ix} + e^{-ix}},$ $= -i \frac{e^{2ix} - 1}{e^{2ix} + 1},$ $\sin x = \frac{\sinh ix}{i},$ $\cos x = \cosh ix,$ $\tan x = \frac{\tanh ix}{i}.$																								
<p>v2.01 ©1994 by Steve Seiden sseiden@acm.org <a href="http://www.csc.lsu.edu/~seiden">http://www.csc.lsu.edu/~seiden</a></p>	<table><tr><th><math>\theta</math></th><th><math>\sin \theta</math></th><th><math>\cos \theta</math></th><th><math>\tan \theta</math></th></tr><tr><td>0</td><td>0</td><td>1</td><td>0</td></tr><tr><td><math>\frac{\pi}{6}</math></td><td><math>\frac{1}{2}</math></td><td><math>\frac{\sqrt{3}}{2}</math></td><td><math>\frac{\sqrt{3}}{3}</math></td></tr><tr><td><math>\frac{\pi}{4}</math></td><td><math>\frac{\sqrt{2}}{2}</math></td><td><math>\frac{\sqrt{2}}{2}</math></td><td>1</td></tr><tr><td><math>\frac{\pi}{3}</math></td><td><math>\frac{\sqrt{3}}{2}</math></td><td><math>\frac{1}{2}</math></td><td><math>\sqrt{3}</math></td></tr><tr><td><math>\frac{\pi}{2}</math></td><td>1</td><td>0</td><td><math>\infty</math></td></tr></table> <p>... in mathematics you don't under- stand things, you just get used to them. - J. von Neumann</p>	$\theta$	$\sin \theta$	$\cos \theta$	$\tan \theta$	0	0	1	0	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{\pi}{2}$	1	0	$\infty$	
$\theta$	$\sin \theta$	$\cos \theta$	$\tan \theta$																							
0	0	1	0																							
$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$																							
$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1																							
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$																							
$\frac{\pi}{2}$	1	0	$\infty$																							

## Theoretical Computer Science Cheat Sheet

## Number Theory

The Chinese remainder theorem: There exists a number  $C$  such that:

$$C \equiv r_1 \pmod{m_1}$$

$$\vdots \quad \vdots \quad \vdots$$

$$C \equiv r_n \pmod{m_n}$$

if  $m_i$  and  $m_j$  are relatively prime for  $i \neq j$ .

Euler's function:  $\phi(x)$  is the number of positive integers less than  $x$  relatively prime to  $x$ . If  $\prod_{i=1}^n p_i^{e_i}$  is the prime factorization of  $x$  then

$$\phi(x) = \prod_{i=1}^n p_i^{e_i-1} (p_i - 1).$$

Euler's theorem: If  $a$  and  $b$  are relatively prime then

$$1 \equiv a^{\phi(b)} \pmod{b}.$$

Fermat's theorem:

$$1 \equiv a^{p-1} \pmod{p}.$$

The Euclidean algorithm: if  $a > b$  are integers then

$$\gcd(a, b) = \gcd(a \bmod b, b).$$

If  $\prod_{i=1}^n p_i^{e_i}$  is the prime factorization of  $x$  then

$$S(x) = \sum_{d|x} d = \prod_{i=1}^n \frac{p_i^{e_i+1} - 1}{p_i - 1}.$$

Perfect Numbers:  $x$  is an even perfect number iff  $x = 2^{n-1}(2^n - 1)$  and  $2^n - 1$  is prime.

Wilson's theorem:  $n$  is a prime iff

$$(n-1)! \equiv -1 \pmod{n}.$$

Möbius inversion:

$$\mu(i) = \begin{cases} 1 & \text{if } i = 1. \\ 0 & \text{if } i \text{ is not square-free.} \\ (-1)^r & \text{if } i \text{ is the product of } r \text{ distinct primes.} \end{cases}$$

If

$$G(a) = \sum_{d|a} F(d),$$

then

$$F(a) = \sum_{d|a} \mu(d) G\left(\frac{a}{d}\right).$$

Prime numbers:

$$\begin{aligned} p_n &= n \ln n + n \ln \ln n - n + n \frac{\ln \ln n}{\ln n} \\ &\quad + O\left(\frac{n}{\ln n}\right), \\ \pi(n) &= \frac{n}{\ln n} + \frac{n}{(\ln n)^2} + \frac{2!n}{(\ln n)^3} \\ &\quad + O\left(\frac{n}{(\ln n)^4}\right). \end{aligned}$$

## Graph Theory

## Definitions:

*Loop* An edge connecting a vertex to itself.

*Directed* Each edge has a direction.

*Simple* Graph with no loops or multi-edges.

*Walk* A sequence  $v_0 e_1 v_1 \dots e_\ell v_\ell$ .

*Trail* A walk with distinct edges.

*Path* A trail with distinct vertices.

*Connected* A graph where there exists a path between any two vertices.

*Component* A maximal connected subgraph.

*Tree* A connected acyclic graph.

*Free tree* A tree with no root.

*DAG* Directed acyclic graph.

*Eulerian* Graph with a trail visiting each edge exactly once.

*Hamiltonian* Graph with a cycle visiting each vertex exactly once.

*Cut* A set of edges whose removal increases the number of components.

*Cut-set* A minimal cut.

*Cut edge* A size 1 cut.

*k-Connected* A graph connected with the removal of any  $k-1$  vertices.

*k-Tough*  $\forall S \subseteq V, S \neq \emptyset$  we have  $k \cdot c(G-S) \leq |S|$ .

*k-Regular* A graph where all vertices have degree  $k$ .

*k-Factor* A  $k$ -regular spanning subgraph.

*Matching* A set of edges, no two of which are adjacent.

*Clique* A set of vertices, all of which are adjacent.

*Ind. set* A set of vertices, none of which are adjacent.

*Vertex cover* A set of vertices which cover all edges.

*Planar graph* A graph which can be embedded in the plane.

*Plane graph* An embedding of a planar graph.

$$\sum_{v \in V} \deg(v) = 2m.$$

If  $G$  is planar then  $n - m + f = 2$ , so

$$f \leq 2n - 4, \quad m \leq 3n - 6.$$

Any planar graph has a vertex with degree  $\leq 5$ .

## Notation:

$E(G)$  Edge set

$V(G)$  Vertex set

$c(G)$  Number of components

$G[S]$  Induced subgraph

$\deg(v)$  Degree of  $v$

$\Delta(G)$  Maximum degree

$\delta(G)$  Minimum degree

$\chi(G)$  Chromatic number

$\chi_E(G)$  Edge chromatic number

$G^c$  Complement graph

$K_n$  Complete graph

$K_{n_1, n_2}$  Complete bipartite graph

$r(k, \ell)$  Ramsey number

## Geometry

Projective coordinates: triples  $(x, y, z)$ , not all  $x, y$  and  $z$  zero.

$$(x, y, z) = (cx, cy, cz) \quad \forall c \neq 0.$$

Cartesian Projective

$$(x, y) \quad (x, y, 1)$$

$$y = mx + b \quad (m, -1, b)$$

$$x = c \quad (1, 0, -c)$$

Distance formula,  $L_p$  and  $L_\infty$  metric:

$$\sqrt{(x_1 - x_0)^2 + (y_1 - y_0)^2},$$


$$[|x_1 - x_0|^p + |y_1 - y_0|^p]^{1/p},$$

$$\lim_{p \rightarrow \infty} [|x_1 - x_0|^p + |y_1 - y_0|^p]^{1/p}.$$

Area of triangle  $(x_0, y_0)$ ,  $(x_1, y_1)$  and  $(x_2, y_2)$ :

$$\frac{1}{2} \text{abs} \begin{vmatrix} x_1 - x_0 & y_1 - y_0 \\ x_2 - x_0 & y_2 - y_0 \end{vmatrix}.$$

Angle formed by three points:



$$\cos \theta = \frac{(x_1, y_1) \cdot (x_2, y_2)}{l_1 l_2}.$$

Line through two points  $(x_0, y_0)$  and  $(x_1, y_1)$ :

$$\begin{vmatrix} x & y & 1 \\ x_0 & y_0 & 1 \\ x_1 & y_1 & 1 \end{vmatrix} = 0.$$

Area of circle, volume of sphere:

$$A = \pi r^2, \quad V = \frac{4}{3} \pi r^3.$$

If I have seen farther than others, it is because I have stood on the shoulders of giants.

– Issac Newton

## Theoretical Computer Science Cheat Sheet

 $\pi$ 

Wallis' identity:

$$\pi = 2 \cdot \frac{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdots}{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdots}$$

Brouncker's continued fraction expansion:

$$\frac{\pi}{4} = 1 + \frac{1^2}{2 + \frac{3^2}{2 + \frac{5^2}{2 + \frac{7^2}{2 + \cdots}}}}$$

Gregory's series:

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \cdots$$

Newton's series:

$$\frac{\pi}{6} = \frac{1}{2} + \frac{1}{2 \cdot 3 \cdot 2^3} + \frac{1 \cdot 3}{2 \cdot 4 \cdot 5 \cdot 2^5} + \cdots$$

Sharp's series:

$$\frac{\pi}{6} = \frac{1}{\sqrt{3}} \left( 1 - \frac{1}{3^1 \cdot 3} + \frac{1}{3^2 \cdot 5} - \frac{1}{3^3 \cdot 7} + \cdots \right)$$

Euler's series:

$$\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \cdots$$

$$\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \frac{1}{9^2} + \cdots$$

$$\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \frac{1}{5^2} - \cdots$$

## Partial Fractions

Let  $N(x)$  and  $D(x)$  be polynomial functions of  $x$ . We can break down  $N(x)/D(x)$  using partial fraction expansion. First, if the degree of  $N$  is greater than or equal to the degree of  $D$ , divide  $N$  by  $D$ , obtaining

$$\frac{N(x)}{D(x)} = Q(x) + \frac{N'(x)}{D(x)},$$

where the degree of  $N'$  is less than that of  $D$ . Second, factor  $D(x)$ . Use the following rules: For a non-repeated factor:

$$\frac{N(x)}{(x-a)D(x)} = \frac{A}{x-a} + \frac{N'(x)}{D(x)},$$

where

$$A = \left[ \frac{N(x)}{D(x)} \right]_{x=a}.$$

For a repeated factor:

$$\frac{N(x)}{(x-a)^m D(x)} = \sum_{k=0}^{m-1} \frac{A_k}{(x-a)^{m-k}} + \frac{N'(x)}{D(x)},$$

where

$$A_k = \frac{1}{k!} \left[ \frac{d^k}{dx^k} \left( \frac{N(x)}{D(x)} \right) \right]_{x=a}.$$

The reasonable man adapts himself to the world; the unreasonable persists in trying to adapt the world to himself. Therefore all progress depends on the unreasonable.  
– George Bernard Shaw

## Calculus

Derivatives:

$$1. \frac{d(cu)}{dx} = c \frac{du}{dx}, \quad 2. \frac{d(u+v)}{dx} = \frac{du}{dx} + \frac{dv}{dx}, \quad 3. \frac{d(uv)}{dx} = u \frac{dv}{dx} + v \frac{du}{dx},$$

$$4. \frac{d(u^n)}{dx} = nu^{n-1} \frac{du}{dx}, \quad 5. \frac{d(u/v)}{dx} = \frac{v \left( \frac{du}{dx} \right) - u \left( \frac{dv}{dx} \right)}{v^2}, \quad 6. \frac{d(e^{cu})}{dx} = ce^{cu} \frac{du}{dx},$$

$$7. \frac{d(c^u)}{dx} = (\ln c) c^u \frac{du}{dx}, \quad 8. \frac{d(\ln u)}{dx} = \frac{1}{u} \frac{du}{dx},$$

$$9. \frac{d(\sin u)}{dx} = \cos u \frac{du}{dx}, \quad 10. \frac{d(\cos u)}{dx} = -\sin u \frac{du}{dx},$$

$$11. \frac{d(\tan u)}{dx} = \sec^2 u \frac{du}{dx}, \quad 12. \frac{d(\cot u)}{dx} = -\csc^2 u \frac{du}{dx},$$

$$13. \frac{d(\sec u)}{dx} = \tan u \sec u \frac{du}{dx}, \quad 14. \frac{d(\csc u)}{dx} = -\cot u \csc u \frac{du}{dx},$$

$$15. \frac{d(\arcsin u)}{dx} = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}, \quad 16. \frac{d(\arccos u)}{dx} = \frac{-1}{\sqrt{1-u^2}} \frac{du}{dx},$$

$$17. \frac{d(\arctan u)}{dx} = \frac{1}{1+u^2} \frac{du}{dx}, \quad 18. \frac{d(\operatorname{arccot} u)}{dx} = \frac{-1}{1+u^2} \frac{du}{dx},$$

$$19. \frac{d(\operatorname{arcsec} u)}{dx} = \frac{1}{u\sqrt{1-u^2}} \frac{du}{dx}, \quad 20. \frac{d(\operatorname{arccsc} u)}{dx} = \frac{-1}{u\sqrt{1-u^2}} \frac{du}{dx},$$

$$21. \frac{d(\sinh u)}{dx} = \cosh u \frac{du}{dx}, \quad 22. \frac{d(\cosh u)}{dx} = \sinh u \frac{du}{dx},$$

$$23. \frac{d(\tanh u)}{dx} = \operatorname{sech}^2 u \frac{du}{dx}, \quad 24. \frac{d(\coth u)}{dx} = -\operatorname{csch}^2 u \frac{du}{dx},$$

$$25. \frac{d(\operatorname{sech} u)}{dx} = -\operatorname{sech} u \tanh u \frac{du}{dx}, \quad 26. \frac{d(\operatorname{csch} u)}{dx} = -\operatorname{csch} u \coth u \frac{du}{dx},$$

$$27. \frac{d(\operatorname{arcsinh} u)}{dx} = \frac{1}{\sqrt{1+u^2}} \frac{du}{dx}, \quad 28. \frac{d(\operatorname{arccosh} u)}{dx} = \frac{1}{\sqrt{u^2-1}} \frac{du}{dx},$$

$$29. \frac{d(\operatorname{arctanh} u)}{dx} = \frac{1}{1-u^2} \frac{du}{dx}, \quad 30. \frac{d(\operatorname{arccoth} u)}{dx} = \frac{1}{u^2-1} \frac{du}{dx},$$

$$31. \frac{d(\operatorname{arcsech} u)}{dx} = \frac{-1}{u\sqrt{1-u^2}} \frac{du}{dx}, \quad 32. \frac{d(\operatorname{arccsch} u)}{dx} = \frac{-1}{|u|\sqrt{1+u^2}} \frac{du}{dx}.$$

Integrals:

$$1. \int cu \, dx = c \int u \, dx, \quad 2. \int (u+v) \, dx = \int u \, dx + \int v \, dx,$$

$$3. \int x^n \, dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1, \quad 4. \int \frac{1}{x} \, dx = \ln x, \quad 5. \int e^x \, dx = e^x,$$

$$6. \int \frac{dx}{1+x^2} = \arctan x, \quad 7. \int u \frac{dv}{dx} \, dx = uv - \int v \frac{du}{dx} \, dx,$$

$$8. \int \sin x \, dx = -\cos x, \quad 9. \int \cos x \, dx = \sin x,$$

$$10. \int \tan x \, dx = -\ln |\cos x|, \quad 11. \int \cot x \, dx = \ln |\cos x|,$$

$$12. \int \sec x \, dx = \ln |\sec x + \tan x|, \quad 13. \int \csc x \, dx = \ln |\csc x + \cot x|,$$

$$14. \int \arcsin \frac{x}{a} \, dx = \arcsin \frac{x}{a} + \sqrt{a^2 - x^2}, \quad a > 0,$$

## Theoretical Computer Science Cheat Sheet

## Calculus Cont.

15.  $\int \arccos \frac{x}{a} dx = \arccos \frac{x}{a} - \sqrt{a^2 - x^2}, \quad a > 0,$
16.  $\int \arctan \frac{x}{a} dx = x \arctan \frac{x}{a} - \frac{a}{2} \ln(a^2 + x^2), \quad a > 0,$
17.  $\int \sin^2(ax) dx = \frac{1}{2a}(ax - \sin(ax) \cos(ax)),$
18.  $\int \cos^2(ax) dx = \frac{1}{2a}(ax + \sin(ax) \cos(ax)),$
19.  $\int \sec^2 x dx = \tan x,$
20.  $\int \csc^2 x dx = -\cot x,$
21.  $\int \sin^n x dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x dx,$
22.  $\int \cos^n x dx = \frac{\cos^{n-1} x \sin x}{n} + \frac{n-1}{n} \int \cos^{n-2} x dx,$
23.  $\int \tan^n x dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x dx, \quad n \neq 1,$
24.  $\int \cot^n x dx = -\frac{\cot^{n-1} x}{n-1} - \int \cot^{n-2} x dx, \quad n \neq 1,$
25.  $\int \sec^n x dx = \frac{\tan x \sec^{n-1} x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x dx, \quad n \neq 1,$
26.  $\int \csc^n x dx = -\frac{\cot x \csc^{n-1} x}{n-1} + \frac{n-2}{n-1} \int \csc^{n-2} x dx, \quad n \neq 1,$
27.  $\int \sinh x dx = \cosh x,$
28.  $\int \cosh x dx = \sinh x,$
29.  $\int \tanh x dx = \ln |\cosh x|,$
30.  $\int \coth x dx = \ln |\sinh x|,$
31.  $\int \operatorname{sech} x dx = \arctan \sinh x,$
32.  $\int \operatorname{csch} x dx = \ln \left| \tanh \frac{x}{2} \right|,$
33.  $\int \sinh^2 x dx = \frac{1}{4} \sinh(2x) - \frac{1}{2} x,$
34.  $\int \cosh^2 x dx = \frac{1}{4} \sinh(2x) + \frac{1}{2} x,$
35.  $\int \operatorname{sech}^2 x dx = \tanh x,$
36.  $\int \operatorname{arcsinh} \frac{x}{a} dx = x \operatorname{arcsinh} \frac{x}{a} - \sqrt{x^2 + a^2}, \quad a > 0,$
37.  $\int \operatorname{arctanh} \frac{x}{a} dx = x \operatorname{arctanh} \frac{x}{a} + \frac{a}{2} \ln |a^2 - x^2|,$
38.  $\int \operatorname{arccosh} \frac{x}{a} dx = \begin{cases} x \operatorname{arccosh} \frac{x}{a} - \sqrt{x^2 + a^2}, & \text{if } \operatorname{arccosh} \frac{x}{a} > 0 \text{ and } a > 0, \\ x \operatorname{arccosh} \frac{x}{a} + \sqrt{x^2 + a^2}, & \text{if } \operatorname{arccosh} \frac{x}{a} < 0 \text{ and } a > 0, \end{cases}$
39.  $\int \frac{dx}{\sqrt{a^2 + x^2}} = \ln \left( x + \sqrt{a^2 + x^2} \right), \quad a > 0,$
40.  $\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \arctan \frac{x}{a}, \quad a > 0,$
41.  $\int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin \frac{x}{a}, \quad a > 0,$
42.  $\int (a^2 - x^2)^{3/2} dx = \frac{x}{8} (5a^2 - 2x^2) \sqrt{a^2 - x^2} + \frac{3a^4}{8} \arcsin \frac{x}{a}, \quad a > 0,$
43.  $\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a}, \quad a > 0,$
44.  $\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right|,$
45.  $\int \frac{dx}{(a^2 - x^2)^{3/2}} = \frac{x}{a^2 \sqrt{a^2 - x^2}},$
46.  $\int \sqrt{a^2 \pm x^2} dx = \frac{x}{2} \sqrt{a^2 \pm x^2} \pm \frac{a^2}{2} \ln \left| x + \sqrt{a^2 \pm x^2} \right|,$
47.  $\int \frac{dx}{\sqrt{x^2 - a^2}} = \ln \left| x + \sqrt{x^2 - a^2} \right|, \quad a > 0,$
48.  $\int \frac{dx}{ax^2 + bx} = \frac{1}{a} \ln \left| \frac{x}{a+bx} \right|,$
49.  $\int x \sqrt{a+bx} dx = \frac{2(3bx-2a)(a+bx)^{3/2}}{15b^2},$
50.  $\int \frac{\sqrt{a+bx}}{x} dx = 2\sqrt{a+bx} + a \int \frac{1}{x\sqrt{a+bx}} dx,$
51.  $\int \frac{x}{\sqrt{a+bx}} dx = \frac{1}{\sqrt{2}} \ln \left| \frac{\sqrt{a+bx} - \sqrt{a}}{\sqrt{a+bx} + \sqrt{a}} \right|, \quad a > 0,$
52.  $\int \frac{\sqrt{a^2 - x^2}}{x} dx = \sqrt{a^2 - x^2} - a \ln \left| \frac{a + \sqrt{a^2 - x^2}}{x} \right|,$
53.  $\int x \sqrt{a^2 - x^2} dx = -\frac{1}{3} (a^2 - x^2)^{3/2},$
54.  $\int x^2 \sqrt{a^2 - x^2} dx = \frac{x}{8} (2x^2 - a^2) \sqrt{a^2 - x^2} + \frac{a^4}{8} \arcsin \frac{x}{a}, \quad a > 0,$
55.  $\int \frac{dx}{\sqrt{a^2 - x^2}} = -\frac{1}{a} \ln \left| \frac{a + \sqrt{a^2 - x^2}}{x} \right|,$
56.  $\int \frac{x dx}{\sqrt{a^2 - x^2}} = -\sqrt{a^2 - x^2},$
57.  $\int \frac{x^2 dx}{\sqrt{a^2 - x^2}} = -\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin \frac{x}{a}, \quad a > 0,$
58.  $\int \frac{\sqrt{a^2 + x^2}}{x} dx = \sqrt{a^2 + x^2} - a \ln \left| \frac{a + \sqrt{a^2 + x^2}}{x} \right|,$
59.  $\int \frac{\sqrt{x^2 - a^2}}{x} dx = \sqrt{x^2 - a^2} - a \arccos \frac{a}{|x|}, \quad a > 0,$
60.  $\int x \sqrt{x^2 \pm a^2} dx = \frac{1}{3} (x^2 \pm a^2)^{3/2},$
61.  $\int \frac{dx}{x \sqrt{x^2 + a^2}} = \frac{1}{a} \ln \left| \frac{x}{a + \sqrt{a^2 + x^2}} \right|,$

## Theoretical Computer Science Cheat Sheet

## Calculus Cont.

$$\begin{aligned}
62. \int \frac{dx}{x\sqrt{x^2 - a^2}} &= \frac{1}{a} \arccos \frac{a}{|x|}, \quad a > 0, & 63. \int \frac{dx}{x^2\sqrt{x^2 \pm a^2}} &= \mp \frac{\sqrt{x^2 \pm a^2}}{a^2 x}, \\
64. \int \frac{x dx}{\sqrt{x^2 \pm a^2}} &= \sqrt{x^2 \pm a^2}, & 65. \int \frac{\sqrt{x^2 \pm a^2}}{x^4} dx &= \mp \frac{(x^2 + a^2)^{3/2}}{3a^2 x^3}, \\
66. \int \frac{dx}{ax^2 + bx + c} &= \begin{cases} \frac{1}{\sqrt{b^2 - 4ac}} \ln \left| \frac{2ax + b - \sqrt{b^2 - 4ac}}{2ax + b + \sqrt{b^2 - 4ac}} \right|, & \text{if } b^2 > 4ac, \\ \frac{2}{\sqrt{4ac - b^2}} \arctan \frac{2ax + b}{\sqrt{4ac - b^2}}, & \text{if } b^2 < 4ac, \end{cases} \\
67. \int \frac{dx}{\sqrt{ax^2 + bx + c}} &= \begin{cases} \frac{1}{\sqrt{a}} \ln \left| 2ax + b + 2\sqrt{a}\sqrt{ax^2 + bx + c} \right|, & \text{if } a > 0, \\ \frac{1}{\sqrt{-a}} \arcsin \frac{-2ax - b}{\sqrt{b^2 - 4ac}}, & \text{if } a < 0, \end{cases} \\
68. \int \sqrt{ax^2 + bx + c} dx &= \frac{2ax + b}{4a} \sqrt{ax^2 + bx + c} + \frac{4ac - b^2}{8a} \int \frac{dx}{\sqrt{ax^2 + bx + c}}, \\
69. \int \frac{x dx}{\sqrt{ax^2 + bx + c}} &= \frac{\sqrt{ax^2 + bx + c}}{a} - \frac{b}{2a} \int \frac{dx}{\sqrt{ax^2 + bx + c}}, \\
70. \int \frac{dx}{x\sqrt{ax^2 + bx + c}} &= \begin{cases} \frac{-1}{\sqrt{c}} \ln \left| \frac{2\sqrt{c}\sqrt{ax^2 + bx + c} + bx + 2c}{x} \right|, & \text{if } c > 0, \\ \frac{1}{\sqrt{-c}} \arcsin \frac{bx + 2c}{|x|\sqrt{b^2 - 4ac}}, & \text{if } c < 0, \end{cases} \\
71. \int x^3 \sqrt{x^2 + a^2} dx &= \left(\frac{1}{3}x^2 - \frac{2}{15}a^2\right)(x^2 + a^2)^{3/2}, \\
72. \int x^n \sin(ax) dx &= -\frac{1}{a}x^n \cos(ax) + \frac{n}{a} \int x^{n-1} \cos(ax) dx, \\
73. \int x^n \cos(ax) dx &= \frac{1}{a}x^n \sin(ax) - \frac{n}{a} \int x^{n-1} \sin(ax) dx, \\
74. \int x^n e^{ax} dx &= \frac{x^n e^{ax}}{a} - \frac{n}{a} \int x^{n-1} e^{ax} dx, \\
75. \int x^n \ln(ax) dx &= x^{n+1} \left( \frac{\ln(ax)}{n+1} - \frac{1}{(n+1)^2} \right), \\
76. \int x^n (\ln ax)^m dx &= \frac{x^{n+1}}{n+1} (\ln ax)^m - \frac{m}{n+1} \int x^n (\ln ax)^{m-1} dx.
\end{aligned}$$

## Finite Calculus

Difference, shift operators:

$$\Delta f(x) = f(x+1) - f(x),$$

$$\mathbb{E} f(x) = f(x+1).$$

Fundamental Theorem:

$$f(x) = \Delta F(x) \Leftrightarrow \sum f(x) \delta x = F(x) + C.$$

$$\sum_a^b f(x) \delta x = \sum_{i=a}^{b-1} f(i).$$

Differences:

$$\Delta(cu) = c\Delta u, \quad \Delta(u+v) = \Delta u + \Delta v,$$

$$\Delta(uv) = u\Delta v + \mathbb{E} v \Delta u,$$

$$\Delta(x^n) = nx^{n-1},$$

$$\Delta(H_x) = x^{-1}, \quad \Delta(2^x) = 2^x,$$

$$\Delta(c^x) = (c-1)c^x, \quad \Delta\binom{x}{m} = \binom{x}{m-1}.$$

Sums:

$$\sum cu \delta x = c \sum u \delta x,$$

$$\sum (u+v) \delta x = \sum u \delta x + \sum v \delta x,$$

$$\sum u \Delta v \delta x = uv - \sum \mathbb{E} v \Delta u \delta x,$$

$$\sum x^n \delta x = \frac{x^{n+1}}{n+1}, \quad \sum x^{-1} \delta x = H_x,$$

$$\sum c^x \delta x = \frac{c^x}{c-1}, \quad \sum \binom{x}{m} \delta x = \binom{x}{m+1}.$$

Falling Factorial Powers:

$$x^{\underline{n}} = x(x-1) \cdots (x-n+1), \quad n > 0,$$

$$x^{\underline{0}} = 1,$$

$$x^{\underline{n}} = \frac{1}{(x+1) \cdots (x+|n|)}, \quad n < 0,$$

$$x^{\overline{n+m}} = x^{\overline{m}}(x-m)^{\underline{n}}.$$

Rising Factorial Powers:

$$x^{\overline{n}} = x(x+1) \cdots (x+n-1), \quad n > 0,$$

$$x^{\overline{0}} = 1,$$

$$x^{\overline{n}} = \frac{1}{(x-1) \cdots (x-|n|)}, \quad n < 0,$$

$$x^{\overline{n+m}} = x^{\overline{m}}(x+m)^{\underline{n}}.$$

Conversion:

$$x^{\underline{n}} = (-1)^n (-x)^{\overline{n}} = (x-n+1)^{\overline{n}}$$

$$= 1/(x+1)^{\overline{-n}},$$

$$x^{\overline{n}} = (-1)^n (-x)^{\underline{n}} = (x+n-1)^{\underline{n}}$$

$$= 1/(x-1)^{\underline{-n}},$$

$$x^n = \sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^{\underline{k}} = \sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (-1)^{n-k} x^{\overline{k}},$$

$$x^{\underline{n}} = \sum_{k=1}^n \left[ \begin{matrix} n \\ k \end{matrix} \right] (-1)^{n-k} x^k,$$

$$x^{\overline{n}} = \sum_{k=1}^n \left[ \begin{matrix} n \\ k \end{matrix} \right] x^k.$$

$$\begin{aligned}
x^1 &= x^{\underline{1}} &= x^{\overline{1}} \\
x^2 &= x^{\underline{2}} + x^{\underline{1}} &= x^{\overline{2}} - x^{\overline{1}} \\
x^3 &= x^{\underline{3}} + 3x^{\underline{2}} + x^{\underline{1}} &= x^{\overline{3}} - 3x^{\overline{2}} + x^{\overline{1}} \\
x^4 &= x^{\underline{4}} + 6x^{\underline{3}} + 7x^{\underline{2}} + x^{\underline{1}} &= x^{\overline{4}} - 6x^{\overline{3}} + 7x^{\overline{2}} - x^{\overline{1}} \\
x^5 &= x^{\underline{5}} + 15x^{\underline{4}} + 25x^{\underline{3}} + 10x^{\underline{2}} + x^{\underline{1}} &= x^{\overline{5}} - 15x^{\overline{4}} + 25x^{\overline{3}} - 10x^{\overline{2}} + x^{\overline{1}} \\
x^{\overline{1}} &= x^1 &x^{\underline{1}} &= x^1 \\
x^{\overline{2}} &= x^2 + x^1 &x^{\underline{2}} &= x^2 - x^1 \\
x^{\overline{3}} &= x^3 + 3x^2 + 2x^1 &x^{\underline{3}} &= x^3 - 3x^2 + 2x^1 \\
x^{\overline{4}} &= x^4 + 6x^3 + 11x^2 + 6x^1 &x^{\underline{4}} &= x^4 - 6x^3 + 11x^2 - 6x^1 \\
x^{\overline{5}} &= x^5 + 10x^4 + 35x^3 + 50x^2 + 24x^1 &x^{\underline{5}} &= x^5 - 10x^4 + 35x^3 - 50x^2 + 24x^1
\end{aligned}$$

## Theoretical Computer Science Cheat Sheet

## Series

Taylor's series:

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2}f''(a) + \dots = \sum_{i=0}^{\infty} \frac{(x-a)^i}{i!} f^{(i)}(a).$$

Expansions:

$$\begin{aligned} \frac{1}{1-x} &= 1 + x + x^2 + x^3 + x^4 + \dots = \sum_{i=0}^{\infty} x^i, \\ \frac{1}{1-cx} &= 1 + cx + c^2x^2 + c^3x^3 + \dots = \sum_{i=0}^{\infty} c^i x^i, \\ \frac{1}{1-x^n} &= 1 + x^n + x^{2n} + x^{3n} + \dots = \sum_{i=0}^{\infty} x^{ni}, \\ \frac{x}{(1-x)^2} &= x + 2x^2 + 3x^3 + 4x^4 + \dots = \sum_{i=0}^{\infty} ix^i, \\ x^k \frac{d^n}{dx^n} \left( \frac{1}{1-x} \right) &= x + 2^n x^2 + 3^n x^3 + 4^n x^4 + \dots = \sum_{i=0}^{\infty} i^n x^i, \\ e^x &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \dots = \sum_{i=0}^{\infty} \frac{x^i}{i!}, \\ \ln(1+x) &= x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \dots = \sum_{i=1}^{\infty} (-1)^{i+1} \frac{x^i}{i}, \\ \ln \frac{1}{1-x} &= x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \dots = \sum_{i=1}^{\infty} \frac{x^i}{i}, \\ \sin x &= x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots = \sum_{i=0}^{\infty} (-1)^i \frac{x^{2i+1}}{(2i+1)!}, \\ \cos x &= 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots = \sum_{i=0}^{\infty} (-1)^i \frac{x^{2i}}{(2i)!}, \\ \tan^{-1} x &= x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \dots = \sum_{i=0}^{\infty} (-1)^i \frac{x^{2i+1}}{(2i+1)}, \\ (1+x)^n &= 1 + nx + \frac{n(n-1)}{2}x^2 + \dots = \sum_{i=0}^{\infty} \binom{n}{i} x^i, \\ \frac{1}{(1-x)^{n+1}} &= 1 + (n+1)x + \binom{n+2}{2}x^2 + \dots = \sum_{i=0}^{\infty} \binom{i+n}{i} x^i, \\ \frac{x}{e^x - 1} &= 1 - \frac{1}{2}x + \frac{1}{12}x^2 - \frac{1}{720}x^4 + \dots = \sum_{i=0}^{\infty} \frac{B_i x^i}{i!}, \\ \frac{1}{2x}(1 - \sqrt{1-4x}) &= 1 + x + 2x^2 + 5x^3 + \dots = \sum_{i=0}^{\infty} \frac{1}{i+1} \binom{2i}{i} x^i, \\ \frac{1}{\sqrt{1-4x}} &= 1 + x + 2x^2 + 6x^3 + \dots = \sum_{i=0}^{\infty} \binom{2i}{i} x^i, \\ \frac{1}{\sqrt{1-4x}} \left( \frac{1 - \sqrt{1-4x}}{2x} \right)^n &= 1 + (2+n)x + \binom{4+n}{2}x^2 + \dots = \sum_{i=0}^{\infty} \binom{2i+n}{i} x^i, \\ \frac{1}{1-x} \ln \frac{1}{1-x} &= x + \frac{3}{2}x^2 + \frac{11}{6}x^3 + \frac{25}{12}x^4 + \dots = \sum_{i=1}^{\infty} H_i x^i, \\ \frac{1}{2} \left( \ln \frac{1}{1-x} \right)^2 &= \frac{1}{2}x^2 + \frac{3}{4}x^3 + \frac{11}{24}x^4 + \dots = \sum_{i=2}^{\infty} \frac{H_{i-1} x^i}{i}, \\ \frac{x}{1-x-x^2} &= x + x^2 + 2x^3 + 3x^4 + \dots = \sum_{i=0}^{\infty} F_i x^i, \\ \frac{F_n x}{1 - (F_{n-1} + F_{n+1})x - (-1)^n x^2} &= F_n x + F_{2n} x^2 + F_{3n} x^3 + \dots = \sum_{i=0}^{\infty} F_{ni} x^i. \end{aligned}$$

Ordinary power series:

$$A(x) = \sum_{i=0}^{\infty} a_i x^i.$$

Exponential power series:

$$A(x) = \sum_{i=0}^{\infty} a_i \frac{x^i}{i!}.$$

Dirichlet power series:

$$A(x) = \sum_{i=1}^{\infty} \frac{a_i}{i^x}.$$

Binomial theorem:

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k.$$

Difference of like powers:

$$x^n - y^n = (x-y) \sum_{k=0}^{n-1} x^{n-1-k} y^k.$$

For ordinary power series:

$$\alpha A(x) + \beta B(x) = \sum_{i=0}^{\infty} (\alpha a_i + \beta b_i) x^i,$$

$$x^k A(x) = \sum_{i=k}^{\infty} a_{i-k} x^i,$$

$$\frac{A(x) - \sum_{i=0}^{k-1} a_i x^i}{x^k} = \sum_{i=0}^{\infty} a_{i+k} x^i,$$

$$A(cx) = \sum_{i=0}^{\infty} c^i a_i x^i,$$

$$A'(x) = \sum_{i=0}^{\infty} (i+1) a_{i+1} x^i,$$

$$xA'(x) = \sum_{i=1}^{\infty} i a_i x^i,$$

$$\int A(x) dx = \sum_{i=1}^{\infty} \frac{a_{i-1}}{i} x^i,$$

$$\frac{A(x) + A(-x)}{2} = \sum_{i=0}^{\infty} a_{2i} x^{2i},$$

$$\frac{A(x) - A(-x)}{2} = \sum_{i=0}^{\infty} a_{2i+1} x^{2i+1}.$$

Summation: If  $b_i = \sum_{j=0}^i a_j$  then

$$B(x) = \frac{1}{1-x} A(x).$$

Convolution:

$$A(x)B(x) = \sum_{i=0}^{\infty} \left( \sum_{j=0}^i a_j b_{i-j} \right) x^i.$$

God made the natural numbers;  
all the rest is the work of man.  
– Leopold Kronecker



## Theoretical Computer Science Cheat Sheet


## Series

Expansions:

$$\begin{aligned} \frac{1}{(1-x)^{n+1}} \ln \frac{1}{1-x} &= \sum_{i=0}^{\infty} (H_{n+i} - H_n) \binom{n+i}{i} x^i, \\ x^{\overline{n}} &= \sum_{i=0}^{\infty} \left[ \begin{matrix} n \\ i \end{matrix} \right] x^i, \\ \left( \ln \frac{1}{1-x} \right)^n &= \sum_{i=0}^{\infty} \left[ \begin{matrix} i \\ n \end{matrix} \right] \frac{n! x^i}{i!}, \\ \tan x &= \sum_{i=1}^{\infty} (-1)^{i-1} \frac{2^{2i} (2^{2i} - 1) B_{2i} x^{2i-1}}{(2i)!}, \\ \frac{1}{\zeta(x)} &= \sum_{i=1}^{\infty} \frac{\mu(i)}{i^x}, \\ \zeta(x) &= \prod_p \frac{1}{1 - p^{-x}}, \\ \zeta^2(x) &= \sum_{i=1}^{\infty} \frac{d(i)}{x^i} \quad \text{where } d(n) = \sum_{d|n} 1, \\ \zeta(x) \zeta(x-1) &= \sum_{i=1}^{\infty} \frac{S(i)}{x^i} \quad \text{where } S(n) = \sum_{d|n} d, \\ \zeta(2n) &= \frac{2^{2n-1} |B_{2n}|}{(2n)!} \pi^{2n}, \quad n \in \mathbb{N}, \\ \frac{x}{\sin x} &= \sum_{i=0}^{\infty} (-1)^{i-1} \frac{(4^i - 2) B_{2i} x^{2i}}{(2i)!}, \\ \left( \frac{1 - \sqrt{1-4x}}{2x} \right)^n &= \sum_{i=0}^{\infty} \frac{n(2i+n-1)!}{i!(n+i)!} x^i, \\ e^x \sin x &= \sum_{i=1}^{\infty} \frac{2^{i/2} \sin \frac{i\pi}{4}}{i!} x^i, \\ \sqrt{\frac{1 - \sqrt{1-x}}{x}} &= \sum_{i=0}^{\infty} \frac{(4i)!}{16^i \sqrt{2} (2i)!(2i+1)!} x^i, \\ \left( \frac{\arcsin x}{x} \right)^2 &= \sum_{i=0}^{\infty} \frac{4^i i!^2}{(i+1)(2i+1)!} x^{2i}. \end{aligned}$$

$$\begin{aligned} \left( \frac{1}{x} \right)^{\overline{-n}} &= \sum_{i=0}^{\infty} \left\{ \begin{matrix} i \\ n \end{matrix} \right\} x^i, \\ (e^x - 1)^n &= \sum_{i=0}^{\infty} \left\{ \begin{matrix} i \\ n \end{matrix} \right\} \frac{n! x^i}{i!}, \\ x \cot x &= \sum_{i=0}^{\infty} \frac{(-4)^i B_{2i} x^{2i}}{(2i)!}, \\ \zeta(x) &= \sum_{i=1}^{\infty} \frac{1}{i^x}, \\ \frac{\zeta(x-1)}{\zeta(x)} &= \sum_{i=1}^{\infty} \frac{\phi(i)}{i^x}, \end{aligned}$$

## Escher's Knot



## Stieltjes Integration

If  $G$  is continuous in the interval  $[a, b]$  and  $F$  is nondecreasing then

$$\int_a^b G(x) dF(x)$$

exists. If  $a \leq b \leq c$  then

$$\int_a^c G(x) dF(x) = \int_a^b G(x) dF(x) + \int_b^c G(x) dF(x).$$

If the integrals involved exist

$$\int_a^b (G(x) + H(x)) dF(x) = \int_a^b G(x) dF(x) + \int_a^b H(x) dF(x),$$

$$\int_a^b G(x) d(F(x) + H(x)) = \int_a^b G(x) dF(x) + \int_a^b G(x) dH(x),$$

$$\int_a^b c \cdot G(x) dF(x) = \int_a^b G(x) d(c \cdot F(x)) = c \int_a^b G(x) dF(x),$$

$$\int_a^b G(x) dF(x) = G(b)F(b) - G(a)F(a) - \int_a^b F(x) dG(x).$$

If the integrals involved exist, and  $F$  possesses a derivative  $F'$  at every point in  $[a, b]$  then

$$\int_a^b G(x) dF(x) = \int_a^b G(x) F'(x) dx.$$

## Cramer's Rule

If we have equations:

$$a_{1,1}x_1 + a_{1,2}x_2 + \cdots + a_{1,n}x_n = b_1$$

$$a_{2,1}x_1 + a_{2,2}x_2 + \cdots + a_{2,n}x_n = b_2$$

$$\vdots \quad \quad \quad \vdots$$

$$a_{n,1}x_1 + a_{n,2}x_2 + \cdots + a_{n,n}x_n = b_n$$

Let  $A = (a_{i,j})$  and  $B$  be the column matrix  $(b_i)$ . Then there is a unique solution iff  $\det A \neq 0$ . Let  $A_i$  be  $A$  with column  $i$  replaced by  $B$ . Then

$$x_i = \frac{\det A_i}{\det A}.$$

Improvement makes strait roads, but the crooked roads without Improvement, are roads of Genius.

– William Blake (The Marriage of Heaven and Hell)

00	47	18	76	29	93	85	34	61	52
86	11	57	28	70	39	94	45	02	63
95	80	22	67	38	71	49	56	13	04
59	96	81	33	07	48	72	60	24	15
73	69	90	82	44	17	58	01	35	26
68	74	09	91	83	55	27	12	46	30
37	08	75	19	92	84	66	23	50	41
14	25	36	40	51	62	03	77	88	99
21	32	43	54	65	06	10	89	97	78
42	53	64	05	16	20	31	98	79	87

The Fibonacci number system:  
Every integer  $n$  has a unique representation

$$n = F_{k_1} + F_{k_2} + \cdots + F_{k_m},$$

where  $k_i \geq k_{i+1} + 2$  for all  $i$ ,  
 $1 \leq i < m$  and  $k_m \geq 2$ .

## Fibonacci Numbers

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, ...

Definitions:

$$F_i = F_{i-1} + F_{i-2}, \quad F_0 = F_1 = 1,$$

$$F_{-i} = (-1)^{i-1} F_i,$$

$$F_i = \frac{1}{\sqrt{5}} \left( \phi^i - \hat{\phi}^i \right),$$

Cassini's identity: for  $i > 0$ :

$$F_{i+1}F_{i-1} - F_i^2 = (-1)^i.$$

Additive rule:

$$F_{n+k} = F_k F_{n+1} + F_{k-1} F_n,$$

$$F_{2n} = F_n F_{n+1} + F_{n-1} F_n.$$

Calculation by matrices:

$$\begin{pmatrix} F_{n-2} & F_{n-1} \\ F_{n-1} & F_n \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^n.$$