


Advanced Diagnostics and Management of Thyroid Nodules



Ultrasonographic Evaluation, Bethesda Risk Stratification, and Targeted Therapeutic
Strategies

Author: BA Medicales Editorial Team

Publisher: BA Medicales Medical Series (www.BAMedicales.com)

Published: September 2026 | Clinical Reference E-Book

Copyright & Publishing Information

Publisher: BA Medicales Medical Series (www.BAMedicales.com)

Publication Date: September 2026

Format: Digital Clinical E-Book & Reference Guide

Copyright Notice: © 2026 BA Medicales Medical Series. All rights reserved. No part of this publication may be reproduced or transmitted without prior permission.

About BA Medicales (www.BAMedicales.com)

BA Medicales is an established medical-learning platform dedicated to advancing accurate, relevant, and source-aware healthcare education across Indonesia and internationally. Founded on 20 November 2024 by **Dr. dr. Bob Andinata, Sp.B., Subsp. Onk(K)**, BA Medicales serves as an integrated learning bridge connecting understandable public health literacy with rigorous professional clinical education.

Editorial Scope & Core Mission:

- *Oncology & Surgical Foundation:* Initial editorial depth centers on surgical oncology, neoplasia, tumor biology, and thyroid pathology.
- *Multi-Audience Reach:* Delivers tailored knowledge modules for the general public, general practitioners, medical specialists, and multidisciplinary healthcare teams.
- *Flexible Modalities:* Programs are executed through online webinars, digital reference libraries, hands-on offline workshops, and hybrid clinical discussions.
- *Source Integrity:* Every educational release maintains explicit source transparency and evidence-based grounding.

PUBLISHER DISCLAIMER: This e-book is published as a clinical reference module by BA Medicales Medical Series (www.BAMedicales.com) for educational and clinical reference purposes. It does not replace individualized clinical judgment or direct patient care.

Foreword from BA Medicales

Thyroid nodules represent one of the most frequently encountered endocrine conditions in daily surgical and oncological practice. While the vast majority of detected nodules are benign, accurate diagnostic differentiation is crucial to ensure early identification of malignant cases while avoiding unnecessary invasive surgical procedures.

At **BA Medicales Medical Series** (www.BAMedicales.com), our commitment is to provide clear, actionable, and evidence-grounded clinical guides. This module synthesizes high-yield ultrasonographic criteria (ACR TI-RADS), cytological classification standards (Bethesda System), and modern therapeutic pathways (PEA, RFA, and systemic targeted therapies).

How to Use This E-Book

This clinical e-book is structured into three main sections to facilitate efficient point-of-care review:

1. **Preliminaries:** Overview of clinical scope, publisher credentials, and navigation structure.
2. **Core Content Chapters:** In-depth analysis of laboratory screening, high-frequency ultrasonography, Bethesda category risk stratification, and non-surgical ablation vs. targeted therapy.
3. **Reference & Back Matter:** Rapid executive takeaways, standard medical glossary, and publisher background.

Look out for **Clinical Pearl** callout boxes throughout the text, which highlight common diagnostic pitfalls (such as ultrasound artifact misinterpretations).

ONLINE PLATFORM ACCESS: Visit www.BAMedicales.com to access supplementary video lectures, clinical case discussions, and interactive learning tools.

Table of Contents

Section / Chapter Title	Page
1. Preliminaries (Cover, Copyright, Foreword, Table of Contents)	Pages 1 – 4
2. Introduction & Prologue: Epidemiology and Diagnostic Scope	Page 4
3. Chapter 1: Ultrasonographic Evaluation & Imaging Artifacts	Page 5
4. Chapter 2: Bethesda Classification & Surgical Indications	Page 6
5. Chapter 3: Minimally Invasive Ablation & Targeted Therapeutics	Page 7
6. Reference & Back Matter: Executive Summary, References, Glossary & Publisher Bio	Pages 8 – 9

Introduction & Prologue

Thyroid nodule management requires a structured diagnostic workflow combining endocrinological, radiological, and cytopathological parameters. Serum **FT4 and sensitive TSH (TSHs)** remain the initial baseline laboratory tests. Ultrasonography serves as the primary non-invasive imaging modality. Although most detected thyroid nodules are benign, the estimated overall risk of malignancy (ROM) ranges between **7% and 15%**.

Accurate diagnostic assessment relies on high-resolution equipment and standardized reporting protocols. This document lays out the key criteria established in clinical practice.

Chapter 1: Ultrasonographic Evaluation & Imaging Artifacts

Optimum Wave Frequency: For effective ultrasonographic examination of superficial structures such as the thyroid gland, a probe frequency range of **10–15 MHz** is required to achieve high spatial resolution.

ACR TI-RADS Classification Advantages: The points-based ACR TI-RADS framework categorizes nodules based on composition, echogenicity, shape, margin, and echogenic foci. A key strength of ACR TI-RADS is that it achieves **0% unclassifiable nodules** and significantly reduces unnecessary fine-needle aspiration (FNA) biopsies.

Pitfalls in Artifact Recognition: Clinicians must strictly distinguish true microcalcifications from benign imaging artifacts. Specifically, **large comet tail artifacts** or high-echogenicity foci arising from the posterior wall of tiny cysts can be misidentified as **punctate echogenic foci (PEF)**.

Because PEF carries a heavy **3-point score** in ACR TI-RADS, misclassifying these benign artifacts incorrectly inflates a nodule's risk category from **TR2 (Benign)** to **TR4 (Moderately Suspicious)**, leading to inappropriate biopsy orders.




Figure 1.1: Ultrasound Scan of Thyroid Gland Using 10–15 MHz Probe & Artifact Distinction.

DIAGNOSTIC ALERT: Punctate Echogenic Foci (PEF) adds 3 points to ACR TI-RADS. Always check for posterior acoustic features to rule out comet tail artifacts before scoring.

Chapter 2: Bethesda Classification & Surgical Indications

Fine-Needle Aspiration Biopsy (FNAB) results are standardized using the Bethesda System for Reporting Thyroid Cytopathology. Each category corresponds to a specific Risk of Malignancy (ROM) and clinical management directive.

Bethesda Category IV (Follicular Neoplasm / Suspicious for Follicular Neoplasm):

- **Risk of Malignancy (ROM):** Estimated at **25–40%**.
- **Surgical Requirement:** A **surgical lobectomy** is required to obtain a definitive diagnostic determination.
- **Clinical Rationale:** Cytology alone cannot distinguish between a benign follicular adenoma and a malignant follicular thyroid carcinoma. Definitive diagnosis requires histopathology demonstrating vascular or capsular invasion, which cannot be evaluated on FNAB cytology alone. Therefore, clinical observation is insufficient.

Bethesda System Category & ROM Summary Table

Bethesda Category	Diagnostic Classification	Risk of Malignancy (ROM)	Recommended Clinical Management
Category I	Nondiagnostic / Unsatisfactory	1 – 4%	Repeat FNAB with USG guidance
Category II	Benign	0 – 3%	Clinical & USG follow-up
Category III	AUS / FLUS	5 – 15%	Repeat FNAB or molecular testing
Category IV	Follicular Neoplasm	25 – 40%	Surgical Lobectomy
Category V	Suspicious for Malignancy	50 – 75%	Near-total thyroidectomy or lobectomy
Category VI	Malignant	97 – 99%	Total Thyroidectomy

Chapter 3: Therapeutic Interventions & Targeted Therapies

Management of thyroid nodules is tailored according to pathological diagnosis, nodule composition, and clinical stage.

1. Non-Surgical Ablation Procedures for Benign Nodules:

For benign nodules that are symptomatic or cosmetically disturbing, non-surgical minimally invasive techniques offer high efficacy:

- **Percutaneous Ethanol Ablation (PEA):** The procedure of choice for predominantly **cystic benign nodules**.
- **Radiofrequency Ablation (RFA):** Preferred thermal ablation modality for **solid benign thyroid nodules**.

2. Targeted Systemic Therapy for Well-Differentiated Thyroid Carcinoma (DTC):

In advanced, progressive, or radioiodine-refractory well-differentiated thyroid carcinoma, multikinase inhibitor targeted drugs are indicated:

- **Sorafenib:** Oral multikinase inhibitor targeted for advanced progressive DTC.
- **Lenvatinib:** Oral multikinase inhibitor demonstrating significant progression-free survival benefits in refractory DTC.



Figure 3.1: Treatment Pathways for Benign Ablation (PEA/RFA) and Targeted Systemic Therapy (Sorafenib/Lenvatinib).

Executive Summary & Clinical Takeaways

- 1. Initial Laboratory Evaluation:** Always check baseline serum FT4 and sensitive TSH (TSHs).
- 2. USG Technical Standard:** Use high-frequency ultrasound transducers (10–15 MHz) for thyroid imaging.
- 3. Avoid Artifact Misinterpretation:** Do not mistake comet tail artifacts or cyst wall foci for punctate echogenic foci (PEF). PEF adds 3 points and falsely elevates TR2 to TR4.
- 4. Bethesda IV Surgical Mandate:** Follicular neoplasms carry a 25–40% ROM; surgical lobectomy is required for histopathological vascular/capsular evaluation.
- 5. Minimally Invasive Ablation:** Utilize Percutaneous Ethanol Ablation (PEA) for cystic benign nodules and Radiofrequency Ablation (RFA) for solid benign nodules.
- 6. Targeted Therapeutics:** Administer multikinase inhibitors Sorafenib or Lenvatinib for advanced, radioiodine-refractory well-differentiated thyroid carcinoma.

References & Primary Sources

- 1. Primary Source Document 1: SOAL KAHOOT.docx** — Medical Assessment Module on Superficial Ultrasonography, Artifact Recognition, Benign Ablation Procedures (PEA/RFA), and Targeted Therapeutics.
- 2. Primary Source Document 2: SOAL LMS.docx** — Clinical Risk Stratification Matrix, Bethesda System Criteria, and Laboratory Marker Evaluation.
- 3. Official Platform Reference: BA Medicales Medical Series** — www.BAMedicales.com (About BA Medicales Platform & Medical Education System).

Medical Glossary

Term / Abbreviation	Clinical Definition & Meaning
ACR TI-RADS	American College of Radiology Thyroid Imaging Reporting and Data System.
Bethesda System	Standardized cytopathological reporting system for thyroid FNAB.
FNAB	Fine-Needle Aspiration Biopsy.
PEA	Percutaneous Ethanol Ablation (minimally invasive non-surgical procedure).
PEF	Punctate Echogenic Foci (high-risk ultrasound feature worth 3 points).
RFA	Radiofrequency Ablation (thermal ablation for solid benign nodules).
TSHs	Sensitive Thyroid-Stimulating Hormone.

About the Author & Publisher: BA Medicales

BA Medicales Editorial Team operates under **BA Medicales Medical Series** (www.BAMedicales.com) to produce publication-grade, source-grounded clinical references and educational resources.

Founded on 20 November 2024 by **Dr. dr. Bob Andinata, Sp.B., Subsp. Onk(K)**, BA Medicales is built on a strong foundation in surgical oncology. The platform connects healthcare professionals, medical students, and the general public through structured learning, clinical case exchanges, and reliable medical references.

For inquiries, educational collaborations, or to explore additional clinical modules, visit: **www.BAMedicales.com**