PSETs Landing Page*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTEX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH. Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) Nth assigned PSET PDF of a Yale course with course number CODE is:

https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf

where TYPE is stats or math. Similarly, to access my solution for this PSET you can go to:

https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email anish.lakkapragada@yale.edu.

^{*}Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

MATH 241 PSET 3

September 16, 2024

1.

a) We can compute the probability Fred completes the project on time, given that he completes his first milestone on time as:

$$P(A_3|A_1) = P(A_3|A_2, A_1)P(A_2|A_1) + P(A_3|A_2^c, A_1)P(A_2^c|A_1)$$

Because A_3 and A_1 are conditionally independent given A_2 and A_2^c , $P(A_3|A_2, A_1) = P(A_3|A_2) = 0.8$ and $P(A_3|A_2^c, A_1) = P(A_3|A_2^c) = 0.3$. We are also know that $P(A_2|A_1) = 0.8$ and $P(A_2^c|A_1) = 1 - P(A_2^c|A_1) = 0.2$. Thus, $P(A_3|A_1) = 0.8(0.8) + 0.3(0.2) = 0.7$.

We now compute the probability Fred completes the project on time, given that he completes his first milestone late as:

$$P(A_3|A_1^c) = P(A_3|A_2, A_1^c)P(A_2|A_1^c) + P(A_3|A_2^c, A_1^c)P(A_2^c|A_1^c)$$

Given $P(A_3|A_2, A_1^c) = P(A_3|A_2) = 0.8$, $P(A_3|A_2^c, A_1^c) = P(A_3|A_2^c) = 0.3$, $P(A_2|A_1^c) = 0.3$ and $P(A_2^c|A_1^c) = 1 - P(A_2|A_1^c) = 0.7$, we get that $P(A_3|A_1^c) = 0.8(0.3) + 0.3(0.7) = 0.45$.

b) The probability Fred will finish his project on time is given by $P(A_3)$. Given $P(A_1) = 0.75$ and $P(A_3|A_1) = 0.7$, $P(A_3|A_1^c) = 0.45$ from (a), we can compute $P(A_3)$ as:

$$P(A_3) = P(A_3|A_1)P(A_1) + P(A_3|A_1^c)P(A_1^c)$$

$$P(A_3) = 0.7(0.75) + 0.45(1 - P(A_1))$$

$$P(A_3) = 0.7(0.75) + 0.45(0.25)$$

$$P(A_3) = 0.6375$$

2.

We are given P(A) = 1 and we have to prove that for any B where P(B) > 0, P(A|B) = 1. The proof is below.

$$P(B) = P(B \cap A) + P(B \cap A^c)$$

Note that $P(B \cap A) = P(B|A)P(A) = P(B)$ as event A is guaranteed to happen. Thus:

$$P(B) = P(B) + P(B \cap A^c)$$
$$P(B \cap A^c) = 0$$

Because $P(A^c|B) = \frac{P(B \cap A^c)}{P(B)}$ and we know P(B) > 0, $P(A^c|B) = 0$. Thus, we have proven $P(A|B) = 1 - P(A^c|B) = 1$ given P(A) = 1 and for any B where P(B) > 0.

3.

b) We compute $P(G|A^c)$ below.

$$P(G|A^c) = \frac{P(A^c|G)P(G)}{P(A^c)} = \frac{(1 - P(A|G))g}{P(A^c|G)P(G) + P(A^c|G^c)P(G^c)}$$

$$P(G|A^c) = \frac{(1 - p_1)g}{(1 - P(A|G))g + (1 - P(A|G^c))(1 - g)} = \frac{(1 - p_1)g}{(1 - p_1)g + (1 - p_2)(1 - g)}$$

c) Note that because A and B are conditionally independent given G or G^c , $P(B|A^c, G) = P(B|G)$ and $P(B|A^c, G^c) = P(B|G^c)$. We compute $P(B|A^c)$ below.

$$P(B|A^{c}) = P(B|A^{c}, G)P(G|A^{c}) + P(B|A^{c}, G^{c})P(G^{c}|A^{c})$$

$$P(B|A^{c}) = P(B|G)P(G|A^{c}) + P(B|G^{c})(1 - P(G|A^{c}))$$

$$P(B|A^{c}) = p_{1}P(G|A^{c}) + p_{2}(1 - P(G|A^{c}))$$

$$P(B|A^{c}) = p_{2} + \frac{(p_{1} - p_{2})(1 - p_{1})g}{(1 - p_{1})g + (1 - p_{2})(1 - g)}$$

4.

a) Let us define events A and B as the event that the sample goes to labs A and B, respectively. Let us define C as the event in which the patient has the disease conditionitis. and the events + and - as the events in which the patient tested positive and negative, respectively.

Given these definitions, P(C) = p, $P(A) = P(B) = \frac{1}{2}$, $P(+|C,A) = a_1$, $P(-|C^c,A) = a_2$, $P(+|C,B) = b_1$, and $P(-|C^c,B) = b_2$. We compute P(C|+) below.

$$P(C|+) = \frac{P(+|C)P(C)}{P(+)} = p\frac{P(+|C,A)P(A) + P(+|C,B)P(B)}{P(+)} = p\frac{(a_1 + b_1)}{2P(+)}$$

We now compute P(+) below. Note that events (C, A) and (C, B) are entirely independent as the patient having conditionitis has no relation to which lab their sample is tested at.

$$P(+) = P(+|C,A)P(C \cap A) + P(+|C^c,A)P(C^c \cap A) + P(+|C,B)P(C \cap B) + P(+|C^c,B)P(C^c \cap B)$$

$$= a_1P(A)P(C) + (1 - a_2)P(C^c)P(A) + b_1P(C)P(B) + (1 - b_2)P(C^c)P(B)$$

$$= \frac{p(a_1 + b_1)}{2} + \frac{(1 - a_2)(1 - p)}{2} + \frac{(1 - b_2)(1 - p)}{2}$$

As such, given P(+), we can compute P(C|+) as:

$$P(C|+) = \frac{p(a_1 + b_1)}{p(a_1 + b_2) + (1 - p)[2 - a_2 - b_2]}$$

b) We compute P(A|+) below:

$$P(A|+) = \frac{P(+|A)P(A)}{P(+)}$$

$$= \frac{P(+|A,C)P(C) + P(+|A,C^c)P(C^c)}{2P(+)}$$

$$= \frac{pa_1 + (1 - a_2)(1 - p)}{2P(+)}$$

Using our calculation for P(+) in (a), we get our final answer:

$$P(A|+) = \frac{pa_1 + (1 - a_2)(1 - p)}{p(a_1 + b_1) + (1 - p)[2 - a_2 - b_2]}$$

5.

a) Let us define M as the event that the mother has the disease and C_1, C_2 as the events that the first and second child have the disease, respectively. Given these definitions, $P(M) = \frac{1}{3}$, $P(C_1|M) = P(C_2|M) = \frac{1}{2}$, and $P(C_1|M^c) = P(C_2|M^c) = 0$. We compute the probability neither children has the condition given by $P(C_1^c \cap C_2^c)$ below:

$$P(C_1^c \cap C_2^c) = P(C_1^c \cap C_2^c | M)P(M) + P(C_1^c \cap C_2^c | M^c)P(M^c)$$

Note that C_1 and C_2 are conditionally independent given M. As such,

$$P(C_1^c \cap C_2^c) = \frac{P(C_1^c|M)P(C_2^c|M)}{3} + P(C_1^c \cap C_2^c|M^c)P(M^c)$$

$$= \frac{(1 - P(C_1|M))(1 - P(C_2|M))}{3} + P(C_1^c \cap C_2^c|M^c)P(M^c)$$

$$= \frac{1}{12} + P(C_1^c \cap C_2^c|M^c)P(M^c)$$

We also know that events C_1, C_2 will not occur given M^c . Thus $P(C_1^c \cap C_2^c | M^c) = 1$:

$$P(C_1^c \cap C_2^c) = \frac{1}{12} + (1 - P(M))$$
$$P(C_1^c \cap C_2^c) = \frac{1}{12} + \frac{2}{3}$$
$$P(C_1^c \cap C_2^c) = \frac{3}{4}$$

c) We compute $P(M|C_2^c \cap C_1^c)$ below, given that $P(C_1^c \cap C_2^c) = \frac{3}{4}$ from (a). Note again that C_1 and C_2 are conditionally independent given M.

$$\begin{split} P(M|C_2^c \cap C_1^c) &= \frac{P(C_2^c \cap C_1^c|M)P(M)}{P(C_1^c \cap C_2^c)} \\ &= \frac{4(1-P(C_1|M))(1-P(C_2|M))}{9} = \frac{4}{2*2*9} \\ P(M|C_2^c \cap C_1^c) &= \frac{1}{9} \end{split}$$

6. Anish Lakkapragada. I worked independently.