

Discretionary Note

Anish Krishna Lakkapragada

IF YOU USE THIS FILE TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH. Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

CONTENT STARTS ON NEXT PAGE.

To access the general instructions for this repository head [here](#).

Math 226- HW 6 Due: October 14 by Midnight

1. (24 points) Let A be $m \times n$, B be $n \times k$, P, Q be $n \times n$ matrices
 - a) (10 points) Show that $(AB)^t = B^t A^t$. (Use the definition of matrix multiplication)
 - b) (6 points) Show that P is invertible then P^t is invertible, and $(P^t)^{-1} = (P^{-1})^t$
 - c) (8 points) Show that if PQ is invertible then $(PQ)^t$ is invertible and $((PQ)^t)^{-1} = (P^{-1})^t (Q^{-1})^t$.
2. (11 points) Let $A = [\alpha_{ij}]_{n \times n}$. Trace of A is defined as $Tr(A) = \sum_{i=1}^n \alpha_{ii}$.
 - a) (7 points) Let $B = [\beta_{ij}]_{n \times n}$, prove that $Tr(AB) = Tr(BA)$.
 - b) (4 points) Use part a) to show if A and B are similar matrices then $Tr(A) = Tr(B)$
3. (8 points) Show that if A is $n \times 1$ matrix, and B is $1 \times n$ matrix then the $n \times n$ matrix AB has rank at most 1. Conversely if C is any $n \times n$ matrix having rank 1, then there exist $n \times 1$ matrix A , and $1 \times n$ matrix B such that $C = AB$.
4. (14 points) In \mathbb{R}^2 , let L be the line $y = mx$, where $m \neq 0$, and T be reflection of \mathbb{R}^2 about L . (We did something similar to the following logic in the class for projection operator)
 - a) (2 points) Find a vector $v_1 \in \mathbb{R}^2$ such that $T(v_1) = v_1$, and a vector $v_2 \in \mathbb{R}^2$ such that $T(v_2) = -v_2$
 - b) (2 points) Let $\beta' = \{v_1, v_2\}$ where v_1, v_2 are the vectors that you found in part a. Find the matrix representation for $[T]_{\beta'}$.
 - c) (6 points) Let $\beta = \{e_1, e_2\}$. Find $Q = [I_{\mathbb{R}^2}]_{\beta}^{\beta'}$, and $Q^{-1} = [I_{\mathbb{R}^2}]_{\beta'}^{\beta}$.
 - d) (4 points) Find an expression for T using $[T]_{\beta} = Q^{-1}[T]_{\beta'}Q$.